

Title	Variation in parasitism of intertidal invertebrates, with a focus on trematodes on the southwest of Ireland
Authors	Geraghty, Amy-Ciara
Publication date	2018
Original Citation	Geraghty, A-C. 2018. Variation in parasitism of intertidal invertebrates, with a focus on trematodes on the southwest of Ireland. PhD Thesis, University College Cork.
Type of publication	Doctoral thesis
Rights	© 2018, Amy-Ciara Geraghty http://creativecommons.org/ licenses/by-nc-nd/3.0/
Download date	2025-08-27 00:28:05
Item downloaded from	https://hdl.handle.net/10468/6627



University College Cork, Ireland Coláiste na hOllscoile Corcaigh Variation in parasitism of intertidal invertebrates, with a focus on trematodes on the southwest of Ireland

Amy-Ciara Geraghty, B.Sc., MRes.



A thesis submitted in fulfilment of the requirements for the degree of Doctor of Philosophy to the National University of Ireland, Cork.

May 2018

School of Biological, Earth and Environmental Sciences.

Head of School: Professor Sarah Culloty

Research supervisors: Drs Rob McAllen, Ruth Ramsay and Prof. Sarah Culloty.

Table of Contents

DECLARATION2
ACKNOWLEDGEMENTS
GENERAL ABSTRACT4
CHAPTER 1: General Introduction5
CHAPTER 2: Factors affecting parasite infection of three common intertidal gastropods on rocky shores of differing water quality on the coast of Ireland
CHAPTER 3: Distribution of parasites of the European Limpet (<i>Patella vulgata</i>) in a marine reserve
CHAPTER 4: A case study on eutrophication and macroparasite infections with the intertidal invertebrates, the green shore crab (<i>Carcinus maenas</i>) and ragworm (<i>Hediste diversicolor</i>) in three estuaries
CHAPTER 5: Parasites and pathologies of the blue mussel (<i>Mytilus</i> spp.): the role of host stress in susceptibility
CHAPTER 6: General Discussion
APPENDIX

Declaration

This thesis has not been submitted for another degree either at University College Cork or elsewhere. The thesis is my own work and any aspect of the work contributed by colleagues is specified in the acknowledgments. This thesis may be available for consultation within the university library.

Signed: Omy Geraghty

Amy-Ciara Geraghty

Acknowledgements

It is with immense gratitude I acknowledge the help and support of my supervisors Prof Sarah Culloty, Drs Rob McAllen and Ruth Ramsay. Without their support, constructive criticism and editing this dissertation would not have been possible. I am also grateful to Drs Sharon Lynch and Tom Kelly for their guidance and advice in relation to laboratory protocols and statistical analysis.

Funding was provided by the Thomas Crawford-Hayes Fellowship from University College Cork. Thanks must be given to Declan O'Donnell and Patrick Graham from Irish National Parks and Wildlife Service, for granting research permits R1-4/14 & R06-10/1 to allow work in boundaries of Lough Hyne Marine Reserve. I am grateful for the contributions of my colleagues in University College Cork to Chapter 4. In the chapter, water analysis and bird surveys were carried out by Xie Quishi and Vincent Murphy, respectively.

I would also like to thank postgraduate and undergraduate students for their assistance and support over the course of the project; namely, Gregory Beecher, Babette Brooker, Mary-Catherine Gallagher, Grace Flannery, Cian Luck, Emer Morgan, Amy O'Reilly, Enya O'Reilly, William O'Shea and Darren Reidy. I am extremely grateful to the technical staff of University College Cork; Elizabeth Cotter, Luke Harman, and Allen Whitaker. In particular, I am grateful to Luke for all the heavy lifting, rowing and help beyond the call of duty in the field.

Finally, I have to thank my family. Mom, Dad, Amber and David thank you so much for your unending support. It is the least I can do, but I promise not to talk about parasites over dinner ever again.

GENERAL ABSTRACT

Parasites are ubiquitous in marine ecosystems, are sensitive to a range of environmental conditions, share characteristics of traditional bioindicators, and can be indicators of environmental stress. To investigate the role of environmental stressors on parasite communities, specifically trematodes, parasitological surveys of marine invertebrates were conducted across spatial and temporal scales on the coast of Ireland. The potential impact of stressors on parasite abundance, diversity and ability to infect new hosts was investigated.

The response of parasite communities and trematodes to environmental stressors varied in relation to habitat, stressor, host and parasite taxa surveyed. Water quality classifications were defined by the Water Framework Directive and used as a proxy for environmental stress. From May 2014 to April 2015, parasite communities of the common periwinkle Littorina littorea, the dogwhelk Nucella lapillus and the common limpet *Patella vulgata*, were sampled on three shores of differing water classifications and Lough Hyne Marine Reserve, Co. Cork. Large variation in parasite richness and abundance was observed between gastropod species, months and sites sampled. Higher occurrence and intensities of trematode infections in P. vulgata and L. littorea were associated with the site of good water quality. From June to August 2015, the Ragworm, Hediste diversicolor, and green shore crab Carcinus maenas were surveyed in three estuaries of differing enrichment levels. Occurrence and intensities of trematode taxa were recorded in C. maenas and H. diversicolor in the site of lower water quality. The increased abundance of trematodes, higher concentration of nutrient parameters and larger bird communities recorded, suggested that eutrophication may encourage trematode infection.

The potential of statutory protection, to encourage parasite communities was assessed by a yearlong survey (May 2013-April 2014) of subtidal *Mytilus* spp. populations and a short survey (June 2015) of intertidal *P. vulgata* populations in and around Lough Hyne. Parasite communities of both invertebrate taxa were similar in abundance and composition within and outside of Lough Hyne marine reserve. The limited exchange between outer waters and Lough Hyne, suggested that the reserve is unlikely to facilitate parasite taxa abundance. To investigate the observed low prevalence of trematodes in subtidal *Mytilus* spp., a series of laboratory trials was undertaken to assess the role of host stress on susceptibility to trematodes. However, the study was inconclusive due to a high level of natural trematode infection.



Fig. 1 Word clouds generated from the following chapters, excluding citations and references, with (a) words with a minimum frequency of 10 and (b) a minimum frequency of 100. The size of a words indicates their frequency of occurrence with larger sizes representing words with a higher frequency.

CHAPTER 1: General Introduction

The oceans cover the majority of this planet and contain an estimated ~2.2 million species (Mora et al., 2011). Yet before 91% of these marine species can be described, they are being lost with mainly unknown consequences (Mora et al., 2011). Marine biodiversity is lost at increasing rates and can be measured in extinctions, population depletions and community homogenisation (Sala et al., 2006, Worm et al., 2006, Crowe et al., 2015). This loss is largely due to the widespread impacts of human activities on the marine environment; to the extent that no area has been left unaffected (Halpern et al., 2008, Crowe et al., 2015). The impacts can be as a result of a combination of stressors, such as coastal development, overexploitation, pollution and climate change (Jackson et al., 2001, Halpern et al., 2008). The loss of biodiversity impacts on the functioning of marine ecosystems and their ability to provide ecosystem goods and services, on which society depends (Duarte 2000, Hooper et al., 2005, Worm et al., 2006, Beaumont et al., 2007). These ecosystem goods and services are thought to be best supported by diverse biological communities (Worm et al., 2006). Consequently, biodiversity loss may jeopardize future generations' food security, livelihoods and safety, which can have implications for a growing population of 7 billion and especially for the estimated 1,200,000,000 people living within 100km of the coast (Small et al., 2003). Society's health could be affected as changes may lead to increases in the occurrence and severity of disease or parasitic infection (Ward & Lafferty, 2004, Ostfeld & Keesing, 2000).

Marine ecosystem health (MEH) is a concept created to illustrate the state of the marine ecosystem (Constanza 1992, Marcogliese, 2005). MEH has attracted increasing scientific interest in the past decade (e.g. Halpern *et al.*, 2012). Health has been reported to be an easily understood, but poorly suited analogy for the state of a

marine ecosystem (Constanza 1992, Rapport *et al.*, 1998, Belaoussoff *et al.*, 2003). This is because (a) ecosystems are not organisms regulated by physiological processes, (b) the ambiguity of the term "health" and (c) the difficultly in attaining a quantifiable measure of "health" (Wicklum *et al.*, 1995, Belaoussoff *et al.*, 1998). MEH has varied definitions in the literature, ranging from the wellbeing of a coupled human-natural system or a reflection of the condition of the marine environment from the perspective of adverse effects caused by anthropogenic activities (Constanza 1992, Knap *et al.*, 2002, Strain *et al.*, 2002, Halpern *et al.*, 2012). Furthermore, MEH is utilised synonymously with ecosystem integrity, stability, functioning or ecological quality in the literature (Muniz *et al.*, 2011). Assessing whether an ecosystem is "healthy" is difficult in relation to assessing whether it is impacted or not by anthropogenic activities. This is carried out on a case-by-case basis or by inferring the status of an ecosystem from the state of sentinel organisms (Bremner *et al.*, 2003, Well, 2003). The majority of MEH assessments, and this thesis, adopt the latter approach.

Sentinel organisms or environmental indicators are taxa or communities which often reflect environmental conditions. Changes in the physiology, behaviour or presence of sentinel organisms or composition of communities within an ecosystem can indicate stress in an ecosystem (Goodsell *et al.*, 2009). Sentinel organisms or indicators can increase the generality of findings and the speed at which declines in MEH are detected. Molluscan communities have been commonly used as sentinel taxa as they are well studied, have wide geographic distributions and are abundant in marine ecosystems (Stewart *et al.*, 2008, Goodsell *et al.*, 2009, Atalah & Crowe, 2012). Mussel species have been preferentially utilised as sentinel organisms on a global scale due to the taxon's wide geographic distribution, abundance, persistence, sedentary nature, and ability to concentrate metals and organic contaminants (Hellou *et al.*, 2003, Kim *et al.*, 2008, Marigómez *et al.*, 2013, Brenner *et al.*, 2014). In particular, *Mytilus*

spp. have been widely used in large biomonitoring program e.g. "Mussel Watch", to investigate regional patterns in ecosystem health in relation to heavy metals, pollution and parasites (Goldberg, 1975, Kim et al., 2006; 2008). Other studies have considered an index, a health status index (HSI) which has been used successfully to track the health of *Mytilus* spp. after the Prestige Oil Spill and to assess levels of pollution along Greek coasts (Raftopoulou & Dimitriadis 2010, Marigómez et al., 2013). Pollution biomarkers in the digestive gland of Mytilus galloprovincialis and HSI collected revealed high stress conditions in locations sampled (Raftopoulou & Dimitriadis 2010). HSI and Bioeffects Assessment Index (BAI) were among the five integrated biomarker indexes used in Galicia and the Bay of Biscay to simplify the interpretation of ecosystem stress in relation to the pollution of the marine habitats (Marigómez et al., 2013). Gastropods have also been utilised as sentinel species. The common periwinkle Littorina littorea (Linnaeus, 1758) and the dogwhelk, Nucella lapillus (Linnaeus, 1758), have been commonly used in the detection of environmental tributyltin (TBT) by recording the occurrence of intersex and imposex in the prosobranchs (e.g. Gibbs et al., 1987, Harding et al., 1992, Oehlmann et al., 1998). The tissue and radula of the common limpet *Patella vulgata* have been used to monitor heavy metal pollution in the Atlantic (Miramand and Bentley, 1992, Davies et al., 2005). The Ragworm Hediste diversicolor (OF Müller, 1776) has been used as a sensitive bioindicator of heavy pollution (Scaps, 2002, references therein). The Green shore crab Carcinus maenas (Linnaeus, 1758) has been used to test the impact of toxicants and as a bioindicator in the monitoring of heavy metal contaminants (Rodrigues & Pardal, 2014).

The response of sentinel organisms to stressors or anthropogenic impacts can be altered by parasites, which are organisms dependent on, or that benefit from another organism, a host (Rohde, 2002, Lefevre *et al.*, 2009). In molluscs, trematode infection

has resulted in castration, endocrine disruption and increased mortality in hosts. When trematode infection and pollution co-occur, decreased respiration, metabolism, food consumption and cardiac activity has been recorded in bivalves (Morley, 2010). Multiple reviews call for the need to include parasites in pollution studies. Parasites can alter the host's response to a pollutant or accumulate the pollutant in its own tissues (Marcogliese & Pietrock, 2011, Sures *et al.*, 2017).

Parasites as indicators of MEH

Parasites are initially a counterintuitive measure of ecosystem health. However, parasites have many of the characteristics of more traditional sentinel organisms. Parasites are ubiquitous, considered to be the most common lifestyle on earth, and can be highly sensitive to environmental conditions (Mouritsen and Poulin, 2002, Clausen et al., 2008, Dobson et al., 2008). Changes in environmental conditions may affect the complex host-parasite relationship. Parasite infection may increase, if host abundance or susceptibility increases, or decrease, if the infected hosts experience increased mortality rates (Sures et al., 2017). Parasites with complex lifecycles, free living stages or that intermittently survive external to the host, may parallel the response of free living biota, as both are in direct contact with the environment (Mackenzie, 1999, Pietrock and Marcogliese, 2003, Vidal-Martinez et al., 2010, Aguirre-Macedo et al., 2011, Sures et al., 2017). In aquatic environments, the free living stages of parasitic worms, helminths, are known to be sensitive to a range of abiotic factors such as, temperature, salinity, light, pH, water pressure, and surrounding biotic communities (Pietrock and Marcogliese, 2003, Thieltges et al., 2008, Welsh *et al.*, 2014). Parasite mediated effects are recognised to influence a range of ecosystem functions and food web structures (Lafferty, 1997, Marcogliese, 2005, Thompson et al., 2005, Hudson et al., 2006). From an ecological perspective, relevant information on ecosystem state has been obtained by identifying and analysing parasite communities (Marcogliese 2005, Vidal-Martinez *et al.*, 2010, Holmes 2014, Sures *et al.*, 2017).

Beyond the health of their individual hosts, parasite communities can reveal ecosystem stress, changes in host populations, food web structure and ecosystem functioning in response to ecological disturbances (Marcogliese 2001; 2005). In terrestrial ecosystems, Hudson *et al.*, (2006) suggested that a "healthy" ecosystem is one that contains a richness and an abundance of parasite taxa (Marcogliese, 2005). This contradicts past ideas of "stressed" ecosystems or hosts where negative interactions like parasitism were expected to increase in line with decreasing physiological health (Odum, 1985).

The ability of parasite taxa to reflect ecosystem state is varied, and is likely context and taxa dependent (Khan & Thulin, 1991, Sures *et al.*, 2017). Thus, when selecting potential parasites to assess MEH or environmental state, detailed knowledge of life histories, phylogenetic lineages, response to different stressors and the host's immune system is needed (Vidal-Martínez, 2010). The biology and life histories of parasite taxa affect their suitability as indicators and sensitivity to environmental stressors (Marcogliese, 2005, Blanar *et al.*, 2009). In particular, parasites with free-living stages and that require multiple hosts to complete their lifecycle are sensitive to ecosystem disturbance (Sures *et al.*, 2017, references therein).

Trematodes, i.e.flukes, are a class within the phylum Platyhelminthes (flatworms) contains the class Trematoda that subdivides into subclasses Aspidogastrea and Digenea. Digenea is the larger subclass; it contains 77 families, 18,000 species and has high rates of species descriptions still occurring (Poulin 2014, Costello 2016). It

is one of the most successful groups of parasitic trematodes (Cribb et al., 2003, Olson et al., 2003). Digenean trematodes are predominately endoparasites of vertebrates with complex lifecycles that usually require two invertebrate intermediate hosts and a final vertebrate host (heteroxenous) (Fig. 1). Eggs are produced through sexual reproduction of adult worms in the vertebrate host. Eggs are released with host faeces and hatch into free living ciliated infective stages (miracidia) either in the water column or after consumption by the first intermediate host (Sousa, 1994, Esch et al., 2001). In the first intermediate host, usually a mollusc, miracidia reach a speciesspecific tissue or site of infection and transform into asexual reproductive bodies (sporocysts). Sporocysts form additional sporocysts or another larval stage with a mouth and primitive gut (rediae). Rediae can ingest host tissue and other larval stages of trematode species (Kuris 1990, Esch et al., 2001). Within sporocysts and rediae, another free living infective stage (cercariae) develop. Cercariae, once mature, are released into the water column. Cercariae are usually short lived, do not feed, and contain limited reserves of energy (Thieltges & Rick, 2006, Haas et al., 2008). In the water column, cercariae can actively seek out second intermediate hosts or suitable substrates to encyst within or on (metacercariae) (Haas 2003, Haas et al., 2008). Ingestion of metacercariae by the final vertebrate host completes the lifecycle of the trematode (Esch et al., 2001, Haas et al., 2008).

Larval digenean trematodes are capable of infecting a compatible molluscan host by evading the host's internal defense system through a variety of approaches. The trematode larvae can inhibit a host's immune response through the downregulation of hemocyte activity, the release of secretory-excretory products which block or destroy hemocyte receptors or mask or mimic molecular signals that promote hemocyte defense responses (Adema & Loker, 1997).



Fig. 1. Schematic of generalised 3-host lifecycle of marine digenean trematodes. Adult worms live in the final host, generally a vertebrate. Eggs are shed into the water column with host faeces. The eggs then hatch into ciliated organisms, miracidia, in the water column or within the second intermediate host, usually a mollusc. In the first intermediate host, miracidia develop into sporocysts which asexually produce rediae or more sporocysts. Large numbers of free living infective stages, cercariae are released from sporocysts or rediae and encyst in second intermediate hosts or on substrate. Upon consumption by a final host these cysts, metacercariae, transform into adult worms. Please note schematic is not to scale.

Trematodes as indicators of MEH

Parasitic trematodes, in particular have been suggested as suitable indicators of "ecosystem health" (Marcogliese 2005, Sures et al., 2017). Since trematodes require multiple hosts to complete their lifecycles, any perturbance to their transmission or hosts will likely result in a decline in trematode abundance. Trematode communities are suggested to inform on the health of the host organism, its trophic position within the food web and the presence in the ecosystem of any other organisms that are involved in their life cycles (Marcogliese 2005b). Trematodes that utilise invertebrates as primary intermediate hosts, can alter host shell growth, zonation on the shore, behaviour, attachment strength and the fecundity of their gastropod first intermediate hosts (Davies & Knowles, 2001, McCarthy et al., 2004, O'Dwyer et al., 2014a; 2014b). Despite the higher occurrence of metacercariae relative to sporocysts, the effects of trematodes on second intermediate hosts are less well studied (Sousa, 1991, Mouritsen & Poulin 2002). Trematodes that utilise bivalves as second intermediate hosts can affect the host's size, respiration rate and movement (Mouritsen & Poulin, Stier et al., 2015). Laboratory studies suggest that heavy infections of 2002a, metacercariae can negatively affect an invertebrate hosts' survival, movement and respiration rate (Stier et al., 2015).

Meta-analyses carried out by Lafferty (1997), Blanar *et al.*, (2009) and Vidal-Martinez *et al.*, (2010) found that trematodes were consistently and negatively affected by a wide range of pollutants: pesticides, polycyclic aromatic hydrocarbons (PAHs), heavy metals, sewage, pulp and paper waste. However, the response of trematodes to eutrophication differed between reviews (Lafferty, 1997, Blanar *et al.*, 2009, Vidal-Martinez *et al.*, 2010). In reviews conducted by Lafferty (1997) and Vidal-Martinez *et al.*, (2010) eutrophication encouraged trematode infection, while Blanar *et al.*,

(2009) found the reverse. Furthermore, larval free-living stages of trematodes, cercariae, are sensitive to environmental conditions (Pietrock & Marcogliese, 2003). Trematodes have been proposed as indicators of biodiversity as the abundance of larval stages have previously been found to predict the abundance of their final reptilian (Byers *et al.*, 2011) and avian hosts (Huspeni & Lafferty, 2004, Hechinger and Lafferty, 2005, Shea *et al.*, 2012).

Larval trematode communities are also affected by host habitat quality (Cross et al., 2001; 2005, Marcogliese 2005a, Palm et al., 2009). Poor habitat quality is suggested to negatively affect trematodes by impairing the infection success of their free living infective stages (Cross et al., 2001; 2005). Littorina littorea when acting as a first intermediate host of trematodes in a polluted environment produced less viable cercariae (Cross et al., 2001; 2005). Trematode cercariae in hosts in poor quality habitats, accumulated heavy metal pollutants and had impaired swimming ability and longevity (Cross et al., 2001; 2003; 2005). Trematode communities in disturbed or impacted habitats require time to recover to pre-impact levels (Huspeni & Lafferty, 2004, Morley & Lewis, 2006, Aguirre-Macedo et al., 2011). Trematode communities of gastropods Lymnaea stagnalis (Linnaeus, 1758) significantly declined during and after the restoration of a canal in the United Kingdom (Morley & Lewis, 2006). In a restored Californian salt marsh, trematode communities of the Californian horn snail, Cerithideopsis californica (Haldeman, 1840), in impacted sites took 6 years to reach abundances and richness recorded in reference sites (Huspeni & Lafferty, 2004). Similarly, it took several years after a hurricane, for trematode communities of the horn snail Cerithideopsis pliculosa (Menke, 1829), in the Yucatán Peninsula, Mexico to reach pre-impact abundances (Aguirre-Macedo et al., 2011). In Northeast American salt marshes, anthropogenic factors: road density, nitrogen and metal contaminants, had a greater effect on the abundance of trematode communities of *Tritia obsoleta* (Say, 1822) than the abundance of parasites' final hosts (Altman & Byers, 2014). Altman & Byers, (2014) speculated that these factors affected trematode communities via the abundance of their final fish hosts.

Trematodes commonly harboured within mullet species *Liza aurata* (Risso, 1810) and *Liza ramada* (Risso, 1827), were absent from a heavily polluted Mediterranean estuary (Dzikowski *et al.*, 2003). Previous work on the variation of trematode communities in relation to ecological quality has focused on fish (Williams & Mackenzie, 2003, Marcogliese, 2005, Sures *et al.*, 2017). However, utilising fish as hosts in parasitological surveys can be problematic due to their large spatial ranges, limited availability for particular species and the ethical dilemmas of utilising destructive techniques on vertebrates (Khan & Thulin, 1991, Jackson *et al.*, 2001, Costello *et al.*, 2016). Low mobility and restricted diets of hosts are factors that may generate stability and predictability in their parasite communities by contributing to regular and stable recruitment rates by different parasite species (Gonzalez and Poulin, 2005). Thus, in this thesis, the invertebrate-trematode relationship was considered the most appropriate host-parasite model to evaluate ecological quality and the role of surrounding ecological communities.

Trematode infection can have ecosystem-level effects, the extent of which is likely dependent on the abundance of the trematode, the induced alterations of the host and the host population size (Wood *et al.*, 2007, Clausen *et al.*, 2008, Mouritsen & Poulin, 2010). In the North Sea, the common periwinkle, *Littorina littorea*, when utilised as a primary intermediate host by trematodes, consumed significantly less algae relative to uninfected specimens. Thus, the structure of macroalgal communities in microcosm experiments was found to significantly differ between populations of trematode-infected and uninfected *L. littorea* (Clausen *et al.*, 2008). Trematode-induced changes

in intermediate hosts have implications for host population ecology, community interactions, nutrient flow between habitats and food web structure (Lefevre *et al.*, 2009). For example, in Californian estuaries, the killifish *Fundulus parvipinnis*, Girard 1854, infected with metacercariae were 10-30 times more likely to be consumed by avian predators (Lafferty, 2008). Behavioural changes in the New Zealand cockle *Austrovenus stutchburyi* (Wood, 1828) as a trematode second intermediate host resulted in decreased sediment disturbance, increased surface structural complexity and increased larval trematode abundance (Mourisen & Poulin, 2010).

Present knowledge of trematodes of intertidal biota in Irish coastal waters

On a global scale, parasitological studies on charismatic or commercial vertebrate species greatly outnumber those on invertebrate species and generally focus on a single host/parasite species (de Montudouin *et al.*, 2000, Thieltges *et al.*, 2006, Bordes & Morand 2009). Studies on marine microparasites outnumber those on marine macroparasites species (Orias *et al.*, 1978, Byrne *et al.*, 1999, O'Connell & Fives 2004, Lynch *et al.*, 2014). Work in an Irish context has had a similar focus, but has utilised a wide array of techniques, histological, genomic and macroscopic, to detect macro-and microparasites e.g. Fermer *et al.*, 2011a; 2011b, Callaway *et al.*, 2012, Lynch *et al.*, 2012; 2014, Firth *et al.*, 2017. In particular, the microparasites of commercial bivalves the Pacific oyster *Magallana gigas* (Thunberg, 1793), mussel *Mytilus* spp. (Linnaeus, 1758) and European flat oyster *Ostrea edulis* (Linnaeus, 1758) have received much attention in the literature e.g. Culloty *et al.*, 1999, Lynch *et al.*, 2010, Lynch *et al.*, 2012.

On the southwest coast of Ireland, intertidal bivalves were the biota most commonly screened for macroparasites (Fermer *et al.*, 2011, Lynch *et al.*, 2012, Morgan *et al.*, 2012) and gastropods (Prinz*et al.*, *et al.*, 2010). In these studies, trematodes were found to be a major component of macroparasite communities of common invertebrates on rocky shores (Prinz *et al.*, 2010) and of the bivalve, the edible cockle *Cerastoderma edule* (Linnaeus, 1758) in soft sediment shores (Fermer *et al.*, 2011a; 2011b). In the north of Ireland, trematodes have been recorded in several intertidal gastropods (Copeland *et al.*, 1987, Field & Irwin 1999, McCarthy *et al.*, 1999; 2000; 2002, Cross *et al.*, 2005), acorn barnacles *Semibalanus balanoides* (Linnaeus, 1767) (Irwin & Irwin 1980) and herring gulls *Larus argentatus*, Pontoppidan, 1763, (Irwin & Prentice 1976). Firth *et al.*, (2017) screened the common limpet *Patella vulgata* (Linnaeus, 1758) from rocky shores on the east and west coast of Ireland. Larval trematodes were abundant in *P. vulgata* for the majority of sites sampled.

OUTLINE

The present study focused on gaining more knowledge of the macroparasites of common marine invertebrate species in Irish coastal waters and relating them to the current known environmental parameters associated with the sites. The study aimed to evaluate if parasite communities, with a focus on trematodes, of biota screened varied in response to ecosystem quality as defined by policy directive classifications (Chapter 2 & 4), levels of statutory protection (Chapter 2, 3 & 5) or nutrient enrichment (Chapter 4).

Chapter 2 describes a year-long survey of the parasite communities of the common periwinkle *Littorina littorea*, common limpet, *Patella vulgata*, and dogwhelk *Nucella lapillus*. Gastropods were sampled from three rocky shores of differing Water Framework Directive classifications as well as from a marine reserve, Lough Hyne, on the Southwest of Ireland. It was hypothesised that shores of higher status would contain richer and more abundant parasite communities.

In **Chapter 3**, parasite communities of the common limpet *Patella vulgata* were surveyed within and outside the marine reserve Lough Hyne. The study aimed to investigate the potential of the marine reserve to act a reservoir of marine parasites due to its sheltered coastline and statutory protection. It was hypothesised that a higher occurrence and diversity of parasites would be observed within the boundaries of the marine reserve.

Chapter 4 examines parasite communities occurring in the European green shore crab, *Carcinus maenas*, and Ragworm, *Hediste diversicolor*, in three different estuaries of different nutrient status classifications according the Water Framework

Directive and characterised by physio-chemical parameters. Avian communities were surveyed at each estuary to estimate the number of potential role of final hosts of trematode taxa recorded.

In **Chapter 5**, the parasitological communities of subtidal blue mussels *Mytilus* spp. within and outside of Lough Hyne marine reserve are described. Also, the susceptibility of *Mytilus* spp., to the cercariae of the Digenean trematode *Echinostephilla patellae* was investigated following previous work identifying the bivalve as an intermediate host. Host stress was considered in terms of differing *Mytilus* spp. habitats, exposure to the dinoflagellate *Karenia mikimotoi* and simulated nutrient enrichment.

Chapter 6 provides conclusions of the study and reviews the results of preceding chapters, evaluates their significance and suggests future work.

REFERENCES

- Aarab N, Pampanin DM, Naevdal A, Oysaed KB, Gastaldi L, Bechmann RK (2008) Histopathology alterations and histochemistry measurements in mussel, *Mytilus edulis* collected offshore from an aluminium smelter industry (Norway). *Marine Pollution Bulletin*, 57, 569-574.
- Adema CM, Loker E (1997) Specifity and Immunobiology of Larval Digenean-Snail Associations. In Advances in trematode biology, Fried B and Graczyk TK (eds.). CRC Press, Boca Raton, Florida, 229–265.
- Aguirre-Macedo ML, Vidal-Martinez VM, Lafferty KD (2011) Trematode communities in snails can indicate impact and recovery from hurricanes in a tropical coastal lagoon. *International Journal for Parasitology*, **41**, 1403-1408.
- Altman I, Byers JE (2014) Large-scale spatial variation in parasite communities influenced by anthropogenic factors. *Ecology*, **95**, 1876-1887.
- Atalah J, Crowe TP (2010) Combined effects of nutrient enrichment, sedimentation and grazer loss on rock pool assemblages. *Journal of Experimental Marine Biology and Ecology*, **388**, 51-57.
- Atalah J, Crowe TP (2012) Nutrient enrichment and variation in community structure on rocky shores: The potential of molluscan assemblages for biomonitoring. *Estuarine Coastal and Shelf Science*, **99**, 162-170.
- Beaumont NJ, Austen MC, Atkins JP, Burdon D, Degraer S, Dentinho TP, Derous S, Holm P, Horton T, van Ierland E, Marboe AH, Starkey DJ, Townsend M, Zarzycki T (2007) Identification, definition and quantification of goods and services provided by marine biodiversity: Implications for the ecosystem approach. *Marine Pollution Bulletin*, 54, 253-265.
- Belaoussoff S, Kevan PG (1998) Toward an ecological approach for the assessment of ecosystem health. *Ecosystem Health*, **4**, 4-8.
- Blanar CA, Munkittrick KR, Houlahan J, MacLatchy DL, Marcogliese DJ (2009) Pollution and parasitism in aquatic animals: A meta-analysis of effect size. *Aquatic Toxicology*, **93**, 18-28.
- Bordes F, Morand S (2009) Parasite diversity: an overlooked metric of parasite pressures? *Oikos*, **118**, 801-806.
- Bremner J, Frid C, Rogers SI (2003) Assessing marine ecosystem health: the longterm effects of fishing on functional biodiversity in North Sea benthos. Aquatic Ecosystem Health and Management: Special Issue, Barometers of Aquatic Ecosystem Health and Integrity, 6, 131-137.

- Brenner M, Broeg K, Frickenhaus S, Buck BH, Koehler A (2014) Multi-biomarker approach using the blue mussel (*Mytilus edulis* L.) to assess the quality of marine environments, Season and habitat-related impacts. *Marine Environmental Research*, **95**, 13-27.
- Byers JE, Altman I, Grosse AM, Huspeni TC, Maerz JC (2011) Using Parasitic Trematode Larvae to Quantify an Elusive Vertebrate Host. *Conservation Biology*, **25**, 85-93.
- Byrne CJ, Holland C, Tully O (1999) Metazoan parasite community structure of sea trout on the west coast of Ireland. *Journal of Fish Biology*, **55**, 127-134.
- Callaway R, Shinn AP, Grenfell SE, Bron JE, Burnell G, Cook EJ, Crumlish M, Culloty S, Davidson K, Ellis RP, Flynn KJ, Fox C, Green DM, Hays GC, Hughes AD, Johnston E, Lowe CD, Lupatsch I, Malham S, Mendzil AF, Nickell T, Pickerell T, Rowley AF, Stanley MS, Tocher DR, Turnbull JF, Webb G, Wootton E, Shields RJ (2012) Review of climate change impacts on marine aquaculture in the UK and Ireland. *Aquatic Conservation-Marine and Freshwater Ecosystems*, 22, 389-421.
- Clausen KT, Larsen MH, Iversen NK, Mouritsen KN (2008) The influence of trematodes on the macroalgae consumption by the common periwinkle *Littorina littorea. Journal of the Marine Biological Association of the United Kingdom*, **88**, 1481-1485.
- Constanza R (1992) *Toward an operational definition of health.*, Island Press, Washington, D.C.
- Copeland MR, Montogomery WI, Hana REB (1987) Ecology of a digenean infection, *Cercaria patellae* in *Patella vulgata* near Portavogie Harbour, Northern Ireland. *Journal of Helminthology*, **61**, 315-328.
- Costello MJ (2016) Parasite Rates of Discovery, Global Species Richness and Host Specificity. *Integrative and Comparative Biology*, **56**, 588-599.
- Cribb TH, Bray RA, Olson PD, Timothy D, Littlewood J (2003) Life cycle evolution in the Digenea: a new perspective from phylogeny. *Advances in Parasitology*, 54, 197-254.
- Cross MA, Irwin SWB, Fitzpatrick SM (2001) Effects of heavy metal pollution on swimming and longevity in cercariae of *Cryptocotyle lingua* (Digenea: Heterophyidae). *Parasitology*, **123**, 499-507.
- Cross MA, Irwin SWB, Fitzpatrick SM (2005) Effects of host habitat quality on the viability of *Cryptocotyle lingua* (Trematoda : Digenea) cercariae. *Parasitology*, **130**, 195-201.

- Crowe T, Austen M, Frid C (2015) Human Impacts on Biodiversity, Functioning and Services Introduction. *Marine Ecosystems: Human Impacts on Biodiversity*, *Functioning and Services*, 3-20.
- Crowe TP, Thompson RC, Bray S, Hawkins SJ (2000) Impacts of anthropogenic stress on rocky intertidal communities. *Journal of Aquatic Ecosystem Stress and Recovery*, **7**, 273-297.
- Culloty SC, Novoa B, Pernas M, Longshaw M, Mulcahy MF, Feist SW, Figueras A (1999) Susceptibility of a number of bivalve species to the protozoan parasite *Bonamia ostreae* and their ability to act as vectors for this parasite. *Diseases* of Aquatic Organisms, **37**, 73-80.
- Davies MS, Knowles AJ (2001) Effects of trematode parasitism on the behaviour and ecology of a common marine snail (*Littorina littorea* (L.)). *Journal of Experimental Marine Biology and Ecology*, **260**, 155-167.
- Davies, M. S., Proudlock, D. J., and Mistry, A. (2005). Metal Concentrations in the Radula of the Common Limpet, *Patella vulgata* L., from 10 Sites in the UK. *Ecotoxicology*, 14, 465–475.
- de Montudouin X, Kisielewski I, Bachelet G, Desclaux C (2000) A census of macroparasites in an intertidal bivalve community, Arcachon Bay, France. *Oceanologica Acta*, **23**, 453-468.
- Dobson A, Lafferty KD, Kuris AM, Hechinger RF, Jetz W (2008) Homage to Linnaeus: How many parasites? How many hosts? *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 11482-11489.
- Duarte CM (2000) Marine biodiversity and ecosystem services: an elusive link. Journal of Experimental Marine Biology and Ecology, **250**, 117-131.
- Dzikowski R, Paperna I, Diamant A (2003) Use of fish parasite species richness indices in analyzing anthropogenically impacted coastal marine ecosystems. *Helgoland Marine Research*, **57**, 220-227.
- Esch GW, Curtis LA, Barger MA (2001) A perspective on the ecology of trematode communities in snails. *Parasitology*, **123**, 57-75.
- Feis ME, Thieltges DW, Olsen JL, de Montaudouin X, Jensen KT, Bazairi H, Culloty SC, Luttikhuizen PC (2015) The most vagile host as the main determinant of population connectivity in marine macroparasites. *Marine Ecology Progress* Series, 520, 85-99.
- Fermer J, Culloty SC, Kelly TC, O'Riordan RM (2011a) Manipulation of Cerastoderma edule burrowing ability by Meiogymnophallus minutus metacercariae? Journal of the Marine Biological Association of the United Kingdom, 91, 907-911.

- Fermer J, Culloty SC, Kelly TC, O'Riordan RM (2011b) Parasitological survey of the edible cockle *Cerastoderma edule* (Bivalvia) on the south coast of Ireland. *Journal of the Marine Biological Association of the United Kingdom*, **91**, 923-928.
- Field L, Irwin S (1999) Digenean larvae in *Hydrobia ulvae* from Belfast Lough (Northern Ireland) and the Ythan Estuary (north-east Scotland). *Journal of the Marine Biological Association of the United Kingdom*, **79**, 431-435.
- Firth L, Grant L, Crowe T, Ellis J, Wiler C, Convery C, O'Connor N (2017) Factors affecting the prevalence of the trematode parasite *Echinostephilla patellae* (Lebour, 1911) in the limpet *Patella vulgata*. *Journal of Experimental Marine Biology and Ecology*, **492**, 99-104.
- Fleming LE, Broad K, Clement A, Dewailly E, Elmir S, Knap A, Pomponi SA, Smith S, Gabriele HS, Walsh P (2006) Oceans and human health: Emerging public health risks in the marine environment. *Marine Pollution Bulletin*, **53**, 545-560.
- Galaktionov KV, Dobrovolskij AA (2003) The Trematode Life Cycle as a System of Adaptations. In: *The Biology and Evolution of Trematodes*. Springer, 215-317.
- Gibbs P, Pascoe P, Burt G (1988) Sex change in the female dog-whelk, *Nucella lapillus*, induced by tributyltin from antifouling paints. *Journal of the Marine Biological Association of the United Kingdom*, **68**, 715-731.
- Girard C (1854). Descriptions of new fishes collected by Dr. A. L. Heermann, naturalist attached to the Survey of the Pacific Railroad Route, under Lt. R. S. Williamson, U.S.A.
- Goldberg ED (1975) The Mussel Watch. A First Step in Global Monitoring. *Marine Pollution Bulletin*, **6**, 111-113.
- González MT, R Poulin T (2005) Spatial and temporal predictability of the parasite community structure of a benthic marine fish along its distributional range. *International Journal for Parasitology*, **35**, 1369–1377.
- Gonzalez MT, R Poulinr (2005) Spatial and temporal predictability of the parasite community structure of a benthic marine fish along its distributional range. *International Journal for Parasitology*, **35**, 1369-1377.
- Goodsell PJ, Underwood AJ, Chapman MG (2009) Evidence necessary for taxa to be reliable indicators of environmental conditions or impacts. *Marine Pollution Bulletin*, **58**, 323-331.
- Haas W (2003) Parasitic worms: strategies of host finding, recognition and invasion. *Zoology*, **106**, 349-364.

- Haas W, Beran B, Loy C (2008) Selection of the host's habitat by cercariae: from laboratory experiments to the field. *The Journal of Parasitology*, **94**, 1233-1238.
- Haldeman SS, (1840). A monograph of the Limniades and other freshwater univalve shells of North America. Philadelphia, J. Dobson 1, (unnumbered page)
- Halpern BS, Longo C, Hardy D, McLeod KL, Samhouri JF, Katona SK, Kleisner K, Lester SE, O'Leary J, Ranelletti M, Rosenberg AA, Scarborough C, Selig ER, Best BD, Brumbaugh DR, Chapin FS, Crowder LB, Daly KL, Doney SC, Elfes C, Fogarty MJ, Gaines SD, Jacobsen KI, Karrer LB, Leslie HM, Neeley E, Pauly D, Polasky S, Ris B, St Martin K, Stone GS, Sumaila UR, Zeller D (2012) An index to assess the health and benefits of the global ocean. *Nature*, **488**, 615-620.
- Halpern BS, Walbridge S, Selkoe KA, Kappel CV, Micheli F, D'Agrosa C, Bruno JF, Casey KS, Ebert C, Fox HE, Fujita R, Heinemann D, Lenihan HS, Madin EMP, Perry MT, Selig ER, Spalding M, Steneck R, Watson R (2008) A global map of human impact on marine ecosystems. *Science*, **319**, 948-952.
- Harding MJC, Bailey SK, Davies IM (1992) TBT imposex survey of the North Sea. UK Department of the Environment, Contract PECD 7/8/214, Annex 4: Germany (Scottish Fisheries Working Paper No. 13/92), 1-7.
- Hechinger RF, Lafferty KD (2005) Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1059-1066.
- Hechinger RF, Lafferty KD (2005) Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1059-1066.
- Hechinger RF, Wood AC, Kuris AM (2011) Social organization in a flatworm: trematode parasites form soldier and reproductive castes. *Proceedings of the Royal Society of London B: Biological Sciences*, **278**, 656-665.
- Hellou J, Law RJ (2003) Stress on stress response of wild mussels, *Mytilus edulis* and *Mytilus trossulus*, as an indicator of ecosystem health. *Environmental Pollution*, **126**, 407–416.
- Holmes JC (2014) Environmental parasitology: parasites as bioindicators of ecosystem health. In: *Parasitism: the Diversity and Ecology of Animal Parasites, 2nd Edition*. Cambridge Univ Press, Cambridge, pp. 459-476.
- Hooper DU, Chapin FS, Ewel JJ, Hector A, Inchausti P, Lavorel S, Lawton JH, Lodge DM, Loreau M, Naeem S, Schmid B, Setala H, Symstad AJ, Vandermeer J, Wardle DA (2005) Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, **75**, 3-35.

- Hudson PJ, Dobson AP, Lafferty KD (2006) Is a healthy ecosystem one that is rich in parasites? *Trends in Ecology & Evolution*, **21**, 381-385.
- Huspeni TC, Hechinger RF, Lafferty KD (2005) Trematode parasites as estuarine indicators: Opportunities, applications, and comparisons with conventional community approaches. *Estuarine Indicators*, (ed Bortone, SA) CRC Press, 297-314.
- Huspeni TC, Lafferty KD (2004) Using larval trematodes that parasitize snails to evaluate a saltmarsh restoration project. *Ecological Applications*, **14**, 795-804.
- Irwin SWB, Irwin BC (1980) The distribution of metacercariae of *Maritrema arenaria* (Digenea: Microphallidae) in the barnacle *Balanus balanoides* at three sites on the east coast of Northern Ireland. *Journal of the Marine Biological* Association of the United Kingdom, **60**, 959-862.
- Irwin SWB, Prentice HJ (1976) The parasitic fauna in the digestive tracts of three herring gulls, *Larus argentatus*, taken from Strangford Lough. *Irish Naturalists' Journal*, **18**, 281-282.
- Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH, Cooke R, Erlandson J, Estes JA, Hughes TP, Kidwell S, Lange CB, Lenihan HS, Pandolfi JM, Peterson CH, Steneck RS, Tegner MJ, Warner RR (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293, 629-638.
- Jennings S, Kaiser MJ (1998) The Effects of Fishing on Marine Ecosystems. In: Advances in Marine Biology (ed by JHS Blaxter). Academic Press, 201-352.
- Khan RA, Thulin J (1991) Influence of pollution on Parasites of Aquatic Animals. *Advances in Parasitology*, **30**, 201-238.
- Kim Y, Powell EN, Wade TL, Presley BJ (2008) Relationship of parasites and pathologies to contaminant body burden in sentinel bivalves: NOAA Status and Trends 'Mussel Watch' Program. *Marine Pollution Bulletin*, 65, 101–127.
- Knap A, Dewailly E, Furgal C, Galvin J, Baden D, Bowen RE, Depledge M, Duguay L, Fleming LE, Ford T, Moser F, Owen R, Suk WA, Unluata U (2002) Indicators of ocean health and human health: Developing a research and monitoring framework. *Environmental Health Perspectives*, **110**, 839-845.
- Kollien A (1996) *Cercaria patellae* Lebour, 1911 developing in *Patella vulgata* is the cercaria of *Echinostephilla patellae* (Lebour, 1911) n. comb. (Digenea, Philophthalmidae). *Systematic Parasitology*, **34**, 11-25.
- Kuris A (1990) Guild structure of larval trematodes in molluscan hosts: prevalence, dominance and significance of competition.gi In: *Parasite communities: patterns and processes*. Springer 69-100.

- Lafferty KD (1997) Environmental parasitology: What can parasites tell us about human impacts on the environment? *Parasitology Today*, **13**, 251-255
- Lafferty KD (2008) Ecosystem consequences of fish parasites. *Journal of Fish Biology*, **73**, 2083-2093.
- Lafferty KD, Allesina S, Arim M, Briggs CJ, De Leo G, Dobson AP, Dunne JA, Johnson PTJ, Kuris AM, Marcogliese DJ, Martinez ND, Memmott J, Marquet PA, McLaughlin JP, Mordecai EA, Pascual M, Poulin R, Thieltges DW (2008) Parasites in food webs: the ultimate missing links. *Ecology Letters*, **11**, 533-546.
- Lafferty KD, Holt RD (2003) How should environmental stress affect the population dynamics of disease? *Ecology Letters*, **6**, 654-664.
- Lebour M, (1911) A Review of the British Marine Cercariae. *Parasitology*, **4**, 416-456.
- Lefevre T, Lebarbenchon C, Gauthier-Clerc M, Misse D, Poulin R, Thomas F (2009) The ecological significance of manipulative parasites. *Trends in Ecology & Evolution*, **24**, 41-48.
- Linnaeus C (1758) Tomus I. Systema naturae per regna tria naturae :secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis10. Holmiae, *Laurentii Salvii*, **1-4**, 1-824.
- Linnaeus C (1767) Systema naturae per regna tria naturae: secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. *Laurentii Salvii*, **1-2**, 1-1327.
- Lynch S, Abollo E, Ramilo A, Cao A, Culloty SC, Villalba A (2010) Observations raise the question if the Pacific oyster, *Crassostrea gigas*, can act as either a carrier or a reservoir for *Bonamia ostreae* or *Bonamia exitiosa*. *Parasitology*, 137, 1515-1526.
- Lynch S, Carlsson J, Reilly AO, Cotter E, Culloty SC (2012) A previously undescribed ostreid herpes virus 1 (OsHV-1) genotype detected in the pacific oyster, *Crassostrea gigas*, in Ireland. *Parasitology*, **139**, 1526-1532.
- Lynch S, Morgan E, Carlsson J, Mackenzie C, Wooton E, Rowley A, Malham S, Culloty S (2014) The health status of mussels, *Mytilus* spp., in Ireland and Wales with the molecular identification of a previously undescribed haplosporidian. *Journal of Invertebrate Pathology*, **118**, 59-65.
- Mackenzie (1999) Parasites as Pollution Indicators in Marine Ecosystems: a Proposed Early Warning System. *Marine Pollution Bulletin*, **38**, 955–959.

- Marcogliese DJ (2001) Implications of climate change for parasitism of animals in the aquatic environment. *Canadian Journal of Zoology*, **79**, 1331-1352.
- Marcogliese DJ (2005) Parasites of the superorganism: Are they indicators of ecosystem health? *International Journal for Parasitology*, **35**, 705–716.
- Marcogliese DJ (2016) The Distribution and Abundance of Parasites in Aquatic Ecosystems in a Changing Climate: More than Just Temperature. *Integrative and Comparative Biology*, **56**, 611-619.
- Marcogliese DJ, Pietrock M (2011) Combined effects of parasites and contaminants on animal health: parasites do matter. *Trends in Parasitology*, **27**, 123-130.
- Marcogliese, D.J., Cone, D.K., (1997). Parasite communities as indicators of ecosystem stress. *Parassitologia*, **39**, 227–232.
- Marcogliese, DJ (2004) Parasites: Small Players with Crucial Roles in the Ecological Theater. *EcoHealth*, **1**, 151-164.
- Marigómez I, Garmendia L, Soto M, Orbea A, Izagirre U, Cajaraville MP (2013) Marine ecosystem health status assessment through integrative biomarker indices: a comparative study after the Prestige oil spill "Mussel Watch". *Ecotoxicology*, 22, 486-505.
- McCarthy HO, Fitzpatrick SM, Irwin SWB (2004) Parasite alteration of host shape: a quantitative approach to gigantism helps elucidate evolutionary advantages. *Parasitology*, **128**, 7-14.
- McCarthy HO, Irwin SWB, Fitzpatrick SM (1999) *Nucella lapillus* as a paratenic host for *Maritrema arenaria*. *Journal of Helminthology*, **73**, 281-282.
- McCarthy T, Hassett D (1993) *Cryptocotyle lingua* (Creplin)(Digenea: Heterophyidae) and other parasites of a coastal otter *Lutra lutra* (L.). *The Irish Naturalists' Journal*, **24**, 280-282.
- Menke KT (1829) Verzeichniss der ansehnlichen Conchylien-Sammlung des Freiherrn von der Malsburg. *Pyrmont*, **6**, 1-123.
- Miramand, P. and Bentley, D. (1992). Heavy metal concentrations in two biological indicators (*Patella vulgata* and *Fucus serratus*) collected near the French nuclear fuel reprocessing plant of La Hague. *Science of the Total Environment*, 111, 135–149.
- Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B (2011) How Many Species Are There on Earth and in the Ocean? *PLoS Biology*, **9**, 1-8.
- Morgan E, O'Riordan RM, Kelly TC, Culloty SC (2012) Influence of disseminated neoplasia, trematode infections and gametogenesis on surfacing and mortality in the cockle *Cerastoderma edule*. *Diseases of Aquatic Organisms*, **98**, 73-84.

- Morley NJ (2010) Interactive effects of infectious diseases and pollution in aquatic molluscs. *Aquatic Toxicology*, **96**, 27-36.
- Morley NJ, Lewis JW (2006) Anthropogenic pressure on a moll uscan-trematode community over a long-term period in the Basingstoke Canal, UK, and its implications for ecosystem health. *Ecohealth*, **3**, 269-280.
- Mouritsen KN, Poulin R (2002) Parasitism, climate oscillations and the structure of natural communities. *Oikos*, **97**, 462-468.
- Mouritsen, KN, Poulin, R (2002) Parasitism, community structure and biodiversity in intertidal ecosystems. *Parasitology*, **124**, 101-117.
- Müller OF, (1776) Zoologiae Danicae Prodromus seu Animalium Daniae et Norvegiae indigenarum characteres, nomina, et synonyma imprimis popularium. *Hafniae*, *Typiis Hallageriis*, **32**, 1-217.
- Muniz P, Venturini N, Hutton M, Kandratavicius N, Pita A, Brugnoli E, Burone L, García-Rodríguez F (2011) Ecosystem health of Montevideo coastal zone: A multi approach using some different benthic indicators to improve a ten-yearago assessment. *Journal of Sea Research*, 65, 38-50.
- O'Connell MP, Fives JM (2004) Helminth communities of the lesser sandeel *Ammodytes tobianus* L. off the west coast of Ireland. *Journal of Parasitology*, **90**, 1058-1061.
- Odum EP (1985) Trends Expected in Stressed Ecosystems. *Bioscience*, **35**, 419-422.
- O'Dwyer K, Kamiya T, Poulin R (2014a) Altered microhabitat use and movement of littorinid gastropods: the effects of parasites. *Marine Biology*, **161**, 437-445.
- O'Dwyer K, Lynch A, Poulin R (2014b) Reduced attachment strength of rocky shore gastropods caused by trematode infection. *Journal of Experimental Marine Biology and Ecology*, **458**, 1–5.
- Oehlmann JBauer B, Minchin D, Schulte-Oehlmann Y, Fioroni P, Markert B (1998) Imposex in *Nucella lapillus* and intersex in *Littorina littorea*: interspecific comparisons of two TBT induced effects and their geographical uniformity. In: O'Riordan, R.M., Burnell, G.M., Davies, M.S., Ramsay, N.F. (Eds.), *Aspects of littorinid Biology*, 199–213.
- Olson PD, Cribb TH, Tkach VV, Bray RA, Littlewood DTJ (2003) Phylogeny and classification of the Digenea (Platyhelminthes : Trematoda). *International Journal for Parasitology*, **33**, 733-755.
- Orias J, Noble E, Alderson G (1978) Parasitism in some East Atlantic bathypelagic fishes with a description of *Lecithophyllum irelandeum* sp. (Trematode). *Journal of Parasitology*, **64**, 49-51.

- Ostfeld R, Keesing F (2000) The function of biodiversity in the ecology of vectorborne zoonotic diseases. *Canadian Journal of Zoology-Revue Canadienne De Zoologie*, **78**, 2061-2078.
- Palm HW, Ruckert S (2009) A new approach to visualize ecosystem health by using parasites. *Trends in Ecology & Evolution*, **105**, 539-553.
- Pietrock M, Marcogliese DJ (2003) Free-living endohelminth stages: at the mercy of environmental conditions. *Trends in Parasitology*, **19**, 293-299.
- Pontoppidan E, (1763) Den Danske atlas eller Konge-Riget Dannemark, med dets naturlige egenskaber, elementer, indbyggere, værter, dyr og andre affødninger, dets gamle tildragelser og nærværende omstændingheder i alle provintzer, stæder, kirker, slotte og herre-gaarde. Forestillet ved en udførlig landsbeskrivelse, saa og oplyst med dertil forfærdigede land-kort over enhver provintz, samt ziret med stædernes prospecter, grund-ridser, og andre merkværdige kaabber-stykker. *Efter Søy-kongelig allernaadigst befalning*, 1,1-723.
- Poulin R (1999) The functional importance of parasites in animal communities: many roles at many levels? *International Journal for Parasitology*, **29**, 903-914.
- Poulin R (2006) Global warming and temperature-mediated increases in cercarial emergence in trematode parasites. *Parasitology*, **132**, 143-151.
- Poulin R (2014) Parasite biodiversity revisited: frontiers and constraints. *International Journal for Parasitology*, **44**,581-589.
- Poulin R (2016). Parasite biodiversity revisited: frontiers and constraints. *International Journal for Parasitology*, **113**, 3-10.
- Poulin R, Blasco-Costa I, Randhawa HS (2016) Integrating parasitology and marine ecology: Seven challenges towards greater synergy. *Journal of Sea Research*, 113, 3-10.
- Poulin R, Mouritsen KN (2006) Climate change, parasitism and the structure of intertidal ecosystems. *Journal of Helminthology*, **80**, 183-191.
- Prinz K, Kelly T, O'Riordan R, Culloty S (2010a) Occurrence of macroparasites in four common intertidal molluscs on the south coast of Ireland. *Marine Biodiversity Records*, **3**, 89-96.
- Prinz K, Kelly T, O'Riordan R, Culloty S (2011) Factors influencing cercarial emergence and settlement in the digenean trematode *Parorchis acanthus* (Philophthalmidae). *Journal of the Marine Biological Association of the United Kingdom*, **91**, 1673-1679.

- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2009) Non-host organisms affect transmission processes in two common trematode parasites of rocky shores. *Marine Biology*, **156**, 2303-2311.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010b) Infection of *Mytilus edulis* by the trematode *Echinostephilla patellae* (Digenea: Philophthalmidae). *Journal of Helminthology*, **84**, 193-198.
- Raftopoulou EK, Dimitriadis VK (2010) Assessment of the health status of mussels *Mytilus galloprovincialis* along Thermaikos Gulf (Northern Greece): An integrative biomarker approach using ecosystem health indices. *Ecotoxicology, and Environmental Safety*, **73**, 1580-1587.
- Rapport DJ, Constanza R, McMichael AJ (1998) Assessing ecosystem health. Trends in Ecology & Evolution, 13, 397-402.
- Rees FG (1934) Cercaria patellae Lebour, 1911, and its effect on the digestive gland and gonads of Patella vulgata. Proceedings of the Zoological Society of London, 104, 45-53.
- Risso A (1810). Ichthyologie de Nice ou histoire naturelle des poissons du département des Alpes-Maritimes. *Schoell*, 1-344.
- Risso A (1827). Histoire naturelle des principales productions de l'Europe Méridionale et particulièrement de celles des environs de Nice et des Alpes Maritimes. Paris, Levrault, **3**, 1-480.
- Rodrigues ET, Pardal MA (2014) The crab *Carcinus maenas* as a suitable experimental model in ecotoxicology. *Environment International*, **70**, 158-182.
- Rohde K (2002) Ecology and biogeography of marine parasites. *Advances in Marine Biology*, **43**, 1-86.
- Sala E, Knowlton N (2006) Global marine biodiversity trends. *Annual Review of Environment and Resource*, **31**, 93-122.
- Say T (1822) An account of some of the marine shells of the United States. *Journal of the Academy of Natural Sciences of Philadelphia*, **2**, 1-325.
- Scaps P (2002) A review of the biology, ecology and potential use of the common ragworm *Hediste diversicolor* (OF Muller) (Annelida : Polychaeta). *Hydrobiologia*, **470**, 203-218.
- Shea J, Kersten J, Puccia CM, Stanton AT, Stiso SN, Helgeson ES, Back EJ (2012) The use of parasites as indicators of ecosystem health as compared to insects in freshwater lakes of the Inland Northwest. *Ecological Indicators*, **13**, 184– 188.

- Small C, Nicholls RJ (2003) A global analysis of human settlement in coastal zones. *Journal of Coastal Research*, **19**, 584-599.
- Sousa WP (1991) Can models of soft-sediment community structure be complete without parasites? *American Zoologist*, **31**, 821-830.
- Sousa WP (1994) Patterns and processes in communities of helminth parasites. *Trends in Ecology & Evolution*, **9**, 52-57.
- Stewart JR, Gast RJ, Fujioka RS, Solo-Gabriele HM, Meschke JS, Amaral-Zettler LA, del Castillo E, Polz MF, Collier TK, Strom MS, Sinigalliano CD, Moeller PDR, Holland AF (2008) The coastal environment and human health: microbial indicators, pathogens, sentinels and reservoirs. *Environmental Health*, 7, 14.
- Stier T, Drent J, Thieltges DW (2015) Trematode infections reduce clearance rates and condition in blue mussels *Mytilus edulis*. *Marine Ecology Progress Series*, **529**, 137-144.
- Strain PM, Macdonald RW (2002) Design and implementation of a program to monitor ocean health. *Ocean & Coastal Management*, **45**, 325-355.
- Sures B (2008) Host-parasite interactions in polluted environments. *Journal of Fish Biology*, **73**, 2133-2142.
- Sures B, Nachev M, Selbach C, Marcogliese DJ (2017) Parasite responses to pollution: what we know and where we go in 'Environmental Parasitology'. *Parasites & Vectors*, **10**, 1-19.
- Thieltges DW (2006) Effect of infection by the metacercarial trematode *Renicola* roscovita on growth in intertidal blue mussel *Mytilus edulis*. Marine Ecology Progress Series, **319**, 129-134.
- Thieltges DW, Bordalo MD, Hernandez AC, Prinz K, Jensen KT (2008) Ambient fauna impairs parasite transmission in a marine parasite-host system. *Parasitology*, **135**, 1111-1116.
- Thieltges DW, Krakau M, Andresen H, Fottner S, Reise K (2006) Macroparasite community in molluscs of a tidal basin in the Wadden Sea. *Helgoland Marine Research*, **60**, 307-316.
- Thieltges DW, Rick J (2006) Effect of temperature on emergence, survival and infectivity of cercariae of the marine trematode *Renicola roscovita* (Digenea: Renicolidae). *Diseases of Aquatic Organisms*, **73**, 63-68.
- Thieltges DW, Wegner KM (2016) Ecology and evolution of marine parasites and diseases: International NIOZ-AWI symposium, 10-14 March 2014, Texel, The Netherlands. *Journal of Sea Research*, **113**, 1-3.

- Thieltges DW, Krakau, M, Andresen, H., Fottner, S and Reise, K (2006). Macroparasite community in molluscs of a tidal basin in the Wadden Sea. *Helgoland Marine Research*, **60**, 307-316.
- Thompson RC, Crowe TP, Hawkins SJ (2002) Rocky intertidal communities: past environmental changes, present status and predictions for the next 25 years. *Environmental Conservation*, **29**, 168-191.
- Thompson RM, Mouritsen KN, Poulin R (2005) Importance of parasites and their life cycle characteristics in determining the structure of a large marine food web. *Journal of Animal Ecology*, **74**, 77-85.
- Thunberg CP, (1793). Tekning och Beskrifning på en stor Ostronsort ifrån Japan. Kongliga Vetenskaps Academiens Nya Handlingar, **14**, 140-142.
- Vidal-Martinez VM, Pech D, Sures B, Purucker ST, Poulin R (2010) Can parasites really reveal environmental impact? *Trends in Parasitology*, **26**, 44-51.
- Ward JR, Lafferty KD (2004) The elusive baseline of marine disease: Are diseases in ocean ecosystems increasing? *PLoS Biology*, **2**, 542-547.
- Wells PG (2003) Assessing health of the Bay of Fundy concepts and framework. *Marine Pollution Bulletin*, **46**, 1059-1077.
- Welsh JE, van der Meer J, Brussaard CPD, Thielges DW (2014) Inventory of organisms interfering with transmission of a marine trematode. *Journal of the Marine Biological Association of the United Kingdom*, **10**, 1-6.
- Wicklum D, Davies RW (1995) Ecosystem health and integrity? *Canadian Journal of Botany*, **7**, 997-1000.
- Williams HH, Mackenzie K (2003) Marine parasites as pollution indicators: an update. *Parasitology*, **126**, 27-41.
- Wood CL, Byers JE, Cottingham KL, Altman I, Donahue MJ, Blakeslee AMH (2007) Parasites alter community structure. *Proceedings of the National Academy of Sciences of the United States of America*, **22**, 9335-9339.

- Wood W, (1828) A supplement to the Index testaceologicus; or a catalogue of shells, British and foreign. London, **4**, 1- 59.
- Worm B, Barbier EB, Beaumont N, Duffy JE, Folke C, Halpern BS, Jackson JB, Lotze HK, Micheli F, Palumbi SR (2006) Impacts of biodiversity loss on ocean ecosystem services. *Science*, **314**, 787-790.
- Zimmer RK, Fingerut JT, Zimmer CA (2009) Dispersal pathways, seed rains, and the dynamics of larval behavior. *Ecology*, **90**, 1933-1947.

CHAPTER 2: Factors affecting parasite infection of three common intertidal gastropods on rocky shores of differing water quality on the coast of Ireland

ABSTRACT

Impacts of anthropogenic activities on marine ecosystems are addressed by legislation to protect and monitor coastal waters. Sentinel organisms have been widely used to indicate anthropogenic disturbance on rocky shores. Parasites on rocky shores are a common, but understudied, occurrence. Parasitic trematodes are a large component of macroparasite communities of intertidal fauna and have been proposed as indicators of ecosystem quality. This study investigated the variation in the abundance and richness of parasites, with a focus on trematodes, of intertidal invertebrates in relation to differing Water Framework Directive classifications. For a year, intertidal gastropods, the common periwinkle Littorina littorea, dog whelk Nucella lapillus, and common limpet Patella vulgata were sampled from three rocky shores with differing classifications. *Littorina littorea* and *P. vulgata* were sampled from a site of high water quality Lough Hyne Marine Reserve. Sampling of L. littorea and P. vulgata was restricted to bimonthly events as stipulated in a National Parks and Wildlife licence to minimise potential detrimental effects. Intertidal biota surrounding three host species were recorded, to evaluate potential disrupters of trematode transmission in the field. In total, 3,973 gastropods were screened and parasite taxa from four phyla were recorded: Annelida, Arthropoda, Ciliophora and Platyhelminthes. Nine trematode taxa from seven families were recorded. Parasite taxa were common and varied greatly over the temporal scales, host species and sites considered. There was no clear pattern in parasite taxa richness between sites related to WFD quality classification. Ciliates and cestodes were less common in the site of high quality. On shores of moderate quality, the highest and lowest number of trematode taxa, the highest and lowest prevalence of primary infections of L. littorea and the lowest prevalence of trematode metacercariae in L. littorea and P. vulgata were recorded. Water quality alone was not
considered a determining factor for richness, prevalence or intensities of parasites recorded.

2.1. INTRODUCTION

Parasites are an understudied but ubiquitous component of ecosystems (Marcogliese 2005, Poulin 1999; 2014, Poulin *et al.*, 2016), estimated to constitute 40% of all known species and form up to 75% of key links in food webs (Dobson *et al.*, 2008, Lafferty *et al.*, 2008). Parasites in contribute biodiversity and productivity to aquatic ecosystems (Poulin, 1999, Mouritsen & Poulin, 2002a). Parasite abundance and dynamics are affected by anthropogenic impacts (Lafferty 1997, Sures, 2008, Sures *et al.*, 2017). Meta-analyses revealed an association between environmental impact and parasite abundance (Lafferty 1997, Sures, 2008, Sures *et al.*, 2017). The magnitude of the association was attributed to the wide range of anthropogenic stressors, as well as to the inherent variability of parasite responses (Lafferty 1997).

Rich or abundant parasite communities are proposed to be indicative of the health or good status of an ecosystem rather than its decline (Marcogliese 2005, Hudson *et al.*, 2006). A heavily impacted or frequently disturbed ecosystem may inhibit the transmission of parasites with free living stages (Lafferty 1997, Sures, 2008, Sures *et al.*, 2017). Trematodes require multiple hosts to complete their life cycle and contain free living stages sensitive to a range of environmental stressors (Lafferty 1997, Pietrock & Mouritsen, 2003, Sures *et al.*, 2017). In the Yucatán, Peninsula, Mexico trematode communities of gastropods took several years to reach pre-impact levels following a hurricane in the area (Aguirre-Macedo *et al.*, 2011). The restoration of a Californian salt marsh, resulted in significant increases in the abundance of trematode communities of the California horn snail, *Cerithideopsis californica*. After six years, the occurrence of trematodes was equivalent to reference zones (Huspeni & Lafferty 2005). This recovery period is likely due to the time required to successfully transmit to multiple hosts.

Abundance of trematodes in invertebrates has been associated with increased water quality. In Ireland, water quality was not recorded to effect the diversity and abundance of parasites harboured by *Mytilus* spp. screened (Lynch *et al.*, 2014). Cercariae of *Cryptocotyle lingua* (Creplin, 1825) in *Littorina littorea* in poor quality water, due to high concentrations of heavy metal pollutants, were observed to have impaired swimming and longevity. This effect of environmental stress would have implications for their ability to find and infect the following host (Cross *et al.*, 2001a; 2003; 2005). In the Mondego Estuary, Portugal an increase in water quality, due to decreased nutrient loading led to heavy infection of the isopod *Cyathura carinata* (Kröyer, 1847) by trematode metacercariae (Ferreira *et al.*, 2007).

In Europe, two key directives for assessing coastal waters are the Water Framework Directive (WFD, 2006/60/EC) and the European Marine Strategy Framework Directive (MSFD, 2008/56/EC). Both the WFD and the MSFD require the development of cost effective monitoring tools to assess ecological status (ES) of European marine waters. In Ireland ES are assessed by the Environmental Protection Agency (EPA) and assigned to one of the five ES categories set out under the WFD, (high, good, moderate, poor and bad). Coastal bodies are assigned to quality classes by the EPA according to biological quality elements, physico-chemical parameters, environmental quality standards and hydromorphological quality elements (Clabby *et al.*, 2008, O'Boyle *et al.*, 2010, Bradley *et al.*, 2015).

The transmission of trematodes could be affected by the (not mutually exclusive) mechanisms of predation, physical disturbance, hyperparasitism, toxic exudates and the infection of unsuitable hosts (Thieltges *et al.*, 2008). Surrounding biota can act as an alternative host or a "decoy" to free living trematode stages. The decoy mechanism has been found in the field in the Wadden Sea with a number of intertidal bivalves. Mouritsen *et al.*, (2003) and Thieltges and Reise (2007) found that cockles *Austrovenus stutchburyi* and *Cerastoderma edule* (Linnaeus, 1758) in close proximity or surrounded by rings of *Mytilus* spp., harboured fewer metacercariae due to the increased infection of the nearby mussels. Laboratory studies have shown that a range

of rocky shore biota can physically disturb and consume trematode cercariae, which impairs the trematode's ability to reach or infect to their second intermediate hosts (Thieltges *et al.*, 2008, Prinz *et al.*, 2009, Studer *et al.*, 2013, Welsh *et al.*, 2014).

Intertidal gastropods have been used in biomonitoring programmes, are key host species for trematodes and influential consumers in the rocky intertidal (Werding 1969, Thompson *et al.*, 2002, Prinz *et al.*, 2010a, O'Connor *et al.*, 2011, Atalah and Crowe, 2012). *Littorina littorea*, *Patella vulgata* and *Nucella lapillus* were chosen as the hosts for this study due to their Atlantic-wide distributions, use in biomonitoring programmes and differing ecological roles of an omnivore grazer, macro-grazer and predator, respectively (Moore 1937, Crothers 1985, Crowe *et al.*, 2000, Jenkins *et al.*, 2001b, Atalah & Crowe, 2010). The aims of the research were to examine the effect of (a) site related water quality classifications according to WFD on parasite communities and (b) investigate the effect of surrounding biotic communities. To account for seasonal variation and host effects, the study was conducted for a year and parasitism in gastropod populations was recorded monthly or bimonthly in some sites.

2.2 MATERIALS AND METHODS

2.2.1 Study Sites

In April 2014, preliminary surveys of seven rocky shores of differing WFD ecological classifications were carried out. The intertidal zone of Cusheen (51°31'47.65"N, 9°31'36.30"W), Cuskinny bay (51°51'26.9"N 8°15'59.0"W), Lough Hyne (51°30'1.55"N, 9°18'22.06"W), Sandycove (51°40'34.20"N, 8°31'24.30"W), Tranabo (51°29'52.7"N 9°17'07"W), Whitegate (51°50'18.11"N, 8°13'45.72"W) and Whiddy Island (51°40'53.5"N 9°30'20.9"W) were surveyed with the methodology detailed in Section 2.2.2 page 40. Shores were chosen to allow comparison between different WFD classifications and sufficient host species populations to support year-long monthly sampling. At the time of surveying only classifications based on data from 2007-2009 were available. No coastal water body was classified as of "poor" or "bad" quality by the Environmental Protection Agency, (EPA) (O'Boyle *et al.*, 2010). From the seven shores, two "moderate" shores, one "good" and one marine reserve of "high" quality were chosen to sample.

Cusheen was a shore of "moderate" quality status. The rocky shoreline had a gentle slope that graduated into sand and mud in the subtidal zone. The sheltered south-west facing rocky shore is situated in Roaringwater bay within a Special Area of Conservation (SAC) (O'Boyle *et al.*, 2010). Lough Hyne is a marine reserve, covers an area of 0.8 km² and is an extremely sheltered shore with restricted tidal exchange with outer waters, due to the dimensions of the connecting channel. Lough Hyne was not evaluated by the EPA but is considered to have high status water quality (Jessopp *et al.*, 2010, Lynch *et al.*, 2014). Sandycove is a semi-exposed south facing shore in Kinsale and is considered to be of "good" quality status by the EPA. Whitegate is a west facing bay on the east side of Cork harbour and is classified as "moderate" quality by the EPA. Whitegate bay is within the boundaries of a Special Protection Area (SPA), contains Ireland's only oil refinery which experienced an oil spill in 1997 (EPA, 1997).



Fig. 1 Map of study sites on the southwest of Ireland (a) Cusheen, (b) Sandycove, (c) Lough Hyne & (d) Whitegate. Adapted from Environmental Protection Agency Water Quality shapefile[®] Ordnance Survey Ireland and Government of Ireland protected by the Copyright Act 1963, Copyright (Amendment) Act 1987, Copyright and Related Rights Act 2000 and EU Copyright and Database Directives.

2.2.2 Field collections

To account for temporal variation a minimum of 30 each of L. littorea, N. lapillus and *P. vulgata* were sampled monthly from Cusheen, Sandycove and Whitegate from May 2014 to April 2015. From May 2014 to March 2015 30 each of L. littorea and P. vulgata were sampled every second month from Lough Hyne as permitted by the National Parks and Wildlife Service. Host densities were calculated per m² from counts within randomised 0.25 m 2 quadrats on the low shore. Low shore was sampled to allow comparison e with Lough Hyne's small tidal range, build on the work of Prinz et al., (2010b) and facilitate the detection of trematodes. Gastropods with shell heights greater than 10 mm were sampled from exposed rock at low shore (Byers et al., 2015). Smaller and sexually immature specimens are less likely be to harbour parasite taxa (Moore, 1937, Lauckner, 1980; 1983). Low shore of the intertidal was considered to the zone between the mean tidal level (MTL) and to mean of the low water spring tides (MLWS). To minimise the variation in microhabitat of gastropods, rock pools were excluded from sampling. Intertidal biota known to affect the transmission of trematodes were recorded, (see Results Table 3) (Thieltges et al., 2008, Prinz et al., 2009, Studer et al., 2013, Welsh et al., 2014). Barnacle species were classified on the SACFOR abundance scale according to Crisp & Southward (1958) and modified by Hiscock (1981) with S representing super abundant, A: abundant, C: common, F: frequent, O: occasional, and R: rare (SACFOR). The presence of polychaetes, decapod crabs, anemones and non-host gastropods was noted. Algal species were recorded as percentage cover in 0.25 m^2 quadrats.



Fig. 2 Schematic diagram of sampling programme for (a) the monthly collection of *Littorina littorea, Nucella lapillus* and *Patella vulgata* and (b) bimonthly sampling of *N. lapillus* and *P. vulgata* from marine reserve Lough Hyne. Asterisks denote the subsampled data from Cusheen, Sandycove and Whitegate for comparison with Lough Hyne

2.2.3 Processing

In the laboratory encrusting biota were removed from shells and discarded. This was done to avoid interference of encrusting biota with gastropod trematodes (Prinz et al., 2009, Welsh et al., 2014) and to avoid including parasites of encrusting biota in counts. Gastropods were weighed to the nearest 0.01 g on an electronic balance (OHAUS Adventurer[®] Pro balance model AV3102). Prior to dissection, shell height, length, and width were measured to the nearest 0.05 mm with Vernier callipers (Draper® 4817P) (see Fig. 2). Shells were cracked or removed from gastropods by first severing the abductor muscle. A second incision was then made to sever the cerebral ganglion, then gastropods were sexed: L. littorea according to Graham (1969), Gibbs et al., (1988) and Reid et al., (1996), N. lapillus according to Bettin et al., (1996), and P. vulgata according to Orton (1919) and Blackmore (1969). Patella vulgata is a sequential hermaphrodite where male gonads develop prior to changing to female gonads (protandry). Patella vulgata was classified as female with the presence of greenish eggs, male with orange sperm, or hermaphroditic with both eggs and sperm. If neither were present, limpets were classified as undifferentiated (Orton, 1919, Blackmore, 1969). The visceral mass and foot tissues of *P. vulgata* were separated prior to compression. Whole live mounts were compressed between 1/4 inch thick glass slides (Kompressorium 23 x 5 cm Hauptner-Herberholz[®]) and examined using transmitted light with a stereomicroscope (Nikon SMZ645) and compound microscope (Nikon Alphaphot-2 YS2-H) at magnifications of x8 and x40 respectively. Parasitological terminology followed that of Bush et al., (1997). Prevalence (P) was defined as the percentage of infected hosts and intensity (Ins) was defined as the average number of parasites within infected hosts. Helminth parasites were counted individually, whereas only the presence or absence of ciliates, cestodes, copepods, mites, nematodes, polychaetes parasite taxa were recorded. This was due to the inability to distinguish individuals for counts. Images were taken with the compound microscope Nikon ECLIPSE 80i and NIS-Elements Microscope Imaging Software.



Fig. 3 Illustrations of morphometric measurements taken from (a) *Littorina littorea* and *Nucella lapillus*, modified from Reid (1996), and (b) & (c) *Patella vulgata* modified from Orton (1919). SH represents Shell height, SL Shell length and SW Shell width.

2.2.4 Parasite identification

Parasite taxa were identified by morphology according to Nicoll (1907), Stunkard (1950), Crewe (1951), James (1968a), Werding (1969), Lauckner (1980), MacKinnon & Burt (1983), Kollien (1996) and Buckland-Nicks *et al.*, (2013). Cestode taxa were identified by the lack of an alimentary canal and the presence of the protoscolex (precursor to attachment sucker) and cercomer (caudal appendage). Larval trematodes were distinguished according to their size and if present; acetabula (suckers), colouration, excretory systems, tail, stylet and eyespots. For metacercariae the cyst wall or collar spines, again if present, were also used to distinguish between taxa. As has been done in previous studies *Cercaria parvicaudata* (Stunkard and Shaw, 1931) and *Renicola roscovita* (Stunkard, 1932) primary intermediate infections were grouped as *Renicola* spp. due to the subjectivity of differentiating between the colours yellow and orange. *Himasthla* and *Gymnophallus* species were also grouped in their respective genera. Trematode identification was confirmed to the family level by Drs David Thieltges and Xavier de Montaudouin (pers. comm.).

2.2.5 Statistical Analysis.

Analysis was carried out in R studio version 1.0.136 (R Core Team 2015). Parasitology terms follows Bush *et al.*, (1997). Statistical significance was considered with P<0.05. When comparing with the marine reserve, Lough Hyne, a subset of data from the three shores was used (Fig.3). Normality of distributions was assessed first visually and then formally by the Shapiro–Wilk test. Normality of distributions was assessed first visually and then formally by the Shapiro–Wilk test. Homoscedasticity was determined using a Levene's Test for Equality of Variances. Prior to analysis, non-normal data were transformed; count data were transformed via square root, proportional data transformed via arcsine and continuous data were transformed via log. If transformation failed to normalise the distribution of data, non-parametric tests were carried out.

2.2.5.1 Summary statistics

Normally distributed data was summarised with an arithmetic mean and standard deviation. For parasite prevalence, non-normally distributed data, 95% Clopper Pearson Confidence intervals have been found to be conservative and appropriate measure of variation (Reiczigel 2003, Reiczigel *et al.*, 2008). 95% Clopper Pearson Confidence were calculated with the r package "PropCIs" where presence/absence of parasite taxa were considered to be success/fail of a Bernoulli experiment. Parasite intensity typically has highly aggregated (right-skewed) frequency distribution and so was summarised by a median and the range of infection (Rozsa *et al.*, 2000).

2.2.5.2 Effect of quality

Gastropod host densities and morphometric measurements were compared with the parametric one-way analysis of variance (ANOVA) and a post hoc Tukey Honest Significant Differences (Tukey HSD) test. (Underwood 1997). ANOVA and Tukey HSD were carried out with the package "car" and "stats", respectively.

Comparisons of parasite prevalences and intensities were done with nonparametric test a Pearson's chi square with Yates continuity correction and Kruskal-Wallis chi square, respectively. If expected values were too low (<5) for a Pearson's chi square, a Fisher's Exact test was used. Posthoc comparisons were made with the nonparametric Pairwise Wilcoxon Rank Sum Tests and carried out with the r package "stats".

2.2.5.3 Effect of surrounding biota

Pearson Correlations were used to investigate relationships between parasite taxa their host morphometrics, abundance surrounding biota and co-occurring parasite taxa. The effects of month, site and size of gastropods on the composition of parasite communities were investigated using non-metric multi-dimensional scaling (nMDS) and non-parametric one-factorial or two-factorial PERMANOVA procedures based on Jaccard similarity indices of non-transformed prevalence or intensity data. The effect of sampling effort (number of quadrats or hosts) on the number of parasite species across all sites, separately for each gastropod species was evaluated by Pearson correlations. was used for statistical comparison with the marine reserve.

2.3. RESULTS

2.3.1 Host morphometrics

The smallest *L. littorea* sampled were from Lough Hyne (F=25.43, df=3, P<0.0001) (Table 1). Significantly larger specimens (F=84.62, df=2, P<0.0001) and higher densities (F=25.43, df=2, P<0.0001) of *L. littorea* were sampled from and recorded in Sandycove (Table 1). Larger *N. lapillus* specimens were sampled from Whitegate and Cusheen relative to Sandycove (F=10.19, df=2, P<0.0001), while significantly higher densities of *N. lapillus* were recorded in Sandycove relative to Whitegate (F=5.18, df=2, P<0.001). Larger *P. vulgata* were sampled from Whitegate relative to the other sites (F=13.42 df=2, P<0.0001). Significantly higher densities of *P. vulgata* were recorded in Sandycove relative to other sites (F=133.4, df=2, P<0.0001). The lowest density of *P. vulgata* recorded was in Lough Hyne (F=125.5, df=3, P<0.0001).

Host	Site						
	Cusheen	Lough Hyne	Sandycove	Whitegate			
Littorina littorea							
n	498	195	400	479			
Shell height	18.32 (± 4.51)	17.11 (± 5.00)	24.39 (±5.70)	21.68 (± 10.10)			
Density	42 (± 46)	19 (± 34)	17 (± 36)	16 (± 27)			
Nucella lapillus							
n	273	NA	301	382			
Shell height	28.47 (± 5.83)	NA	26.15 (±5.17)	28.51 (± 5.65)			
Density	9 (± 10)	NA	$10 (\pm 15)$	13 (± 16)			
Patella vulgata							
n	429	216	410	390			
Shell height	13.47 (± 4.91)	12.05 (± 4.29)	12.14 (±5.78)	14.28 (± 6.14)			
Density	58 (± 42)	41 (±32)	62 (± 54)	24 (± 23)			

Table 1. The size and density $(\bar{x}\pm SD)$ per m² of gastropod host screened at each location and for all months combined 2014/15.

2.3.2 Parasite taxa

Of the host species screened, five macroparasite taxa were recorded; halacarid mites, the burrowing polychaete *Polydora ciliata* (Johnston, 1838), harpacticoid copepods, the cestode *Ophryocotyle insignis*, Loennberg, 1890, larval nematodes and nine genera of juvenile trematodes. One microparasite taxon was recorded; ciliates were recorded and were predominantly in *L. littorea* and *P. vulgata* (Table 1). Prevalence of parasite infection significantly differed between gastropod species (χ^2 =1007.2, df=2, P<0.0001) (Fig 3 (a), (b)). *Patella vulgata* and *N. lapillus* had respectively the highest and lowest prevalence of infection (P<0.0001) (Fig.4). Co-infection between parasitic taxa was significantly more common in *P. vulgata* than in other gastropod hosts screened (χ^2 =531.7, df=2, P<0.0001) (Fig. 4). Ciliates, most likely the species *Leiotrocha patellae* (Cúenot, 1891), and trematodes were the two most commonly cooccurring parasite taxa in *P. vulgata*. The prevalence of ciliates and trematodes in *P. vulgata* were positively correlated (r=0.09, P<0.05).

2.3.2.1 Non-Trematode parasite taxa

Ciliates were the second most common parasite taxa recorded, 16% of all gastropods screened harboured ciliates (Fig. 4). The polychaete *P. ciliata* occurred in all hosts at all sites and had an overall prevalence of 10% (Fig.4). Halacarid mites occurred in all host species. The cestode *O. insignis* solely occurred in *P. vulgata* and only in 3% of specimens screened. *Patella vulgata*, relative to the other host species, were significantly more infected by copepods (χ^2 =79.91, df=2, P<0.001), ciliates (χ^2 =189.81, df=2, P<0.0001), mites (χ^2 =657.68, df=2, P<0.001). Nematode prevalence was low (<1%) and did not differ between host species screened.



Fig. 4 (a) Overall prevalence of parasite taxa in the three gastropod hosts screened per site for all months combined. C represents Cusheen, S Sandycove and W Whitegate. (b) Overall prevalence of parasite taxa in three gastropod hosts from each site with bimonthly dataset for comparison with Lough Hyne Marine Reserve, LH. NS represents not sampled, as *Nucella lapillus* was not sampled from Lough Hyne due to insufficient population size.

2.3.2.2 Trematodes

Trematodes were the most common parasite recorded in the study for each host species (Fig. 4). The nine trematode taxa recorded were from seven families Echinostomatidae, Heterophyidae Renicolidae, Philophthaimidae, Notocotylidae, Opecoelidae, and Fellodistomidae. *Littorina littorea* had the richest trematode community of all hosts screened (Table 2). *Littorina littorea* were mainly second intermediate hosts for trematodes, with 7.69% infected with metacercariae and 3.65% with rediae or sporocysts. *Nucella lapillus* had the lowest richness and prevalence of trematode taxa with only 2.31% infected. *Patella vulgata* was the most commonly infected host and, overall, 41% harboured trematodes. *Echinostephilla patellae* (Fig. 6(b)) and *Renicola* spp. (Fig. 6(c)) metacercariae dominated trematode communities of *P. vulgata* and *L. littorea*, respectively (Table 2). Co-infection between two trematode species was rare and was found only in the form of *Renicola* spp. metacercariae and *Cryptocotyle lingua* sporocysts in *L. littorea*.

Parasite taxa in both *L. littorea* and *P. vulgata* varied between months sampled (Figs. 7&8). In all sites sampled, temporal variation in primary trematode infection in *L. littorea* was not statistically significant (Fig. 8). *Echinostephilla patellae* prevalence significantly differed between months for Cusheen (χ^2 =54.82, df=11, P<0.0001), Lough Hyne (χ^2 =21.90, df=5, P<0.001), Sandycove (χ^2 =45.57, df=11, P<0.0001) and Whitegate (χ^2 =73.23, df=11, P<0.0001).

Chapter 2

Table 2. The prevalence and range (min-max) of intensity in brackets of trematode taxa recorded in gastropod species screened. Calculated for all sampling events in 2014/15 combined. NS denotes not-sampled as populations of *Nucella lapillus* in Lough Hyne were insufficient to allow yearlong sampling

Host species	Trematode species	Cusheen	Lough Hyne	Sandycove	Whitegate	Total prevalence
SPOROCYST						
(Primary intermediate	hosts)					
Littorina littorea	<i>Paramonostomum chabaudi</i> ^{*1} (Van Strydonck, 1965)	0.00	0.00	0.50	0.00	0.20
	Cryptocotyle lingua	0.87	3.76	1.49	1.41	1.63
	Himasthla spp. (Dietz, 1909)	0.40	0.47	0.00	0.00	0.27
	Podocotyle atomon (Rudolphi, 1802)	0.22	0.47	1.24	3.74	1.56
	Renicola spp.	0.66	0.00	0.00	0.00	0.07
Nucella lapillus	Parorchis acanthus (Nicoll, 1906) Nicoll, 1907	0.00	NS	0.33	0.00	0.00
METACERCARIAE						
(Second intermediate h	losts)					
L. littorea	Renicola spp. (Stunkard, 1932)	15.72 (1-875)	3.76 (1-5)	6.70 (1-133)	1.17 (1-8)	7.13
	Echinostephilla patellae	0.93 (1-8)	0.47 (1)	0.75 (1)	0.47 (1-6)	0.68
	Himasthla spp.	0.00	0.47(1)	0.00	0.00	0.06
N. lapillus	Renicola spp.	1.20 (1-2)	NS	3.96 (1-12)	0.21(1)	1.26
-	E. patellae	1.12 (1-4)	NS	0.00	0.00	0.31
Patella vulgata	E. patellae	48.60 (1-88)	26.61 (1-35)	47.60 (1-131)	26.59 (1-44)	40.11
0	<i>Gymnophallus</i> spp. ^{*2} (sensu, Crewe, 1951)	0.25 (103)	0.00	0.00	0.00	0.07
	Renicola spp.	0.79 (1)	0.00	0.00	0.28 (1)	0.28
ADULT WORM	~~	• •				
N. lapillus	Proctoeces maculatus ^{*3} (Looss, 1901)	0.44 (1-2)	NS	0.00	0.00	0.10
P. vulgata		0.23 (1)	0.00	0.00	0.00	0.07

*1 formally *Cercaria lebouri* (Van Strydonck, 1965)
*2 "Cercariae B"

*3 formally *Proctoeces buccini* (Loos-Frank, 1969)

Chapter 2



Fig. 5. Photomicrographs of trematode cercariae of the following taxa, scale cars in brackets (a) *Cryptocotyle lingua* (100 μ m) (b) *Podocotyle atomon* (100 μ m), (c) *Himasthla* spp. (100 μ m), (d) *Parorchis acanthus* (200 μ m), (e) *Renicola* spp. (100 μ m) and (f) *Paramonostomum chabaudi* (100 μ m)

Chapter 2



Fig. 6. Photomicrographs of helminth taxa recorded in the with study, scale bar measurements provided in brackets. (a) metacercariae of *Gymnophallus* spp. (50µm) reported as "Cercaria B" by Crewe (1951) (b) metacercariae of *Echinostephilla patellae* from Prinz *et al.*, (2009) (50µm) (c) metacercariae *Renicola* spp. (200µm) (d) cysticercoid of *Ophryocotyle insignis* (200µm) (e) Adult worm of *Proctoeces maculatus* Removed from *Nucella lapillus* (1000µm) (f) Juvenile stage of a nematode (200µm).



Fig. 7 Temporal variation in parasite taxa in *Patella vulgata* in (a) Cusheen, (b) Lough Hyne, (c) Sandycove and (d) Whitegate. NS represent not sampled as Lough Hyne was only sampled & surveyed every second month. Error bars are 95% Clopper Pearson Confidence Intervals.



Fig. 8 Temporal variation in primary intermediate infection of *Littorina littorea* in (a) Cusheen, (b) Lough Hyne, (c) Sandycove and (d) Whitegate. NS represent not sampled as Lough Hyne was only sampled & surveyed every second month. Error bars are 95% Clopper Pearson Confidence Intervals.

2.3.3 Effect of water quality

2.3.3.1 Non-trematode parasites

Prevalence of ciliates in *P. vulgata* significantly differed between non-reserve sites $(\chi^2=27.37, df=2, P<0.001)$ and the marine reserve $(\chi^2=36.64, df=3, P<0.001)$. Significantly lower prevalence of ciliates was recorded in the sites of high quality, Lough Hyne and Sandycove, relative to the site of moderate quality, Whitegate (Fig. 4). The prevalence of *P. ciliata* in the shells of hosts differed significantly between sites for *L. littorea* ($\chi^2=16.31$, df=2, P=1), *N. lapillus* ($\chi^2=39.47$, df=2, P<0.001) and *P. vulgata* ($\chi^2=29.08$, df=2, P<0.001).

Cestode prevalence in *P. vulgata* differed between the three sites (χ^2 =53.56, df=2, P<0.0001) and the marine reserve (χ^2 =29.91, df=3, P<0.001). Highest (7.40%) and lowest (0.03%). cestode prevalences were recorded in the sites of moderate quality. Copepod prevalence in *P. vulgata* did differ between sites (χ^2 =5.89, df=2, P=0.05) and the marine reserve (χ^2 =5.88, df=3, P=0.05). Halacarid mites occurred in all hosts and significantly higher prevalences were recorded in the site of good quality, relative to sites of moderate quality (χ^2 =12.80, df=2, P<0.05). Nematode prevalence was low for all hosts (<1%) and did not differ between sites or gastropod hosts.

2.3.3.2 Trematode parasites

Highest (n=7) and lowest (n=4) trematode richness were both recorded in sites of moderate quality. Seven trematode taxa were record in Cusheen, six in Sandycove, five in Lough Hyne and four in Whitegate (Table 2). Highest and lowest prevalences of trematodes were also recorded in sites of moderate quality, Cusheen and Whitegate, respectively (Fig. 4) (χ^2 =99.56, df=2, P<0.0001). This was also the case with the subsampled dataset that was used for comparison with Lough Hyne (χ^2 =77.34, df=3, P<0.0001). Trematode prevalence in *L. littorea* and *P. vulgata* significantly differed between the four sites (χ^2 =77.39, df=3, P<0.0001). The prevalence of trematodes significantly differed between sites for all hosts, *L. littorea* (χ^2 =23.91, df=2, P<0.001),

N. lapillus (χ^2 =14.48, df=2, P<0.001) and *P. vulgata* (χ^2 =27.37, df=2, P<0.001). Trematodes were more common in *N. lapillus* collected in Cusheen (P<0.001) and Sandycove (P<0.05) relative to Whitegate but overall had low prevalences (Fig. 4). *Nucella lapillus* is not discussed in further detail below as the gastropod species was rarely parasitised and by a limited number of trematode taxa (Table 2).

Littorina littorea was the most common primary intermediate host of trematode taxa (Table 2). Trematode richness in *L. littorea* did not significantly differ between sites. The highest richness (n=4) and lowest (n=2) number of trematode taxa were observed in sites of moderate quality (Table 2). The prevalence of primary intermediate infection in L. littorea did differ between sites, but the difference was only statistically significant between sites of moderate quality (χ^2 =6.80, df=2, P<0.05) (Fig. 9). Paramonostomum chabaudi (Fig 5 (f)) was significantly more prevalent in the site of good quality relative to moderate quality sites (χ^2 =6.45, df=2, P<0.05). *Podocotyle* atomon (Fig 5 (b)) was significantly less prevalent in the site of good quality $(\chi^2=12.30, df=2, P<0.05)$. Renicola spp. (Fig 5 (e)) and C. lingua (Fig 5 (a)) primary infections of L. littorea were more common in the site of good quality (Table 2) but did not significantly differ. Primary infection of L. littorea were grouped by the trematode's second intermediate host did vary in relation to quality status (Fig. 9). The highest number of groups were recorded in Sandycove and the lowest number in Whitegate (Fig. 9). Trematode taxa that infect molluscs were significantly more prevalent in Cusheen (χ^2 =51.59, df=3, P<0.0001). Trematode taxa that encyst, external to host tissues, on hard substrates were only recorded in Sandycove (Fig 9).



Fig. 9 The prevalence of trematode primary intermediate infections in *Littorina littorea* classified by their second intermediate host required in their lifecycle. The four primary intermediate host groups were fishes, crustaceans, molluscs, and hard substrates. Values were based on pooled total infections of all *L. littorea* per site.

The occurrence of trematode metacercariae in *L. littorea* significantly differed between sites of moderate quality, with lower prevalence recorded in Whitegate relative to Cusheen (χ^2 =51.38, df=3, P<0.0001). *Renicola* spp. metacercariae were the most common infection of *L. littorea* utilised as second intermediate host (Table 2). Prevalence of *Renicola* spp. metacercariae significantly differed between sites, with the highest and lowest prevalences recorded in sites of moderate quality (χ^2 =57.20, df=2, P<0.001) (Fig. 9). Prevalence of trematodes in *L. littorea* was higher in Cusheen relative to Sandycove (P<0.05) and Whitegate (P<0.001). When compared with Lough Hyne, highest and lowest prevalence of *Renicola* spp. was recorded in sites of moderate quality. Prevalence of trematodes in *L. littorea* in the marine reserve did not differ from other sites.

Patella vulgata was the most common second intermediate host for all sites and months sampled. *Echinostephilla patellae* prevalence differed between sites $(\chi^2=56.68, df=2, P<0.001)$ and Lough Hyne $(\chi^2=56.68, df=3, P<0.001)$ (Table 2 & Fig.

10). When compared with Lough Hyne the most significant differences were seen between high and moderate quality (P<0.001).



Fig. 10 (a) Prevalence (black) and median intensity (grey) of *Echinostephilla patellae* metacercariae in *Patella vulgata*. (b) Prevalence (black) and median intensity (grey) of *Renicola* spp. metacercariae in *Littorina littorea*. Error bars are 95% Clopper-Pearson 95% Confidence Intervals (CI). Prevalence and intensity estimated from all sampling events. Note the differing y-axes between (a) and (b).



2.3.4 Effect of local biotic factors and trematode infection

Echinostephilla patellae in *Patella vulgata* and *Renicola* spp. in *L. littorea* were the most common trematode taxa recorded in the study. For this reason, subsequent analysis in relation to local biotic factors focused on *E. patellae* and *Renicola* spp. metacercariae. Higher densities of *P. vulgata* were associated with higher prevalences of *E. patellae* in Sandycove (r=0.12, df=408, P<0.05). The positive association *Echinostephilla patellae* prevalence and *P. vulgata* density was observed in the three remaining sites but was not statistically significant. The size of *P. vulgata* also positively correlated with *E. patellae* prevalence in Sandycove (r=0.24, df=438, P<0.0001) and Whitegate (r=0.25, df=359, P<0.0001). Higher intensities of *E. patellae* metacercariae were recorded in larger *P. vulgata* (r=0.15, df=1444, P<0.001).

In all sites sampled, there was a negative correlation between *E. patellae* intensity with *P. vulgata* density, but this was statistically insignificant. In Cusheen, where *Mytilus* spp. were common (Table 3), the intensity of *E. patellae* significantly and negatively correlated with the bivalve's density (r=-0.13, df=391, P<0.05). The negative correlation was found in all remaining sites but was not statistically significant. Perennial alga percentage cover negatively correlated *E. patellae* intensity with but was only statistically significant in Lough Hyne (r=-0.20, df=212, P<0.05). Increased coverage of ephemeral algal coverage negatively correlated with *P. vulgata* shell height (r=-0.07, P<0.05).

Renicola spp. metacercariae positively correlated with their host *L. littorea* density (r=0.10, P<0.001). No significant correlations were found between the remaining ambient biota and *Renicola* spp. metacercariae in *L. littorea*. More trematodes were recorded in larger *L. littorea*. *Cryptocotyle lingua* positively correlated with *L. littorea* shell height (r=0.10, P<0.001), width (r=0.09, P<0.001), whole weight (r=0.09, P<0.001), tissue weight (r=0.08, P<0.001) and negatively with *L. littorea* density (r=-0.33, P<0.001). Prevalence of *C. lingua* positively correlated with ephemeral algal species percentage cover (r=0.14, P<0.001). It can be seen from the nMDS that infection status of *P. vulgata* had a stronger effect on the grouping than site (Fig 11 a & b).

Table 3 Recorded biota for in study sites in 0.25 m 2 quadrats from May 2014 until April 2015. Abundance was calculated from a total of 117 quadrats in Cusheen, 49 in Lough Hyne, 118 in Sandycove and 111 in Whitegate. Units are represented as 1 is presence/absence, 2 is SACFOR, 3 is counts per 1 m 2 , and 4 is percentage cover.

Biota recorded	Functional group	Unit	Cusheen	Lough Hyne	Sandycove	Whitegate
Actinia equina	Anemone	1	0.06	0.03	0.27	0.02
(Linnaeus, 1758)						
Anemonia	Anemone	1	0.01	0.01	0.17	0.00
viridis (Forsskål,						
1775)						
Austrominius	Barnacle	2	0.69	0.22	0.00	0.64
modestus (Darwin,						
1854)						
Chthamalus	Barnacle	2	0.31	0.02	0.02	0.15
montagui						
Southward, 1976						
Chthamalus	Barnacle	2	0.02	0.01	0.00	0.03
stellatus (Poli,						
1791)						
Semibalanus	Barnacle	2	0.13	0.00	0.00	0.03
balanoides						
(Linnaeus, 1767)						
Cancer pagurus	Decapod	1	0.00	0.02	0.03	0.02
Linnaeus, 1758	crab					
Carcinus maenas	Decapod	1	0.02	0.02	0.03	0.21
_	crab					
Pagurus sp.	Decapod	1	0.01	0.00	0.03	0.03
Fabricius, 1775	crab					
Littorina littorea	Host	1	0.75	0.19	0.54	0.65
	gastropod		0 <i></i>	0.00	0	0.67
Nucella lapillus	Host	3	0.65	0.09	0.62	0.65
	gastropod	2	0.05	0.07	0.07	0.74
Patella vulgata	Host	3	0.95	0.37	0.86	0.76
T (gastropod	4	0.01	0.04	0.00	0.01
<i>Ectocarpus</i> sp.	Ephemeral	4	0.01	0.04	0.00	0.01
Lyngbye, 1819	alga	4	0.00	0.11	0.14	0.11
Ulva spp. 1752	Ephemeral	4	0.00	0.11	0.14	0.11
Linnaeus, 1753	alga	2	0.72	0.05	0.02	0.07
<i>Myttilus</i> spp.	Paratenic	3	0.72	0.05	0.03	0.07
Cildente ann	nost Deveterrie	1	0.40	0.12	0.59	0.57
Gibbula spp.	Paratenic	1	0.49	0.12	0.38	0.57
KISSO, 1820	nost Dereterie	1	0.07	0.02	0.06	0.11
Lillorina fabalis	raratemic bost	1	0.07	0.05	0.00	0.11
(1011011, 1823) Littoring obtugate	Dorotonio	1	0.23	0.03	0.25	0.43
(Linnaeus 1758)	host	1	0.23	0.05	0.23	0.43
(Linnacus, 1730)	11051					

Table 3 Continued						
Biota recorded	Functional	Unit	Cusheen	Lough	Sandycove	Whitegate
	group			Hyne	-	_
Littorina saxatilis	Paratenic	1	0.42	0.06	0.04	0.05
(Olivi, 1792)	host					
Ascophyllum	Perennial	4	0.26	0.23	0.02	0.17
nodosum	alga					
(Linnaeus) Le						
Jolis, 1863						
Fucus serratus	Perennial	4	0.09	0.01	0.49	0.75
Linnaeus, 1753	alga					
Fucus spiralis	Perennial	4	0.01	0.04	0.08	0.03
Linnaeus, 1753	alga					
Fucus vesiculosus	Perennial	4	0.21	0.16	0.21	0.17
Linnaeus, 1753	alga					
<i>Laminaria</i> sp.	Perennial	4	0.00	0.01	0.02	0.03
	alga					
Spirobranchus sp.	Polychaete	1	0.07	0.07	0.22	0.41
Blainville, 1818						
<i>Spirorbis</i> sp.	Polychaete	1	0.10	0.15	0.36	0.57
Daudin, 1800						
Chondrus sp.	Turf alga	4	0.03	0.01	0.14	0.05
Stackhouse, 1797						
Cladophora sp.	Turf alga	4	0.05	0.20	0.02	0.01
Kützing, 1843						
Mastocarpus sp.	Turf alga	4	0.03	0.02	0.25	0.07
Kützing, 1843						



Fig. 11 Nonmetric multi-dimensional scaling (nMDS) plots comparing the parasite species composition in (a) *Littorina littorea* and (b) *Patella vulgata* among the four rocky shores in SW Ireland sampled 2014-2015. Plots are based on Jaccard similarities and presence/ absence data. Stress values of all non-metric multidimensional scaling ordinations (nMDS) were below 0.2 and so were considered to represent data accurately in the ordination.



Fig. 12 Nonmetric multi-dimensional scaling (nMDS) plots of (a) *Littorina littorea* and (b) *Patella vulgata* among the four rocky shores in SW Ireland sampled 2014-2015. Symbols denote sites sampled and colours deonote the presence (black) and absence (grey) of trematode metacercariae of (a) *Renicola* spp. or (b) *Echinostephilla patellae*. Plots are based on Jaccard similarities and presence/ absence data. Stress values of all non-metric multidimensional scaling ordinations (nMDS) were below 0.2 and so were considered to represent data accurately in the ordination.

2.4. DISCUSSION

This study illustrates the complexity of the response of recorded parasite taxa to quality. There was no clear pattern in parasite taxa richness between sites related to WFD water quality classification. Ciliates and cestodes were less common in the sites of high quality. Highest trematode occurrence and richness were observed in the sites of moderate quality. The functional groups of trematodes utilising *L. littorea* as a primary intermediate host also varied in relation to quality classifications, with higher richness found in the site of good quality. However, the highest and lowest prevalence of primary intermediate infection of *Littorina littorea* were observed on shores of moderate quality. Trematode taxa *Echinostephilla patellae* and *Renicola* spp. were more prevalent on shores of good quality.

Surrounding intertidal biota were expected to decrease intensity of trematode metacercariae, as this has been demonstrated in multiple laboratory trials (Hopper et al., 2008, Thieltges et al., 2008, Prinz et al., 2009, Studer et al., 2013, Welsh et al., 2014). The effect of biota on trematode infection may be context specific. The association between host density and prevalence of *E. patellae* and *Renicola* spp. was not consistent between sites or hosts, P. vulgata and L. littorea respectively. Furthermore, the effect of surrounding biota on trematode transmission would likely depend on the abundance of the biota. In this study, the correlation between decreased E. patellae intensity and increased number of Mytilus spp. was only significant on the shore where the bivalve was abundant. A possible reason for the decreased intensity of E. patellae in P. vulgata associated with patches of Mytilus spp. is that E. patellae cercariae may be physically disturbed in the water column or infect the bivalve instead of P. vulgata (Thieltges et al., 2008, Prinz et al., 2009, Welsh et al., 2014). Intertidal Mytilus spp. have been recoded to serve as second intermediate hosts for Echinostephilla patellae (Prinz et al., 2009; 2010) and can physically interfere with the swimming and subsequent infection of cercariae through the action of filter feeding (Prinz et al., 2009). Patella vulgata and N. lapillus were the most and least parasitised host species of the study. Nucella lapillus was predominately parasitised by the

burrowing polychaete, *P. cilata* but rarely by other parasite taxa. This difference is likely be due to the immunocompetence of the gastropods and differing encounter rates between the gastropods and their parasite taxa. *Nucella lapillus* within the size range sampled and geographic region of the study, have been recorded to be commonly infected by a range of parasite taxa (Ebling *et al.*, 1964, Crothers, 1985, Prinz*et al.*, *et al.*, 2010). Final hosts of trematodes are considered a determining factor in trematode distribution and prevalence (Kube *et al.*, 2002, Hechinger & Lafferty, 2005, Feis *et al.*, 2015). Thus, the observed difference in trematode occurrence in *N. lapillus* relative to *P. vulgata* may be due to differing encounter rates with avian final hosts. Evidence to support this suggestion is the preference of the Eurasian oystercatcher *Haematopus ostralegus* (Linnaeus, 1758), a final host of the trematode *Echinostephilla patellae*, to consume *Patella* spp. over *N. lapillus*. The preference is attributed to the unpalatable secretions that *N. lapillus* produces to bore into molluscs' shells (Feare, 1970, Kollien, 1996, Hawkins, 1999).

Littorina littorea was parasitized by the highest number of trematode taxa. Its trematode community structure was similar to that of the Atlantic and Wadden Sea; *C. lingua* was the dominant species utilsing *L. littorea* as a primary intermediate host. In terms of abundance it is followed by *Renicola* spp., *Himasthla* spp., *P. chabaudi* and *P. atomon* (James 1968a, Thieltges *et al.*, 2006, Prinz *et al.*, 2010a). A higher prevalence of mollusc and crustacean infecting primary intermediate infections was recorded in Cusheen and Whitegate. This, may be due to the higher abundance of the trematodes' *Renicola* spp., and *P. atomon* second intermediate hosts, *C. maenas* and *Mytilus* spp., observed on Cusheen and Whitegate, respectively.

The recorded abundance of trematodes from the families Echinostomatidae (e.g. *E. patellae, Renicola* spp) and Heterophyidae (*C. lingua*), and low number of co-infections could be due to competitive interactions between trematode taxa within the host. Taxa within these families have rediae, "a soldier caste", that consume heterospecifics and conspecifics from other infections (Hechinger *et al.*, 2011, Garcia-Vedrenne *et al.*, 2016; 2017). The "soldier" caste of treamtode rediae were observed

within primary intermediate hosts and have been recorded to differ from reproductive rediae in their morphology, spatial distribution, and behaviour (Hechinger *et al.*, 2011). *Echinostephilla patellae* and *C. lingua* have redial stages, however, competitive interactions between their rediae and other trematode infections have yet to be observed (Kollien 1996, Rees 1934, Stunkard 1930). This study provides a conservative estimate of the trematodes *Paramonostomum chabaudi* and *Parorchis acanthus* as their metacercariae enycyst in the external environment and were not observed.

Echinostephilla patellae was the most common trematode of the study. Prevalences and intensities of *E. patellae* recorded in this study were higher than in previous work (see Table 4). This is likely due to the different shore heights sampled, a different examination method and the inclusion of metacercariae in counts (see Table 4). This study and work by Prinz *et al.*, 2010a examined the whole body of *P. vulgata* while Firth *et al.*, 2017 largely examined gastric smears and so may have encountered more parasites as more tissue was examined. In contrast to this study, Firth *et al.*, 2017 and Prinz *et al.*, 2010a did not include metacercariae in their prevalence calculations of *E. patellae*.

No primary infections of *E. patellae* were recorded in this study, but they were observed on other shores during the time period (pers. obs.). The absence of primary infection *E. patellae* was likely due to the fact that *P. vulgata* outside of rock pools from the low shore were sampled. Previous work has recorded higher prevalence of *E. patellae* primary infections in rock pools (Crewe, 1951, Thomas, 1965, Copeland *et al.*, 1987). However, the opposite pattern has been reported by Kollien, (1996) and some shores sampled by Firth *et al.*, (2017).

	11	υ	υ		
Location	Method	Life stage	n	Prevalence	Reference
Great Britain	Dissection	Redial	3,631 3,777	4%	Crewe, 1947, 1951
Inner Farne, Northumberland, UK	Dissection	Redial	1,750	13.5%	Thomas, 1965
Cardigan Bay, Southwest UK	Dissection	Redial	503	3.77%	James 1968a
Portavogie Harbour, Northern Ireland.	Squash preparations	Redial	200	91.5%	Copeland et al., 1987
St. Andrews, east coast Scotland, UK	Macroscopic examination and dissection	Redial	1,160	32%	Kollien, 1996
	Dissection	Metacercariae	400	67%	Kollien, 1996
West Ireland Clare Island	Macroscopic examination	Redial	3,350	12%	Delany 2000
Southwest coast Ireland	Squash preparations	Redial	500	3.6+2.4%	Prinz <i>et al.,</i> 2010a
East and West coast Ireland	Visual examination & Gastric smears	Redial	1,580	28%	Firth <i>et al.,</i> 2017

Table 4 Summary of results of previous studies and methods of assessing *Echinostephilla patellae* abundance in *Patella* spp. arranged in chronological order

Prevalence and intensity of trematode taxa varied between sampling events but not in relation to the season. Temporal variation in *E. patellae* metacercariae prevalence or intensity did not exhibit a distinct peak but was higher in the winter months. No seasonal peak in prevalence of trematodes in *L. littorea* utilised as a primary or secondary host was observed. Similar patterns have been recorded in metacercariae of other trematodes (Sindermann & Farrin, 1962, Robson & Williams, 1970, Martin, 1971, Lauckner, 1980). Patterns in seasonal variation of *E. patellae* or *Renicola* spp. metacercariae could be confounded by pre-existing infections from multiple years within a single gastropod. Nikolaev *et al.*, (2006) found that metacercariae of trematodes within superfamily Echinostomatidae in *Mytilus edulis* were viable years after initial infection.

The first record of the adult worm *Proctoeces* sp. in *P. vulgata* and *N. lapillus* in Ireland was also of note. *Proctoeces* spp. have previously recorded in *Patella* spp. in the Mediterranean (Le Pommelet *et al.*, 1997) and in *N. lapillus* in New England had low prevalence but high intensities of adult worms in the kidneys (Pondick, 1983). In this study *Proctoeces* occurred at one site, Cusheen, which had a high density of *Mytilus* sp., the preferred primary host of *Proctoeces* sp. (Lang & Deunis, 1976).

Trematode taxa recorded in the study are considered relatively harmless to humans (Loos-Frank 1967, Loos-Frank 1969). To the author's knowledge, the practice of consuming uncooked marine invertebrate taxa screened in this study is rare in Ireland. Thus, food-borne infection by trematode taxa recorded is improbable but possible. Infection by *Cryptocotyle lingua*, may occur via the consumption of the its second intermediate host, bony fish, rather than its primary intermediate host *L. littorea* (Cummins *et al.*, 2002, Stunkard 1930). *Cryptocotyle lingua* infects fish-consuming-mammals and has been detected in coastal otters *Lutra lutra* in Ireland (McCarthy *et al.*, 1993) and humans in Greenland and Alaska (Babbott *et al.*, 1961, Rausch *et al.*, 1967). Infection was not considered to be harmful and can be avoided with sufficient cooking or freezing of fish, prior to consumption (Babbott *et al.*, 1961, Rausch *et al.*, 1967, BIOHAZ 2010, Borges *et al.*, 2015).

The distribution of trematode taxa, recorded in this study, varied over time and in relation to quality. This study found site-specific effects of biotic communities on the intensity of trematode metacercariae. To provide baselines or accurate predictions of trematodes to future changes, the taxa must be studied in the field or under biologically realistic conditions.

Potential future changes in intertidal communities such as climate change and associated stressors may further impact on gastropod: trematode dynamics (Thompson *et al.*, 2002, Poulin, 2006, Poulin & Mouritsen, 2007). One such change is the rise in global ocean temperature of 0.10 °C, from the surface to a depth of 700m, since 1961 with a predicated maximum increase of 4 °C by the year 2100 is (Callaway *et al.*,

2012). Increases in temperature have already occurred in the seas around Ireland, as rises of 0.3-0.4 °C per decade have been recorded since the 1980s (Dunne et al., 2013). In an Irish context, sea temperature, mean annual temperatures of 1–1.6°C and storm severity are all predicted to increase (Nolan, 2015). The predicted changes may facilitate the transmission of larval stages of trematodes. Metanalysis by Poulin (2006) indicated that a 10°C increase could results in up to a 200-fold increase of cercarial emergence from first intermediate hosts. The emergence of cercariae vary within the same infection, and between trematode taxa and first intermediate host taxa (Poulin, 2006). However, the predications on the response of parasite taxa to predicated changes have largely focused on the effect of temperature on a single trematode taxon (Poulin 2006, Marcogliese 2016). Thus, accurate predictions of parasite taxa's response to predicated climate change are difficult to make, due to the gaps in the knowledge, the inherent complexity of the host-parasite relationship and the neglect of environmental stressors outside of predicted anthropogenic warming (Harvell et al., 1999, Ward & Lafferty 2004, Poulin, 2006, Marcogliese, 2016). In particular the relationships need to be evaluated in relation to abundant trematode such as Echinostephilla patellae, further work should consider its ecology and the effect of its metacercariae on P. vulgata.
REFERENCES

- Aguirre-Macedo ML, Vidal-Martinez VM, Lafferty KD (2011) Trematode communities in snails can indicate impact and recovery from hurricanes in a tropical coastal lagoon. *International Journal for Parasitology*, **41**, 1403-1408.
- Altman I, Byers JE (2014) Large-scale spatial variation in parasite communities influenced by anthropogenic factors. *Ecology*, **95**, 1876-1887.
- Atalah J, Crowe TP (2010) Combined effects of nutrient enrichment, sedimentation and grazer loss on rock pool assemblages. *Journal of Experimental Marine Biology and Ecology*, **388**, 51-57.
- Babbott FL, Gordon JE, Frye WW (1961) Intestinal parasites of man in Arctic Greenland. *American Journal of Tropical Medicine and Hygiene*, **10**, 185-190.
- Bettin C, Oehlmann J, Stroben E (1996) TBT-induced imposex in marine neogastropods is mediated by an increasing androgen level. *Helgolander Meeresuntersuchungen*, **50**, 299-317.
- BIOHAZ (2010) Scientific Opinion on risk assessment of parasites in fishery products. *European Food Safety Authority Journal*, **8**, 1-91.
- Blackmore DT (1969) Studies of *Patella vulgata* L. I. Growth, reproduction and zonal distribution. *Journal of Experimental Marine Biology and Ecology*, **3**, 200– 213.
- Blainville H de (1818). Mémoire sur la classe des Sétipodes, partie des Vers à sang rouge de M. Cuvier, et des Annélides de M. de Lamarck. *Bulletin des Sciences, par la Société Philomatique de Paris*, **1818**, 78-85.
- Borges JN, Skov J, QZM B, Moller L, Kania P, Santos C, Buchmann K (2015) Viability of *Cryptocotyle lingua* metacercariae from Atlantic cod (*Gadus morhua*) after exposure to freezing and heating in the temperature range from -80 °C to 100 °C. *Food Control*, **50**, 371-377.
- Bradley C, Byrne C, Craig M, Free G, Gallagher T, Kennedy B, Little R, Lucey J, Mannix A, McCreesh P, McDermott G, McGarrigle M, Longphuirt N, S. OB, S., Plant C, Tierney D, Trodd W, Webster P, Wilkes R, Wynne C (2015) Water Quality in Ireland 2010-2012. (eds Byrne C, Fanning A).
- Burrows MT, Harvey R, Robb L (2008) Wave exposure indices from digital coastlines and the prediction of rocky shore community structure. *Marine Ecology Progress Series*, **353**, 1-12.
- Bush AO, Lafferty KD, Lotz JM, Shostak AW (1997) Parasitology Meets Ecology on Its Own Terms: Margolis *et al.*, Revisited. *The Journal of Parasitology*, **83**, 575-583.

- Byers JE, Malek AJ, Quevillon LE, Altman I, Keogh CL (2015) Opposing selective pressures decouple pattern and process of parasitic infection over small spatial scale. *Oikos*, **124**, 1511-1519.
- Callaway R, Shinn AP, Grenfell SE, Bron JE, Burnell G, Cook EJ, Crumlish M, Culloty S, Davidson K, Ellis RP, Flynn KJ, Fox C, Green DM, Hays GC, Hughes AD, Johnston E, Lowe CD, Lupatsch I, Malham S, Mendzil AF, Nickell T, Pickerell T, Rowley AF, Stanley MS, Tocher DR, Turnbull JF, Webb G, Wootton E, Shields RJ (2012) Review of climate change impacts on marine aquaculture in the UK and Ireland. *Aquatic Conservation-Marine and Freshwater Ecosystems*, **22**, 389-421.
- Clabby KJ, Bradley C, Craig M, Daly D, Lucey J, McGarrigle M, O'Boyle S, Tierney D, Bowman J (2008) Water Quality in Ireland 2004-2006. Environmental Protection Agency, Johnstown Castle, Co. Wexford, Ireland.
- Coleman RA, Goss-Custard JD, Durell S, Hawkins SJ (1999) Limpet *Patella* spp. consumption by oystercatchers *Haematopus ostralegus*: a preference for solitary prey items. *Marine Ecology Progress Series*, **183**, 253-261.
- Copeland MR, Montogomery WI, Hana REB (1987) Ecology of a Digenean infection, *Cercaria patellae* in *Patella vulgata* near Portavogie Harbour, Northern Ireland. *Journal of Helminthology*, **61**, 315-328.
- Coustau C, Renaud F, Delay B, Robbins I, Mathieu M (1991) Mechanisms involved in parasitic castration: In vitro effects of the trematode *Prosorhynchus squamatus* on the gametogenesis and the nutrient storage metabolism of the marine bivalve mollusc *Mytilus edulis*. *Experimental Parasitology*, **73**, 36-43.
- Creplin FCH, (1825) Observationes de entozois. Octavo Gryphiswaldiae.
- Crewe W, (1951) The occurrence of *Cercaria patellae* Lebour (Trematoda) and its effects on the host; with notes on some other helminth parasites of british limpets. *Parasitology*, **41**, 15-22.
- Crisp DJ, & Soutward, AJ. (1958). The distribution intertidal organisms along the coasts of the English Channel. *Journal of the Marine Biological Association of the United Kingdom*, **37**, 157-208.
- Cross MA, Irwin SWB, Fitzpatrick SM (2001) Effects of heavy metal pollution on swimming and longevity in cercariae of *Cryptocotyle lingua* (Digenea : Heterophyidae). *Parasitology*, **123**, 499-507.
- Cross MA, Irwin SWB, Fitzpatrick SM (2005) Effects of host habitat quality on the viability of *Cryptocotyle lingua* (Trematoda : Digenea) cercariae. *Parasitology*, **130**, 195-201.
- Crothers JH (1985) Dog-whelks: an introduction to the biology of *Nucella lapillus*. *Field Studies*, **6**, 291-360.

- Crowe TP, Thompson RC, Bray S, Hawkins SJ (2000) Impacts of anthropogenic stress on rocky intertidal communities. *Journal of Aquatic Ecosystem Stress and Recovery*, **7**, 273-297.
- Cúenot L. (1891) Infusoires commensaux des ligies, patelles et arenicole. *Revue Biologique du Nord de la France*, **4**,81-89.
- Cummins V, Coughlan S, McClean O, Connolly N, Mercer J, Burnell G (2002) An assessment of the potential for the sustainable development of the edible periwinkle, *Littorina littorea*, industry in Ireland. Marine Institute, Ireland/
- Darwin C (1854) A Monograph on the Sub-Class Cirripedia with Figures of All the Species. The Balanidae, (or Sessile Cirripedia) *The Ray Society, London*, **1-8**, 1-684.
- Daudin, FM (1800) Receuil de mémoires et de notes sur des espèces inédites ou peu connues de Mollusques, de vers et de zoophytes. *Fuchs & Treuttel et Wurtz. Paris*, **18**, 19-50.
- Delany J (2000) Ecology of patallid limpets: aspects of the ecology and reproduction of the two patellid limpets in mid-shore tidepools on Clare Island, Co. Mayo. Doctor of Philosophy Thesis, Department of Zoology, and Animal Ecology. University College Cork, Cork, Ireland.
- Dietz E (1909) Die Echinostomiden der Vogel. Zoologischer Anzeiger, 34, 180-192.
- Dobson A, Lafferty KD, Kuris AM, Hechinger RF, Jetz W (2008) Homage to Linnaeus: How many parasites? How many hosts? *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 11482-11489.
- Dunne S, Hanafin J, Lynch P, McGrath R, Nishimura E, Nolan P, Venkata Ratnam J, Semmler T, Sweeney C, Varghese S, Wang S (2013). STRIVE Report Series No. 27. Ireland in a Warmer World –Scientific Predictions of the Irish Climate in the twenty-first century. 2001-CDM2. Environmental Protection Agency, Environmental Protection Agency, Wexford, Ireland.
- EPA (1997) Oil spillage from Irish Refining Plc, Whitegate, Co. Cork. Environmental Protection Agency, Ardcavan, Wexford, Ireland. 1- 34.
- Evans DW, Irwin SWB, Fitzpatrick SM (1997) Metacercarial encystment and in vivo cultivation of *Cercaria lebouri* Stunkard 1932 (Digenea: Notocotylidae) to adults identified as *Paramonostomum chabaudi* van Strydonck 1965. *International Journal for Parasitology*, **27**, 1299-1304.
- Fabricius JC (1775) Systema entomologiae, sistens insectorum classes, ordines, genera, species, adiectis, synonymis, locis, descriptionibus, observationibus. *Flensburg and Leipzig*, **1-32**, 1-832.

- Feare C (1970) Aspects of the ecology of an exposed shore population of dogwhelks *Nucella lapillus* (L.). *Oecologia*, **5**, 1-18.
- Feis ME, Thieltges DW, Olsen JL, de Montaudouin X, Jensen KT, Bazairi H, Culloty SC, Luttikhuizen PC (2015) The most vagile host as the main determinant of population connectivity in marine macroparasites. *Marine Ecology Progress* Series, 520, 85-99.
- Ferreira SM, Brandao A, Baeta A, Neto JM, Lillebo AI, Jensen KT, Pardal MA (2007) Effects of restoration management on the estuarine isopod *Cyathura carinata*: mediation by trematodes and habitat change. *Marine Biology*, **151**, 109-118.
- Firth L, Grant L, Crowe T, Ellis J, Wiler C, Convery C, O'Connor N (2017) Factors affecting the prevalence of the trematode parasite *Echinostephilla patellae* (Lebour, 1911) in the limpet *Patella vulgata*. *Journal of Experimental Marine Biology and Ecology*, **492**, 99-104.
- Firth LB, Crowe TP (2008) Large-scale coexistence and small-scale segregation of key species on rocky shores. *Hydrobiologia*, **614**, 233-241.
- Forskål P (1775) Descriptiones animalium, avium, amphibiorum, piscium, insectorum, vermium, quae in itenere orientali observavit. *Hafniae: Möller.* 1-34, 1-164.
- Galaktionov KV, Dobrovolskij AA (2003) The Trematode Life Cycle as a System of Adaptations. In: *The Biology and Evolution of Trematodes*. Springer, pp. 215-317.
- Galli P, Crosa G, Mariniello L, Ortis M, D'Amelio S (2001) Water quality as a determinant of the composition of fish parasite communities. *Hydrobiologia*, **452**, 173-179.
- Garcia-Vedrenne A, Quintana A, DeRogatis A, Dover C, Lopez M, Kuris A, RF H (2017) Trematodes with a reproductive division of labour: heterophyids also have a soldier caste and early infections reveal how colonies become structured. *International Journal for Parasitology*, **47**, 41-50.
- Garcia-Vedrenne A, Quintana A, DeRogatis M, Martyn K, Kuris A, Hechinger R (2016) Social Organization in Parasitic Flatworms-Four Additional Echinostomoid Trematodes Have a Soldier Caste and One Does Not. *Journal* of Parasitology, **102**, 11-20.
- Gibbs P, Pascoe P, Burt G (1988) Sex change in the female dog-whelk, *Nucella lapillus*, induced by tributyltin from antifouling paints. *Journal of the Marine Biological Association of the United Kingdom*, **68**, 715-731.

Grahame J (1969) Shedding of Penis in *Littorina littorea*. Nature, 221, 976.

- Granovitch AI, Mikhailova NA (2004) Rocky shore trematodes of the west coast of Sweden: distribution and life cycle strategies. *Acta Parasitologica*, **49**, 228-236.
- Harvell C, Kim K, Burkholder J, Colwell R, Epstein PR, Grimes D, Hofmann E, Lipp E, Osterhaus A, Overstreet RM (1999) Emerging marine diseases-climate links and anthropogenic factors. *Science*, **285**, 1505-1510.
- Hawkins SJ (1999) Limpet *Patella* spp. consumption by oyster-catchers *Haematopus* ostralegus: a preference for solitary prey items. Marine Ecology Progress Series, **183**, 253-261.
- Hechinger RF, Lafferty KD (2005) Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1059-1066.
- Hechinger RF, Wood AC, Kuris AM (2011) Social organization in a flatworm: trematode parasites form soldier and reproductive castes. *Proceedings of the Royal Society of London B: Biological Sciences*, **278**, 656-665.
- Hiscock K (1981) The rocky shore ecology of Sullom Voe. *Proceedings of the Royal Society of Edinburgh Section B*, **80**, 219-240.
- Hopper JV, Poulin R, Thieltges DW (2008) Buffering role of the intertidal anemone Anthopleura aureoradiata in cercarial transmission from snails to crabs. Journal of Experimental Marine Biology and Ecology, **367**,153-156.
- Huspeni TC, Hechinger RF, Lafferty KD (2005) Trematode parasites as estuarine indicators: Opportunities, applications, and comparisons with conventional community approaches. *Estuarine Indicators*, (ed Bortone, SA) CRC Press, 297-314.
- Intergovernmental Panel on Climate Change (IPCC) (2007) The Physical Science Basis. Contribution of Working Group 1 to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Solomon,S, Qin D, Manning M, Averyt K, Tignor M, Miller HL, (Editors). Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1-996.
- James B (1968a) The distribution and keys of species in the family Littorinidae and of their digenean parasites, in the region of Dale, Pembrokeshire. *Field Studies*, 2, 615-650.
- James B (1968b) The occurrence of larval Digenea in ten species of intertidal prosobranch molluscs in Cardigan Bay. *Journal of Natural History*, **2**, 37-41.

- Jenkins SR, Arenas F, Arrontes J, Bussell J, Castro J, Coleman RA, HawkinsRA, Kay S, Martínez B, Oliveros, Roberts MJ, Sousa S, Thompson RC,. Hartnoll RG (2001) European-scale analysis of seasonal variability in limpet grazing activity and microalgal abundance. *Marine Ecology Progress Series*, **211**, 193-203.
- Jenkins SR, Hartnoll RG (2001) Food supply, grazing activity and growth rate in the limpet *Patella vulgata* L.: a comparison between exposed and sheltered shores. *Journal of Experimental Marine Biology and Ecology*, **258**, 123-139.
- Jessopp M, McAllen R, O'Halloran J, Kelly T (2010) Nutrient and Ecosystem Dynamics in Ireland's Only Marine Nature Reserve (NEIDIN). In: *Science, Technology, Research and Innovation for the Environment (STRIVE)*. Environmental Protection Agency, Co. Wexford, Ireland, 1-57.
- Jessopp M, Mulholland O, McAllen R, Johnson M, Crowe T, Allcock A (2008) Coastline configuration as a determinant of structure in larval assemblages. *Marine Ecology Progress Series*, **352**, 67-75.
- Johnston, G (1838) Miscellanea Zoologica Ariciadae. *Magazine of Zoology and Botany*, **2**, 63-73.
- Khan RA, Thulin J (1991) Influence of pollution on Parasites of Aquatic Animals. *Advances in Parasitology*, **30**, 201-238.
- Kollien A (1996) *Cercaria patellae* Lebour, 1911 developing in *Patella vulgata* is the cercaria of *Echinostephilla patellae* (Lebour, 1911) n. comb. (Digenea, Philophthalmidae). *Systematic Parasitology*, **34**, 11-25.
- Kröyer H (1847). Karcinologiste Bidrag. Naturhistorisk Tidsskrift, Kjobenhavn, 2, 366-346.
- Kube J, Kube S, Dierschke V (2002) Spatial and Temporal variations in the Trematode Component Community of the Mudsnail *Hydrobia ventrosa* in Relation to the Occurrence of Waterfowl as Definitive Hosts. *The Journal of Parasitology*, 88, 1075-1086.
- Kützing FT (1843) Phycologia generalis oder Anatomie, Physiologie und Systemkunde der Tange. Mit 80 farbig gedruckten Tafeln, gezeichnet und gravirt vom Verfasser. **1&2**, 1-458.
- Lafferty KD (1997) Environmental parasitology: What can parasites tell us about human impacts on the environment? *Parasitology Today*, **13**, 251-255.
- Lafferty KD, Allesina S, Arim M, Briggs CJ, De Leo G, Dobson AP, Dunne JA, Johnson PTJ, Kuris AM, Marcogliese DJ, Martinez ND, Memmott J, Marquet PA, McLaughlin JP, Mordecai EA, Pascual M, Poulin R, Thieltges DW (2008) Parasites in food webs: the ultimate missing links. *Ecology Letters*, **11**, 533-546.

- Lambert WJ, Corliss E, Sha J, Smalls J (2012) Trematode Infections in *Littorina littorea* on the New Hampshire Coast. *Northeastern Naturalist*, **19**, 461-474.
- Lang WH, Deunis EA (1976) Morphology and seasonal incidence of infection of Proctoeces maculatus (Looss, 1901) Odhner, 1911 (Trematoda) in Mytilus edulis L. Ophelia, 15, 65-75.
- Lauckner G (1980) Diseases of Mollusca: Gastropoda In: *Diseases of Marine Animals*. (ed by Kinne O), John Wiley & Sons, New York. **1**, 311–424.
- Lauckner G (1983) Introduction, Bivalvia to Scaphopoda In: *Diseases of Marine Animals.* (ed by Kinne O), John Wiley & Sons **2**, 477-897.
- Le Pommelet E, Bartoli P, Silan P (1997) Biodiversite des gènes et autres helminthes intestinaux des rougets: synthese pour *Mullus surmuletus* (Linnaeus, 1758) et *M. barbatus* (L., 1758). *Annales des Sciences Naturelles*, **18**, 117-133.
- Linnaeus C (1753) Species plantarum, exhibentes plantas rite cognitas ad genera relatas cum differentiis specificis, nominibus trivialibus, synonymis selectis, locis natalibus, secundum systema sexuale digestas. *Holmiæ Impensis Laurentii Salvii*, **1753**.
- Linnaeus C (1758) Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata. *Laurentius Salvius: Holmiae*, **2**, 1-824.
- Linnaeus C (1767). Systema naturae per regna tria naturae: secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio duodecima. 1. Regnum Animale. *Holmiae, Laurentii Salvii, Laurentii Salvii*, 1&2, 1-1327.
- Loennberg E (1890) Helminthologische Beobachtungen von der Westküste Norwegens. Bihang til Kongliga Svenska Vetenskaps-akademiens Handlingar, 16, 1-47.
- Loos-Frank B (1967) Experimentelle Untersuchungen über Bau, Entwicklung und Systematik der Himasthlinae (Trematoda, Echinostomatidae) des Nordseeraumes. Zeitschrift für Parasitenkunde, **28**, 299-351.
- Loos-Frank B (1969) Studies on Gymnophallid Trematodes of North Sea. I. Alternative Life-Cycles of *Gymnophallus choledochus* Odhner 1900. *Zeitschrift Fur Parasitenkunde*, **32**, 135-156.
- Looss A (1901) Ueber einige Distomen der Labriden des Triester Hafens Zentralblatt für Bakteriologie Parasitenkunde, **29**, 398-405.

- Lynch S, Morgan E, Carlsson J, Mackenzie C, Wooton E, Rowley A, Malham S, Culloty S (2014) The health status of mussels, *Mytilus* spp., in Ireland and Wales with the molecular identification of a previously undescribed haplosporidian. *Journal of Invertebrate Pathology*, **118**, 59-65.
- Lyngbye HC (1819) Tentamen hydrophytologiae danicae continens omnia hydrophyta cryptogama Daniae, Holsatiae, Faeroae, Islandiae, Groenlandiae hucusque cognita, systematice disposita, descripta et iconibus illustrata, adjectis simul speciebus norvegicis. *Hafniae*, **32**, 1-248.
- MacKenzie K, (1999) Parasites as pollution indicators in marine ecosystems: a proposed early warning system. *Marine Pollution Bulletin* **38**, 955-959.
- Mackinnon BM & Burt MD (1983): Polymorphism of microtriches in the cysticercoid of *Ophryocotyle insignis* Lönnberg,1890 from the limpet *Patella vulgata Candian Journal of Zoology*, **61**, 1062-1070.
- Marcogliese DJ (2005) Parasites of the superorganism: Are they indicators of ecosystem health? *International Journal for Parasitology*, **35**, 705–716.
- Marcogliese DJ (2016) The Distribution and Abundance of Parasites in Aquatic Ecosystems in a Changing Climate: More than Just Temperature. *Integrative and Comparative Biology*, **56**, 611-619.
- Marcogliese DJ, Pietrock M (2011) Combined effects of parasites and contaminants on animal health: parasites do matter. *Trends in Parasitology*, **27**, 123-130.
- Marcogliese DJ (2016) The Distribution and Abundance of Parasites in Aquatic Ecosystems in a Changing Climate: More than Just Temperature. *Integrative and Comparative Biology*, **56**, 611-619.
- Marine Strategy Framework Directive, (2008) Directive 2008/56/EC of the European Parliament and the Council of 17 June 2008 Establishing a Framework for Community Action in the Field of Marine Environmental Policy (Marine Strategy Framework Directive). Official Journal of European Communities, 164, 19-40.
- Martin WE (1971) Larval stages of renicolid trematodes. *Transactions of the American Microscopical Society*, **90**, 188–194.
- McCarthy T, Hassett D (1993) *Cryptocotyle lingua* (Creplin) (Digenea: Heterophyidae) and other parasites of a coastal otter *Lutra lutra* (L.). *The Irish Naturalists' Journal*, **24**, 280-282.
- Moore HB (1937) The Biology of *Littorina littorea*. Part I. Growth of the Shell and Tissues, Spawning, Length of Life and Mortality. *Journal of the Marine Biological Association of the United Kingdom*, **21**, 721-742.

- Mouritsen KN, McKechnie S, Meenken E, Toynbee J, Poulin R (2003) Spatial heterogeneity in parasite loads in the New Zealand cockle: the importance of host condition and density. *Journal of the Marine Biological Association of the United Kingdom*, **83**, 307-310.
- Mouritsen KN, Poulin R (2002) Parasitism, community structure and biodiversity in intertidal ecosystems. *Parasitology*, **124**, 101-117.
- Nicoll W, (1907) *Parorchis acanthus*, the type of a new genus of trematodes. *Quarterly Journal of Microscopical Science*, **51**, 345-355.
- Nikolaev KE, Sukhotin AA, Galaktionov KV (2006) Infection patterns in White Sea blue mussels Mytilus edulis of different age and size with metacercariae of *Himasthla elongata* (Echinostomatidae) and *Cercaria parvicaudata* (Renicolidae). *Diseases of Aquatic Organisms*, **71**, 51-58.
- Nolan P (2015) Ensemble of regional climate model projections for Ireland. Report no. 159, Environmental Protection Agency, Environmental Protection Agency, Wexford, Ireland.
- O'Boyle R, Wilkes R, McDermott G (2010) Quality of Estuarine and Coastal Waters 2007-2009. In: McGarrigle M, Lucey J, Ó Cinnéide M (eds) Water Quality in Ireland 2007-2009. Environmental Protection Agency Ireland.
- O'Connor NE, Donohue I, Crowe TP, Emmerson MC (2011) Importance of consumers on exposed and sheltered rocky shores. *Marine Ecology Progress Series*, **443**, 65-75.
- Olivi G (1792) Zoologia Adriatica, ossia catalogo ragionato degli animali del golfo e della lagune di Venezia. *Bassano* **9**, 1-32.
- Orton JH (1919) Sex-Phenomena in the Common Limpet: (*Patella vulgata*). *Nature*, **104**, 373-374.
- Ostfeld R, Keesing F (2000) The function of biodiversity in the ecology of vectorborne zoonotic diseases. *Canadian Journal of Zoology*, **78**, 2061-2078.
- Pietrock M, Marcogliese DJ, (2003) Free-living endohelminth stages: at the mercy of environmental conditions. *Trends in Parasitology*, **19**, 293-299.
- Poli GS (1791) Testacea utriusque Siciliae eorumque historia et anatome. Parme, 1.
- Pondick JS (1983) The Geographical Distribution of an Adult Trematode, *Proctoeces maculatus*, in the Gastropod *Nucella lapillus* from New England. *Proceedings of the Helminthological Society of Washington*, **50**, 174-176.

- Poulin R (1999) The functional importance of parasites in animal communities: many roles at many levels? *International Journal for Parasitology*, **29**, 903-914.
- Poulin R (2006) Global warming and temperature-mediated increases in cercarial emergence in trematode parasites. *Parasitology*, **132**, 143-151.
- Poulin R, Blasco-Costa I, Randhawa HS (2016) Integrating parasitology and marine ecology: Seven challenges towards greater synergy. *Journal of Sea Research*, 113, 3-10.
- Poulin R, Mouritsen KN (2006) Climate change, parasitism and the structure of intertidal ecosystems. *Journal of Helminthology*, **80**, 183-191.
- Poulin R, Mouritsen KN (2007) Climate change, parasitism and the structure of intertidal ecosystems. *Journal of Helminthology*, **80**, 183-191.
- Prinz K, Kelly T, O'Riordan R, Culloty S (2010a) Occurrence of macroparasites in four common intertidal molluscs on the south coast of Ireland. *Marine Biodiversity Records*, **3**, 89-96.
- Prinz K, Kelly T, O'Riordan R, Culloty S (2011) Factors influencing cercarial emergence and settlement in the digenean trematode *Parorchis acanthus* (Philophthalmidae). *Journal of the Marine Biological Association of the United Kingdom*, **91**, 1673-1679.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2009) Non-host organisms affect transmission processes in two common trematode parasites of rocky shores. *Marine Biology*, **156**, 2303-2311.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010b) Infection of *Mytilus edulis* by the trematode *Echinostephilla patellae* (Digenea: Philophthalmidae). *Journal of Helminthology*, **84**, 193-198.
- R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <u>http://www.R-project.org/</u>
- Rausch RL, Scott EM, Rausch VR (1967) Helminths in eskimos in Western Alaska, with particular reference to *Diphyllobothrium* infection and anaemia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **61**, 351-357.
- Rees FG (1934) Cercaria patellae Lebour, 1911, and its effect on the digestive gland and gonads of Patella vulgata. Proceedings of the Zoological Society of London, 104, 45-53.
- Reiczigel J (2003) Confidence intervals for the binomial parameter: some new considerations. *Statistics in Medicine*, **22**, 611-621.

- Reiczigel J, Abonyi-Tóth Z, Singer J (2008) An exact confidence set for two binomial proportions and exact unconditional confidence intervals for the difference and ratio of proportions. *Computational Statistics and Data Analysis*, **52**, 5046-5053.
- Reiczigel, J and Rózsa, L Quantitative Parasitology. 3.0 edn 2005.
- Reid DG, Rumbak E, Thomas RH (1996) DNA, morphology and fossils: Phylogeny and evolutionary rates of the gastropod genus *Littorina*. *Philosophical Transactions of the Royal Society B*, **351**, 877-895.
- Risso A (1826) Histoire naturelle des principales productions de l'Europe Méridionale et particulièrement de celles des environs de Nice et des Alpes Maritimes, **4**, 1-439.
- Robson E, Williams I (1970) Relationships of Some Species of Digenea with the Marine Prosobranch *Littorina littorea* (L.) I. The Occurrence of Larval Digenea in *L. littorea* on the North Yorkshire Coast. *Journal of Helminthology*, 44, 153-168.
- Rozsa L, Reiczigel J, Majoros G (2000) Quantifying parasites in samples of hosts. *Journal of Parasitology*, **86**, 228-232.
- Rudolphi CA, (1809) Entozoorum sive vermium intestinalium historia naturalis. Amsterdam. 2, 1-145.
- Silva ACF, Hawkins SJ, Boaventura DM, Thompson RC (2008) Predation by small mobile aquatic predators regulates populations of the intertidal limpet *Patella vulgata* (L.). *Journal of Experimental Marine Biology and Ecology*, **367**, 259-265.
- Silva ACF, Hawkins SJ, Clarke KR, Boaventura DM, Thompson RC (2010) Preferential feeding by the crab Necora puber on differing sizes of the intertidal limpet Patella vulgata. Marine Ecology Progress Series, 416, 179-188.
- Sindermann CJ, Farrin AE (1962) Ecological studies of *Cryptocotyle lingua* (Trematoda: Heterophyidae) whose larvae cause" pigment spots" of marine fish. *Ecology*, **43**, 69-75.
- Southward AJ (1976) On the taxonomic status and distribution of *Chthamalus stellatus* (Cirripedia) in the northeast Atlantic region: with a key to the common intertidal barnacles of Britain. *Journal of the Marine Biological Association of the United Kingdom*, **56**, 1007–1028.
- Stuankard, HW (1930) The life history *Cryptocotyle lingua* (Creplin), with notes on the physiology of the metacercariae. *Journal of Morphology*, **50**, 143-191.

- Studer A, Kremer L, Nelles J, Poulin R, Thieltges DW (2013) Biotic interference in parasite transmission: Can the feeding of anemones counteract an increased risk of parasitism in amphipods at higher temperature? *Journal of Experimental Marine Biology and Ecology*, 445, 116-119.
- Stunkard H W (1930) The life history of *Cryptocotyle lingua* (Creplin), with notes on the physiology of the metacercariae. *Journal of Morphology*, **50**, 143-191.
- Stunkard H, Shaw CR (1931) The effect of dilution of sea water on the activity and longevity of certain marine cercariae, with descriptions of two new species. *The Biological Bulletin*, **61**, 242-271.
- Stunkard HW (1932) Some larval trematodes from the coast in the region of Roscoff, Finistère. *Parasitology*, **24**, 321–343.
- Sures B (2008) Host-parasite interactions in polluted environments. *Journal of Fish Biology*, **73**, 2133-2142.
- Sures B, Nachev M, Selbach C, Marcogliese DJ (2017) Parasite responses to pollution: what we know and where we go in 'Environmental Parasitology'. *Parasites & Vectors*, **10**, 1-19.
- Thieltges D, Jensen K, Poulin R, (2008) The role of biotic factors in the transmission of free-living endohelminth stages. *Parasitology*, **135**, 407-426.
- Thieltges DW (2006) Effect of infection by the metacercarial trematode *Renicola* roscovita on growth in intertidal blue mussel *Mytilus edulis*. Marine Ecology Progress Series, **319**, 129-134.
- Thieltges DW, Bordalo MD, Hernandez AC, Prinz K, Jensen KT (2008) Ambient fauna impairs parasite transmission in a marine parasite-host system. *Parasitology*, **135**, 1111-1116.
- Thieltges DW, Krakau M, Andresen H, Fottner S, Reise K (2006) Macroparasite community in molluscs of a tidal basin in the Wadden Sea. *Helgoland Marine Research*, **60**, 307-316.
- Thieltges DW, Rick J (2006) Effect of temperature on emergence, survival and infectivity of cercariae of the marine trematode *Renicola roscovita* (Digenea: Renicolidae). *Diseases of Aquatic Organisms*, **73**, 63-68.
- Thomas M (1965) Observations on the occurrence of *Cercaria patella* Lebour in *Patella vulgata* L. on the Inner Farne. *Transactions of the Natural History Society of Northumberland*, **15**, 140-146.
- Thompson RC, Crowe TP, Hawkins SJ (2002) Rocky intertidal communities: past environmental changes, present status and predictions for the next 25 years. *Environmental Conservation*, **29**, 168-191.

- Turton W (1825) Description of some new British shells. *Zoological Journal*, **2**, 361-367.
- Underwood AJ (1997) Experiments in Ecology: their logical design and interpretation using analysis of variance. *Cambridge University Press, Cambridge*, 1-504.
- van Strydonck D, (1965). Esptces du genre *Paramonostomum*. Ltihe 1909 (fam. Notocotylidae, Ltihe 1909). *Annales de Parasitologie*, **40**, 141-148.
- Vidal-Martinez VM, Pech D, Sures B, Purucker ST, Poulin R (2010) Can parasites really reveal environmental impact? *Trends in Parasitology*, **26**, 44-51.
- Ward JR, Lafferty KD (2004) The elusive baseline of marine disease: Are diseases in ocean ecosystems increasing? *PLoS Biology*, **2**, 542-547.
- Water Framework Directive (2000) 2000/60/EC of the European Parliament and of the Council establishing a framework for the Community action in the field of water policy". *Official Journal of European Communities*, **371**, 1-72.
- Welsh JE, van der Meer J, Brussaard CPD, Thielges DW (2014) Inventory of organisms interfering with transmission of a marine trematode. *Journal of the Marine Biological Association of the United Kingdom*, **10**, 1-6.
- Werding B (1969) Morphologie, Entwicklung und Ökologie digener Trematoden-Larven der Strandschnecke *Littorina littorea*. *Marine Biology*, **3**, 306-333.

CHAPTER 3: Distribution of parasites of the European Limpet (*Patella vulgata*) in a marine reserve

ABSTRACT

Marine reserves are instrumental in the implementation of successful marine conservation or commercial fisheries management. In the literature to date, some work has been done on the dynamics of pathogens and parasites within marine reserves. Marine reserves can encourage parasitic infection, as within their boundaries, biota can attain larger sizes and densities, which can favour increased parasite load and enhanced transmission. This study was conducted to examine whether parasite communities of a common grazer, the limpet Patella vulgata, varied inside and outside a marine reserve in Southwest Ireland. The roles of coastal configuration, host density, and the location of populations inside and outside of the reserve were considered in relation to distribution of parasites in P. vulgata. Parasitism was commonly observed inside and outside the reserve. Over half of P. vulgata screened were infected by one or more of five parasite taxa from four phyla: Annelida, Arthropoda, Ciliophora and Platyhelminthes. The most common parasite taxa were trematode metacercariae of Echinostephilla patellae and the ciliate Leiotrocha patellae. Large variation in parasite distribution over small spatial scales was recorded, but generally, higher occurrence and burdens of trematode metacercariae were recorded in sheltered sites and with slower water currents, indicating the significance of water movements in parasite transmission and retention. The observed similarity in parasite communities within and outside the reserve is likely to result from the wide dispersal range of final avian hosts, which extends to all sites sampled. This study indicates that other site-specific local factors such as host density, coastal configuration and host interaction have a greater influence on parasites than location within a reserve alone.

3.1 INTRODUCTION

Parasitic taxa are generally ignored in conservation biology, despite their ubiquity and potential ecological relevance (Poulin 1999, Dobson *et al.*, 2008). The neglect of parasites has been attributed to a general distaste for and the difficulty in studying cryptic taxa (Windsor 1995, Nichols *et al.*, 2011, Gómez *et al.*, 2013). Parasites are increasingly recognised to play a role in community structure and functioning (Marcogliese 2004, Poulin & Mouritsen, 2006). The effect of anthropogenic impacts, such as declining biodiversity and predicted climatic change, on marine parasites is largely unknown (Ward *et al.*, 2004, Marcogliese, 2016, Thieltges *et al.*, 2016). Despite this, there is speculation that the extent and severity of infection by marine parasites will increase in response to predicted changes in global warming, host range and abundance (Poulin 2006, Poulin & Mouritsen, 2006).

Marine protected areas (hereafter MPAs) have a remit to protect biodiversity or provide a refuge for marine biota from commercial harvesting and anthropogenic impacts. MPAs are a common management tool in conservation biology and are defined by the IUCN as "(a) clearly defined geographical space, recognised, dedicated and managed, through legal or other effective means, to achieve the long-term conservation of nature with associated ecosystem services and cultural values" (Day *et al.*, 2012). The most restricted or protected area is a reserve, within which no extractive activities are permitted (Lester and Halpern, 2008). Within a reserve's boundaries, a species' abundance or biomass can rapidly increase (Gell & Roberts, 2003, Byers, 2005, Lester & Halpern, 2008). This may benefit local fisheries if the increasingly abundant or larger sized individuals of commercially exploited species leave the boundaries of the reserve, an effect known as "spillover", and can then be harvested from nearby unprotected areas (Gell and Roberts, 2003).

Generally, the effects of marine reserves on commercial, especially on vertebrate species, are well understood (Murray et al., 1999, Gell & Roberts, 2003, Halpern et al., 2009). Some attention has been paid to non-commercial or invertebrate species (Lubchenco et al., 2003, Fenberg et al., 2012) and parasite dynamics within marine reserves (Wootton et al., 2012). Within a reserve, a host species could increase in abundance, density, and size (Gell & Roberts, 2003, Byers, 2005, Lester & Halpern, 2008). According to classic epidemiological theory, these changes may favour successful transmission of parasites or disease outbreak (Kermack & McKendrick, 1927). This has occurred in terrestrial systems (Lebarbenchon et al., 2006). The increased abundance of a parasite within the protected area can result in a parasite "jumping" to new host species, that gather in large numbers e.g. shorebirds (Lebarbenchon et al., 2006). To ensure conservation of the marine reserves' target species it is critical to understand parasite dynamics within their boundaries and whether these dynamics can inhibit its objectives. A marine reserve could act as a parasite reservoir if a parasite is permanently maintained within the reserve's boundaries and is able to transmit to nearby host populations or "spillover" into the surrounding area (Haydon et al., 2002).

At present, work on pathogens in MPAs is fragmented, with a limited number of locations (n=10) and taxa (n=25) considered (Table 1). The limited number of empirical studies on parasites in MPAs, make generalisations of parasites within reserves difficult and have provided conflicting results (see Table 1). The potential for a marine reserve to facilitate parasitic infections is likely dependent on several factors, the life history of the parasite taxa, their hosts, and the characteristics and location of the reserve itself. The location of the reserve, its water currents and degree of exposure may affect parasite communities within its boundaries. In the Northern Hemisphere at Lundy Island Marine Reserve, UK, European *Homarus gammarus* (Linnaeus, 1758) experienced a higher incidence of shell disease within protected areas (Wootton *et al.*, 2012, Davies *et al.*, 2015). While *Jasus edwardsii* (Hutton, 1875) experienced a lower incidence of tail fan necrosis in New Zealand marine reserves (Freeman & MacDiarmid, 2009). Work, in the Great Barrier Reef and Mediterranean Sea, found that protected areas had partial or no effect on coral

pathologies (Ternengo *et al.*, 2009, Lamb *et al.*, 2016). In France, monogenean communities of the White Seabream *Diplodus sargus* (Linnaeus, 1758) did not differ between protected and unprotected areas (Sasal *et al.*, 2004).

Table 1 List of studies generated by the database ISI web of Science TM for studies on parasites or pathogens within MPAs or marine reserves. Magnitude of effect of the reserve (if any) on parasite taxa an increase in infection parameters is represented as +, a decrease as – and no effect as N.

Parasite	Host	Site	Effect of reserve	Magnitude	Reference
Acanthocephalan	Bucchich's goby	Cerbère-	Individuals within protected	+	Sasal et al.,
three species of	Gobius bucchichii	Banyuls marine	zones were on average		1996
trematode,	Steindachner, 1870	reserve, in the	larger, older and had higher		
nematode.	n=396	southeast of	prevalence of parasites.		
		France			
The monogenean	White seabream		No modification in global	+	Sasal <i>et al.</i> ,
Lamellodiscus	Diplodus sargus		parasite community. Only		2004
spp., Johnston &	(Linnaeus, 1758)		one, L. elegans, increased its		
11egs, 1922	n=104		abundance in the protected		
			area relative to fished zones.		
Bacterial wasting	Purple sea urchin n=27, 456	Southern	Disease was rare at reserve	-	Behrens and
disease.	Strongylocentrotus	Californian	sites but common at fished		Lafferty,
	purpuratus (Stimpson, 1857)	rocky reef	sites.		2004
	Red sea urchin n=19, 338				
	Mesocentrotus franciscanus				
	(A. Agassiz, 1863)				
Bacteriae	Abalone Haliotis spp	NA-(Modelled	A deterministic model of	т	McCallum et
Rickettsia spn	(Linnaeus 1758)	Californian	microparasite infection	,	al_{2005}
Menensia spp.	(Enniceds, 1750)	ecosystem)	found that the yield of		un, 2005
		•••••	abalone decreased in		
			response to <i>Rickettsia</i> spp.		
			1 TT		

Table 1 Continued

Parasite	Host	Site	Effect of reserve		Reference
Monogenean and Digenean trematodes, cestodes, nematodes, acanthocenhalans	Bicolor chromis n=25 Chromis margaritifer Fowler, 1946 Blackbar devil n=50, Plectroglyphidodon dickii (Liénard, 1830)	Northern Pacific	Generally, parasite richness, intensity and prevalence were higher for all fish species in the non-fished site.	+	Lafferty <i>et</i> <i>al.</i> , 2008
and copepods.	Arc-Eye Hawkfish n=49 Paracirrhites arcatus (Cuvier, 1829) Whitecheek Tang n=26 Acanthurus nigricans (Linnaeus, 1758) Two-spot red snapper n=38 Lutjanus bohar (Forsskål, 1775)				
Handling-related bacterial infection.	Spiny lobster n=7, 466 Jasus edwardsii	New Zealand marine reserve	Lobsters outside the reserve were significantly more affected by tail fan necrosis than lobsters within the reserve.	-	Freeman and MacDiarmid, 2009
Skeletal eroding band, brown band disease and growth anomalies.	Reef building coral Acroporidae and Poritidae	Republic of Palau, located in the western Pacific Ocean	Skeletal eroding band was 2- fold higher within MPA.	+	Page <i>et al.</i> , 2009

Table 1 Continued

Parasite	Host	Site	Effect of reserve		Reference
Parasite Aspidogastrean, Digenean and monogean trematodes, cestodes, nematodes, acanthocephalans and crustaceans. Fungal shell disease.	HostCommon two-bandedseabream Diplodus vulgaris(Geoffroy Saint-Hilaire,1817) n=72Striped red mullet n=68Mullus surmuletusLinnaeus, 1758Common pandora n=47Pagellus erythrinus,(Linnaeus, 1758)Forkedbeard n=40Phycis phycis, (Linnaeus,1766)Red scorpionfish n=57Scorpaena scrofa Linnaeus,1758Peacock wrasse n=58Symphodus tinca (Linnaeus,	Site Bonifacio Strait Marine Reserve, Mediterranean Sea	Parasite distributions were not influenced by the protection status of the site. parasite life traits history remained a fundamental determinant in the community composition	N	Reference Ternengo <i>et</i> <i>al.</i> , 2009
Fungal shell disease.	1758) European lobsters n=666 <i>Homarus gammarus</i> (Linnaeus, 1758)	Lundy Island, United Kingdom	Increased host abundance, injury and shell disease detected in the no take zone of the reserve.	+	Wootton <i>et al.</i> , 2012

Table 1 Continued

Parasite	Host	Site	Effect of reserve		Reference
Digenean trematode <i>Proctoeces lintoni</i> (Siddiqi & Cable, 1960)	Keyhole limpet n=202 <i>Fissurella latimarginata</i> G. B. Sowerby I, 1835 Intertidal mussels n=2,509 <i>Perumytilus purpuratus</i> (Lamarck, 1819) Clingfish n=71 <i>Sicyases sanguineus</i> Müller & Troschel, 1843	Central Chile	Higher number of infection recorded in the MPAs in limpets and mussels	+	Loot <i>et al.</i> , 2005
P. lintoni Gill parasites Maxillopoda, Malacostraca, Pinnotherid pea crab, Monogenean and Digenean trematodes.	<i>F. latimarginata</i> n=59 Bilagay n=100 <i>Cheilodactylus variegatus</i> Valenciennes, 1833 Jerguilla n=120 <i>Aplodactylus punctatus</i> Valenciennes, 1832 Sea urchin n=149 <i>Loxechinus albus</i> (Molina, 1782) <i>F. latimarginata</i> n=123	Central Chile	Higher number of eggs recorded in the MPAs One significant effect for one parasite species; <i>Encotyllabe</i> sp. (Diesing, 1850) from the gills of <i>C. variegatus</i>	+	Loot <i>et al.</i> , 2008, Wood <i>et al.</i> , 2013
P. lintoni	<i>F. latimarginata</i> n=558		Increased parasitism within the marine reserves	+	Aldana <i>et</i> <i>al.</i> , 2013

Table 1 Continued

Parasite	Host	Site		Effect of reserve	Reference
Black band disease, skeletal eroding band, brown, band disease, white syndromes, atramentous necrosis or growth anomalies.	Reef-building corals n=36, 104	Great Reef Park	Barrier Marine	The marine reserves were only partially effective in mitigating disease prevalence	- Lamb <i>et al.</i> , 2016

Conflicting results can even be found within the same protected area (Table 1). *Perumytilus purpuratus* and *F. crassa* were more commonly infected with the trematode *P. lintoni* in Chilean MPAs (Loot *et al.*, 2005). However, further work in the areas found no difference in the parasite communities of *F. latimarginata*) and sea urchins *L. albus* between protected and unprotected zones (Wood *et al.*, 2013).

Coastal configuration is known to affect shore exposure, water currents and intertidal communities (Jessopp *et al.*, 2008, Burrows, 2012). Wave exposure is known to be a determining factor in the structure of intertidal communities on rocky shores (Burrows (Burrows *et al.*, 2008). Increasing exposure of rocky shores leads to a shift from macroalgae to filter feeders in intertidal communities (Thompson *et al.*, 2002). Changes in potential host communities will likely change respective parasite communities.

This study was undertaken to examine parasite infection of a non-commercial invertebrate species in a marine reserve. Lough Hyne is the only statutory marine reserve in the Republic of Ireland. Parasites have been previously recorded within the reserve, in the form of parasitic copepods and barnacles in decapod crabs (Renouf 1932, Costello & Myers, 1989, Minchin, 1997, De Grave & Holmes, 1998, Lyons, 2011). Trematodes have been recorded in barnacles, *Chthamalus montagui* and *C. stellatus* (O'Riordan *et al.*, 1992), decapod crabs (Lyons, 2011), bivalves, *Mytilus* spp. (Lynch *et al.*, 2014) and sea squirts *Ascidiella aspersa* (Müller, 1776) (Lynch *et al.*, 2016). However, the present study aimed to explicitly compare parasite infections within and outside the reserve.

Patella vulgata was chosen as model species to screen as a host for parasite abundance, as it frequently harbours parasite taxa in Ireland (Copeland *et al.*, 1987, Prinz *et al.*, 2010a, Firth *et al.*, 2017). Furthermore, it is a keystone grazer and an abundant gastropod on rocky shores, including those within Lough Hyne (Little *et al.*, 1988, Crowe *et al.*, 2000, Moore *et al.*, 2007). *Patella vulgata* also has high site fidelity, has

a foraging range of up to 1 metre (Wright & Hartnoll, 1981) and is in close contact with the substrate (Orton, 1929, Little *et al.*, 1988). These factors may enable the detection of variation in parasite communities over a small spatial scale. Although, *P. vulgata* is not commercially harvested in Ireland, on a global scale, many patellid limpet stocks are overexploited (Borges *et al.*, 2015). Simulated exploitation of *P. vulgata* induced changes in the life history of the gastropod. *Patella vulgata* populations at two locations in southwest England had the largest individuals systematically removed for 18 months. In areas of lower density of large individuals sex change was induced at smaller sizes. indicating that the gastropod may be susceptible to harvesting (Borges *et al.*, 2015).

Lough Hyne is sheltered, has restricted tidal exchange with outer waters and can retain and self-seed with invertebrate larvae (Johnson *et al.*, 1995, Jessopp *et al.*, 2007, Jessopp & McAllen, 2008). Larvae of the saddle oyster *Anomia ephippium* and the crustacean *Austrominius modestus* were, imported into the marine reserve and were more abudnant withithan in adjacent shores (Jessopp & McAllen, 2007; 2008). It is speculated that the unusual coastal configuration of Lough Hyne, in combination with less anthropogenic disturbance could facilitate parasitic communities of *P. vulgata* within Lough Hyne. It was hypothesised that a greater incidence and richness of parasite taxa would occur within the boundaries of the reserve.

3.2. MATERIALS AND METHODS

3.2.1 Sites

The study area encompassed ten rocky shores along the southwest coast of Ireland (see Table 2). Toe Head Bay was the most easterly site and the Western shore of Lough Hyne marine reserve was the most westerly (Fig. 1). Lough Hyne, covers an area of 0.8km² and has a limited tidal range (ca. 1m) (Ebling *et al.*, 1960). Lough Hyne has restricted tidal exchange with outer waters due to the dimensions of the connecting

channel. The channel, called the "rapids", is approximately 5m deep and 25m wide. Within the channel, an area rapidly shallows, creating a sill. During a flood tide, water must overcome the sill to enter Lough Hyne. Currents in the "rapids" reach speeds greater than 3ms^{-1} during the flood tide (Bassindale *et al.*, 1948, Bassindale *et al.*, 1957, Ebling *et al.*, 1960). It takes an estimated 41 days for water to be flushed from Lough Hyne (Johnson *et al.*, 1995). Water enters the reserve and moves northward in an anticlockwise gyre, slowing as it reaches the west side of Lough Hyne (Bassindale *et al.*, 1957). Water continues to slow as it leaves the Southern basin and moves towards the North basin (Bassindale *et al.*, 1957, Greenwood *et al.*, 2001, Bell & Barnes, 2002). The other sites, Tralispean, Tragumna and Toe Head Bay, were chosen to compare with Lough Hyne, as they are close, but not connected, to the reserve (Fig. 1). The low shore was sampled at all sites to allow comparison with the reserve as intertidal biota are concentrated within in its limited tidal range (Little *et al.*, 1988, Little, 1991).



Fig. 1 Map showing ten sites along the south coast of Ireland where 30 *Patella vulgata* (b) were sampled from in June 2015. 1 Lough Hyne North, 2 Lough Hyne East, 3 Castle Island, 4 Lough Hyne South, 5 Lough Hyne West, 6 Barloge, 7 Tranabo, 8 Tralispean, 9 Tragumna and 10 Toe Head Bay. Adapted from Environmental Protection Agency Water Quality shapefile.

Table 2.	Descriptions of study	y sites (modified from	Bassindale et al., 195	7 & Ebling <i>et al.</i> ,	1960), with Ballar	tine scale added	(Ballantine,	1961)
	1 2	/	,	U /	//		· · · · ·	

Site	GPS	Ballantine Scale	Site description
1 Northern Shore	51°30'21''N 9°18'12''W	5	Steep rock that continues into sublittoral boulders. The average effective depth of the north shore is 12m. At this shore, there are slow currents in the inflow with speeds of 0.02-2ms ⁻¹ . Water in the North Basin originates from the South Basin of Lough Hyne.
2 Eastern Shore	51°30'14.6"N 9°17'54.6" W	5	The shore comprises of steep rocks sloping with an angle of 45° and with some boulders.
3 Castle Island	51°30'06'0''N 9°18'02.5'' W	6	The shoreline and slope varies greatly on the island. The south side of the island has vertical rock and steep intertidal rock ledges. The north side of the island has moderately (c. 45°) sloping rocks which continue into boulders and mud in the subtidal.
4 Southern Shore	51°29'57''N, 9°17'53'W	7	The slope of the shoreline varies from gentle to near vertical on the South shore of Lough Hyne. In the subtidal the slope is generally gentle. The substrate is composed of gravel and boulders before giving away to mud. The shore experiences powerful currents with speeds of 6-8ms ^{-1.}
5 Western Shore	51°30'02''N 9°18'22''W	7-8	Steep rock in the intertidal, gently sloped rocks and small stones below low water in the subtidal. At points, it is heavily shaded by overhanging trees. Inflowing water has been recorded to have speeds of 4-6 ms ⁻¹ as it moves from the South Basin towards the North Basin. The Western Basin contains the deepest point (48m) within Lough Hyne.
6 Barloge	51°29'37.3"N 9°17'45.4''W	6-7	Barloge bay is small and separated by a promontory. The shoreline consists of steeply sloping rock, stones and gravel.
7 Tranabo	51°29'52.7"N 9°17'07"W		The shore is a wide south facing bay connects to Lough Hyne at high water. It takes just under a day for the bay to flush.
8 Tralispean	51°29'54.4"N 9°15'54.1"W	7	Tralispean is a small sheltered bay with slate and gravel as sediment. On the shore there is substantial freshwater runoff across the beach.
9 Tragumna	51°29'54.4"N 9°15'54.1"W	7	Adjacent to Tralispean, it is a small, sandy south/south-west facing beach. Tragumna's shoreline is gently sloping with steep irregular boulders.
10 ToeHead Bay	51°29'49.3"N 9°14'08.2"W	7	The shore is a wide south facing beach with extensive irregular rocky outcrops and boulders.

3.2 Sampling

Patella vulgata were collected from each site 29/06/2015-04/07/2015 (n=30 per site). The sampling period was chosen to coincide with known seasonal peaks of macroparasites of *P. vulgata* (Prinz *et al.*, 2010b). Three limpets were collected from each of ten 0.25m² quadrats. Quadrats were haphazardly placed at low shore. In each quadrat, the number of *P. vulgata*, the total number of all *Patella* spp. and percentage cover of macroalgal species *Ascophyllum nodosum*, *Fucus* spp. and *Ulva* spp. were recorded. Limpets found in rock pools, on vertical rock faces and measuring under <10mm in shell length were recorded but not sampled. In total, 300 *P. vulgata* were screened. Salinity was measured at each site with a handheld refractometer (Atago ATC S/Mill-E).

3.2.3 Processing

Patella vulgata were processed immediately after collection. *Patella vulgata* were processed and sexed as specified in Chapter 2, Section 2.2.3, page 43.

3.2.4 Statistical analysis

The factors were Site (fixed, orthogonal, 10 levels) and marine reserve (fixed, nested in location, 2 levels). Analysis of prevalence and median intensity was done by following the approach of recommended by Rozsa *et al.*, (2000), carried out in R studio version 1.0.136 (R Core Team 2015) and detailed further in Chapter 2 Section 5.2 page 44. Differences in prevalence between locations and gonadal stages were tested with a Pearson's chi square with Yates continuity correction. If expected values were too low (<5) for a Pearson's chi square, a Fisher's Exact test was used. Intensities were compared with a non-parametric Kruskal Wallis test if transformation failed to normalise distributions. Normality of distributions was assessed first visually and then formally by the Shapiro–Wilk test. Variation among locations in host characteristics, namely size, weight, and density, were tested with a one-way analysis of variance (ANOVA). Significant results were investigated with Tukey's Honest Significant Differences (Tukey HSD) or pairwise comparisons following an ANOVA or Kruskal Wallis test. Host weight and shell heights were log transformed, while parasite counts and host density were square root transformed. Parametric tests, Pearson correlations, ANOVAs and Post-hoc tests were carried out in R studio with the packages stats and car (R Core Team, 2017). Statistical significance was considered at P<0.05.

3.2.5 Parasite identification

Parasite identification and terminology follows that of that of previous specification in Chapter 2, Section 2.2.3, page 43 & Section 2.2.4 page 44.

3.3 RESULTS

3.3.1 Host morphometrics

The mean shell height of sampled limpets was 17.24 (±3.69, SD) mm. The largest limpets were found in Toe Head Bay and Lough Hyne East (See Table 3), while the smallest individuals were found in Barloge (F=10.92, df=9, P<0.001). The mean density of *P. vulgata* for all sites combined was 48 (±43, SD) gastropods per m² sampled quadrats. Density ranged from 18 to 96 limpets per m². Highest and lowest densities of limpets were both found in the reserve, on the south (Tukey HSD, P<0.001) and north shore, respectively (Tukey HSD, P<0.001). The majority of limpets, 229 of the 300 processed, had undifferientated gonads (Fig. 2). Gonadal

stages significantly differed between sites (χ^2 =53.10, df=18, P<0.001) with the highest proportion of undifferientated gonads found at Toe Head Bay (Fig. 2).

Table 3 *Patella vulgata* population characteristics (n=30 per site) and site salinity in June 2015 in the form of \bar{x} (±SD).

Site	Host Density per	Shell height	Shell width	Foot length	Whole Weight	Tissue Weight	Salinity
	m^2	mm	mm	mm	g	g	
1 Lough Hyne North	18 (±5)	16.84 (±3.87)	40.57 (±6.88)	22.65 (±4.60)	13.30 (±6.83)	4.76 (±2.13)	32 (±5.25)
2 Lough Hyne East	36 (±15)	19.94 (±3.20)	46.27 (±6.66)	24.93 (±2.52)	21.25 (±6.56)	8.10 (±5.16)	34 (±1.41)
3 Castle Island	38 (±16)	17.86 (±2.99)	44.55 (±6.53)	24.07 (±3.91)	16.89 (±7.96)	5.75 (±2.51)	32 (±4.97)
4 Lough Hyne South	87 (±31)	16.89 (±3.40)	44.09 (±5.00)	24.14 (±3.80)	15.65 (±6.18)	5.55 (±1.94)	35 (±0.00)
5 Lough Hyne West	31 (±47)	16.32 (±3.15)	45.50 (±5.20)	24.33 (±4.45)	16.94 (±7.34)	6.10 (±2.25)	34 (±1.08)
6 Barloge	20 (±4)	14.83 (±3.25)	39.79 (±3.62)	21.81 (±2.29)	10.74 (±3.64)	3.81 (±1.39)	35 (±0.82)
7 Tranabo	27 (±9)	17.24 (±2.68)	44.00 (±4.42)	26.56 (±3.95)	17.63 (±6.29)	9.45 (±15.65)	28 (±3.68)
8 Tralispean	65 (±15)	19.19 (±3.00)	43.46 (±4.35)	25.44 (±5.67)	15.79 (±7.00)	6.87 (±2.54)	35 (±0.00)
9 Tragumna	96 (±81)	14.68 (±3.95)	35.03 (±5.41)	21.70 (±7.63)	8.27 (±3.49)	3.24 (±3.17)	34 (±1.41)
10 Toe Head Bay	60 (±32)	18.15 (±4.09)	40.97 (±6.13)	23.51 (±4.56)	14.10 (±8.18)	5.22 (±2.92)	35 (±0.00)

Chapter 3



Fig. 2 Sex of the protandrous *Patella vulgata* sampled at each site on the southwest coast of Ireland June 2015.

3.2 Parasite taxa

Of the 300 limpets examined, 52% harboured one or more parasite taxa from four phyla: Annelida, Arthropoda, Ciliophora, or Platyhelminthes. Trematodes were the most prevalent taxa with a total prevalence of 49%. All trematode infections were in the form of metacercariae; an encysted life stage which, upon consumption by a vertebrate, metamorphoses into an adult worm. Trematodes were followed in abundance by ciliates (27%), a cestode *Ophryocotyle insignis* (3%), mites (3%), and a burrowing polychaete *Polydora ciliata* (2%). Five parasitic taxa were recorded within and outside the reserve (Fig. 3). No increase in parasite diversity or intensity of infection was observed within the marine reserve relative to shores outside (Figs 3(b) & 4 (a)) and (Table 4).

Echinostephilla patellae was found in 48% of limpets and occurred at every site sampled except the south shore of Lough Hyne. *Echinostephilla patellae* prevalence and intensity ranged from 7-93% and 1-223 metacercariae per infected limpet (Fig. 3). *Echinostephilla patellae* prevalence significantly differed between sites (χ^2 =114.82, df=9, P<0.001) but not between protected and unprotected zones (χ^2 =2.81, df =1, P =0.09) (see Fig 4). High prevalences of *E. patellae* were found in Lough Hyne North (93%) and Toe Head Bay (93%) (Table 4). Low prevalences of *E. patellae* were found in Lough Hyne West (7%). The parasite species was absent from limpets sampled from Lough Hyne South (Table 4). Highest intensities of *E. patellae* were found in Barloge with 223 metacercariae found in one *P. vulgata* (Fig 4 (a)). When grouped by the Ballainte scale, significantly higher intensities of *E. patellae* were observed in sheltered shores (χ^2 =57.74, df=2, P<0.00) (Fig 4(b)).

The remaining trematode infections (1%) were *Gymnophallus* spp., from the family Gymnophallidae. *Gymnophallus* spp. occurred only outside of the reserve in Tragumna. Prevalence of ciliates, most likely the species *Leiotrocha patellae* significantly differed between sites (χ^2 =18.01, df=9, P<0.05). Highest prevalence of *L. patellae* was found in Castle Island (43%) and lowest in North Shore (7%). The prevalence of the cestode *O. insignis* varied from 7-58% and 2-114 cysts per limpet found on Toe Head Bay and the North and South shores of Lough Hyne. The polychaete, *Polydora ciliata*, occurred within the reserve on the West shore of Lough Hyne and in Barloge (Fig 2 (b)). *Polydora ciliata* was found in the shells of 3% of limpets at both sites.

	Site	Echinostephilla patellae	Gymnophallus spp.	Ophryocotyle insignis
L	Lough Hyne North	30 (6-158)	0	58 (28-87)
2	Lough Hyne East	46 (3-213)	0	0
3	Castle Island	16 (1-192)	0	0
4	Lough Hyne South	0	0	36 (4-114)
5	Lough Hyne West	2 (1-2)	0	0
6	Barloge	18 (1-223)	0	0
7	Tranabo	4 (1-7)	0	0
8	Tralispean	16.5 (2-60)	0	0
9	Tragumna	24 (3-67)	3 (1-67)	0
10	Toe Head Bay	22 (2-130)	0	7 (2-23)

Table 4. Median intensity and range of infection of helminth parasites of *Patella vulgata* sampled at each site (n=30)





Fig. 3. (a). Map of study sites; 1 Lough Hyne North, 2 Lough Hyne East, 3 Castle Island, 4 Lough Hyne South, 5 Lough Hyne West, 6 Barloge, 7 Tranabo, 8 Tralispean, 9 Tragumna, 10 Toe Head. Pie charts show the proportion of the 30 *P. vulgata* harbouring macroparasite taxa (black) and not parasitised (white). (b) Prevalence of parasitic taxa found in *Patella vulgata* sampled. Error bars are upper 95% Clopper-Pearson confidence intervals. Map adapted from Environmental Protection Agency Water Quality shapefile.

3.3.3 Parasite and host dynamics

Echinostephilla patellae had an overall prevalence of 48% and was the dominant parasite in this study, so further analysis focused on this species. *Echinostephilla patellae* prevalence significantly differed between gonadal stages (χ^2 =53.1, df=2, P<0.001). A higher prevalence of *E. patellae* was found in *P. vulgata* with undifferientated gonads. The largest proportion of undifferentiated gonads was found on Toehead Bay (Fig.2). However, *E. patellae* intensity did not differ between gonadal stages (F=1.008, df=2, P>0.05). *Echinostephilla patellae* prevalence and intensity were negatively and significantly correlated with *Patella* spp. density (r=0.27, df=297, P<0.001) and (r=-0.22, df=297, P<0.001) repsectively. *Echinostephilla patellae* prevalence and intensity correlated with limpet size (r=0.12, df=298, P<0.05) and (r=0.13, df=298, P<0.05). *Echinostephilla patellae* prevalence did not significantly differ with other host morphometrics, namely, shell width, whole organism or wet tissue weight (Table 2).





Fig. 4 The median intensity of *Echinostephilla patellae* metacercariae per sampling site and averaged (a) over protected and unprotected zones or (b) the sampling sites' Ballantine scale values.
3.4 DISCUSSION

In the current study, parasite prevalence, intensity and richness in *P. vulgata* did not vary in relation to the level of legal protection of a site. Consequently, other factors must determine the distributions recorded. Determining factors in macroparasites recorded are likely to be local site factors such as water movement and host population's characteristics (Hechinger *et al.*, 2007, Byers *et al.*, 2008).

Coastal configuration is known to play a role in the retention of water in bays. It also affects the structure of benthic invertebrate communities and therefore is likely to affect the parasite communities recorded in this study. In the water column, cercariae may be functionally similar to benthic lecithotrophic larvae (Crewe, 1951, Kollien, 1996, Zimmer *et al.*, 2009). Cercariae recorded in this study are similarly sized (<1mm) to benthic invertebrate larvae of previous work in the area (Crewe, 1951, Kollien, 1996, Jessopp, 2007, Jessopp & McAllen, 2008). Cercariae are also soft bodied, free living, and short lived (<24 hours) (Kollien, 1996, Pietrock & Marcogliese, 2003). This stage is sensitive to a range of environmental conditions and has a suite of behaviours which increases the probability of contacting an appropriate second intermediate host (Pietrock & Marcogliese, 2003, Sukhdeo & Sukhdeo 2004, Thieltges, 2006, Zimmer *et al.*, 2009).

Evidence for the role of configuration and degree of trematode infection is illustrated through *E. patellae*. Previous work on the role of wave exposure regarding the prevalence of *E. patellae* cercarial infection is contradictory with sheltered (Thomas, 1965) and exposed shores (James 1968) in British waters. In the present study, higher prevalences of *E. patellae* tended to be recorded on shores with slower currents, gentle slopes and long water retention times namely, Barloge, Castle Island, the North and East shore in Lough Hyne and Toe Head Bay outside of the reserve. However, this pattern was not statistically significant.

The highest intensity of *E. patellae* was recorded in Barloge. This is likely due to slow moving water and low host densities. Under laboratory conditions, *Patella* spp. infected by *E. patellae*, release a pulse of cercariae, up to 500, after immersion (Crewe, 1951). At the beginning of a flood tide, when water in Barloge is practically still (Bassindale *et al.*, 1958), so a mass release of cercariae may occur in an area of low host density. Potentially, multiple cercariae could reach and infect the limited number of *Patellae* was recorded on shores with steep slopes and exposure to faster water currents. Prevalence of *E. patellae* was low on Western shore and absent on the Southern shore of Lough Hyne. The absence of *E. patellae* on the Southern shore is likely due to turbulent currents at the "rapids". In this channel, significant mortality of invertebrate larvae occurs (Bassindale *et al.*, 1948, Jessopp, 2007). The turbulence probably has a similar effect on *E. patellae* cercariae.

In sheltered shores cercariae may persist for longer in the water column. This increases the probability of cercariae finding a suitable host and decreases the probability of cercariae being dislodged during the infection process. Lough Hyne is an extremely sheltered habitat with limited tidal exchange between outer waters due to its unusual coastal configuration. Evidence for the degree of wave exposure is provided by Swedish rocky shores where increasing wave exposure coincided with low trematode prevalence in *Littorina saxatilis* (Granovitch *et al.*, 2004).

Echinostephilla patellae prevalence or intensity were not correlated with host morphometrics observed in this study. Greater prevalence of *E. patellae* metacercariae was recorded in *P. vulgata* with undifferentiated gonads. However, no significant difference in *E. patellae* intensities was recorded between gonadal stages. Thus, higher prevalence in undifferentiated gonads is likely due to the large proportion of *P. vulgata* with undifferentiated gonads in populations sampled, rather than a preference of *E. patellae*. The proportions of gonadal stages in populations agree with

histological work on a *P. vulgata* population on a moderately exposed shore. It was found that most *P. vulgata* sampled had undifferentiated gonads in June and matured in December (McCarthy *et al.*, 2008).

The composition of the parasite community and its variation over small spatial scales agreed with previous records for Ireland (Copeland *et al.*, 1987, Prinz *et al.*, 2010a) and the UK (Crewe, 1951, Thomas, 1965). Parasite communities in *P. vulgata*, like other molluscan hosts, were dominated by trematodes, while other metazoans, such as the cestode *O. insignis*, were less common (Sousa, 1994, Poulin, 2001, Prinz *et al.*, 2010a). Moreover, species richness and occurrence were similar to *P. vulgata* parasite communities recorded a year previously (Chapter 2) for the Western shore and Tralispean.

There was a notable absence of trematode life stages preceding metacercariae, the cercariae, and the taxon "Cercaria B" (Crewe, 1951). These absences may be due to sampling the low shore at sites, which was chosen in order for sites outside the Lough to be comparative due to the restricted tidal range in Lough Hyne. High levels of cercarial infections have been recorded at the mid tide level of rocky shores in previous studies (Thomas, 1965, Prinz *et al.*, 2010b) due to the foraging preference of parasites' avian hosts. For *E. patellae* and for the cestode *O. insignis* the oystercatcher, *Haematopus ostralegus* is a recorded final host, (Burt, 1962, Kollien, 1996), and for *Gymnophallus* spp. gulls, *Larus* spp., are considered likely final hosts (Crewe, 1951). While the mobility and behaviour of avian hosts play a major role in the distribution of the species (Hechinger *et al.*, 2005, Feis *et al.*, 2015), they were not surveyed during the study. Local site factors favouring gull final hosts have been found to the major determinant of trematode prevalence in *Littorina* spp. in New England (Hechinger *et al.*, 2005, Feis *et al.*, 2015). The small spatial scale (<10km) of the present study means that all sites sampled were within the dispersal range of final avian hosts. However,

only low numbers of *H. ostralegus* have been observed in the reserve (Kelly, T pers. comm., 2015).

This study found that *P. vulgata* populations within the reserve do not function as a parasite reservoir per the definition provided by Haydon et al., (2002). No effect of the reserve was detected in relation to parasite or host populations. Turbulent currents at the "rapids" makes it likely that cercariae from inside the reserve will not transfer or "spillover" into P. vulgata populations outside of the reserve. Hosts within a reserve do not automatically become resistant to infection or act as a reservoir for their parasites. Caution is urged when drawing the conclusion that a marine reserve could not act as a reservoir or amplify parasite dynamics. Lough Hyne, although well studied, may not be representative of other marine reserves due to its sheltered nature and limited tidal exchange with outer waters. In particular, the limited tidal range may have negative implications for the available space and time spent by potential avian hosts on the shores foraging. It was designated because of its biological diversity and varied habitats contained in a small area (Bell & Barnes., 2002). However, recent MPA designations have specific criteria e.g. the restoration of commercial species or habitat. Proposed networks of MPAs will have a high degree of connectivity due to increased water movement and strategic locations. These MPAs may have greater potential to facilitate parasite transmission and act as reservoirs (Fenberg et al., 2012).

Multiple factors drive disease or parasite development. The ecological consequences of recorded taxa merit further work due to the ecological relevance of *P. vulgata* and the ubiquity of parasite taxa. In general, this is also true of parasite taxa of other invertebrate species. It is necessary to consider non-commercial species or neglected parasite taxa in MPAs. This is in order to avoid a panglossian view of MPAs and to ensure their conservation objectives are achieved.

REFERENCES

- Agassiz A. (1863). List of the echinoderms sent to different institutions in exchange for other specimens, with annotations. *Bulletin of the Museum of Comparative Zoölogy at Harvard College*, **1**, 17-28.
- Ballantine W (1961) A Biologically-defined Exposure Scale for the Comparative Description of Rocky Scores. *Field Studies Journal*, **1**, 1-19.
- Bassindale R, Davenport E, Ebling FJ, Kitching JA, Sleigh MA, Sloane JF (1957) The ecology of the Lough Ine rapids with special reference to water currents: VI. Effects of the rapids on the hydrography of the South Basin. *Journal of Ecology*, 45, 879-900.
- Bassindale R, Ebling FJ, Kitching JA, Purchon RD (1948) The ecology of the Lough Ine Rapids with special reference to water currents. *Journal of Ecology*, **36**, 305-322.
- Bell J, Barnes D (2002) The Relationship between Sedimentation, Flow Rates, Depth and Time at Lough Hyne Marine Nature Reserve. *The Irish Naturalists' Journal*, 27, 106-116.
- Blackmore DT (1969) Studies of *Patella vulgata* L. I. Growth, reproduction and zonal distribution. *Journal of Experimental Marine Biology and Ecology*, **3**, 200-213.
- Borges CDG, Hawkins SJ, Doncaster CP, Crowe TP (2015) Effects of simulated human exploitation of a key grazer, *Patella vulgata*, on rocky shore assemblages. *Marine Ecology Progress Series*, **533**, 163-176.
- Burrows MT (2012) Influences of wave fetch, tidal flow and ocean colour on subtidal rocky communities. *Marine Ecology Progress Series*, **445**, 193-207.
- Burrows MT, Harvey R, Robb L (2008) Wave exposure indices from digital coastlines and the prediction of rocky shore community structure. *Marine Ecology Progress Series*, **353**, 1-12.
- Burt MDB (1962) A contribution to the knowledge of the cestode genus *Ophryocotyle* Friis, 1870. *Journal of the Linnean Society London Zoology*, **44**, 645-668.
- Byers JE (2005) Marine reserves enhance abundance but not competitive impacts of a harvested nonindigenous species. *Ecology*, **86**, 487-500.

- Byers JE, Blakeslee AMH, Linder E, Cooper AB, Maguire TJ (2008) Controls of spatial variation in the prevalence of trematode parasites infecting a marine snail. *Ecology*, **89**, 439-451.
- Copeland MR, Montogomery WI, Hana REB (1987) Ecology of a Digenean infection, *Cercaria patellae* in *Patella vulgata* near Portavogie Harbour, Northern Ireland. *Journal of Helminthology*, **61**, 315-328.
- Costello MJ, Myers AA (1989) Observations on the parasitism of *Aora gracilis* (Bate) (Amphipoda) by *Sphaeronella leuckartii* Salensky (Copepoda), with a review of amphipod-*Sphaeronella* associations. *Journal of Natural History* **23**, 81-91.
- Crewe W (1947) Helminth Parasites of Limpets. Nature, 159, 238-238.
- Crewe W (1951) The occurrence of *Cercaria patellae* Lebour (Trematoda) and its effects on the host; with notes on some other helminth parasites of british limpets. *Parasitology*, **41**, 15-22.
- Crowe TP, Thompson RC, Bray S, Hawkins SJ (2000) Impacts of anthropogenic stress on rocky intertidal communities. *Journal of Aquatic Ecosystem Stress and Recovery*, **7**, 273-297.
- Cuvier G, Valenciennes A (1829) Histoire naturelle des poissons. Tome troisième. Suite du Livre troisième. Des percoïdes à dorsale unique à sept rayons branchiaux et à dents en velours ou en cardes. F. G. Levrault, Paris. **3**, 1-500.
- Cuvier G, Valenciennes A (1833) Histoire naturelle des poissons. Tome neuvième. Suite du livre neuvième. Des Scombéroïdes. **9**,1-512.
- Davies CE, Johnson AF, Wootton EC, Greenwood SJ, Clark KF, Vogan CL, Rowley AF (2015) Effects of population density and body size on disease ecology of the European lobster in a temperate marine conservation zone. *ICES Journal of Marine Science Journal du Conseil*, **75**, 128-138.
- Day J, Dudley N, Holmes G, Laffoley D, Stolton S, Wells S (2012) Guidelines for applying the IUCN Protected Area Management Categories to Marine Protected Areas., Gland, Switzerland, IUCN (International Union for Conservation of Nature).
- De Grave S, Holmes JMC (1998) The distribution of marine Isopoda (Crustacea) in Lough Hyne. Biology and Environment *Proceedings of the Royal Irish Academy* **98B**, 23-30.

- de Montaudouin X, Wegeberg AM, Jensen KT, Sauriau PG (1998) Infection characteristics of *Himasthla elongata* cercariae in cockles as a function of water current. *Diseases of Aquatic Organisms*, **34**, 63-70.
- Diesing CM (1850) Encotyllabe. Systema Helminthum, 1, 290-427.
- Dobson A, Lafferty KD, Kuris AM, Hechinger RF, Jetz W (2008) Homage to Linnaeus: How many parasites? How many hosts? *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 11482-11489.
- Ebling FJ, Sleigh MA, Sloane JF, Kitching JA (1960) The Ecology of Lough Ine: VII. Distribution of Some Common Plants and Animals of the Littoral and Shallow Sublittoral Regions. *Journal of Ecology*, **48**, 29-53.
- Feis ME, Thieltges DW, Olsen JL, de Montaudouin X, Jensen KT, Bazairi H, Culloty SC, Luttikhuizen PC (2015) The most vagile host as the main determinant of population connectivity in marine macroparasites. *Marine Ecology Progress* Series, 520, 85-99.
- Fenberg PB, Caselle JE, Claudet J, Clemence M, Gaines SD, Antonio Garcia-Charton J, Goncalves EJ, Grorud-Colvert K, Guidetti P, Jenkins SR, Jones PJS, Lester SE, McAllen R, Moland E, Planes S, Sorensen TK (2012) The science of European marine reserves: Status, efficacy, and future needs. *Marine Policy*, 36,1012-1021.
- Fingerut JT, Zimmer CA, Zimmer RK (2003) Patterns and processes of larval emergence in an estuarine parasite system. *Biological Bulletin*, **205**, 110-120.
- Firth L, Grant L, Crowe T, Ellis J, Wiler C, Convery C, O'Connor N (2017) Factors affecting the prevalence of the trematode parasite *Echinostephilla patellae* (Lebour, 1911) in the limpet *Patella vulgata*. *Journal of Experimental Marine Biology and Ecology*, **492**, 99-104.
- FJ Ebling, Kitching JA, Muntz L, Taylor CM (1964) The Ecology of Lough Ine: XIII. Experimental Observation of the destruction of *Mytilus* edulis and *Nucella lapillus* by crabs. Journal *of Animal Ecology*, **33**, 73-82.
- Forsskål, PS (1775) Descriptiones animalium avium, amphibiorum, piscium, insectorum, vermium; quae in itinere orientali observavit. *Post mortem auctoris edidit Carsten Niebuhr Hauniae*, **1-20**, 1-34.
- Fowler HW (1946) A Collection of Fishes Obtained in the Riu Kiu Islands by Captain Ernest R. Tinkham A.U.S. *Proceedings of the Academy of Natural Sciences of Philadelphia*, **98**, 123-218.

- Fredensborg BL, Mouritsen KN, Poulin R (2004) Intensity-dependent mortality of *Paracalliope novizealandiae* (Amphipoda, Crustacea) infected by a trematode: experimental infections and field observations. *Journal of Experimental Marine Biology and Ecology*, **311**, 253-265.
- Freeman DJ, MacDiarmid AB (2009) Healthier lobsters in a marine reserve: effects of fishing on disease incidence in the spiny lobster, *Jasus edwardsii*. *Marine and Freshwater Research*, **60**, 140-145.
- Gell FR, Roberts CM (2003) Benefits beyond boundaries: the fishery effects of marine reserves. *Trends in Ecology & Evolution*, **18**, 448-455.
- Geoffroy Saint-Hilaire E (1817) Poissons du Nil, de la mer Rouge et de la Méditerranée. In: Description de l'Egypte ou recueil des observations et des recherches qui ont été faites en Égypte pendant l'expedition de l'Armée français, publié par les ordres de sa Majesté-L'Empereur Napoléon le Grand. (Imprimerie Impériale). Paris. *Histoire Naturelle*. **1**, 18-27.
- Gómez A, Nichols E (2013) Neglected wild life: Parasitic biodiversity as a conservation target. *International Journal for Parasitology: Parasites and Wildlife*, **2**, 222–227.
- Granovitch AI, Mikhailova NA (2004) Rocky shore trematodes of the west coast of Sweden: distribution and life cycle strategies. *Acta Parasitologica*, **49**, 228-236.
- Greenwood A, O'Riordan RM, Barnes DKA (2001) Seasonality and vertical zonation of zooplankton in a semi-enclosed sea lough. *Journal of the Marine Biological Association of the United Kingdom*, **81**, 213-220.
- Halpern BS, Lester SE, Kellner JB (2009) Spillover from marine reserves and the replenishment of fished stocks. *Environmental Conservation*, **36**, 268-276.
- Haydon DT, Cleaveland S, Taylor LH, Laurenson MK (2002) Identifying reservoirs of infection: A conceptual and practical challenge. *Emerging Infectious Diseases*, **8**, 1468-1473.
- Hechinger RF, Lafferty KD (2005) Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal Society B*, **272**, 1059-1066.
- Hechinger RF, Lafferty KD, Huspeni TC, Brooks AJ, Kuris AM (2007) Can parasites be indicators of free-living diversity? Relationships between species richness and the abundance of larval trematodes and of local benthos and fishes. *Oecologia*, **151**, 82-92.

- Hutton F (1875) Descriptions of Two New Species of Crustacea from New Zealand. *Transactions and Proceedings of the Royal Society of New Zealand*, **7**, 279–80.
- James B (1968) The occurrence of larval Digenea in ten species of intertidal prosobranch molluscs in Cardigan Bay. *Journal of Natural History*, **2**, 37-41.
- Jessopp M (2007) The quick and the dead: larval mortality due to turbulent tidal transport. *Journal of the Marine Biological Association of the United Kingdom*, **87**, 675-680.
- Jessopp M, McAllen R (2008) Go with the flow: tidal import and export of larvae from semi-enclosed bays. *Hydrobiologia*, **606**, 81-92.
- Jessopp M, McAllen R, O'Halloran J, Kelly T (2010) Nutrient and Ecosystem Dynamics in Ireland's Only Marine Nature Reserve (NEIDIN). In: *Science, Technology, Research and Innovation for the Environment (STRIVE).* Environmental Protection Agency, Co. Wexford, Ireland, 1-57.
- Jessopp M, Mulholland O, McAllen R, Johnson M, Crowe T, Allcock A (2008) Coastline configuration as a determinant of structure in larval assemblages. *Marine Ecology Progress Series*, **352**, 67-75.
- Johnson M, Costello MJ, Odonnell D (1995) The nutrient economy of a marine inlet: Lough Hyne, South West Ireland. *Ophelia*, **41**, 137-151.
- Kermack W, McKendrick A (1927) A contribution to the mathematical theory of epidemics. I. *Proceedings of the Royal Society of London A*, **115**, 138-141.
- Kollien A (1996) *Cercaria patellae* Lebour, 1911 developing in *Patella vulgata* is the cercaria of *Echinostephilla patellae* (Lebour, 1911) n. comb. (Digenea, Philophthalmidae). *Systematic Parasitology*, **34**, 11-25.
- Lamarck JBM (1819) Histoire naturelle des animaux sans vertèbres. Tome sixième, 1re partie. Paris, Verdière, 1815-22.
- Lamb J, Wenger AS, Devlin MJ, Ceccarelli DM, Williamson DH, Willis BL (2016) Reserves as tools for alleviating impacts of marine disease. *Philosophical Transactions of the Royal Society* B. **371**, 1-11.
- Lauckner G (1980) Diseases of Mollusca: Gastropoda. John Wiley & Sons, New York 1.
- Lebarbenchon C, Poulin R, Gauthier-Clerc M, Thomas F (2006) Parasitological consequences of overcrowding in protected areas. *Ecohealth*, **3**, 303-307.

- Lefevre T, Lebarbenchon C, Gauthier-Clerc M, Misse D, Poulin R, Thomas F (2009) The ecological significance of manipulative parasites. *Trends in Ecology & Evolution*, **24**, 41-48.
- Lester SE, Halpern BS (2008) Biological responses in marine no-take reserves versus partially protected areas. *Marine Ecology Progress Series*, **367**, 49-56.
- Liénard F (1839) Poissons. In: L. Bouton (ed) (Dixième) Rapport Annuel sur les travaux de la Société d'Histoire Naturelle de l'île Maurice. *Rapport Annuel sur les travaux de la Société d'Histoire Naturelle de l'Ile Maurice*, **10**, 31-47.
- Linnaeus C (1758) Tomus I. Systema naturae per regna tria naturae :secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis10. Holmiae, *Laurentii Salvii*, **1-4**, 1-824.
- Linnaeus C (1766) Systema naturae sive regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. *Laurentii* Salvii, Holmiae, 1, 1-532.
- Linnaeus C (1767) Systema naturae per regna tria naturae: secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. *Laurentii* Salvii, **1-2**, 1-1327.
- Little C (1991) Ecology of the rocky intertidal zone. The ecology of Lough Hyne Proceedings of a conference 4-5 September, **1990**, 53-60.
- Little C Williams, Morritt D, Perrins J, Stirling P (1988) Foraging behaviour of *Patella* vulgata L. in an Irish sea lough. Journal of Experimental Marine Biology and Ecology, **120**, 1-21.
- Loot G, Aldana M, Navarrete SA (2005) Effects of human exclusion on parasitism in intertidal food webs of central Chile. *Conservation Biology*, **19**, 203-212.
- Lubchenco J, Palumbi SR, Gaines SD, Andelman S (2003) Plugging a hole in the ocean: The emerging science of marine reserves. *Ecological Applications*, **13**, 3-7.
- Lynch S, Darmody G, O'Dwyer K, Gallagher M, Nolan S, McAllen R, Culloty S (2016) Biology of the invasive ascidian *Ascidiella aspersa* in its native habitat: Reproductive patterns and parasite load. *Estuarine, Coastal and Shelf Science*, 181, 249-255.

- Lynch S, Morgan E, Carlsson J, Mackenzie C, Wooton E, Rowley A, Malham S, Culloty S (2014) The health status of mussels, *Mytilus* spp., in Ireland and Wales with the molecular identification of a previously undescribed haplosporidian. *Journal of Invertebrate Pathology*, **118**, 59-65.
- Lyons L (2011) Aspects of the biology of decapods in South West Ireland. Doctor of Philosophy Thesis, University College Cork, Cork, Ireland.
- Marcogliese DJ (2004) Parasites: Small Players with Crucial Roles in the Ecological Theater. *EcoHealth*, **1**, 151-164.
- Marcogliese DJ (2016) The Distribution and Abundance of Parasites in Aquatic Ecosystems in a Changing Climate: More than Just Temperature. *Integrative and Comparative Biology*, **56**, 611-619.
- McCarthy M, Woosnam P, Culloty SC (2008) Histological investigation of the reproductive cycles of the limpets *Patella vulgata* and *Patella ulyssiponensis*. *Marine Biology*, **153**, 871-877.
- Minchin D (1997) The influence of the parasitic cirripede Sacculina carcini on its brachyuran host Carcinus maenas within its home range. Centre for Research on Introduced Marine Pests Technical Report, **11**, 76-80.
- Molina GI (1782). Saggito sulla storia naturale del Chili. 4, Animali del Chili. Bolonga, 367, 175-308.
- Moore P, Thompson R, Hawkins S (2007) Effects of grazer identity on the probability of escapes by a canopy-forming macroalga. *Journal of Experimental Marine Biology and Ecology*, **344**, 170–180.
- Müller J,Troschel FH (1843) Beiträge zur Kenntniss der natürlichen Familien der Fische. Archiv für Naturgeschichte, **9**, 292-330.
- Müller OF, (1776) Zoologiae Danicae Prodromus seu Animalium Daniae et Norvegiae indigenarum characteres, nomina, et synonyma imprimis popularium. *Hafniae*, *Typiis Hallageriis*, **32**, 1-217.
- Murray SN, Ambrose RF, Bohnsack JA, Botsford LW, Carr MH, Davis GE, Dayton PK, Gotshall D, Gunderson DR, Hixon MA (1999) No-take reserve networks: sustaining fishery populations and marine ecosystems. *Fisheries* 24, 11-25.
- Nichols E, Gomez A (2011) Conservation education needs more parasites. *Biological Conservation.* **144**, 937-941.

- O'Riordan RM, Myers AA, Cross TF (1992) Brooding in the intertidal barnacles *Chthamalus stellatus* (Poli) and *Chthamalus montagui* Southward in southwestern Ireland. *Journal of Experimental Marine Biology and Ecology*, **164**, 135-145.
- Orton JH (1929) Observation on *Patella vulgata*. Part III. Habitat and habits. *Journal* of the Marine Biological Association of the United Kingdom, **16**, 277-288.
- Pietrock M, Marcogliese DJ (2003) Free-living endohelminth stages: at the mercy of environmental conditions. *Trends in Parasitology*, **19**, 293-299.
- Poulin R (1999) The functional importance of parasites in animal communities: many roles at many levels? *International Journal for Parasitology*, **29**, 903-914.
- Poulin R (2001) Interactions between species and the structure of helminth communities. *Parasitology*, **122**, 3-11.
- Poulin R (2006) Global warming and temperature-mediated increases in cercarial emergence in trematode parasites. *Parasitology*, **132**, 143-151.
- Poulin R, Mouritsen K (2006) Climate change, parasitism and the structure of intertidal ecosystems. *Journal of Helminthology*, **80**, 183-191.
- Prinz K, Kelly TC, O'Riordan RM, Culloty S (2010b) Temporal variation in prevalence and cercarial development of *Echinostephilla patellae* (Digenea, Philophthalmidae) in the intertidal gastropod *Patella vulgata*. Acta Parasitologica, 55, 39-44.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2009) Non-host organisms affect transmission processes in two common trematode parasites of rocky shores. *Marine Biology* 156, 2303-2311.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010a) Occurrence of macroparasites in four common intertidal molluscs on the south coast of Ireland. *Marine Biodiversity Records*, 3, 89-96.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010c) Infection of *Mytilus edulis* by the trematode *Echinostephilla patellae* (Digenea: fPhilophthalmidae). *Journal of Helminthology*, **84**, 193-198.
- R Core Team 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <u>http://www.R-project.org/.</u>

- Rees PG (1934) *Cercaria patellae* Lebour, 1911, and its Effect on the Digestive Gland and Gonads of *Patella vulgata*. *Proceedings of the Zoological Society of London*, **104**, 45-53.
- Renouf LPW (1932) Xantho incisus Leach and Cancer pagurus L., infested with Sacculina carcini Thompson, on the south coast of Co. Cork. Annals & Magazine of Natural History, **10**, 132-133.
- Rózsa L, Reiczigel J, Majoros G (2000) Quantifying parasites in samples of hosts. *Journal of Parasitology*, **86**, 228-232.
- Sasal P, Desdevises Y, Durieux E, Lenfant P, Romans P (2004) Parasites in marine protected areas: success and specificity of monogeneans. *Journal of Fish Biology*, 64, 370-379.
- Sousa W (1994) Patterns and processes in communities of helminth parasites. *Trends* in Ecology & Evolution, 9, 52-57.
- Sowerby, GB (1835) A Catalogue of the recent species of *Fissurella*. The conchological illustration, 1-8.
- Steindachner F (1870) Ichthyologische Notizen (X) (Schluss) Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften Mathematisch Naturwissenschaftliche *Classe*, **5**, 623-642.
- Stimpson W (1857) On the Crustacea and Echinodermata of the Pacific shores of North America. *Boston Journal of Natural History*, **6**, 444-532
- Sukhdeo MVK, Sukhdeo SC (2004) Trematode behaviours and the perceptual worlds of parasites. *Canadian Journal of Zoology*, **82**, 292-315.
- Ternengo S, Levron C, Mouillot D, Marchand B (2009) Site influence in parasite distribution from fishes of the Bonifacio Strait Marine Reserve (Corsica Island, Mediterranean Sea). *Parasitology Research*, **104**, 1279-1287.
- Thieltges DW (2006) Effect of infection by the metacercarial trematode *Renicola* roscovita on growth in intertidal blue mussel *Mytilus edulis*. Marine Ecology Progress Series, **319**, 129-134.
- Thieltges DW, Wegner KM (2016) Ecology and evolution of marine parasites and diseases: International NIOZ–AWI symposium, 10-14 March 2014, The Netherlands. *Journal of Sea Research*, **113**, 1-2.

- Thomas M (1965) Observations on the occurrence of *Cercaria patella* Lebour in *Patella vulgata* L. on the Inner Farne. *Transactions of the Natural History Society of Northumberland*, **15**, 140-146.
- Thompson RC, Crowe TP, Hawkins SJ (2002) Rocky intertidal communities: past environmental changes, present status and predictions for the next 25 years. *Environmental Conservation*, **29**, 168-191.
- Ward JR, Lafferty KD (2004) The elusive baseline of marine disease: Are diseases in ocean ecosystems increasing? *PLoS Biology*, **2**, 542-547.
- Windsor D (1995) Equal Rights for Parasites. *Conservation Biology*, 9, 1-2.
- Wood CL, Micheli F, Fernandez M, Gelcich S, Castilla JC, Carvajal J (2013) Marine protected areas facilitate parasite populations among four fished host species of central Chile. *Journal of Animal Ecology*, **82**, 1276-1287.
- Wootton EC, Woolmer AP, Vogan CL, Pope EC, Hamilton KM, Rowley AF (2012) Increased Disease Calls for a Cost-Benefits Review of Marine Reserves. *PLoS One*, **7**, 1-8.
- Wright J, Hartnoll R (1981) An energy budget for a population of the limpet Patella vulgata. Journal of the Marine Biological Association of the United Kingdom, 61, 627-646.
- Zimmer RK, Fingerut JT, Zimmer CA (2009) Dispersal pathways, seed rains, and the dynamics of larval behavior. *Ecology*, **90**, 1933-1947.

CHAPTER 4: A case study on macroparasite infections in the intertidal invertebrates, the green shore crab (*Carcinus maenas*) and ragworm (*Hediste diversicolor*) in nutrient enriched estuaries.

ABSTRACT

Eutrophication is a widespread stressor in the marine environment. Eutrophication may encourage parasite taxa by increasing the density of gastropod hosts and enhancing the production of free living infectious stages via increased host survival. This study examined the occurrence of macroparasites in the intertidal invertebrates, Carcinus maenas and Hediste diversicolor, from three estuaries of differing nutrient classifications. Carcinus maenas and H. diversicolor are abundant in estuaries, have differing ecological roles, wide geographic distributions, and previously have been used in biomonitoring programmes. Both invertebrate taxa may be valuable host: parasite models to evaluate the response of parasite communities to eutrophic conditions. Nutrient classifications were provided by the Water Framework Directive and characterised by the parameters, ammonia, ortho-P, total oxidised nitrogen, total phosphorus, total nitrogen, salinity and temperature. To account for potential vertebrate host abundance, bird communities were surveyed once a month, at fixed points within the three selected estuaries. From the 1,002 invertebrates screened, parasitic taxa from the classes Trematoda, Cestoda, Copepoda, Nematoda and Acari were recorded. Digenean trematodes were the most common parasite infection of C. maenas and H. diversicolor. Carcinus maenas harboured few taxa but was heavily infected by the trematode metacercariae of Microphallus spp. This study found a relationship between eutrophication levels and trematode distribution in estuaries. Higher abundances of potential avian hosts and digenean trematodes were recorded in the enriched estuary. Given the predicted increases in eutrophication, and the global

distribution of invertebrate hosts screened, this study has broad ecological significance.

4.1 INTRODUCTION

Estuaries are some of the most productive, diverse, and economically important ecosystems on earth (Hobbie, 2000). Approximately 75% of the world's population are dependent on estuarine and coastal watersheds (Vitousek *et al.*, 1997, Small & Nicholls, 2003). In Ireland, over half of the population resides within 10km of the coastline (Stapleton *et al.*, 2000). Globally, high levels of anthropogenic impacts such as urbanisation, industrialisation, commercial fishing, and intensification of agriculture, have resulted in estuaries becoming some of the most impacted ecosystems (Jackson *et al.*, 2001, Lotze *et al.*, 2006, Airoldi & Beck, 2007). Nitrogen from terrestrial sources, agricultural runoff and sewage discharge, are all considered causal factors for eutrophication in marine ecosystems (Howarth & Marino, 2006). Eutrophication is a chronic widespread stress on marine ecosystems on a global scale (Smith, 2003) and a national scale as over half of coastal waters and estuaries in Ireland are affected by the stress (O'Boyle *et al.*, 2010, Hartnett *et al.*, 2011). The extent and severity of eutrophication will likely increase in response to growing human populations (Smith, 2003, Lotze *et al.*, 2006).

Eutrophication is broadly speculated to encourage parasitic infection directly, by providing nutrients to opportunistic parasite taxa to assimilate and indirectly by changing ecological communities' composition (Lafferty, 1997, McKenzie & Townsend, 2007, Budria, 2017). The facilitation maybe be threshold dependant as low to moderate eutrophication may favour parasite taxa able to infect multiple hosts and those with complex lifecycles that involve benthic invertebrates or fish (Esch, 1971, Galli *et al.*, 2001, Zander & Reimer, 2002, Johnson & Carpenter, 2008). The survival of infected hosts may increase with the availability of resources and the carrying capacity of potential Low to moderate eutrophication can affect hosts (Lafferty & Holt, 2003, Smith & Schindler, 2009). However, severe eutrophication may inhibit parasite infection due to the resulting hypoxia, mass invertebrate

mortalities, blooms of harmful algae and macroalgae (Hull, `1987, Raffaelli, 1999, Zander *et al.*, 2002, Howarth & Marino, 2006). Macroalgal blooms could impair the survival and movement of free living parasite life stages and oxygen deficiency could decrease host abundance and survival (Dušek *et al.*, 1998, Zander & Reimer, 2002, Pietrock & Marcogliese, 2003).

Parasitic trematode life cycles contain free living infective stages which are sensitive to a range of environmental conditions (Pietrock & Marcogliese, 2003). Eutrophic conditions in freshwater ecosystems encouraged trematode infection by increasing the abundance of intermediate hosts and the number of cercariae produced by primary intermediate hosts (Johnson et al., 2007, Johnson & Carpenter, 2008). In the intertidal zone of estuaries and coastal shores, trematodes dominate invertebrate macroparasite communities (Dobson et al., 2008, Kuris et al., 2008, Fermer et al., 2011). In estuarine ecosystems, trematodes collectively, can reach a substantial biomass. In three Californian estuaries, trematode biomass was equivalent to that of birds, fishes, burrowing shrimps, and polychaetes (Kuris et al., 2008). Consequently, evaluating the effect of eutrophication on parasites, such as trematodes, in ecosystems where they This study, aimed to investigate the potential of are abundant, is crucial. eutrophication to encourage macroparasite infection of two invertebrate taxa in estuaries. It was hypothesised that enriched estuaries would experience greater richness and intensity of parasitic infection.

The two invertebrate taxa chosen to be screened in estuaries of differing levels of enrichment were the green shore crab *Carcinus maenas* and polychaete *Hediste diversicolor*. Both invertebrates are abundant in estuaries, have wide physiological tolerances and Atlantic-wide distributions (Crothers, 1967, Taylor *et al.*, 1973, Scaps, 2002). As a decapod crab, *C. maenas* experience a high degree of parasitic infection from a range of macroparasite taxa (Thieltges *et al.*, 2009, Lyons, 2011). Outside of its native range *C. maenas* is found in Pacific North America, southern Australia,

Tasmania, Japan and South Africa (Crothers, 1967, Carlton & Cohen, 2003). The success of *C. maenas* as an invasive is partly attributed to its escape from native parasites (Torchin *et al.*, 2001, Zetlmeisl *et al.*, 2011). Polychaetes are parasitised by a range of helminth taxa (Margolis, 1971; 1973, Vismann, 1990), but are rarely considered in parasitological studies (Margolis, 1971, 1973, Peoples *et al.*, 2012). *Hediste diversicolor* play a role in the bioturbation of sediment, are a common prey of birds and have potential economic value as a bait fishery (Scaps, 2002, Nybakken & Bertness, 2005, Waston *et al.*, 2016).

Bird communities were also surveyed as final hosts are known to play a key role in the distribution of macroparasites (Hechinger & Lafferty, 2005, Levakin *et al.*, 2013, Feis *et al.*, 2015).

4.2. MATERIALS AND METHODS

4.2.1 Study Sites

Three estuaries were sampled on the southwest coast of Ireland; Clonakilty (51°36'24.3"N 8°52'37.5"W), Oysterhaven (51°42'20.8"N 8°27'37.5"W) and Youghal (51°97' 862" N, 007°85' 844" W) (Fig. 1) Clonakilty Bay is a small estuary (approx. 272 ha) with mud and sand flats. It is a proposed Special Protection Area (SPA), a proposed Natural Heritage Area (NHA) and a candidate Special Area of Conservation (SAC). Clonakilty estuary has transient algal mats within the bay and the town sewage inflow is situated at the head of the estuary. Youghal is a moderately-sized (approx. 729 ha), sheltered south-facing estuary. Youghal is an SPA into which the town sewerage system discharges untreated effluent to the harbour via 3 sewage outfalls. Oysterhaven is a small (approx. 100 ha) south facing estuary. Oysterhaven estuary is not an SPA but is adjacent to the Sovereign Islands SPA. The nutrient statuses of Irish estuaries were assigned by the Environmental Protection Agency (EPA) according the

Water Framework Directive (WFD). Estuaries' nutrient statuses were evaluated by the analysis of water parameters ammonia (NH₃), ortho-phosphate (MRP), total nitrogen (TN), total oxidised nitrogen (TON), total phosphorus (TP) as mgL⁻¹, temperature and salinity. Clonakilty was classified as eutrophic by the EPA according to the classifications of WFD This was due to elevated nutrient concentrations and accelerated growth of macroalgae. Youghal and Oysterhaven were classified as "moderate" and "unpolluted" water quality according the WFD (O'Boyle *et al.*, 2010). "Eutrophic" status is awarded when threshold values for nutrients, macroalgal growth and an "undesirable disturbance" are exceeded. "Moderate" status is awarded when two criteria were breached and "unpolluted" status when no criteria were exceeded.

Chapter 4



Fig. 1. Study sites (a) Youghal estuary (b) Oysterhaven estuary (51°42'51.8"N 8°28'29.0"W), middle (51°42'29.5"N 8°27'32.2"W) and outer (51°42'04.5"N 8°27'30.4"W) points and (c) Clonakilty estuary. (i) Inner, (ii) middle and (ii) outer sampling points with sites designated with an arrow. Adapted from Environmental Protection Agency Water Quality shapefile.

127

4.2.2 Sampling

4.2.2.1 Host samples

Carcinus maenas and *H. diversicolor* were sampled at three fixed points within each estuary in June, July and August 2015 (Fig. 1). Sampling points were at the inner, middle and outer zones of the estuary where *C. maenas* was abundant to account for spatial variation in salinity. In Clonakilty estuary, the inner (51°61'932" N, 008°88'103" W), middle (51°36'264"N, 008°52'454"W), and outer (51°36'10.9"N, 8°51'14.8"W) points in were sampled. The inner (51°97'862" N, 007°85'844" W), middle (51°95'881" N, 007°84'951" W) and outer (51°94'641" N, 007°84'174" W) of Youghal estuary. In Oysterhaven, the inner (51°42'51.8"N 8°28'29.0"W), middle (51°42'29.5"N, 8°27'32.2"W) and outer (51°42'04.5"N, 8°27'30.4"W) points. The inner (51°56'46.8"N, 7°50'32.9"W) of Youghal estuary. At low tide, *Carcinus maenas* and *Hediste diversicolor* were haphazardly sampled respectively, by hand or from the top 20 cm of sediment until 30 of each were found.



Fig. 2 Schematic diagram of the sampling programme for intertidal invertebrates and water samples for each estuary. Factors are shown on the left.

4.2.2.2 Potential avian host surveys

Each month, point censuses were taken of bird populations at the mid points of estuaries sampled (Fig. 3). Within a week of invertebrate sampling the abundance of all species was recorded at the same locations with survey methodology based on that of Bibby *et al.*, (2000). Surveys were conducted during a flood tide, occurred within a week of invertebrate sampling events and lasted an hour and thirty to forty-three minutes. Ambiant temperature and wind speed were taken. Cloud cover was categorised in oktas as defined by the World Meteorological Organisation (WMO); 0 for clear skies, 1–2 for few clouds, 3–4 for scattered clouds, 5–7 for broken clouds, and 8 for overcast skies.



Fig. 3 Schematic diagram of the sampling programme for point censuses of bird populations

4.2.2.3 Water samples

Samples of estuarine water were collected during the flood tide at the three sampling points in each estuary on a monthly basis. At each of these points 500ml of water was collected in opaque plastic containers previously washed once with 10% HCL solution and twice with deionised water. Salinity and temperature of water were measured at the exact location of sampling, prior to the emersion of a container, with a handheld

conductivity meter (Cond 330i/Set 2C20-001). Turbidity was estimated with a secchi tube. For each sample, containers were opened just below the surface (ca. 10cm), in depths of less than 0.5m, away from sediment disturbance and approximately 3 hours after low water. Containers were sealed under water, after at least a minute post immersion, then placed on ice and brought to the laboratory for processing as soon after sampling as possible.

All water analysis was carried out by the accredited Aquatic Services Unit (ASU) laboratory in UCC. The ASU regularly participates in an international laboratory proficiency scheme (Aquacheck) to ensure the quality of the data generated. Water chemistry parameters, Ammonia (mg/L), Ortho-P (mg P/L), Total Oxidised Nitrogen (mg N/L), Total Phosphorus (mg P/L) and Total Nitrogen (mg N/L), were measured using standardised laboratory procedures. Ammonia (Total Ammonia Nitrogen, including both NH₄⁺ and NH₃) was analysed with flow injection analysis (FIA) using the supplied method QuikChem Method 10-107-06-3-D. Ortho-P was analysed manually on filtered samples using the Murphy Riley (1962) molybdate/ascorbic acid method 10-107-04-1-C. TP was analysed manually on unfiltered samples using the Murphy Riley (1962) molybdate/ascorbic acid method after sample digestion. TN was analysed with FIA on unfiltered samples using the supplied method QuikChem Method 10-107-04-1-C after sample digestion.

4.2.4 Processing

Invertebrates were frozen after collection (ca.-18°C), as freezing does not affect size of trematodes' metacercariae (Lepitzki et al., 1994). After defrosting and prior to examination invertebrates were weighed, sexed and in the case of C. maenas, carapace width was measured. Ten of the thirty C. maenas collected were haphazardly subsampled and classified as male or female depending on the number and development of pleopods (Shen, 1935). Hediste diversicolor were classed as female with the presence of large eggs ($\geq 200 \mu m$), male with white masses in the coelom and undifferentiated with the absence of both (Dales 1950). The visceral cavity of C. maenas was excised and examined as trematode metacercariae are concentrated within the area (Torchin et al., 2001, Torchin et al., 2002, Blakeslee et al., 2009). Whole mounts of *H. diversicolor* were examined. All tissue squashes were examined under x40 to x100 magnifications with transmitted light. Parasites were identified based on morphology and descriptions provided by Burt (1962), Crothers (1968), Brock & Lightner (1990), Meyers (1990), Raftos & Cooper (1990), Koie (2000), Pina et al., (2011) and Peoples et al., (2012). Larval trematodes were distinguished according to their colouration, size and if present, acetabula (suckers), excretory systems, stylet and eyespots. For metacercariae, the cyst wall or collar spines, again if present, were also used to distinguish between taxa. Cestode taxa were identified by the lack of an alimentary canal and the presence of the protoscolex (precursor to attachment sucker) and cercomer (caudal appendage). Parasitological terminology and microscopes follow the specifications of Chapter 2 Section 2.2.3 page 43.

4.2.5 Analysis

The water and invertebrate samples were 3 factor analyses. The factors were estuary (fixed, orthogonal, 3 levels), month (fixed, random, 3 levels) and site (random, nested in estuary, 3 levels) (See Fig. 2). The bird point surveys were two factor analyses with

two factors estuary (estuary, fixed, orthogonal, 3 levels) and month (fixed, orthogonal, 3 levels). Analysis was carried out in R studio (R Core Team, 2017).

4.2.5.1 Invertebrate and parasite data analysis

Normality was formally tested by the Shapiro–Wilk test. Homogeneity of variance was then checked with the Cochran test. Prior to analysis, non-normal data were transformed; count data were transformed via square root, proportional data transformed via arcsine and continuous data were transformed via log. If transformation failed to normalise the distribution of data, non-parametric tests were carried out. Host characteristics were compared with a parametric ANOVA test and a post hoc Tukey honest significant difference (Tukey HSD) test. Prevalences between sites, zones, sampling events and sexes were compared with a non-parametric Pearson's Chi-squared test with Yates Correction. Intensity of infection was compared with the non-parametric Kruskal-Wallis rank sum test and a paired Wilcoxon test with Bonferroni correction. Statistical significance was considered at P<0.05.

4.2.5.2 Water samples

Physico-chemical parameters were compared with a parametric ANOVA test and a post hoc Tukey HSD test.

4.2.5.3 Potential avian host surveys

Abundance and species richness were the factors considered in the survey. Richness was compared with the Shannon-Wiener diversity index (H). Pearson correlations between the counts of bird taxa listed as final hosts of trematode taxa recorded and trematode intensity were carried out. N(N-1)/ Σ n(n-1)

4.3. RESULTS

4.3.1 Parasite taxa

In total, five macroparasite taxa were recorded. Trematodes, a cestode, unidentified copepods, nematodes and a halacarid mite were recorded in the 1,002 invertebrates screened (Fig.4). Trematodes dominated the parasitic communities of both host species screened at all sampling events and all sites (Fig. 4). Trematode infections in C. maenas were solely metacercariae of a microphallid, most likely Microphallus primas (Jägerskiöld, 1908). However, the trematode is referred to as Microphallus spp. hereafter as morphologically similar but distinct microphallid species have been recorded in the region in C. maenas (Gibson et al., 2005, Leung et al., 2009, Lyons, 2011) and Pagurus bernhardus (Linnaeus, 1758) (Lynch et al., 2015). This was similar to the approach of Lynch et al., (2015), in the preliminary survey of P. bernhardus (Lynch et al., 2015). Microphallus spp. was the most common parasite recorded in the study with 82% of all C. maenas infected. Hediste diversicolor was parasitised by a range of parasite taxa: trematodes, larval nematodes, cestodes, copepods and mites (Fig. 4a). Trematode infections of H. diversicolor consisted of metacercariae of an echinostomatid, likely Himasthla militaris (Rudolphi, 1802), and a gymnophallid, likely Gymnophallus choledochus (Odhner, 1900). Metacercariae of the echinostomatid trematode (15%) were more common than the gymnophallid (5%). The nematode, Cucullanus sp. (Müller, 1777), and an unidentified copepod harboured by *H. diversicolor* were the next most abundant parasite as both had a prevalence of 2%. The cestode, Ophryocotyle proteus, Friis, 1870, and halacarid mite were the least common taxa with a prevalence of 0.15%. The site richest in parasitic taxa was the enriched site, Clonakilty with 7 parasitic taxa recorded, and was followed by the Oysterhaven (5), and Youghal estuaries (4).



Fig.4. (a) The prevalence of macroparasite taxa in *Hediste diversicolor* in estuaries per site and sampling event in 2015. (b) The prevalence of trematode *Microphallus* spp. metacercariae taxa in *Carcinus maenas* per site and sampling event in 2015. Error bars for prevalence are upper 95% Clopper Pearson Intervals. Median values of metacercariae intensities are plotted on the secondary axis. Note the differing y-axes between (a) and (b).



Fig. 5. Photomicrographs of helminth larvae recorded at three estuaries on the southwest coast of Ireland. (a) Microphallid metacercaria in the two developmental stages (i) and (ii). Note the characteristic thick cyst wall (arrow) and dark gonad and vitellarium tissue (thick arrow); (b) Echinostomatid metacercariae with collar spines (arrow) which are characteristic of the family; (c) Gymnophallid metacercariae, note the lack of cyst wall (arrow) and surrounding orange colouration (thick arrow); (d) *Ophryocotyle* sp. cysticeroid with protoscolex (arrow). Microphallid metacercariae were recorded in *Carcinus maenas*. Helminth larvae (b-d) were recorded in *Hediste diversicolor*. Scale bars denote 100µm.

<u>Carcinus</u> maenas

Microphallus spp. prevalence (χ^2 =54.31, df=2, P<0.0001) and intensity (χ^2 =104.09, df=2, P<0.0001) significantly differed between estuaries. For all sampling events, higher prevalences and intensities of *Microphallus* spp. were recorded in the enriched site, Clonakilty estuary, relative to Oysterhaven and Youghal estuaries (Fig. 4 & 6). The most significant differences in prevalence and intensity occurred between Clonakilty and Youghal estuaries (P<0.0001). Overall, 96%, 92% and 59% of *C. maenas* were infected with *Microphallus* spp. in Clonakilty, Oysterhaven and Youghal estuaries, respectively.

Prevalence of *Microphallus* spp. in *C. maenas* varied within estuaries (see Fig. 4). In Youghal estuary, *Microphallus* spp. was significantly more prevalent in the inner and middle sites relative to the outer site (χ^2 =33.20, df=2, P<0.0001). Intensity of *Microphallus* spp. within estuaries differed in Clonakilty (χ^2 =8.58, df=2, P<0.05) and Oysterhaven (χ^2 =7.31, df=2, P<0.05). Higher intensities of *Microphallus* spp. were recorded the middle site relative to the outer site in the case of Clonakilty (P<0.05) and relative to the inner site in Oysterhaven (P=0.05).

The intensity of *Microphallus* spp. significantly differed between sampling dates only in Clonakilty estuary, (χ^2 =8.5, df=2, P<0.05). Higher intensities were recorded in August relative to June (P<0.05). Prevalence of *Microphallus* spp. in relation to sampling date only differed in Oysterhaven estuary with significantly lower prevalences recorded in August (χ^2 =7.78, df=2, P<0.05). *Microphallus* spp. prevalence did not differ between sampling sites for Clonakilty and Oysterhaven estuaries.

Microphallus spp. prevalence and intensity were positively correlated (r=0.25, df=291, P<0.0001). Prevalence or intensity of *Microphallus* spp. did not differ

between sexes of *C. maenas*. However, *Microphallus* spp. intensity positively correlated with *C. maenas* weight (r=0.21, df=285, P<0.001) and carapace width (r=0.15, df =289, P<0.05).

Hediste diversicolor

Nematodes were found in 1%, 5% and 0.4% of *H. diversicolor* sampled from Clonakilty, Oysterhaven and Youghal, respectively (Fig. 4). 2% of *H. diversicolor* from Oysterhaven and Youghal contained copepods. The least common parasitic taxa in *H. diversicolor* were the cestode, *Ophryocotyle proteus* (Friis, 1870) and the halacarid mite with a prevalence of 0.15%. The cestode *O. proteus* and the halacarid mite were recorded in Clonakilty in two separate *H. diversicolor*.

Echinostomid metacercariae were significantly more prevalent in Clonakilty estuary than other estuaries(χ^2 =84.00, df=2, P<0.001). 32.7% of *H. diversicolor* screened from Clonakilty, 10.8% from Oysterhaven and 3.3% from Youghal were infected by the echinostomid. The intensity of the echinostomid metacercariae was significantly higher in Clonakilty than in Youghal estuary (χ^2 =7.76, df=2, P <0.05). Prevalence of the gymnophallid significantly differed between sites (χ^2 =22.29, df=2, P<0.001). Highest prevalences of the gymnophallid were recorded in Clonakilty with 10.4% of *H. diversicolor* screened infected. Low prevalence was recorded in samples from Oysterhaven (3.3%) and Youghal (2%) estuaries (Fig. 4).

The prevalence of the echinostomid trematode significantly differed between sites sampled in Clonakilty (χ^2 =35.51, df=2, P<0.0001), Oysterhaven (χ^2 =11.01, df=2, P<0.01) and Youghal (χ^2 =31.71, df=2, P<0.0001) estuaries (Fig. 7). The prevalence of the gymnophallid also differed between sites in Clonakilty (χ^2 =23.77, df=2, P<0.001), Oysterhaven (χ^2 =6.13, df=2, P=0.05) and Youghal (χ^2 =15.24, df=2, P<0.001) estuaries (Fig. 8). Prevalence of the echinostomid trematode was lower in

the outer (more coastal) site in Clonakilty, Oysterhaven and Youghal estuaries, relative to the inner (brackish) site. In Clonakilty and Youghal lower prevalence of the gymnophallid was recorded in worms collected from outer sites relative to the inner sites. In Oysterhaven, the echinostomid had lower prevalence and the gymnophallid was absent, in the middle site of the estuary. The intensity of both trematode taxa did not significantly differ between sites within the estuaries.



Fig. 6. Intensity of infection of *Carcinus maenas* by microphallid metacercariae in the sites within the estuaries (a) Clonakilty (b) Oysterhaven and (c) Youghal for the months of June, July and August 2015. Note the differing y-axes.





Fig. 7. Intensity of infection of *Hediste diversicolor* by echinostomid metacercariae in the sites within the estuaries (a) Clonakilty (b) Oysterhaven and (c) Youghal for the months of June, July and August 2015. Note the differing y-axes.



Fig. 8. Intensity of infection of *Hediste diversicolor* by gymnophallid metacercariae in the siteswithin the estuaries (a) Clonakilty (b) Oysterhaven and (c) Youghal for the months of June, July and August 2015. Note the differing y-axes.

Chapter 4

4.3.2 Host morphometrics

Carcinus maenas

Carcinus maenas weight (χ^2 =20.92, df=2, P<0.0001) and carapace width (χ^2 =16.86 df=2, P<0.001) significantly differed between the three estuaries. *Carcinus maenas* sampled from Youghal estuary were larger than those sampled from Clonakilty (P<0.001) and Oysterhaven (P<0.05) estuaries. *Carcinus maenas* morphometrics also significantly varied between sites within Oysterhaven and Youghal estuaries. Larger *C. maenas* were sampled from the middle site in Oysterhaven (χ^2 =8.74, df=2, P<0.05) and Youghal (χ^2 =7.73, df=2, P<0.05) estuaries.

The sex ratio of *C. maenas* populations sampled did not significantly differ between estuaries, sites, or sampling events. The colour morphs of *C. maenas* significantly varied between estuaries (χ^2 =21.07, df=2, P<0.0001). The green colour morph outnumbered the red morph in all estuaries (Table 1). In order of decreasing abundance, the red colour morph was found in Youghal, Oysterhaven and Clonakilty estuaries (Table 1). The occurrence of the red morph of *C. maenas* significantly differed between sitesin Clonakilty (χ^2 =9.50, df=2, P<0.001) and Youghal estuaries (χ^2 =16.80, df=2, P<0.001). The red morph of *C. maenas* was more common in the outer sampling site of Clonakilty estuary. In the case of Youghal estuary, the red morph was absent from the outer site.

Table 1	Carcinus	maenas	morphometrics	averaged	over	all	sampling events a	ıre
presente	d as x̄ (±SI	D). Male	C. maenas is rep	presented b	oy m a	ind t	female crabs by f.	
Parame	eters				Es	tuar	ries	

1 drumeters	Listuaries				
	Clonakilty	Oysterhaven	Youghal		
Weight (g)	10.63 (±7.4)	10.57 (±4.53)	14.03 (± 8.25)		
Carapace width (mm)	34.62 (±7.89)	36.63 (±6.78)	38.36 (±7.86)		
Sex ratio (m:f)	1:0.83	1:1.14	1:0.38		
Colour morph ratio (red: green)	1:32	1:14	1:8		
Hediste diversicolor

The wet weights of *H. diversicolor* did not significantly differ between estuaries (F=2.426, df=2, P>0.05) but did between siteswithin estuaries (see Table 2). In Clonakilty estuary, significantly heavier *H. diversicolor* were recorded in the outer site (χ^2 =37.5, df=2, P<0.0001) relative to the inner and middle sites. In Oysterhaven estuary significantly heavier *H. diversicolor* were sampled from the middle site relative to the inner and outer sites (F=14.89, df=2, P<0.0001). In Youghal estuary heavier *H. diversicolor* were recorded in the outer site relative to the inner and outer sites (χ^2 =11.81, df=2, P<0.05).

Table 2 *Hediste diversicolor* morphometrics averaged over all sampling events are presented as \bar{x} (±SD). *Hediste diversicolor* with undifferentiated gonads are represented by u, males by m and female with f.

Parameters	Estuaries							
	Clonakilty	Oysterhaven	Youghal					
Weight (g)	0.55 (±0.26)	0.53 (±0.33)	0.67 (±0.49)					
Sex ratio (u:m:f)	1:0.18:0.25	1:0.84:0.54	1:0.36:0.26					

4.3.3 Physico-chemical parameters

According to the nutrient parameters measured, Clonakilty estuary was the most nutrient rich estuary (Table 3). Results agreed with the WFD designation. Total phosphorous differed between estuaries (F=7.37, df=2, P<0.01) with significantly higher values recorded in Clonakilty (Tukey HSD, P<0.05). Ortho-phosphate significantly differed between estuaries (F=37.70, df=2, P<0.001) with higher values again recorded in Clonakilty (Tukey HSD, P<0.001). Levels of ortho-phosphate and

total phosphorous were highest in June and July at Clonakilty and exceeded threshold values for WFD classifications for unpolluted waters (O'Boyle *et al.*, 2010). Temperature also significantly differed between estuaries (F=6.02, df=2, P<0.01) with higher temperatures recorded at Clonakilty (Tukey HSD, P<0.01)

	Cl	Clonakilty Oysterhaven				Youghal			Grand	
Values	Inner	Middle	Outer	Inner	Middle	Outer	Inner	Middle	Outer	Total
Ammonia (mgL ⁻¹)	0.23	0.04	0.05	0.04	0.05	0.03	0.05	0.06	0.04	0.07
	(±0.02)	(±0.01)	(±0.02)	(±0.01)	(±0.03)	(± 0.01)	(± 0.01)	(±0.02)	(±0.01)	(± 0.06)
Ortho-P (mgL ⁻¹)	0.11	0.03	0.01	<0.01	<0.01	<0.01	0.01	0.01	0.01	0.02
	(±0.05)	(±0.02)	(±0.01)	(±<0.01)	(±<0.01)	(±<0.01)	(±<0.01)	(±<0.01)	(±<0.01)	(±0.04)
Total Phosphorus (mgL ⁻¹)	0.32	0.13	0.06	0.04	0.05	0.04	0.13	0.03	0.04	0.09
	(±0.22)	(±0.08)	(±0.02)	(±0.02)	(±0.01)	(±0.02)	(±0.09)	(±0.01)	(±0.01)	(±0.11)
Total Nitrogen (mgL ⁻¹)	2.19	1.17	0.76	0.85	0.90	0.75	1.56	0.71	0.62	1.06
	(±0.77)	(±0.18)	(±0.21)	(±0.49)	(±0.36)	(±0.52)	(±0.52)	(±0.20)	(±0.05)	(±0.61)
Total Oxidised Nitrogen (mgL ⁻¹)	0.93	0.12	0.05	0.27	0.14	0.06	0.78	0.24	0.15	0.30
	(±0.34)	(±0.09)	(±0.02)	(±0.31)	(±0.11)	(± 0.05)	(±0.33)	(±0.15)	(±0.10)	(±0.35)
Salinity (%)	25.47	28.07	31.67	28.90	30.87	31.13	21.63	28.77	31.93	28.71
	(±7.80)	(±5.50)	(±2.36)	(±2.38)	(±2.97)	(± 2.32)	(±4.75)	(±3.39)	(±2.69)	(± 4.75)
Turbidity (NTU)	8.67	5.00	6.67	10.33	5.00	5.00	56.67	6.67	6.67	12.30
	(±3.21)	(±<0.01)	(±2.89)	(±9.24)	(±<0.01)	(±<0.01)	(±80.98)	(±2.89)	(±2.89)	(±27.79)
Temperature (°C)	21.90	22.70	18.07	15.63	16.27	16.23	19.20	17.67	19.13	18.53
	(±2.77)	(±2.76)	(±1.62)	(±2.20)	(±0.23)	(±0.81)	(±4.35)	(±4.01)	(±6.03)	(±3.60)

Table 3 Physico-chemical water parameters sampled from three sites within each estuary sampled for June-August 2015. Parameters are presented as \bar{x} (±SD) and n=3 per site within estuaries sampled.

4.3.4 Potential avian host surveys

There was no significant difference in avian richness between estuaries with 25 bird species recorded in Oysterhaven and 24 in both Clonakilty and Youghal (see Table 4). High abundances of birds were found in Clonakilty with 1, 249 individuals recorded, 561 individuals in Youghal and 264 in Oysterhaven. The abundance and distribution of species did vary between months with peaks seen in August. Simpson's Diversity index found Oysterhaven to have the most diverse bird community on average with an index of 1.69 (SD 1.45), followed by Youghal with 9.16 (SD 0.72) and Clonakilty with 6.79 (SD 2.04). Of the potential final hosts for helminth taxa recorded, gulls *Larus* spp. (Linnaeus, 1758), the ruddy turnstone *Arenaria interpres* (Linnaeus, 1758) and the Eurasian oystercatcher *Haematopus ostralegus*, only the abundance *H. ostralegus* significantly differed between estuaries with larger numbers recorded in Clonakilty (F=19.42, df=2, P<0.05).

Bird Taxa	Estuary					
-	Clonakilty	Oysterhaven	Youghal	Total		
Actitis hypoleucos, (Linnaeus, 1758)	1	5	6	12		
Common sandpiper						
Anas spp. (Linnaeus, 1758)	39	23	8	70		
Dabbling ducks						
Ardea cinerea (Linnaeus, 1758)	3	4	11	18		
Grey Heron						
Arenaria interpres	1	0	0	1		
Ruddy turnstone						
Bubulcus ibis (Linnaeus, 1758)	3	7	0	10		
Cattle egret						
Calidris spp. (Merrem, 1804)	16	3	0	19		
Sand pipers						
Carduelis carduelis (Linnaeus, 1758)	0	2	0	2		
European goldfinch						
Charadrius hiaticula (Linnaeus, 1758)	0	0	3	3		
Common ringed plover						
Columba palumbus (Linnaeus, 1758)	0	13	0	13		
Common Woodpigeon						
Corvus spp. (Linnaeus, 1758)	58	40	30	128		
Crow species						
Cygnus olor (Gmelin, 1789)	48	0	0	48		
Mute swan						
Egretta garzetta (Linnaeus, 176	66) 12	3	16	31		
Little egret						
Haematopus ostralegus	101	3	67	171		
Oystercatcher						
Hirundo rustica (Linnaeus, 1758)	0	4	5	9		
Barn swallow						
Larus spp.	532	84	210	826		
Gulls						
Limosa spp. Brisson, 1760	247	27	48	322		
Godwit						
Numenius spp. Moehring, 1758	145	29	74	248		
Curlews						
Phalacrocorax carbo (Linnaeus, 1758)	2	2	14	18		
Great cormorant						
Sterna spp. (Linnaeus, 1758)	0	0	26	26		
Terns						
Tadorna tadorna (Linnaeus, 1758)	11	0	8	19		
Common shelduck						
Tringa spp. (Linnaeus, 1758)	30	15	35	80		
Shanks						
Total number of individuals	1249	264	561	2074		

Table 4 Avian taxa recorded in the three estuaries for the months June-August 2015 summed for all sampling events

Table 5 Correlation between trematode prevalence and final host abundance for all sites and sampling events in Summer 2015 for the three estuaries combined. Asterisk denote level of significance * for P<0.05, ** P<0.001 and *** P<0.0001

Trematode species	Final avian host	Correlation coefficient
Echinostomatid	Anas sp.	-0.13**
	Arenaria interpres	0.16***
	Calidris sp.	0.22***
	Haematopus ostralegus	0.06
	<i>Larus</i> sp.	-0.21
	<i>Limosa</i> sp.	0.36
	<i>Numenius</i> sp.	0.01
	<i>Sterna</i> sp.	-0.43**
	<i>Tringa</i> sp	0.07
Gymnophallid	Anas sp.	0.31***
	Haematopus ostralegus	0.24***
	Larus sp.	0.20***
	Sterna sp.	-0.27***
Microphallus spp.	Anas sp.	0.34***
	Haematopus ostralegus	0.03*
	Larus sp.	0.21***
	<i>Tringa</i> sp.	0.13*

Table 6 Correlation between trematode taxa and physico-chemical properties of estuarine water samples from all sampling points and events in Summer 2015 for all estuaries combined. NA represents not applicable as microphallid metacercariae were recorded in *Carcinus maenas* and echinostomid or gymnophallid metacercariae were recorded in *Hediste diversicolor*. Asterisk denote level of significance * for P<0.05, ** P<0.001 and *** P<0.0001.

	Echinostomatid	Gymnophallid	Microphallid	NH ₃	PO4 ³⁻	TON	TP	TN	Salinity	Turbidity	Temperature
Echinostomatid	1										
Gymnophallid	0.13**	1									
Microphallid	NA	NA	1								
NH ₃	0.02***	-0.15***	-0.15	1							
PO ₄ ³⁻	0.22***	-0.07	-0.15	0.52	1						
TON	0.11***	-0.24***	0.02	0.59	0.31	1					
TP	021***	-0.11***	0.48***	0.21	0.57	0.35	1				
TN	0.17***	-0.20***	0.33***	0.35	0.44	0.62	0.85	1			
Salinity	-0.03	0.02	-0.20 ***	-0.08	-0.03	-0.64	-0.21	-0.35	1		
Turbidity	-0.17***	-0.18***	0.02	0.42	0.40	0.56	0.31	0.28	-0.41	1	
Temperature	0.18***	0.07	0.11	0.25	0.62	0.30	0.56	0.52	-0.26	0.51	1

4.4. DISCUSSION

The study supports the hypothesis that eutrophication encourages macroparasite infection. The mechanisms by which eutrophication facilitates trematode infection are proposed to act at the juvenile stages (Johnson *et al.*, 2008). Excess nutrients are speculated to encourage intermediate host density and/or output of infective stages. No trematode infections preceding metacercariae, i.e. cercarial, were recorded during this study. However, the prevalence and intensity of metacercariae recorded in the enriched site indicates that trematode infections could be facilitated.

Higher abundances of avian hosts, and parasitic taxa were recorded at the eutrophic site, Clonakilty. In this study, trematode prevalence in both hosts was positively correlated with total Nitrogen and total Phosphorous concentrations. Nitrogen in salt marshes has been linked to increased trematode prevalence (Altman *et al.*, 2014). The differences in trematode intensity between estuaries could not be attributed to differences in host morphometrics. *Hediste diversicolor* did not significantly differ in size or between estuaries. Moreover, significantly lower intensity and prevalence of *Microphallus* spp. occurred in Youghal in which significantly larger *C. maenas* were screened. Stronger correlations were found with the proposed hosts for taxa found. Final hosts' distribution and movement are known to play a determining role in the distribution and prevalence of trematodes (Hechinger & Lafferty, 2005, Feis *et al.*, 2015).

Of the two invertebrate host species screened, *C. maenas* was the most commonly infected. Parasite communities were not taxa rich and were dominated by trematode metacercariae. The higher prevalence and intensity of *Microphallus* spp. is indicative of the trematode's ecological relevance and importance. The method of hand collection may produce high estimates of parasitic infection. Heavily infected *C. maenas* can be lethargic due to physiological stress and thus slower and, more easily

sampled by hand (Crothers, 1968, Werner, 2001). As a result, estimates of intensity of trematode infection in *C. maenas* sampled may be slightly higher than in the total population.

In all estuaries trematodes were the predominant helminth with high intensities of Microphallus spp. metacercariae in C. maenas. This finding agrees with the work of Zetlmeisl et al., (2011) and Blakeslee et al., (2015). Richness of macroparasite taxa recorded in C. maenas was lower than previous work in the United Kingdom and New Zealand but was consistent with histological work on the hermit crab, Pagurus bernhardus (Linnaeus, 1758) in Ireland (Stentiford & Feist 2005, Lynch et al., 2015). Notably lacking from C. maenas sampled in this study was the rhizocephalan Sacculina carcini, Thompson, 1836, the turbellarian Fecampia erythrocephala, Giard, 1886, and the acanthocephalan Profilicollis botulus (Van Cleave, 1916). The absence of these taxa is likely due to the habitat and sizes of C. maenas sampled. Sacculina carcini is commonly found in subtidal habitats (Waser et al., 2016). This distribution could be attributed to parasitic manipulation, with castrated or "feminised" C. maenas moving into the subtidal, like gravid females would. The subtidal habitat may facilitate the survival of S. carcini infective stages increasing the likelihood of them reaching and infecting a decapod (Rasmussen 1959, Werner, 2001, Waser et al., 2016). Alternatively, the parasitic taxa could have eluded detection due to their small size or an early stage of infection. According to Kuris et al., (2002), F. erythrocephala is commonly found in small C. maenas with carapaces less than 11mm in width.

Macroparasite communities of *C. maenas* warrant further study, as they play a role in the ecology and physiology of *C. maenas* (Zetlmeisl *et al.*, 2011). Microphallids, in particular *Microphallus similis*, affect the body mass index, energy storage, and immune activation of *C. maenas* in its invasive range (Blakeslee *et al.*, 2015). The escape of *C. maenas* from two thirds of its native parasites is considered key to the crab's invasive success (Torchin *et al.*, 2001). In the invasive range, *C. maenas* is not

only parasitised by fewer taxa but also to a lesser extent than individuals within its native range (Torchin *et al.*, 2001, Blakeslee *et al.*, 2015). The higher intensity of *Microphallus* spp. may have implications for the health of *C. maenas*. Heavy microphallid infections in the crab *Neohelice granulata* (Dana, 1851) resulted in structural damage to the hepatopancreas, necrosis of digestive tubules and impaired physiological storage (Robaldo *et al.*, 1999). Heavy infection by microphallid metacercariae in *C. maenas* in the field impaired the crab's self-righting ability (Blakeslee *et al.*, 2015) and survival under laboratory conditions (Stunkard 1957).

Hediste diversicolor was found to harbour greater parasite richness than *C. maenas* in the three estuaries surveyed. *Hediste diversicolor* was used as a second intermediate host by two trematode taxa (Margolis, 1971; 1973). Polychaetes in New Zealand were primarily infected by a single trematode species in the form of metacercariae (Peoples *et al.*, 2012). High intensities of metacercariae, 18 per worm, impaired the structural integrity of New Zealand polychaetes and led to fragmentation of the worm's body (Peoples *et al.*, 2012). If metacercariae have similar effects on *H. diversicolor*, it could affect their economic value as a bait fishery (Torchin *et al.*, 2001, Peoples *et al.*, 2012).

The effect of transient algal mats in Clonakilty on host-trematode dynamics in estuaries, is likely complex. Algal mats may increase local density of intermediate invertebrate hosts (Hull, 1987, Lewis *et al.*, 2014). Dense mats of vegetation can be important feeding areas for birds that either feed directly on the vegetation or on invertebrates found within the mats (Goss-Custard *et al.*, 2006). Algal mats may favour trematode taxa by increasing the abundance of their common primary intermediate host the Laver spire shell *Peringia ulvae* (Pennant, 1777), formerly *Hydrobia ulvae*. *Peringia ulvae* can colonise and consume transient algal mats that are present in Clonakilty estuary (Soulsby *et al.*, 1982, Hull, 1987). *Peringia ulvae* is also a host for a large range of trematode species including *Microphallus* spp. and echinostomatids (Stunkard, 1957, Deblock 1980). Proximity of trematode hosts is vital

for the taxa's transmission and the increased density of primary intermediate trematode hosts is likely to facilitate high prevalence of *Microphallus* spp. recorded (Fredensborg & Poulin, 2006).

The infective stage preceding the metacercaria, the cercaria, is short lived, free living and mobile. It is usually released into the water column to encounter and infect a suitable host and continue its life cycle. Under laboratory conditions, macroalgae can impair the dispersal and successful infection of cercariae (Welsh *et al.*, 2014). This could decrease the intensites of metacercarial infections as the local abundance of cercariae determines the encounter rates of the trematode with their crustacean intermediate hosts (Thieltges *et al.*, 2009). However, the high intensities of *Microphallus* spp. recorded in Clonakilty suggest that any physical disturbance of infective stages by algae, is overcome by increased densities of gastropod hosts.

Eutrophication from organic sources has been recorded to favour gulls (Raven & Coulson 2001). The effect of algal mats on avian hosts varies in magnitude and in relation to the avian species. Soulsby *et al.*, (1982) suggested that the effect was small and several prey items decreased due to covering or algal decomposition could be compensated by increased densities of the gastropod *P. ulvae*. In Clonakilty estuary, the effect of algal mats on birds is species-specific with black backed gulls, *Larus* marinus, foraging in algal cleared areas (Lewis *et al.*, 2014). Algal mats may also affect the foraging strategy of bird species within the estuary. Californian shorebirds switch from foraging pecks to probes in response to increased coverage by algal mats in eutrophic estuaries (Green *et al.*, 2015). If bird communities in Irish estuaries adopt the same foraging behaviour, this could explain the higher prevalence and richness of *H. diversicolor* in Clonakilty than the less eutrophic estuaries sampled. Algal mats may affect the density of *Hediste diversicolor*. However, the effect may vary as previous work has found that the abundance of *H. diversicolor* was greater (Hull,

1987; Norkko & Bonsdorff, 1996b) and less abundant (Lopes *et al.*, 2000) under algal mats relative to uncovered areas of mudflat.

The occurrence of trematodes in all estuaries surveyed was high. However, greater richness, prevalence and intensity of parasitic taxa was found in the eutrophic estuary of Clonakility. In the case of trematodes, this could be due to an increased output of infective stages by molluscan and/or avian hosts. There was a positive correlation between enrichment parameters, phosphorous and nitrogen, and trematode prevalence. This was a correlative field study, so interpretation and extrapolation of results was conservative. Observed patterns of trematode distributions could be confounded by additional, unmeasured variables e.g. heavy metal concentrations or differing invertebrate host abundances (Mischler et al., 2016). Human infection by trematode taxa in this study is possible but likely harmless. Cercariae from trematodes that utilise birds as final hosts, may encounter and infect the skin of a swimmer. Dermatitis or "swimmers' itch" results from the penetration of cercariae into skin and the subsequent irritation, arrested maturation or death of cercariae (Fraser et al., 2009, Horak et al., 2015). However, the dermatitis is generally temporary, due to the short lifespan of the cercariae (<24hrs) and the immune-physiology of human skin (Fraser et al., 2009, Horak et al., 2015, Thieltges et al., 2006b).

The high prevalence of trematodes in all estuaries sampled is indicative of their ecological relevance. If the hypothesis is correct and the recorded higher trematode intensity is causally related to increased nutrient parameters, there may be implications for the survival and ecological interactions of invertebrates screened. Future work should consider manipulative laboratory trials to test the mechanisms of how eutrophication may facilitate parasitism in estuaries. Laboratory trials by Johnson *et al.*, 2007 with the freshwater snail species composition *Planorbella* spp., *Ribeiroia ondatrae* and . Additionally, the effect of trematode metacercariae on the physiology and behaviour of invertebrate hosts screened warrants investigation. This study reports

the detection of trematode metacercariae in *Hediste diversicolor* and raises the question as to what impact this parasite may be having on populations of the worm in estuaries on the southwest coast of Ireland.

REFERENCES

- Airoldi L, Beck MW (2007) Loss, status and trends for coastal marine habitats of Europe. In: Gibson RN, Atkinson RJA, Gordon JDM (eds) Oceanography and Marine Biology, 45, Book 45.
- Altman I and Byers JE (2014) Large-scale spatial variation in parasite communities influenced by anthropogenic factors. *Ecology*, **95**, 1876-1887.
- Baeta A, Cabral HN, Neto JM, Marques JC, Pardal MA (2005) Biology, population dynamics and secondary production of the green crab *Carcinus maenas* (L.) in a temperate estuary. *Estuarine Coastal and Shelf Science*, **65**, 43-52.
- Bibby C, Burgess N, Hill DA, Mustoe S (2000) Bird Census Techniques. *Academic Press*, London, UK.
- Blakeslee AMH, Keogh CL, Byers JE, Kuris AM, Lafferty KD, Torchin ME (2009) Differential escape from parasites by two competing introduced crabs. *Marine Ecology Progress Series*, **393**, 83-96.
- Blakeslee AMH, Keogh CL, Fowler AE, Griffen BD (2015) Assessing the Effects of Trematode Infection on Invasive Green Crabs in Eastern North America. *PloS One*, **10**, 1-20.
- Blanar CA, Munkittrick KR, Houlahan J, MacLatchy DL, Marcogliese DJ (2009) Pollution and parasitism in aquatic animals: A meta-analysis of effect size. *Aquatic Toxicology*, **93**, 18-28.
- Brock J, Lightner D (1990) Diseases of Crustacea, *Diseases of Marine Animals*. Biologische Anstalt Helgoland, Hamburg Germany.
- Budria A (2017) Beyond troubled waters: the influence of eutrophication on hostparasite interactions. *Functional Ecology*, **31**, 1348-1358.
- Burt MDB (1962) A contribution to the knowledge of the cestode genus *Ophryocotyle* Friis, 1870. *Journal of the Linnean Society London Zoology*, **44**, 645-668.
- Carlton JT, Cohen AN (2003) Episodic global dispersal in shallow water marine organisms: the case history of the European shore crabs *Carcinus maenas* and *C. aestuarii. Journal of Biogeography*, **30**,1809-1820.
- Crothers JH (1967) The biology of the shore crab *Carcinus maenas* (L.). *Field Studies*, **2**,407-434.

- Crothers JH (1968) The biology of the shore crab *Carcinus maenas* (L.): 2. The life of the adult crab. *Field Studies*, **2**,579-614.
- Dales RP (1950) The reproduction and larval development of *Nereis diversicolor* O.
 F. Müller *Journal of the Marine Biological Association of the United Kingdom*, 29, 321-360.
- Dana JD (1851) On the Classification of the msaioid Crustacea or Oxyrhyncha. *American Journal of Scientce and the Arts*, **11**, 425-429.
- Deblock S (1980) Survey of the larval Trematoda parasites of *Hydrobia* (Prosobranches) mollusca of the coasts of France. *Parassitologia*, **22**, 1-105.
- Dobson A, Lafferty KD, Kuris AM, Hechinger RF, Jetz W (2008) Homage to Linnaeus: How many parasites? How many hosts? *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 11482-11489.
- Dušek L, Gelnar M, Šebelová Š (1998) Biodiversity of parasites in a freshwater environment with respect to pollution: metazoan parasites of chub (*Leuciscus cephalus* L.) as a model for statistical evaluation. *International Journal for Parasitology*, 28,1555-1571.
- Esch GW (1971) Impact of ecological succession on the parasite fauna in centrarchids from oligotrophic and eutrophic ecosystems. *American Midland Naturalist*, **86**, 160-168.
- Feis ME, Thieltges DW, Olsen JL, de Montaudouin X, Jensen KT, Bazairi H, Culloty SC, Luttikhuizen PC (2015) The most vagile host as the main determinant of population connectivity in marine macroparasites. *Marine Ecology Progress* Series, 520, 85-99.
- Fermer J, Culloty SC, Kelly TC, O'Riordan RM (2011) Parasitological survey of the edible cockle *Cerastoderma edule* (Bivalvia) on the south coast of Ireland. *Journal of the Marine Biological Association of the United Kingdom*, **91**, 923-928.
- Fraser SJ, Allan SJR, Roworth M, Smith HV, Holme SA (2009) Cercarial dermatitis in the UK. *Clinical and Experimental Dermatology*, **34**, 344-346.
- Fredensborg BL, Poulin R (2006) Parasitism shaping host life-history evolution: adaptive responses in a marine gastropod to infection by trematodes. *Journal of Animal Ecology*, **75**, 44-53.
- Friis GM (1970) En hitil ubeskreven Baendelorme hos Fugle. Videnskabelige Meddelelser Naturhistorisk Forening i København. (aar 1896), **1**, 121-124.

- Galli P, Crosa G, Mariniello L, Ortis M, D'Amelio S (2001) Water quality as a determinant of the composition of fish parasite communities. *Hydrobiologia*, **452**,173-179.
- Giard A (1886) Sur un Rhabdocoele nouveau, parasite et nidulant (Fécampia erythrocephala). Comptes Rendus de l'Académie des Sciences, Paris, 103, 499-501.
- Gibson D, Bray R, Harris E (2005) Host-Parasite Database of the Natural History Museum, London.
- Gmelin JF (1789) Caroli a Linné, systema naturae. 1-2, 501-1032.
- Goss-Custard J, West A, Yates MG, Caldow R, Stillman R, Bardsley L, Castilla J, Castro M, Dierschke V, Durell S, Eichhorn G, Ens B, Exo K, Udayangani-Fernando P, Ferns P, Hockey P, Gill J, Johnstone I, Kalejta-Summers B, Masero J, Moreira F, Nagarajan R, Owens I, Pacheco C, Perez-Hurtado A, Rogers D, Scheiffarth G, Sitters H, Sutherland W, Triplet P, Worrall D, Zharikov Y, Zwarts L, Pettifor R (2006) Intake rates and the functional response in shorebirds (Charadriiformes) eating macro-invertebrates. *Biological Reviews*, 81, 501-529.
- Green L, Blumstein DT, Fong P (2015) Macroalgal Mats in a Eutrophic Estuary Obscure Visual Foraging Cues and Increase Variability in Prey Availability for Some Shorebirds. *Estuaries and Coasts*, **38**, 917-926.
- Hartnett M, Wilson JG, Nash S (2011) Irish estuaries: Water quality status and monitoring implications under the water framework directive. *Marine Policy*, 35, 810-818.
- Hechinger RF, Lafferty KD (2005) Host diversity begets parasite diversity: bird final hosts and trematodes in snail intermediate hosts. *Proceedings of the Royal Society B*, **272**, 1059-1066.
- Hobbie JE (2000) Estuarine science: a synthetic approach to research and practice. Island Press, Washington, DC.
- Horak P, Mikes L, Lichtenbergova L, Skala V, Soldanova M, Brant SV (2015) Avian schistosomes and outbreaks of cercarial dermatitis. *Clinical Microbiology Reviews*, 28, 165-190.
- Howarth RW, Marino R (2006) Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: Evolving views over three decades. *Limnology and Oceanography*, **51**, 364-376.

- Hull SC (1987) Macroalgal mats and species abundance: a field experiment. *Estuarine Coastal and Shelf Science*, **25**, 519-532.
- Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, Bradbury RH, Cooke R, Erlandson J, Estes JA, Hughes TP, Kidwell S, Lange CB, Lenihan HS, Pandolfi JM, Peterson CH, Steneck RS, Tegner MJ, Warner RR (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293, 629-638.
- Jägerskiöld LA, (1908). Kleine Beitrage zur Kenntnis der Vogeltrematode. Zentralblatt für Bakteriologie Parasitenkunde **48**, 302-317.
- Johnson, Chase JM, Dosch KL, Hartson RB, Gross JA, Larson DJ, Sutherland DR, Carpenter SR (2007) Aquatic eutrophication promotes pathogenic infection in amphibians. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 15781-15786.
- Johnson P, Carpenter S (2008) Influence of Eutrophication on Disease in Aquatic Ecosystem: Patterns, Processes, and Prediction. In: *Infectious Disease Ecology* (ed. by Ostfield R, Kessing F, Eviner V). Princeton University Press, Princeton University Press.
- Johnson P, Thieltges DW (2010) Diversity, decoys and the dilution effect: how ecological communities affect disease risk. *Journal of Experimental Biology*, **213**, 961-970.
- Johnson P, Townsend AR, Cleveland CC, Glibert PM, Howarth RW, McKenzie VJ, Rejmankova E, Ward MH (2010) Linking environmental nutrient enrichment and disease emergence in humans and wildlife. *Ecological Applications*, **20**, 16-29.
- Koie M (2000) The life-cycle of the flatfish nematode *Cucullanus heterochrous*. Journal of Helminthology, **74**, 323-328.
- Kuris AM, Hechinger RF, Shaw JC, Whitney KL, Aguirre-Macedo L, Boch CA, Dobson AP, Dunham EJ, Fredensborg BL, Huspeni TC, Lorda J, Mababa L, Mancini FT, Mora AB, Pickering M, Talhouk NL, Torchin ME, Lafferty KD (2008) Ecosystem energetic implications of parasite and free-living biomass in three estuaries. *Nature*, **454**, 515-518.
- Kuris AM, Torchin ME, Lafferty KD (2002) Fecampia erythrocephala rediscovered: prevalence and distribution of a parasitoid of the European shore crab, Carcinus maenas. Journal of the Marine Biological Association of the United Kingdom, 82, 955-960.

- Lafferty KD (1997) Environmental parasitology: What can parasites tell us about human impacts on the environment? *Parasitology Today*, **13**, 251-255.
- Lafferty KD, Holt RD (2003) How should environmental stress affect the population dynamics of disease? *Ecology Letters*, **6**, 654-664.
- Lepitzki D, Scott M, McLaughlin J (1994) Influence of storage and examination methods on the recovery and size of metacercaria of *Cerastoderma edule* (L.) from commercial beds of the lower Thames estuary. *Zeitschrift für Parasitenkunde*, **56**, 1-11.
- Leung TLF, Keeney DB, Poulin R (2009) Cryptic species complexes in manipulative echinostomatid trematodes: when two become six. *Parasitology*, **136**, 241-252.
- Levakin IA, Nikolaev KE, Galaktionov KV (2013) Long-term variation in trematode (Trematoda, Digenea) component communities associated with intertidal gastropods is linked to abundance of final hosts. *Hydrobiologia*, **706**, 103-118.
- Lewis LJ, Kelly TC, Davenport J (2014) Black-tailed Godwits *Limosa limosa islandica* and Redshanks *Tringa totanus* respond differently to macroalgal mats in their foraging areas. *Wader Study Group Bulletin*, **121**, 85-93.
- Linnaeus C, (1758). Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. *Editio decima, reformata. Laurentius Salvius: Holmiae*, **2**, 1-824.
- Looss A, (1907) Zur Kenntnis der Distomen families Hemiuridae (Studies of Distomatidae Haemiuridae family). *Zoologischer Anzeiger*, **31**, 585-620.
- Lotze HK, Lenihan HS, Bourque BJ, Bradbury RH, Cooke RG, Kay MC, Kidwell, SM, Kirby MX, Peterson CH, Jackson JBC (2006) Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science*, **312**, 1806-1809.
- Lynch S, Darmody G, Laide C, Walsh D, Culloty SC (2015) A preliminary health survey of the hermit crab, *Pagurus bernhardus*, on the southwest coast of Ireland. *Journal of Invertebrate Pathology*, **127**, 73-75.
- Lyons L (2011) Aspects of the biology of decapods in South West Ireland. Doctor of Philosophy Thesis, University College Cork, Cork.
- Margolis L (1971) Polychaetes as intermediate hosts of helminth parasites of vertebrates: a review. *Journal of the Fisheries Research Board of Canada*, **28**, 1385-1392.

- Margolis L (1973) Additional notes on polychaetes as intermediate hosts of helminth parasites of vertebrates. *Journal of the Fisheries Research Board of Canada*, **30**, 469-470.
- McKenzie VJ, Townsend AR (2007) Parasitic and infectious disease responses to changing global nutrient cycles. *EcoHealth*, **4**, 384-396.
- Meyers TR (1990) Chapter 3: Diseases of Crustacea, Vol 3. Biologische Anstalt Helgoland. Hamburg Notkestrasse 31, W-2000 Hamburg 52, Germany, Hamburg.
- Mischler J, Johnson , McKenzie VJ, Townsend AR, (2016) Parasite infection alters nitrogen cycling at the ecosystem scale. *Journal of Animal Ecology*, **85**, 817-828.
- Moehring PH (1958) Geslachten der vogelen. 3, 361.
- Müller, OF (1777) Zoologica danica seu animalium 1. 1st Edition. Ghent University, Ghent, Belgium, 1-292.
- Murphy, J, Riley, JP, (1962) A modified single solution method for the determination of phosphate natural water. *Analytica Chimica Acta*, **27**, 31–36. receipt
- Nybakken JW, Bertness MD (2005) Marine biology: an ecological approach. Sixth edition. **1-6**, 1-579.
- O'Boyle R, Wilkes R, McDermott G (2010) Quality of Estuarine and Coastal Waters 2007-2009. In: McGarrigle M, Lucey J, Ó Cinnéide M (eds) Water Quality in Ireland 2007-2009. Environmental Protection Agency Ireland.
- Odhner T, (1900) par *G nereicola* Rebecq et G. Prévot, 1962 en Camargue (France) (Digenea, Gymnophallidae). *Bulletin de la Société Zoologique de France*, **99**, 551–559.
- Pennant T, (1777) British Zoology. Crustacea. Mollusca. Testacea, 4, 1-154.
- Peoples RC, Randhawa HS, Poulin R (2012) Parasites of polychaetes and their impact on host survival in Otago Harbour, New Zealand. *Journal of the Marine Biological Association of the United Kingdom*, **92**, 449-455.Pietrock M, Marcogliese DJ (2003) Free-living endohelminth stages: at the mercy of environmental conditions. *Trends in Parasitology*, **19**, 293-299.

- Pina S, Russell-Pinto F, Rodrigues P (2011) Morphological and molecular study of *Microphallus primas* (Digenea: Microphallidae) metacercaria, infecting the shore crab *Carcinus maenas* from northern Portugal. *Folia Parasitologica*, 58, 48-54.
- R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <u>http://www.R-project.org/</u>
- Raffaelli D (1999) Nutrient enrichment and trophic organisation in an estuarine food web. Acta Oecologica-*International Journal of Ecology*, **20**, 449-461.
- Raftos DA, Cooper EL (1990) Chapter 2: Diseases of Annelida, Vol 3. Biologische Anstalt Helgoland, Hamburg, Germany.
- Rasmussen E (1959) Behaviour of Sacculinized Shore Crabs (*Carcinus maenas* Pennant). *Nature*, **183**, 479-480.
- Raven SJ, Coulson JC (2001) Effects of cleaning a tidal river of sewage on gull numbers: a before-and-after study of the River Tyne, northeast England. *Bird Study*, **48**, 48-58.
- Robaldo R B, Monserrat J, Cousin JCB, Bianchini A (1999) Effects of metacercariae (Digenea: Microphallidae) on the hepatopancreas of *Chasmagnathus granulata* (Decapoda: Grapsidae). *Diseases of Aquatic Organisms*, **37**, 153-157.
- Rudolphi CA (1802) Beobachtungen ber die Eingerweidewurmer. Archiv für Zoologie und Zootomie, **2-3**, 1-125.
- Scaps P (2002) A review of the biology, ecology and potential use of the common ragworm *Hediste diversicolor* (OF Muller) (Annelida : Polychaeta). *Hydrobiologia*, **470**, 203-218.
- Shen CJ (1935) An investigation of the post-larval development of the shore-crab *Carcinus maenas*, with special reference to the external secondary sexual characters. *Proceedings of the Zoological Society of London*, **105**, 1-33.
- Small C, Nicholls RJ (2003) A global analysis of human settlement in coastal zones. *Journal of Coastal Research*, **19**, 584-599.
- Smith VH (2003) Eutrophication of freshwater and coastal marine ecosystems: A global problem. *Environmental Science and Pollution Research*, **10**, 126-139.

- Smith VH, Schindler DW (2009) Eutrophication science: where do we go from here? *Trends in Ecology & Evolution*, **24**, 201-207.
- Soulsby PG, Lowthion D, Houston M (1982) Effects of macroalgal mats on the ecology of intertidal mudflats. *Marine Pollution Bulletin*, **13**,162-166.
- Stapleton L, Lehane M, Toner P (2000) Ireland's environment: a millennium report. Environmental Protection Agency, Environmental Protection Agency, Wexford, Ireland.
- Stentiford GD, Feist SW (2005) A histopathological survey of shore crab (*Carcinus maenas*) and brown shrimp (*Crangon crangon*) from six estuaries in the United Kingdom Journal of Invertebrate Pathology, 88, 136-146.
- Stunkard HW (1957) The Morphology and Life-History of the Digenetic Trematode, *Microphallus similis* (Jägerskiöld, 1900) Baer, 1943. *Biological Bulletin*, **112**, 254-266.
- Taylor EW, Butler PJ, Sherlock PJ (1973) The respiratory and cardiovascular changes associated with the emersion response of *Carcinus maenas* (L.) during environmental hypoxia at three different temperatures. *Journal of Comparative Physiology*, **86**, 95-115.
- Thieltges DW, Fredensborg BL, Poulin R (2009) Geographical variation in metacercarial infection levels in marine invertebrate hosts: parasite species character versus local factors. *Marine Biology*, **156**, 983-990.
- Thieltges DW, Rick J (2006b) Effect of temperature on emergence, survival and infectivity of cercariae of the marine trematode *Renicola roscovita* (Digenea: Renicolidae). *Diseases of Aquatic Organisms*, **73**, 63-68.
- Thompson JV (1836) Natural history and metamorphosis of an anomalous crustaceous parasite of *Carcinus maenas*, the *Sacculina carcini*. *The Entomological Magazine*, **3**, 452-456.
- Torchin M, Lafferty K, Kuris A (2002) Parasites and marine invasions. *Parasitology*, **124**,137-151.
- Torchin ME, Lafferty KD, Kuris AM (2001) Release from parasites as natural enemies: Increased performance of a globally introduced marine crab. *Biological Invasions*, **3**, 333-345.
- Van Cleave, HJ, (1916). *Filicollis botulus* n. sp. with notes on the characteristics of the genus. *Transactions of the American Microscopical Society*, **35**, 131-134.

- Vidal-Martinez VM, Pech D, Sures B, Purucker ST, Poulin R (2010) Can parasites really reveal environmental impact? *Trends in Parasitology*, **26**, 44-51.
- Vismann B (1990) Sulfide detoxification and tolerance in Neries (Hediste) diversicolor and Nereis (Hediste) virens (Annelida: Polychaeta). Marine Ecology Progress Series, **59**, 229-238.
- Vitousek PM, Mooney HA, Lubchenco J, Melillo JM (1997) Human domination of Earth's ecosystems. *Science*, **277**, 494-499.
- Waser AM, Goedknegt MA, Dekker R, McSweeney N, Witte JIJ, van der Meer J, Thieltges DW (2016) Tidal elevation and parasitism: patterns of infection by the rhizocephalan parasite *Sacculina carcini* in shore crabs *Carcinus maenas*. *Marine Ecology Progress Series*, 545, 215-225.
- Waston G, Murray J, Schaefer M, Bonner A (2016) Bait worms: a valuable and important fishery with implications for fisheries and conservation management. *Fish and Fisheries*, **18**, 374-388.
- Welsh JE, van der Meer J, Brussaard CPD, Thielges DW (2014) Inventory of organisms interfering with transmission of a marine trematode. *Journal of the Marine Biological Association of the United Kingdom*, **10**, 1-6.
- Werner M (2001) Prevalence of the parasite *Sacculina carcini* Thompson 1836 (Crustacea, Rhizocephala) on its host crab *Carcinus maenas* (L.) on the west coast of Sweden. *Ophelia*, **55**, 101-110.
- World Meteorological Organization (1983) Guide to Meteorological Practice No. 168.
- Zander CD, Reimer LW (2002) Parasitism at the ecosystem level in the Baltic Sea. *Parasitology*, **124**, 119-135.
- Zetlmeisl C, Hermann J, Petney T, Glenner H, Griffiths C, Taraschewski H (2011) Parasites of the shore crab *Carcinus maenas* (L.): implications for reproductive potential and invasion success. *Parasitology*, **138**, 394-401.

CHAPTER 5: Parasites and pathologies of the blue mussel (*Mytilus* spp.): the role of host stress in susceptibility

ABSTRACT

The following study was undertaken to investigate the relationship between parasite communities of the blue mussel, Mytilus spp. and water quality in areas where the bivalve was found. A year-long survey assessed parasite communities of two subtidal *Mytilus* spp. populations inside and outside a marine reserve, Lough Hyne Co. Cork. It was hypothesised that Mytilus spp. outside the reserve would experience higher prevalence and intensity of parasite infection due to increased exposure to anthropogenic stressors. From the 731 Mytilus spp. screened nine parasite taxa were recorded: four macroparasites and five microparasites, including apicomplexans gregarines, ciliates, prokaryotic inclusions, Rickettsia-like organism /Chlamydia-like organisms, trematodes, nematodes, copepods and a cestode. Copepods and gregarines were found to be the most common macroparasite and microparasite, respectively. Data were analysed in relation to sex, gonadal development stage, condition index and the presence of parasites. Month, sex of Mytilus spp. and site sampled were not important determinants for the observed parasite prevalence or intensity. Prevalence of parasites did not correlate with pathologies recorded. Lipofuscin accumulation in tissues was the most common pathology recorded in *Mytilus* spp. and varied in relation to month of sampling. The low prevalence and intensity of trematode taxa were recorded in both *Mytilus* spp. populations in the field study was investigated with a series of laboratory trials assessing the susceptibility of *Mytilus* spp. to the trematode *Echinostephilla patellae.* To investigate the role of host stress in susceptibility, *Mytilus* spp. were sampled from habitats of differing levels of exposure to air, a bloom of Karenia mikimotoi, and to simulated nutrient enrichment. The findings of the laboratory trials were inconclusive, due to the high level of natural trematode infections of the 240 specimens included in the trials. There was a general, but insignificant trend, of increased E. patellae prevalence in Mytilus spp. exposed to cercariae. It was not possible to distinguish between pre-existing natural and laboratory infections of *E. patellae* due to the inability to age metacercariae based on morphology and current knowledge. Overall, the study suggests *Mytilus* spp. are not negatively affected by recorded parasite taxa and intensity of infection under current environmental conditions and water quality classifications.

5.1. INTRODUCTION

Mussels from the genus *Mytilus* are widely used in biomonitoring programs such as "Mussel watch" (Goldberg, 1975, Borja *et al.*, 2008, Kim *et al.*, 2008). Taxa used in biomonitoring programs are generally well studied, abundant, sedentary, have high tolerances to physiological stresses and wide global distributions. *Mytilus* are distributed from the subarctic to temperate regions, are sessile, resistant to pollution, euryhaline and can accumulate metals at 10^3 to 10^5 higher concentrations than the surrounding environment (Gosling, 2003). *Mytilus* spp. are filter feeders and can also concentrate pollutants, allow them to act as an early warning system of ecosystem decline or anthropogenic impacts (Gosling, 2003).

Mytilus spp. are ecosystem engineers in the rocky intertidal providing habitat through their shells or the dense aggregations that they form. Patches of *Mytilus* spp. can provide refuge and habitat for a wide variety of associated organisms in the rocky intertidal (e.g. Seed, 1996, O'Connor & Crowe 2007; 2008). *Mytilus* spp. are also hosts to a variety of different parasite taxa, including shell boring polychaetes, intestinal copepods, and trematodes (Lauckner, 1983). Work on Irish rocky shores and UK estuaries found trematodes to be common parasites of *Mytilus* spp. (Bignell *et al.*, 2008, Prinz *et al.*, 2010a, Lynch *et al.*, 2014). In the North Atlantic, blue mussels are commonly found to be second intermediate hosts of the trematode genera *Gymnophallus*, *Himasthla* and *Renicola* (Galaktionov*et al.*, 2015). Trematodes that utilise mussels as primary hosts are usually from the Bucephalid family; *Prosoryhnchus squamatus* (Odhner, 1905) and are found predominantly in Northwestern Europe, Britain and Atlantic Canada (Lauckner 1983, Coustau *et al.*, 1991).

Histopathology can assess the pathological, reproductive and metabolic condition of individual *Mytilus* spp. simultaneously. Tissue pathologies in *Mytilus* spp. have been

used in a wide range of studies to evaluate individual populations and their surrounding environment's health (Svärdh *et al.*, 2002, Aarab *et al.*, 2008, Bignell *et al.*, 2008). In response to increasing concentration of contaminants and heavy metals, increased levels of the age pigment (lipofuscin) and inflammatory lesions have been recorded in *Mytilus* spp. (Aarab *et al.*, 2008, Bignell *et al.*, 2011, Brenner *et al.*, 2014). Accumulation of lipofuscin granules, a type of insoluble cellular biomarker, can reflect the sublethal effects of pollutants and cell damage (Riveros *et al.*, 2002; Dimitriadis *et al.*, 2004). Histopathological surveys of *Mytilus* spp. in Ireland, the UK and USA have found the bivalve to commonly harbour trematode taxa (Bignell *et al.*, 2008, Bignell *et al.*, 2011, Lynch *et al.*, 2014, Powell *et al.*, 2015a; 2015b). Studies that consider the whole specimen with dissection or compression techniques have also found this to be the case in the Wadden, North Atlantic, Pacific and Arctic Seas (Thieltges *et al.*, 2006a, Prinz *et al.*, 2010a, Wilson *et al.*, 2013).

Parasite communities of *Mytilus* species have previously been used as indicators for environmental quality (Kim *et al.*, 2008; Sures, 2008, Brenner *et al.*, 2014). Parasite diversity and abundance in mussels have been linked with poor water quality in previous studies in Ireland, Sweden and the UK (Svärdh, 1999; Svärdh and Johannesson, 2002; Bignell *et al.*, 2008, Lynch *et al.*, 2014). Predicted environmental changes have been shown to affect the immuo-competence of *Mytilus* spp. which may affect their susceptibility to parasites (Brenner *et al.*, 2014, Mackenzie *et al.*, 2014a; 2014b). Pollution can decrease the abudnance of parasites depending on its effect on the host and the parasite taxa in question (Sures, 2008, Sures *et al.*, 2017). Also it should be noted that harbouring parasites is not necessarily equated to poor health of individuals or ecosystems (Hudson *et al.*, 2006).

The effect of eutrophication on the susceptibility of *Mytilus* spp. was also of interest. Eutrophication is expected to broadly facilitate trematode infection by increasing the abundance of gastropod hosts and the production of infective stages (Johnson *et al.*, 2010, Sures *et al.*, 2017). Finally, three intertidal populations of *Mytilus* spp. experiencing high, moderate and low exposure to a bloom were sampled from shores increasingly distant) from a bloom of *Karenia mikimotoi* (Miyake & Kominami ex Oda) Gert Hansen & Ø.Moestrup, 2000.

The first aim of this study were to examine if populations of *Mytilus* spp. could be characterised by their histopathology and parasite communities within and outside a marine reserve. Subtidal mussels were sampled as they may readily reflect water quality being constantly immersed and potentially continuously filtering. The survey was carried out over a year to account for seasonal variation in *Mytilus* spp. physiology and reproduction (Bignell *et al.*, 2008).

Laboratory trials were then used to investigate the role of host stress on *Mytilus* spp. and its susceptibility to the trematode *Echinostephilla patellae*. *Echinostephilla patellae* was chosen as it is an abundant trematode on the southwest coast of Ireland and has been experimentally established as a parasite of *Mytilus* spp. (Prinz *et al.,* 2010a; 2010b). The susceptibility of intertidal and subtidal populations of *Mytilus* spp. to *E. patellae* was of interest. The intertidal zone of rocky shores is physiologically stressful, as biota experience large changes in temperature, moisture and physical disturbance by waves. *Mytilus edulis* from or translocated to the intertidal zone have been recorded to have wider physiological tolerances than subtidal *Mytilus* spp. would show a higher occurrence of infection.

5.2. MATERIALS AND METHODS

5.2.1 Parasite Field Survey study sites

The study sites were three mooring lines on the southwest coast of Ireland (see Fig. 1). In Lough Hyne marine reserve, *Mytilus* spp. were sampled in the North Basin $(51^{\circ}30'06'0" \text{ N } 9^{\circ}18'02.5" \text{ W})$. Tranabo bay $(51^{\circ}29'52.7"\text{ N } 9^{\circ}17'07"\text{ W})$ is a small southern facing shore found to the east of the reserve. Cunnamore pier $(51^{\circ}30'21.1"\text{ N } 9^{\circ}25'22.3"\text{ W})$ is located within the small and sheltered Rincolisky Bay enclosed behind Heir Island and Sherkin Island. To have a replicate subtidal population of *Mytilus* spp. to sample, a buoy was deployed ca. 5km to the west of Lough Hyne. However, the buoy failed to recruit *Mytilus* spp.

5.2.1.2 Parasite Field Survey Sampling of *Mytilus* spp.

Mytilus edulis (Linnaeus, 1758) and *Mytilus galloprovincialis* (Lamarck, 1819) hybridise on shores along the southwest coast of Ireland (Gosling *et al.*, 1981; 2008). Distinguishing between the different *Mytilus* species, based on morphology, is problematic as no factor to date can consistently differentiate between the two species. Molecular screening is needed to differentiate between the two species and their hybrids based on the Me15/16 gene (Coughlan & Gosling, 2007). The gene encoding a polyphendic adhesive protein which is a key component in the attachment of mussels to substrates. These markers allow for the reliable estimation of the numbers of parental and hybrid mussel classes within hybrid zones (Inoue *et al.*, 1995). Thus, in this study, the mussels *M. edulis* and *M. galloprovincialis* are grouped into the genus *Mytilus* spp. Coustau *et al.*, (1990) suggested *M. edulis* species were more susceptible to the trematode *Prosorhynchus squamatus*, Odhner, 1905, relative to hybrids of *M. edulis* and *M. galloprovincialis*. However, more recent work has found no significant difference between the species and their hybrid in terms of susceptibility to parasite

infection (Bignell *et al.*, 2008). In Ireland, higher diversity and prevalence of parasites were recorded in *M. edulis* relative to *M. galloprovincialis* but no evidence of increased susceptibility of the species' hybrid was found (Lynch *et al.*, 2014).

Subtidal *Mytilus* spp. were sampled on a monthly basis from April 2013 until April 2014 in Lough Hyne and Tranabo bay. A once off sample was taken from a mooring line in Cunnamore pier, as after sampling the line was cleared of subtidal *Mytilus* spp. For all sampling events of subtidal *Mytilus* spp. specimens were sampled haphazardly by hand from mooring lines in the first 1-2 m below the waters' surface. *Mytilus* spp. <40 mm in shell length were not collected as smaller sizes are generally not parasitised (Bignell *et al.*, 2008; 2011). *Mytilus* spp. were then transported to the laboratory and processed for histology within 48 hours of collection.

5.2.1.3 Parasite Field Survey Methodology

The soft tissue of each mussel was dissected within 24 hours of collection from the fresh individuals. Prior to their dissection, the shell length, width and weight of individual *Mytilus* spp. were measured to the nearest 0.05 mm with vernier callipers. Adductor muscles were severed and soft tissue was removed from the shell. For each *Mytilus* spp. sampled, a section of tissue containing gill, gut, gonad and foot tissue, approximately 1 cm, in width was excised. The tissue section was then fixed in Davidson's solution at 4 °C for 48h as outlined by Shaw and Battle (1957). After 48 h in 70% ethanol, tissue was dehydrated through an ascending ethanol series in a Shadon Tissue Processor Citadell 1000. Tissue was embedded in paraffin wax, cut in 5-7 μ m slices with a Leica RM2235 microtome. Tissue sections were placed on clean standard microscope slides and dried overnight in an oven. Slides were then stained using haematoxylin & eosin in a Leica ST5010 Autostainer XL. Slides were mounted in DPX (Distrene, Plasticiser, Xylene). After a week examined under x 200 or 400 magnifications with transmitted light to ensure the mounting medium was set.

Parasitological terminology and microscopes follows the specifications of Chapter 2 Section 2.2.3 page 43.

5.2.1.4 Parasite Field Survey analysis

Histopathological analysis was carried out according to the protocol outlined by Kim *et al.*, (2006). Parasites such as ciliates, prokaryotic inclusion bodies (PIBS), nematodes, gregarines, cestodes and trematode metacercariae were counted per tissue type. PIBS were counted when a dark staining intracellular inclusion 5-10 μ m in size was seen within a tissue. Pathologies listed in Kim *et al.*, (2006) were qualitatively recorded per tissue type examined. Each cross section of a nematode or a copepod was counted as a single individual. The sex and gonadal stage of specimens were noted, based on scales modified from Seed (1969). Four stages of gonadal maturity were identified: resting, developing, ripe and spawning.

5.2.2 Laboratory trial studies

Three laboratory trials were carried out to investigate the role of host stress on *Mytilus* spp. and its susceptibility to *Echinostephilla patellae* cercariae, collected from its intermediate host *Patella vulgata* Linnaeus, 1758 (see Fig. 2 for experimental designs). Mortality was considered when the shell was gaping and the mantle was unresponsive to mechanical stimulation. Attempts were made to recover cercariae that failed to infect *Mytilus* spp. from tanks. 48 hours after exposure to cercariae whole mounts of *Mytilus* spp. were examined for parasite taxa. Prior to examination shell length, width and weight of individual *Mytilus* spp. were measured to the nearest 0.05 mm with Vernier callipers. Whole tissue mounts were compressed between ¹/₄ inch thick glass slides and examined with microscopes detailed in Chapter 2, Section 2.2.3, page 43. The foot of *Mytilus* spp. was examined separately due to the differing density

of the tissue and recorded preference of *E. patellae* to infect the tissue (Prinz *et al.,* 2009). All parasites encountered were recorded in tissue specific counts.

Trial 1 (T1) investigated the role of differing levels of physiological stress experienced by *Mytilus* spp. in the intertidal and subtidal zone. It was hypothesised that the abiotic conditions of the intertidal zone may be physiologically demanding relative to the subtidal, thus intertidal *Mytilus* spp. would be more susceptible to trematode infection. Trial 2 (T2) investigated three intertidal populations of *Mytilus* spp. experiencing high, moderate and low exposure to a bloom of *Karenia mikimotoi*. Concentrations of *K. mikimotoi* reached 729,000 cells -¹ at the height of the bloom and were roughly 12,000 cells L⁻¹ at the time of sampling. Previous blooms of *K. mikimotoi* in Ireland and Scotland resulted in mass mortality (>90%) of invertebrate populations through a combination of hypoxia and toxicity (Raine *et al.*, 2001, Davidson *et al.*, 2009). *Karenia mikimotoi* produces haemolytic toxins that supress *Mytilus* spp. immune function and influence bivalve growth and mortality (Matsuyama *et al.*, 1999, Hégaret *et al.*, 2011). Thus, *Mytilus* spp. populations that experienced greater exposure due to their higher proximity to the bloom *K. mikimotoi* were hypothesised to be more susceptible to *E. patellae* (Matsuyama *et al.*, 1999).

Trial 3 (T3) investigated the effect of acute host exposure to enriched nutrient conditions on host susceptibility. *Mytilus* spp. in higher nutrient concentrations were hypothesised to be more susceptible to *E. patellae*, due to conditions favouring cercariae survival and infectivity (Zander and Reimer, 2002, Johnson *et al.*, 2010, Johnson & Carpenter, 2008). *Mytilus* spp. were sampled from an intertidal population on the east coast previously sampled and found to be free of *E. patellae* infection by Prinz *et al.*, (2010b).

5.2.2.1 Laboratory trial collection sites

For all trials, *Patella vulgata* were sampled from the intertidal zone of Bullen's Bay, a sheltered rocky shore on the southwest of Ireland ($51^{\circ}38'37.0"N 8^{\circ}33'19.0"W$). For T2 *Mytilus* spp. were collected from the intertidal zone of three rocky shores, the moderately wave exposed Ringaskiddy ($51^{\circ}49'54.4"N$, $8^{\circ}18'04.0"W$), the sheltered shore Cusheen ($51^{\circ}31'43"N$, $9^{\circ}31'34"W$), and the North Shore in Lough Hyne ($51^{\circ}30'21"N$, $9^{\circ}18'12"W$). The distance of the site from highest recorded concentration of *K. mikimotoi* cells was a proxy for decreasing exposure, in order of proximity Cusheen (ca. 15km), Lough Hyne (ca. 30km) and Ringaskiddy (ca. 100km). For T3 *Mytilus* spp. were collected from the moderately exposed rocky shore Blackrock ($53^{\circ}18'00"N 6^{\circ}10'03"W$) Co. Dublin on the east coast of Ireland.

5.2.2.2 Laboratory trial study collection of mussels

Mytilus spp. >4cm in length were sampled haphazardly by hand at low tide from all sites. In November 2015 intertidal and subtidal *Mytilus* spp. were collected in Lough Hyne from the intertidal zone and from mooring lines in the first 1-2m below the waters' surface as described previously in Section 5.2.1. In September 2015 intertidal *Mytilus* spp. were sampled from Cusheen, Lough Hyne and Ringaskiddy for T2. In October 2015 intertidal *Mytilus* spp. were sampled at low tide for T3. 100 intertidal *P. vulgata* were sampled haphazardly by hand at low tide in Bullen's Bay for on the first day of the trial for each run of laboratory trials.



5.2.2.2 Laboratory trial study experimental design

Fig. 2 Schematic diagram of experimental design of the *Mytilus* spp. challenge experiments (a) Trail 1: Host habitat and susceptibility to trematode infection (b) Trial 2: Red tide susceptibility trial (c) Trial 3: Acute exposure nutrient enrichment and host susceptibility trial. Three experimental runs of each trial were carried out.

5.2.2.3 Laboratory trial study experimental protocols

T 1 Host habitat and susceptibility to trematode infection trial

Prior to the commencement of the trial epibionts were removed from *Mytilus* spp. However, *Mytilus* spp. were not cleaned and over the duration of the trial the water in the tanks water was not changed. Intertidal and subtidal *Mytilus* spp. were placed in two plastic aerated tanks (30 X 20 X 20 cm) filled with UV-filtered seawater. Tanks were in were kept within a climate change chamber at 15 °C with 12:12 light: dark photoperiod. *Mytilus* spp. were randomly and individually placed in glass tanks (Fig. 3(c)) with 300 ml of UV-filtered seawater 24 hours prior to exposure to *E. patellae* cercariae collected from *P. vulgata* (Fig. 3(b)). *Mytilus* spp. in assigned tanks had acclimatised for 24 h prior to exposure to cercariae. Each of the three treatments was replicated 4 times per experimental run (Fig. 2a). Water from replicate tanks were filtered with a 20 μ m sieve.

T 2: Red tide susceptibility trial

Prior to the commencement of the trial the epibionts were removed from *Mytilus* spp. However, *Mytilus* spp. were not cleaned and over the duration of the trial and the water in tanks was not changed. *Mytilus* spp. from the three sites sampled were placed in three separate plastic aerated tanks (30 X 20 X 20 cm) filled with UV-filtered seawater and maintained in the same environmental conditions the previous trial. Individual *Mytilus* spp. from Cusheen, Lough Hyne and Ringaskiddy were randomly and individually placed in glass tanks with 300 ml of UV-filtered seawater 24 h prior to exposure to *E. patellae* cercariae collected from *P. vulgata*. Each of the three treatments was replicated 4 times per experimental run. Water from replicate tanks were filtered with a 20 µm sieve.

T 3 Acute exposure nutrient enrichment and host susceptibility trial

Prior to the commencement of the trial the *Mytilus* spp. epibionts were removed from Mytilus spp. However, were not cleaned and over the duration of the trial the in the tanks water was not changed. Mytilus spp. sampled from Blackrock were randomly assigned to different treatments of nutrient concentrations of either 'Low' (7 µM phosphate, 140 µM nitrate and 2.6 µM nitrite), 'Medium' (15 µM phosphate, 278 µM nitrate and 5.14 µM nitrite) or 'High' (30 µM phosphate, 556 µM nitrate and 10.28 µM nitrite) (Fig. 2(c)). Concentrations were chosen based on the work of Malham et al., (2009), varying amouny of Phosphate (PO_4^{3-}), Nitrate (NO_3^{1-}), and Nitrite (NO_2^{-}), were added to high, medium and low treatments see Table 2. In each plastic tank (30 X 20 X 20 cm) 25 Mytilus spp. were kept in a 7.5L of UV treated seawater with added nutrients or without in the case of control tanks. Mytilus spp. were maintained in plastic tanks at the same environmental conditions as T1 and T2 for a week and fed with 9.41ml of Reed Shellfish Diet 1800[®] (approx. 2 billion cells ml⁻¹) twice in that period. Mytilus spp. were then randomly and individually placed in glass tanks with 300ml of UV-filtered seawater 24 h prior to exposure to E. patellae cercariae collected from *P. vulgata*. Water from replicate tanks were filtered with a 20µm sieve.

Table 1. Nutrient levels proposed based on Malham *et al.*, (2009). All values are given as mgL-1

Chemical parameter	High	Medium	Low	Total per run
Sodium Phosphate monobasic dihydrate	2.85	1.42	0.66	4.40
(NaH ₂ PO ₄)				
Sodium Nitrate (NaNO ₃)	517.12	17.24	8.68	543.04
Sodium Nitrite (NaNO ₂)	0.47	0.24	0.12	0.83

Echinostephilla patellae cercariae collection from infected Patella vulgata

Echinostephilla patellae cercariae were obtained from infected *Patella vulgata*, based on work by Prinz *et al.*, (2009; 2010b). Prior to storage in dry darkness for an hour at ca. -4°C, epibionts were removed from the limpets' shells to avoid physical disturbance or impairment of *E. patellae* cercariae (Prinz *et al.*, 2011). *Patella vulgata* were then immersed in 30ml of UV-filtered seawater in plastic petri dishes under bright light (Fig. 3 (a)). *Echinostephilla patellae* cercariae were shed from *P. vulgata* utilised as primary intermediate hosts. One *P. vulgata* utilised as a primary



Fig. 3. Experimental set up of obtaining cercariae (a) Primary intermediate host *Patella vulgata* "sheds" *Echinostephilla patellae* cercariae in 30ml of UV filtered seaweater. (b) *Echinostephilla patellae* cercariae (c) *Mytilus* spp. in aerated glass tanks of 300ml UV-filtered seawater to which cercariae were pippetted. Scale bars indicates (a) 2.25cm and (b) 200µm (c) 9cm.
intermediate host by *E. patellae* was selected from the 100 screened to collect standard doses of cercariae from, either 10 or 20 individuals. *Echinostephilla patellae* cercariae were then pipetted into assigned glass tanks of 300ml UV filtered water (see Fig 1).

5.2.3 Statistical Analysis

Analysis was carried out in R studio version 1.0.136 (R Core Team 2017). Statistical significance was considered with P<0.05. The field study was 2 factor analysis with the location (fixed, orthogonal, 2 levels) and month sampled (fixed, orthogonal, 12 levels) considered factors. The laboratory study was a series of two factor analyses. In T1 the factors were habitat (fixed, orthogonal, 2 levels), and *E. patellae* exposure (fixed, orthogonal, 2 levels) (See Fig. 2 (a)). In T2 location (random, orthogonal, 3 levels) and *E. patellae* exposure (fixed, orthogonal, 2 levels) were the factors considered (See Fig. 2 (b)). In T3 the nutrient level (fixed, orthogonal, 4 levels) and *E. patellae* exposure (fixed, orthogonal, 2 levels) were the factors considered (See Fig. 2 (b)).

5.2.3.1 Parasite data analysis

Differences in parasite prevalences were tested with a Pearson's chi square with Yates continuity correction. If expected values were too low (<5) for a Pearson's chi square, a Fisher's Exact test was used. A Kruskal-Wallis test was carried out to examine differences in parasite intensities between sites, seasons, and sex. Pairwise Wilcoxon Rank Sum Tests was performed if significant differences were found. Parasitology terms follows Bush *et al.*, (1997).

5.2.3.2 Host data analysis

Host size characteristics between populations were tested with an Analysis of variance (one-way ANOVA) in combination with the posthoc test Tukey's honestly significant difference (HSD) test. Prior to the ANOVA being carried out, assumptions of homogeneity of variance and normally distributed data were assessed visually and then formally by the Shapiro–Wilk test. The condition factor based on Fulton's condition index as follows: CF =wet weight/ (shell length)³ × 100. Images were taken with previously cited microscopes and software in Chapter 2, Section 2.2.3, page 43.

5.3. RESULTS

5.3.2.1 Morphometrics of subtidal mussels in field survey

Morphometrics of *Mytilus* spp., sex ratio and sample sizes of subtidal populations sampled are summarised in Table 1. Subtidal *Mytilus* spp., sampled from the North Buoy in Lough Hyne were larger than those sampled in Tranabo (F=8.37, df=1, P<0.0001). No clear seasonal pattern in spawning was detected in either population (Fig. 4).

Table 2. Morphometrics of *Mytilus* spp. populations sampled in Lough Hyne North Buoy and Tranabo Buoy in the form \bar{x} (±SD). CI represents condition index. F:M:U represents the ratio of *Mytilus* spp. with female, male or undifferentiated gonads within the sample screened.

Location	Average Length (mm)	Condition Index	F:M:U	n
Cuanmore pier	61.10 (±11.85)	0.01 (±0.02)	0.37:0.60:0.03	30
North Buoy Lough Hyne	54.35 (±11.79)	0.04 (±0.36)	0.35:0.37:0.27	365
Tranabo	51.10 (±9.33)	0.04 (±0.42)	0.47:0.38:0.15	363



Fig. 4. The percentage of *Mytilus* spp. at each stage of gametogenesis from samples taken from (a) the North Buoy in Lough Hyne and (b) Tranabo bay

5.3.2.2 Histological screening for parasite taxa in field survey

Nine parasite taxa were recorded during the histological study, comprising of four macroparasite and five microparasite taxa. The apicomplexans gregarines were the most common parasite taxa (Table 3). Apicomplexans gregarines were followed in abundance by ciliates, prokaryotic inclusions (PIB), *Rickettsia*-like

organism/*Chlamydia*-like organism (RLO/CLO), trematodes, nematodes, coccidian and a cestode (Table 3).

Table 3. Overall parasite prevalence per site sampled and range (min-max) of intensity in brackets for subtidal populations of *Mytilus* spp. *Rickettsia*-like organism/*Chlamydia*-like organism and prokaryotic inclusion body are represented by RLO/CLO and PIB, respectively.

Parasite taxa	North Buoy	Tranabo	Cuanmore
	Lough Hyne		pier
MICROPARASITE			
Ciliate	8.52 (1-121)	18.73 (1-31)	16.70 (1-2)
Coccidian	1.00 (1)	8.82 (1)	0.00
Gregarine	13.28 (1-124)	31.40 (1-85)	0.00
PIB	10.78 (1-44)	14.33 (1-78)	6.70 (1)
RLO/CLO	9.52 (1-48)	13.50 (1-14)	10.00 (1-4)
MACROPARASITE			
Cestode	0.25 (1)	0.00	0.00
Copepod	4.01 (1-4)	12.5 (1-3)	3.30(1)
Nematode	7.02(1-2)	9.37 (1-2)	0.00
Trematode	9.52 (1-11)	11.85 (1-7)	3.30 (1)

The prevalence of parasite taxa fluctuated over the sampling period (see Fig. 5). *Nematopsis* sp., Schneider, 1892, oocysts (Fig. 6e) were the predominant gregarine infection, the most common parasites at all sites and months sampled. *Nematopsis* sp. were the only parasite taxon to significantly differ between with higher prevalence observed in Tranabo (χ^2 =15.10, df=1, P<0.001). Ciliates were next most abundant taxon and were commonly found on gill filaments. Trematode metacercariae were more common than sporocyst infections. Infection usually occurred within digestive tubules. RLO/CLOs ranged from 1-48 in *Mytilus* spp. and were found predominately in the gills. Higher intensity of RLO/CLOs were seen in males (χ^2 =5.1, df=2, P=0.05). Copepods were mainly found in the digestive tract and to a lesser extent in the gills. It

is likely that *Mytilicola intestinalis*, Steuer, 1902, was the predominant copepod found in the digestive tract. This is due to the location and size of sections of copepod, however characteristics of *M. intestinalis* used to distinguish from the invasive *Mytilicola orientalis* Mori, 1935 such as were not apparent in a cross section. No difference was detected between sexes of *Mytilus* spp. and infection by parasite taxa.



Fig. 5. Parasite prevalence for subtidal populations of *Mytilus* spp. sampled from a buoy in (a) Lough Hyne and (b) Tranabo bay from April 2013 until April 2014.

Coinfection, differing parasite taxa within a host, commonly occurred in screened *Mytilus* spp. Overall, coinfection was recorded in 21% of all *Mytilus* spp. and was significantly more common in *Mytilus* spp. populations in Tranabo bay (27%) than Lough Hyne (15%) (χ^2 =15.48, df=1, P<0.0001). Coinfection was predominately between ciliates and other microparasite taxa for both locations (Table 4). Of the significant correlations between parasite taxa, predominantly negative correlations were found between prokaryotes and helminths.

Table 4. Overall correlations of taxa with other parasite taxa recorded in subtidal populations of *Mytilus* spp. sampled. *Rickettsia*-like organism/*Chlamydia*-like organism and prokaryotic inclusion body are represented by RLO/CLO and PIB, respectively. Asterisk denotes statistical significance: * for P<0.05, ** for <0.001 and *** for <0.0001. ¹ represents helminth taxa

Parasite	Co-occurring Parasite	Correlation coefficient	Overall prevalence of coinfection
Ciliate	Cestode ¹	0.00	0.26
	Copepod	-0.01	0.66
	Gregarine	0.07***	3.95
	Nematode ¹	0.00	0.39
	PIB	0.01	2.50
	Trematode ¹	0.00	2.63
	RLO/CLO	0.01	3.16
Copepod	Cestode ¹	0.06	0.13
	Gregarine	0.40	1.97
	Nematode ¹	0.11***	0.26
	Trematode ¹	0.01	0.39
	RLO/CLO	0.02	0.92
	PIB	-0.01	0.00
Gregarine	Cestode ¹	-0.01	0.13
	Nematode ¹	-0.01	0.39
	Trematode ¹	-0.20***	1.57
	PIB	-0.14	1.18
	RLO/CLO	0.07	1.97
Nematode1	Trematode ¹	0.01	0.39
PIB	Trematode ¹	-0.33***	0.52



Fig. 6 Photomicrographs of parasites recorded in subtidal populations of *Mytilus* spp. sampled in Lough Hyne and Tranabo from April 2013 until April 2014. (a) *Rickettsia*-like organism/ *Chlamydia*-like organism on the gills (RLO/CLO), (b) Ciliate in gill tissue (c) Copepod in digestive tract, (d) Trematode metacercariae in digestive tubule (e) *Nematopsis* oocyst in the connective tissue, (f) Prokaryotic inclusion body (PIB) in gill tissue, (g) Granulocytoma in foot tissue & (h) Lipofuscin accumulation in mantle. (a)-(f) scale bars represent 20µm, 200µm for (g) and 100µm for (h). Haematoxylin and eosin stain.



Fig. 7. Photomicrographs of haplosporidian-like particles recorded in (a) connective tissue, (b) gill tissue, (c) kidney tissue and (d) mantle tissues of subtidal *Mytilus* spp. sampled in Lough Hyne and Tranabo from April 2013 until April 2014. Scale bars represent 200μ m for (a) and 20μ m for (b), (c) and (d). Haematoxylin and eosin stain.

5.3.2.3 Observed pathological conditions in mussels in field survey

Four pathologies were recorded during the study; lipofuscin accumulation, inflammation, granulocytoma and pearl formation. Lipofuscin accumulation was the most common pathology of study and was recorded in just under half of all *Mytilus* spp. examined (Table 5) (Fig. 6(h)). The occurrence of the remaining pathologies did not differ significantly between sites or months sampled.

Table 5. Overall pathology occurrence and correlation with parasite taxa recorded in subtidal populations of *Mytilus* spp. sampled. *Rickettsia*-like organism/*Chlamydia*-like organism is represented by RLO/CLO. Asterisk denotes statistical significance: * for P<0.05, ** for <0.001 and *** for <0.0001.

Pathology	Prevalence	Parasite taxa	Correlation coefficient
Granulocytoma	4.72	Ciliate	-0.52***
		RLO/CLO	-0.45***
		Trematode	-0.50***
		Copepod	-0.29***
		Gregarine	-0.24***
		Lipofuscin	0.11**
Inflammation	11.55	Copepod	-0.31***
		RLO/CLO	-0.66***
		Trematode	-0.24***
		Ciliate	-0.13
Lipofuscin	47.77	Copepod	0.35***
		Trematode	-0.26***
		RLO/CLO	-0.13**
		Ciliate	-0.07*
Pearl formation	0.79		

5.3.3.1 Morphometrics of mussels in laboratory trials

Overall, 350 *Mytilus* spp. were screened during the study, 230 of which were included in the trials T1-3. In all three trials, no mortality was observed.

Table 6. Morphometric data of *Mytilus* spp. included in the trials carried out in October 2015. CI represents Fulton's Condition Index. Data are presented in the form of \bar{x} (±SD)

Trial	n	Shell length (mm)	Whole Weight (g)	CI
1	54	42.30 (±2.89)	12.4 (±2.38)	0.01 (±<0.01)
2	48	34.06 (±8.12)	6.27 (±3.73)	0.01 (±<0.01)
3	128	41.86 (±2.17)	11.78 (±2.30)	0.03 (±0.32)
Total	230	40.48 (±5.40)	10.91 (±3.60)	0.02 (±0.24)

T 1 Intertidal and subtidal Mytilus spp. trial

Mytilus spp. in the laboratory trial were significantly larger in the intertidal than the subtidal habitat (F=12.32, df=1, P<0.001). *Mytilus* spp. morphometrics did not significantly vary within treatment groups between experimental runs.

T 2 Harmful algal bloom

Two *Mytilus* spp. spawned in both the second and third experimental run of T2. The individuals were still included in the analysis as their mantle retracted or shell closed in response to mechanical stimulation. *Mytilus* spp. morphometrics did not vary within treatment groups, between experimental runs or between treatment groups.

T 3 Acute exposure nutrient enrichment and host susceptibility trial

Mytilus spp. morphometrics did not significantly vary within treatment groups, between experimental runs or between treatment groups.

Parasite taxa detected in mussels screened prior in laboratory trials

Macroparasite taxa from Phylum Platyhelminthes and Copepoda were recorded in *Mytilus* spp. Trematode taxa were the most abundance parasite taxa recorded. In addition to natural infections of *E. patellae*, four other trematode taxa; *Renicola* spp., *Microphallus* spp., *Himasthla* spp, and *Proctoeces maculates* (Looss, 1901); were recorded in *Mytilus* spp. included in T1-3 (see Table 7). *Renicola* spp. was the most common taxa harboured by *Mytilus* spp. for all laboratory trials. The copepod *M. intestinalis* and the turbellarian *Urastoma cyprinae* (Graff, 1882) were also recorded

in *Mytilus* spp. (Table 7 & 8). Negative controls of each trial, demonstrate the differences in natural infections between *Mytilus* spp. populations sampled (Table 7 & 8).

Chapter 5

Table 7 Prevalence of trematode taxa in populations of *Mytilus* spp. included in the three trials run in October 2015. Data are presented in the form of prevalence (min-max). – or + indicate whether *Mytilus* spp. was exposed to *Echinostephilla patellae* cercariae. For Trial 1, I represent intertidal and S subtidal *Mytilus* spp. collected in Lough Hyne. For trial 2 Cu represents Cusheen, LH Lough Hyne, R Ringaskiddy. For trial 3 C represents the control, L low nutrient enrichment, M medium enrichment, and H high enrichment treatments.

Trial	Treatment	Ν	Echinostephilla patellae	Renicola spp.	Microphallus spp.	Himasthla spp.	Proctoeces
							maculatus
1	I-	12	91.67 (1-198)	75.00 (1-142)	0.00	0.00	0.00
1	I+	12	75.00 (2-128)	33.33 (1-3)	0.00	8.33 (1-3)	0.00
1	S-	12	16.67 (7-25)	0.00	0.00	8.33 (52)	0.00
1	S+	12	33.33 (1-6)	0.00	0.00	0.00	0.00
2	Cu-	9	33.33 (1-3)	77.78 (3-80)	0.00	88.89 (2-26)	0.00
2	Cu+	9	22.22 (1)	66.67 (3-40)	0.00	77.78 (2-20)	11.11
2	LH-	9	22.22 (1)	55.56 (1-2)	0.00	44.44 (4-38)	0.00
2	LH+	9	55.56 (1-14)	44.44 (1-3)	0.00	1.00 (1-7)	0.00
2	R-	9	22.22 (1)	66.67 (1-2)	0.00	11.11 (4)	0.00
2	R+	9	33.33 (1)	0.00	0.00	33.33 (2-18)	0.00
3	C-	17	5.88 (1)	100 (2-129)	5.88 (1)	0.00	0.00
3	C+	16	6.25 (1)	93.75 (2-231)	0.00	0.00	0.00
3	L-	14	14.29 (1-3)	92.86 (1-94)	0.00	0.00	0.00
3	L+	17	5.88 (3)	100 (3-22)	0.00	11.76 (4-38)	0.00
3	M-	16	12.50 (5)	100 (2-105)	0.00	0.00	0.00
3	M+	16	12.50 (1)	100 (4-177)	6.25 (1)	6.25 (1-7)	0.00
3	H-	15	6.67 (1)	86.67 (3-70)	0.00	13.33 (2-4)	0.00
3	H+	17	11.76 (1-5)	100 (6-208)	0.00	5.88 (1)	0.00
Grand To	tal	230	23.10 (1-198)	73.91 (1-231)	0.87 (1)	17.39 (1-52)	0.43

Table 8 Prevalence of macroparasites other than trematodes and pearl formation from populations of *Mytilus* spp. included in the three trials run in October 2015. – or + indicate whether *Mytilus* spp. was exposed to *Echinostephilla patellae* cercariae. For Trial 1, I represent intertidal and S subtidal *Mytilus* spp. collected in Lough Hyne. For Trial 2, Cu represents Cusheen, LH Lough Hyne, R Ringaskiddy. For Trial 3, *Mytilus* spp. collected from Blackrock were assigned to the control; C, low nutrient enrichment; L, medium enrichment; M, or high enrichment treatments; H.

Trial	Treatment	n	Pearl	Urastoma	Mytilicola
			formation	cyprinae	intestinalis
1	I-	12	0.00	0.00	8.33
1	I+	12	0.00	0.00	33.33
1	S-	12	0.00	0.00	0.00
1	S+	12	0.00	0.00	0.00
2	Cu-	9	0.00	0.00	33.33
2	Cu+	9	0.00	0.00	0.00
2	LH-	9	0.00	0.00	0.00
2	LH+	9	0.00	0.00	0.00
2	R-	9	0.00	0.00	0.00
2	R+	9	0.00	0.00	0.00
3	C-	17	5.88	0.00	0.00
3	C+	16	6.25	0.00	0.00
3	L-	14	21.43	0.00	0.00
3	L+	17	23.53	0.00	0.00
3	M-	16	25.00	0.00	0.00
3	M+	16	18.75	0.00	0.00
3	H-	15	6.67	6.67	0.00
3	H+	17	23.53	0.00	0.00
Grand To	otal	230	9.13	0.43	2.17

For T1 intertidal (n=10) and subtidal (n=10) *Mytilus* spp. were screened prior to laboratory trials. Three parasite taxa were observed, *Himasthla* spp., *Renicola* spp. and *M. intestinalis*. Prevalences of all parasite taxa were higher in *Mytilus* spp. from the intertidal of Lough Hyne; 100% of screened *Mytilus* spp. harboured *Himasthla* spp., 60% harboured *Renicola* spp. and 10% contained *M. intestinalis*. In the subtidal *Mytilus* spp. screened, *Renicola* spp. was the only taxa and a prevalence at 10% was observed. For T2, intertidal *Mytilus* spp. from Cusheen, Lough Hyne and Ringaskiddy were screened prior to the laboratory trial. Prevalences of *Renicola* spp. in *Mytilus* spp. from Ringaskiddy. Higher prevalence of *M. intestinalis* was observed in *Mytilus* spp. from Ringaskiddy (50%) relative to Cusheen (25%). The copepod was absent from *Mytilus* spp. screened from Lough Hyne. In T3, *E. patellae* (12%), *Renicola* spp. (34%), the *Proctoeces maculatus* (Looss, 1901) (1%), copepods (2%) and pearls (18%) were observed in *Mytilus* spp. (n=100) sampled from Blackrock.



Fig. 8 Photomicrographs of helminth taxa recorded during the *Mytilus* spp. trials carried out in October 2015 (a) *Echniostephilla patellae* metacercariae in the bivalve *Mytilus* spp. (b) *Proctoeces maculatus* cercariae (c) Turbellarian *Urastoma cyprinae*. Scale bars indicate 100µm.

T 1 Intertidal and subtidal Mytilus spp. trial

Prevalence and intensity of *E. patellae* did not significantly differ between treatment groups. *Echinostephilla patellae* was significantly more prevalent in intertidal *Mytilus* spp. relative to subtidal *Mytilus* spp. controls ($\chi^2 = 10.74$, df=1, P<0.00) (Table 7). Prevalence of *Renicola* spp. and the copepod *M. intestinalis* were higher in intertidal *Mytilus* spp. than subtidal specimens (Table 7). Negative controls of intertidal and subtidal *Mytilus* spp. had similar prevalences and intensities of trematode taxa and copepods (Table 7).

<u>T 2 Harmful algal bloom</u>

Prevalence and intensity of *E. patellae* did not significantly differ between treatment groups. Prevalence *Renicola* spp. ($\chi^2 = 7.65$, df=2, P<0.05) and *Himasthla* spp. ($\chi^2=8.25$, df=2, P<0.05) significantly differed between controls. Primary intermediate infection by *Proctoeces maculatus* was observed in a negative control of *Mytilus* spp. from Cusheen. Higher prevalences of both trematode taxa were observed in *Mytilus* spp. sampled from Cusheen (Table 7).

T 3 Acute exposure nutrient enrichment and host susceptibility trial

Echinostephilla patellae prevalence and intensity did not differ between controls or within treatments of differing experimental runs. Prevalence and intensity of *E. patellae* also did not significantly differ between different treatment groups. The turbellarian *Urastoma cyprinae* was recorded in one specimen in the negative control of the high treatment. *Renicola* spp. and the pathology of pearl formation were more common in *Mytilus* spp. sampled from Blackrock (Table 7 & 8). *Renicola* spp.

prevalence and intensity were high in all enrichment treatments and did not statistically differ. Differences in the occurrence of pearl in treatments were not statistically significant, despite seeming more common in *Mytilus* spp. exposed to moderately enriched conditions (Table 8).

Overall, coinfection of parasite taxa in *Mytilus* spp. included in all trials was low (1.42%) (Fig. 9). In T3 coinfection was the highest of all the three trials (Fig. 9). Recorded coinfections of macroparasites were predominantly in the form of trematodes with copepods. Coinfections of trematode taxa were in the form of *E*.



Fig. 9 (a) The prevalence of *Mytilus* spp. with no macroparasite taxon (0), harbouring a single parasite taxon (1) or two parasite taxa (2). (b) The prevalence of *Mytilus* spp. harbouring no trematode taxon (0), one trematode taxon (1) or multiple trematode taxa (2, 3). Exposure to *Echinostephilla patellae* cercariae is represented with "+" and controls, to which no cercariae were added, are represented with "-".

patellae, *Himasthla* spp. and *Renicola* spp. Coinfections between *E. patellae* and other trematode metacercariae were positively correlated for T2 and T3. The intensity of *Himasthla* spp. and *E. patellae* in the foot of *Mytilus* spp. were negatively correlated, however it was statistically insignificant (r=-0.24, df=14, P>0.05).

5.4. DISCUSSION

Both the field and laboratory trials recorded a range of largely non-pathogenic parasite taxa in *Mytilus* spp. In the field study, parasite communities or pathologies did not differ between subtidal populations in sites of differing water quality except for a higher occurrence of gregarines in Tranabo bay and a single cestode observed in the North Buoy population. The mooring lines sampled in Lough Hyne and Tranabo are connected by water at high tide. and so subtidal populations of *Mytilus* spp. sampled may have experienced similar exposure to contaminants and parasite taxa. The study would need additional replicates of subtidal *Mytilus* spp. populations to investigate the potential effect of surrounding water quality and parasite communities. An alternative study to address the hypothesis would be a transplant experiment of samples from a common subtidal *Mytilus* spp. population for further investigation into the effect of water quality on parasite communities.

Parasite communities of *Mytilus* spp., screened in the field study were dominated by apicomplexan gregarines, followed by ciliates, prokaryotic inclusions (PIB), *Rickettsia*-like organism/*Chlamydia*-like organism (RLO/CLO), trematodes, nematodes, a coccidian and a cestode. In relation to seasonality of parasites this study agrees with work in UK estuaries (Bignell *et al.*, 2008). Except for the pathology of lipofuscin accumulation, no trends in prevalence of parasites or intensity in relation to month or season were found. Similar patterns have been reported in *M. galloprovincialis* with peaks of lipofuscin accumulation recorded in summer month (Koukouzika *et al.*, 2009). Parasite composition differed slightly from a previous health survey of cultured and wild *Mytilus* spp. (Lynch *et al.*, 2014). In the present study, copepods were the dominant macroparasites followed by trematodes while Lynch *et al.*, (2014) found the reverse. Parasite communities' composition agree with that in the Black Sea and Netherlands (Lauckner 1983).

The field study, recorded lower parasite prevalence and higher intensity extent of infections and pathologies than previous work on *Mytilus* spp. in Irish marine waters (Lynch et al., 2014, Giltrap et al., 2016). This observation could be due to subtidal habitat and immunological vigour of *Mytilus* spp. screened. The fact that *Mytilus* spp. in the field study were sampled from the subtidal zone could explain the low macroparasite prevalences recorded (Buck et al., 2005, Brenner et al., 2014). Although, intertidal molluscs are key hosts for digenean trematodes and other macroparasite taxa, increased distance between potential hosts and current hosts lowers the likelihood of a macroparasite encountering and infecting a potential host (Galaktionov et al., 2003, Buck et al., 2005, Brenner et al., 2014). This is likely the case for trematodes typical intermediate hosts may be restricted to nearshore habitats e.g. *Littorina* spp. abundant in mid- and low intertidal zones but is rare in shallow subtidal habitats (Reid 1996, Carlson et al., 2006). Thus, the further the populations of *Mytilus* spp. are from the substrate or intertidal zone, the lower their encounter rate with preceding trematode hosts and subsequent infection rates would be. In the German Bight, offshore mussels were not infected by trematodes or shell-boring polychaetes(Buck et al., 2005, Brenner et al., 2014). Low trematode prevalence in this study was recorded in subtidal *Mytilus* spp. sampled for laboratory trials the same location which were sampled simultaneously. Higher prevalence and intensity of trematodes were recorded in intertidal *Mytilus* spp. in Lough Hyne intertidal relative to subtidal Mytilus spp. sampled at the same time. Mytilus edulis haemocytes have been found to be more active in phagocytosis and superoxide generation relative to sympatric bivalves the edible cockle Cerastoderma edule and the razor-shell Ensis siliqua (Linnaeus, 1758) (Wootton et al., 2003).

By definition a parasite is an organism that exerts some harm on the host (Thieltges *et al.*, 2013). The extent of harm varies between taxa and life stages and a well-adapted parasite can be intermediately pathogenic or non-pathogenic (Combes, 1997). The pathology of parasite taxa recorded in the field study, trematodes and the turbellarian

U. cyprinae, is dependent on the parasitic life stage and intensity of infection (Lauckner, 1983, Thieltges, 2006, Stier & Thieltges 2015). Trematode sporocyst infections can be highly pathogenic as heavy infections can result in parasitic castration, weakness in valve closure, reduced byssal thread activity and may affect growth (Matthews 1973, Lauckner ,1983, Coustau *et al.*, 1991; 1993, Thieltges 2006, Stier & Thieltges, 2015). The low prevalence of trematode sporocysts in the subtidal specimens in Lough Hyne and of *Proctoeces maculatus* in the intertidal specimens on the east coast suggest that trematode primary infection is not a major threat to the health of either population sampled.

The histological study found no association between local inflammation or pathology and recorded parasite taxa. Gregarines in the form of *Nematopsis* sp. oocysts were the most prevalent parasites for all of the months and both sites sampled. Gregarine infection has been recorded to cause focal and benign inflammation in *Mytilus* spp. (Bower *et al.*, 1994). Ciliates were another highly prevalent microparasite. Ciliates are commensals but can become pathogenic in high intensities or if the host is stressed (Lauckner 1983). The copepod *M. intestinalis* was the most prevalent macroparasite but is also considered to be a commensal (Francisco *et al.*, 2010). The pathology of the copepod is low and generally restricted to mechanical damage of tissue around the site of its hook-like appendages (Moore *et al.*, 1978). A 10-year study in the UK failed to establish *M. intestinalis* as a determining or major contributing factor in bivalve mortality (Davey & Gee 1976). RLO/CLOs have been recorded in a wide range of bivalves and are usually non-pathogenic (Lauckner 1983).

Tissue compression techniques, utilised in the laboratory trials, recorded a smaller range of parasite taxa then histological techniques. In this study, no difference in the prevalence or intensity of the trematode *E. patellae* was recorded during the trials. Although, not statistically significant there seemed to be a general trend of increased *E. patellae* prevalence in *Mytilus* spp. exposed to cercariae. Coinfection of trematode

taxa was common in laboratory trials T2 and T3. The high level of natural trematode infection in the foot tissue of *Mytilus* spp. could have impaired *E. patellae* infection through competitive displacement. *Echinostephilla patellae* cercariae may have been prevented from encysting in the foot by metacercariae as the area was already occupied by *Renicola* spp. and *Himasthla* spp. Pearls were recorded predominantly in the Blackrock population. The formation of such structures in bivalves have been associated with parasitic trematodes, copepods, cestodes, nematopsis oocysts and inorganic irritants (Garner 1872, Sindermann & Rosenfield 1967 Dakin, 1913). In particular previous work on the south coast of Ireland have documented the formation of pearls around metacercariae of *Gymnophallus* spp. in the bivalves *Mytilus* spp. and *Cerastoderma edule* (Fermer *et al.*, 2011, Morgan 2013). However relative to other studies the prevalence and intensities of pearls in *Mytilus* spp. was lower (94%) have been recorded with up to 264 recorded in a single *Mytilus edulis* (Fernandes & Seed 1983, Ambariyanto & Seed 1991).

When the life cycle of *E. patellae* was initially investigated by Kollien (1996), *Mytilus* spp. was not considered in the list of secondary hosts, the gastropod *P. vulgata* was investigated but was exposed to unnaturally high doses of *E. patellae* cercariae. Prinz *et al.*, (2010b) exposed *Mytilus* spp. to varying doses of *E. patellae* and the trematode successfully infected the bivalve's foot. The low doses of 10 cercariae per mussel could be a contributing factor to the failure to detect a statistically significant difference in *E. patellae* prevalence and intensity between *Mytilus* spp. in the trials. In previous work the doses were successful and are likely more biologically realistic (Poulin 2010, Prinz*et al.*, 2010b).

Biomass of *Mytilus* spp. was unlikely a determining factor for the differences in parasite prevalences observed in the laboratory trials. The only difference in size between *Mytilus* spp. specimens was recorded in T1, where larger specimens were sampled from intertidal relative to the subtidal zone of Lough Hyne. This difference

is likely due to sampling methodology, as while intertidal and subtidal differ in size due to their differing growth rates, typically larger specimens are recorded in the subtidal relative to the intertidal zone (Seed, 1969, Page & Hubbard 1987). Subtidal *Mytilus* spp. can grow continuously while intertidal *Mytilus* spp. undergo periods of exposure to air and suppressed feeding activity (Labarta *et al.*, 1997, Tagliarolo *et al.*, 2012). While the size of *Mytilus* spp. is not necessarily related to their age, larger *Mytilus* spp. are likely to have a higher encounter rate with and infection by parasites (Buschbaum & Saier, 2001, Nikolaev *et al.*, 2006). It is acknowledged that there was a difference in holding time of *Mytilus* spp. between runs of laboratory trials. *Mytilus* spp. included in the second and third experimental run of the laboratory trials were held for 48 and 60 hours, respectively, more that individuals in the first run. This may have had an effect, but one which was undetected in parameters measured as there was no significant difference in *Mytilus* spp. mortality, size, spawning or parasite infection between experimental runs.

This study cannot conclusively report the susceptibility of *Mytilus* spp. to *E. patellae* cercariae exposed to environmental stressors simulated or otherwise. The interpretation of results was confounded by natural infection of *Mytilus* spp. Future work, could include either farmed or subtidal *Mytilus* spp. in laboratory trials along with work to investigate the feasibility of estimating the age of *E. patellae* metacercariae according to its morphology. This would allow for the differentiation between pre-existing natural *E. patellae* and experimental infections. The effect of environmental stressor eutrophication on host: parasite dynamics requires further work. The susceptibility of *Mytilus* spp. to parasite taxa in eutrophic conditions is particularly necessary as suspended mussel aquaculture has been proposed as one possible solution to mitigate coastal water quality problems associated with excess nutrients (Edebo *et al.*, 2000, Gren *et al.*, 2009, Lindahl *et al.*, 2005). There is a need to evaluate the risk that parasites taxa recorded in both studies, may pose to *Mytilus* spp. under climate change.

REFERENCES

- Aarab N, Pampanin DM, Naevdal A, Oysaed KB, Gastaldi L, Bechmann RK (2008) Histopathology alterations and histochemistry measurements in mussel, *Mytilus edulis* collected offshore from an aluminium smelter industry (Norway). *Marine Pollution Bulletin*, 57, 569-574.
- Altieri AH (2006) Inducible variation in hypoxia tolerance across the intertidalsubtidal distribution of the blue mussel *Mytilus edulis*. *Marine Ecology Progress Series*, **325**, 295-300.
- Bignell JP, Dodge MJ, Feist SW, Lyons B, Martin PD, Taylor NGH, Stone D, Travalent L, Stentiford GD (2008) Mussel histopathology, and effects of season, disease and species. *Aquatic Biology*, **2**, 1-15.
- Bignell JP, Stentiford GD, Taylor NGH, Lyons BP (2011) Histopathology of mussels (sp.) from the Tamar estuary, UK. *Marine Environmental Research*, **72**, 25-32.
- Borja A, Bricker SB, Dauer DM, Demetriades NT, Ferreira JG, Forbes AT, Hutchings P, Jia X, Kenchington R, Carlos Marques J, Zhu C (2008) Overview of integrative tools and methods in assessing ecological integrity in estuarine and coastal systems worldwide. *Marine Pollution Bulletin*, 56, 1519-1537.
- Bower SM, McGladdery SE, Price IM (1994) Synopsis of infectious diseases and parasites of commercially exploited shellfish. *Annual Review of Fish Diseases*, 4, 1-199.
- Brenner M, Broeg K, Frickenhaus S, Buck BH, Koehler A (2014) Multi-biomarker approach using the blue mussel (*Mytilus edulis* L.) to assess the quality of marine environments, Season and habitat-related impacts. *Marine Environmental Research*, **95**, 13-27.
- Buck BH, Thieltges DW, Walter U, Rosenthal H (2005) Inshore–offshore comparison of parasite infestation in *Mytilus edulis*, implications for open ocean aquaculture. *Journal of Applied Ichthyology*, **21**, 107-113.
- Buschbaum C, Saier B (2001) Growth of the mussel *Mytilus edulis* L. in the Wadden Sea affected by tidal emergence and barnacle epibionts. *Journal of Sea Research*, **45**, 27-36.

- Callaway R, Shinn AP, Grenfell SE, Bron JE, Burnell G, Cook EJ, Crumlish M, Culloty S, Davidson K, Ellis RP, Flynn KJ, Fox C, Green DM, Hays GC, Hughes AD, Johnston E, Lowe CD, Lupatsch I, Malham S, Mendzil AF, Nickell T, Pickerell T, Rowley AF, Stanley MS, Tocher DR, Turnbull JF, Webb G, Wootton E, Shields RJ (2012) Review of climate change impacts on marine aquaculture in the UK and Ireland. *Aquatic Conservation-Marine and Freshwater Ecosystems*, **22**, 389-421.
- Carlson RL, Shulman MJ, Ellis JC (2006) Factors contributing to spatial heterogeneity in the abundance of the common periwinkle *Littorina littorea* (L.) *The Journal of Molluscan Studies*, **72**, 149-156.
- Combes C (1997) Fitness of parasites: Pathology and selection. *International Journal for Parasitology*, **27**,1-10.
- Coughlan B, Gosling E (2007) Genetic structure of Hybrid mussel populations in the west of Ireland: two hypotheses revisited, *Marine Biology*, **150**, 841-852.
- Coustau C, Renaud F, Maillard C, Pasteu N, Delay, B (1991) Differential susceptibility to a trematode parasite among genotypes of the *Mytilus* edulis/galloprovincialis complex. Genetical Research, **57**, 207-212.
- Dakin WJ (1913) Cambridge Manuals of Science and Literature: Pearls. *Cambridge University Press. Cambridge*, **4**, 1-143.
- Daugbjerg N, Hansen G, Larsen J, Moestrup O (2000) Phylogeny of some of the major genera of dinoflagellates based on ultrastructure and partial LSU rDNA sequence data, including the erection of three new genera of unarmoured dinoflagellates. *Phycologia*, **39**, 302-317.
- Davey JT, Gee JM (1976) The occurrence of *Mytilicola intestinalis* Steuer, an intestinal copepod parasite of *Mytilus*, in the south-west of England. *Journal of the Marine Biological Association of the United Kingdom*, **56**, 85-94.
- Davidson K, Miller P, Wilding TA, Shutler J, Bresnan E, Kennington K, Swan S (2009) A large and prolonged bloom of *Karenia mikimotoi* in Scottish waters in 2006. *Harmful Algae*, 8, 349-361.
- Dimitriadis VK, Domouhsidou GP, Cajaraville MP, (2004). Cytochemical and histochemical aspects of the digestive gland cells of the mussel *Mytilus galloprovincialis* (L.) in relation to function. *Journal of Molecular Histology* 35, 501-509.

- Dubois R (1901) Sur la mécanisme de la formation des perles fines dans le *Mytilus* edulis. Les Comptes rendus hebdomadaires des séances de l'Académie des sciences (Paris), **133**, 603-605.
- Dubois R (1903) L'origine des perles chez le Mytilus galloprovincialis. Les Comptes rendus hebdomadaires des séances de l'Académie des sciences (Paris), **136**, 178-179.
- Dubois R (1907) Sur les métamorphoses du Distome parasite des *Mytilus perliers*. Les *Comptes rendus hebdomadaires des séances de l'Académie des sciences* (Paris), **63**, 334-336.
- Dubois R (1909) Contribution a l'étude des perles fines de la nacre et des animaux qui les produisent. *Annales de l'Université de Lyon*, **29**, 1-126.
- Edebo L, Haamer J, Lindahl O, Loo LO, Piriz L (2000) Recycling of macronutrients from sea to land using mussel cultivation. *International Journal of Environment and Pollution*, **13**, 190–207.
- Francisco CJ, Hermida MA, Santos MJ (2010) Parasites and symbionts from *Mytilus* galloprovincialis (Lamark, 1819) (Bivalves, Mytilidae) of the Aveiro estuary Portugal. Journal of Parasitology, 96, 200-205
- Galaktionov KV, Bustnes JO, Bardsen BJ, Wilson JG, Nikolaev KE, Sukhotin AA, Skirnisson K, Saville DH, Ivanov MV, Regel KV (2015) Factors influencing the distribution of trematode larvae in blue mussels *Mytilus edulis* in the North Atlantic and Arctic Oceans. *Marine Biology*, **162**, 193-206.
- Galaktionov KV, Dobrovolskij AA (2003) The trematode life cycle as a system of adaptations. *The Biology and Evolution of Trematodes*. Springer, 215-317.
- Garner R (1872) Experimental Observations on the Appearance of the Lower Forms of Life. *The Lancet*, **8**, 789-791.
- Giltrap M, Ronan J, Tanner C, O'Beirn FX, Lyons BP, Aoidh RM, Rochford H, McHugh B, McGovern E, Wilson J (2016) Application of a weight of evidence approach utilising biological effects, histopathology and contaminant levels to assess the health and pollution status of Irish blue mussels (*Mytilus edulis*). *Marine Environmental Research*, **122**, 33-45.
- Gmelin JF (1789) Caroli a Linné, systema naturae. 1-2, 501-1032.
- Goldberg ED (1975) The Mussel Watch. A First Step in Global Monitoring. *Marine Pollution Bulletin*, **6**, 111-113.

- Gosling E (2003) Bivalve molluscs: biology, ecology and culture. *Oxford: Fishing new books*.
- Gosling E, Wilkins NP (1981) Ecological genetics of the mussels *Mytilus edulis* and *M. galloprovincialis* on Irish coasts. *Marine Ecology Progress Series*, **4**, 221-227.
- Gosling E, Doherty S, Howley N (2008) Genetic characterization of hybrid mussel (*Mytilus*) populations on Irish coasts *Journal of the Marine Biological* Association of the United Kingdom, **88**, 341–34.G
- Graff L (1882). Monographie der Turbellarien I. Rhabdocoelida. Verlag Wilhelm Engelmann, Leipzig, 2, 1-442.
- Gren I, Lindahl O, Lindqvist M (2009) Values of mussel farming for combating eutrophication. *Ecological Engineering*, **35**, 935–945.
- Hégaret H, Da Silva PM, Wikfors GH, Haberkorn H, Shumway SE, Soudant P (2011) In vitro interactions between several species of harmful algae and haemocytes of bivalve molluscs. *Cell Biology and Toxicology*, **27**, 249-266.
- Hudson PJ, Dobson AP, Lafferty KD (2006) Is a healthy ecosystem one that is rich in parasites? *Trends in Ecology & Evolution*, **21**, 381-385.
- mein adhesive protein sequences of *Mytilus edulis*, *M. galloprovincialis* and *M. trossulus*. *Biological Bulletin*, **189**, 370–375.
- Johnson P, Carpenter S (2008) Influence of Eutrophication on Disease in Aquatic Ecosystem: Patterns, Processes, and Prediction. In: *Infectious Disease Ecology* (ed. by Ostfield R, Kessing F, Eviner V). Princeton University Press, Princeton University Press.
- Johnson P, Townsend AR, Cleveland CC, Glibert PM, Howarth RW, McKenzie VJ, Rejmankova E, Ward MH (2010) Linking environmental nutrient enrichment and disease emergence in humans and wildlife. *Ecological Applications*, **20**, 16-29.
- Khan RA, Thulin J (1991) Influence of pollution on Parasites of Aquatic Animals. *Advances in Parasitology*, **30**, 201-238.
- Kim Y, Powell E, Ashton-Alcox K (2006) Histopathology analysis. Histological techniques for marine bivalve molluscs, Update NOAA Technical Memorandum NOS NCCOS 27, 19-52.

- Kim Y, Powell EN, Wade TL, Presley BJ (2008) Relationship of parasites and pathologies to contaminant body burden in sentinel bivalves, NOAA Status and Trends 'Mussel Watch' Program *Marine Pollution Bulletin*, 65, 101–127.
- Kollien A (1996) Cercaria patellae Lebour, 1911 developing in Patella vulgata is the cercaria of Echinostephilla patellae (Lebour, 1911) n. comb. (Digenea, Philophthalmidae). Systematic Parasitology, **34**, 11-25.
- Koukouzika N, Raftopoulou EK, Dimitriadis VK (2009) Seasonal differences of lysosomal, lipid and lipofuscin parameters in the digestive gland of the mussel *Mytilus galloprovincialis. Journal of Molluscan Studies*, **75**, 261-267.
- Labarta U, FernandezReiriz MJ, Babarro JMF (1997) Differences in physiological energetics between intertidal and raft cultivated mussels *Mytilus galloprovincialis*. *Marine Ecology Progress Series*, **152**, 167-173.
- Lamarck JBM (1819) Histoire naturelle des animaux sans vertèbres. Tome sixième, 1re partie. Paris, Verdière, 1815-22.
- Lauckner G (1983) Diseases of Marine Animals, Introduction, Bivalvia to Scaphopoda. Chapter 13. Kinne, O (Ed) *Diseases of Marine Animals*, 2, 477-897.
- Liddell C, Welsh J, van der Meer J, Thieltges D (2017). The effect of dose and frequency of exposure to infectious stages on trematode infection levels in mussels. *Diseases of Aquatic Organisms*, **125**, 85-92.
- Lindahl O, Hart R, Hernroth B, Kollberg S, Loo LO, Olrog L, Rehnstam-Holm AS, Svensson J, Svensson S, Syversen U (2005) Improving marine water quality by mussel farming: A profitable solution for Swedish society. *Ambio*, **34**, 131-138.
- Linnaeus C (1758) Tomus I. Systema naturae per regna tria naturae :secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis10. Holmiae, *Laurentii Salvii*, **1-4**, 1-824.
- Lynch S, Morgan E, Carlsson J, Mackenzie C, Wooton E, Rowley A, Malham S, Culloty S (2014) The health status of mussels, *Mytilus* spp., in Ireland and Wales with the molecular identification of a previously undescribed haplosporidian. *Journal of Invertebrate Pathology*, **118**, 59-65.
- Mackenzie CL, Lynch S, Culloty SC, Malham SK (2014) Future Oceanic Warming and Acidification Alter Immune Response and Disease Status in a Commercial Shellfish Species, *Mytilus edulis* L. *PLoS One*, **9**, 1-12.

- Malham SK, Cotter E, O'Keeffe S, Lynch S, Culloty SC, King JW, Latchford JW, Beaumont AR (2009) Summer mortality of the Pacific oyster, *Crassostrea* gigas, in the Irish Sea: The influence of temperature and nutrients on health and survival. Aquaculture, 287, 128-138.
- Marigómez I, Garmendia L, Soto M, Orbea A, Izagirre U, Cajaraville MP (2013) Marine ecosystem health status assessment through integrative biomarker indices, a comparative study after the Prestige oil spill "Mussel Watch". *Ecotoxicology*, 22, 486-505.
- Marigómez I, Zorita I, Izagirre U, Ortiz-Zarragoitia M, Navarro P, Etxebarria N, Orbea A, Soto M, Cajaraville MP (2013) Combined use of native and caged mussels to assess biological effects of pollution through the integrative biomarker approach. *Aquatic Toxicology*, **136–137**, 32-48.
- Matsuyama Y, Uchida T, Honjo T (1999) Effects of harmful dinoflagellates, *Gymnodinium mikimotoi* and *Heterocapsa circularisquama*, red-tide on filtering rate of bivalve molluscs. *Fisheries Science*, **65**, 248-253.
- Matthews RA (1973) The life-cycle of *Prosorhynchus crucibulum* (Rudolphi, 1819) Odhner, 1905, and a comparison of its cercaria with that of *Prosorhynchus squamatus* Odhner, 1905. *Parasitology*, **66**, 133-164.
- Moore MN, Icarus Allen J, McVeigh A (2006) Environmental prognostics, An integrated model supporting lysosomal stress responses as predictive biomarkers of animal health status. *Marine Environmental Research*, **61**, 278-304.
- Moore MN, Lowe DM, Gee JM (1978) Histopathological effects induced in *Mytilus edulis* by *Mytilicola intestinalis* and the histochemistry of the copepod intestinal cells. *Journal Du Conseil*, **38**, 6-11.
- Morgan E (2013) Synecology in soft sediment bivalves: the influence of arasites, physiological processes, and environmental stressors on health and disease. Doctor of Philosophy Thesis, University College Cork, Cork.
- Mori T (1935) *Mytilicola orientalis*, a new species of parasitic Copepoda. *Zoological Magazine (Dobutsugaku Zasshi)*, **47**, 687-690.
- Nikolaev KE, Sukhotin AA, Galaktionov KV, (2006). Infection patterns in White Sea blue mussels *Mytilus edulis* of different age and size with metacercariae of *Himasthla elongata* (Echinostomatidae) and *Cercaria parvicaudata* (Renicolidae). *Diseases of Aquatic Organisms*, **71**, 51-58.

- O'Connor NE, Crowe TP (2007) Biodiversity among mussels: separating the influence of sizes of mussels from the ages of patches. *Journal of the Marine Biological Association of the United Kingdom*, **87**, 551-557.
- O'Connor NE, Crowe TP (2008) Do mussel patches provide a refuge for algae from grazing gastropods? *Journal of Molluscan Studies*, **74**, 75-78.
- Odhner T (1905). Die Trematoden des arktischen Gebietes. *Fauna Arctica*, **4**, 289-372.
- Page HM, Hubbard DM (1987) Temporal and spatial patterns of growth in mussels *Mytilus edulis* on an offshore platform: relationships to water temperature and food availability. *Journal of Experimental Marine Biology and Ecology*, **111**, 159-179.
- Powell EN, Kim Y (2015a) The Regional Spatial Structure of Parasites and Pathologies in Oysters and Mussels in the United States: 16 Years of Mussel Watch. *Journal of Shellfish Research* **34**, 939-965.
- Powell EN, Kim Y, Bushek D (2015b) Temporal Structure and Trends of Parasites and Pathologies in U.S. Oysters and Mussels: 16 Years of Mussel Watch. *Journal of Shellfish Research* 34, 967-993.
- Prinz K, Kelly T, O'Riordan R, Culloty S (2011) Factors influencing cercarial emergence and settlement in the digenean trematode *Parorchis acanthus* (Philophthalmidae). *Journal of the Marine Biological Association of the United Kingdom*, **91**, 1673-1679.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2009) Non-host organisms affect transmission processes in two common trematode parasites of rocky shores. *Marine Biology*, **156**, 2303-2311.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010a) Occurrence of macroparasites in four common intertidal molluscs on the south coast of Ireland. *Marine Biodiversity Records*, **3**, 89-96.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010b) Infection of *Mytilus edulis* by the trematode *Echinostephilla patellae* (Digenea: Philophthalmidae). *Journal of Helminthology*, **84**, 193-198.
- R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <u>http://www.R-project.org/</u>

- Raine R, S OB, O'Higgins T, White M, Patching J, Cahill B, McMahon T (2001) A satellite and field portrait of a *Karenia mikimotoi* bloom off the south coast of Ireland, August 1998. *Coastal Shellfish-A Sustainable Resource Developments in Hydrobiology*, **160**, 187-193.
- Reid DG (1996) Systematics and evolution of *Littorina*. *The Dorset Press, Dorchester, Dorset, United Kingdom.* 1-463.
- Riveros A, Zúñiga M, Hernandez A, Camaño A, (2002). Cellular biomarkers in native and transplanted populations of the mussel *Perumytilus purpuratus* in the intertidal zones of San Jorge Bay, Antofagasta, Chile. *Archives of Environmental Contamination and Toxicology*, **42**, 303-312.
- Schneider A (1892) Signalement d'un nouvean sporozoaire. *Tablettes Zoologiques*, **2**, 209–210.
- Seed R (1969). The ecology of *Mytilus edulis* L. (Lamellibranchiata) on exposed rocky shores. I. Breeding and settlement. *Oecologia*, (Berlin) **3**, 277-316.
- Seed R (1974). Morphological variations in *Mytilus* from the Irish coasts in relation to the occurrence and distribution of *Mytilus galloprovincialis*, *Cahiers de Biologie Marine*, **15**, 1-25.
- Seed R (1976). Marine Mussels: their ecology and physiology, Cambridge University Press, New York. 13-65.
- Seed R (1996). Patterns of biodiversity in the macro-invertebrate fauna associated with mussel patches on rocky shores. *Journal of the Marine Biological Association of the United Kingdom*, **76**, 203–210.
- Shaw BL, Battle HI (1957) The gross and microscopic anatomy of the digestive tract of the oyster *Crassostrea virginica* (Gmelin, 1791). *Canadian Journal of Zoology*, **35**, 325-347.
- Sindermann CJ (1990) Diseases of Marine Shellfish. Academic Press, Limited, London.
- Sindermann CJ and Rosenfield A (1967) Principal diseases of commercially important marine bivalve mollusca and crustacea. *Fishery Bulletin*, **66**, 335–385.
- Sindermann CJ, Farrin AE (1962) Ecological studies of *Cryptocotyle lingua* (Trematoda: Heterophyidae) whose larvae cause" pigment spots" of marine fish. *Ecology*, **43**, 69-75.

- Steuer A (1902) Mittheilungen aus der K.K. zoologischen Station in Triest. No. 4. Beobachtungen über das Plankton des Triester Golfes im Jahre 1901. *Zoologischer Anzeiger*, **25**, 369-372.
- Stier T Drent J, Thieltges DW (2015) Trematode infections reduce clearance rates and condition in blue mussels *Mytilus edulis*. *Marine Ecology Progress Series*, 529, 137-144.
- Sures B (2008) Host-parasite interactions in polluted environments. *Journal of Fish Biology*, **73**, 2133-2142.
- Sures B, Nachev M, Selbach C, Marcogliese DJ (2017) Parasite responses to pollution: what we know and where we go in 'Environmental Parasitology'. *Parasites & Vectors*, **10**, 1-19.
- Svärdh L (2003) Tissue sampling from live blue mussels, *Mytilus edulis*. A field study from the Swedish west coast. *Journal of Sea Research*, **49**, 221–225.
- Svärdh L, Johanesson K (2002) Incidence of hemocytes and parasites in coastal populations of blue mussels (*Mytilus edulis*) - testing correlations with area, season, and distance to industrial plants. *Journal of Invertebrate Pathology*, 80, 22-28.
- Tagliarolo M, Clavier J, Chauvaud L, Koken M, Grall J (2012) Metabolism in blue mussel: intertidal and subtidal beds compared. *Aquatic Biology*, **17**, 167-180.
- Thieltges DW (2006) Effect of infection by the metacercarial trematode *Renicola* roscovita on growth in intertidal blue mussel *Mytilus edulis*. Marine Ecology Progress Series, **319**, 129-134.
- Wootton EC, Dyrynda EA, Ratcliffe NA (2003) Bivalve immunity, comparisons between the marine mussel (*Mytilus edulis*), the edible cockle (*Cerastoderma edule*) and the razor-shell (*Ensis siliqua*). Fish & Shellfish Immunology, **15**, 195-210.
- Zander CD, Reimer LW (2002) Parasitism at the ecosystem level in the Baltic Sea. *Parasitology*, **124**, 119-135.

CHAPTER 6: General Discussion

To assess an impacted ecosystem, knowledge of pathological conditions and parasites is essential, as it provides valuable insight into the health of host populations and the state of the ecosystem. Data on dominant parasites in multiple host taxa can inform on how parasites and hosts respond to current anthropogenic impacts and aid in the creation of accurate predictions of e.g. a changing climate, increased anthropogenic impact on a specific site in marine waters.

The current study was undertaken to gain knowledge on how the distribution of parasite communities of wild invertebrate taxa on the southwest coast of Ireland varied in response to ecosystem quality. The focus on larval trematodes developed as the taxa are abundant in the intertidal zone and have been proposed, by multiple reviews, as indicators of ecosystem health (Lafferty 1997, Mouritsen & Poulin 2002, Huspeni *et al.*, 2005, Marcogliese 2005, Vidal- *et al.*, 2010, Prinz *et al.*, 2010, Sures *et al.*, 2017).

The observed variation of parasite and trematode communities in relation to ecosystem quality appeared dependent on site specific local factors, host taxa screened, parasite taxa and the definition of quality considered. Ecosystem quality was defined by policy directive classifications in Chapters 2 & 4, and by statutory protection in Chapters 3 & 5. In Chapter 2, metacercariae of the dominant trematodes *Echinostephilla patellae* and *Renicola* spp. were also more prevalent in the site of higher WFD status quality. Trematode primary infection has been suggested as an indicator for water quality (Huspeni *et al.*, 2005). Work in Northern Ireland found that *Cryptocotyle lingua* cercariae in *L. littorea* were negatively affected by poor water quality and had impaired swimming and potentially transmission (Cross *et al.*, 2001; 2003; 2005). In Chapter 2 the prevalence of primary infections in *Littorina littorea* did not differ

between sites but the number of functional groups was higher in the site of good water quality. In Chapter 4, trematode prevalence, intensity and community richness were greater in the site of low water quality. In Chapter 3, parasite communities of Patella *vulgata* were similar within and outside the marine reserve. The heterogeneous distribution of the dominant trematode, E. patellae, within and outside of the marine reserve was proposed to be determined by local site-specific factors, such as wave exposure and local currents rather than the level of protection. In Chapter 5, gregarines were the only parasite taxon harboured by subtidal *Mytilus* spp. that significantly differed between populations sampled as significantly higher prevalences recorded outside of the marine reserve. In Chapter 5, low prevalence of trematode taxa was recorded in both subtidal *Mytilus* spp. populations was likely to be due to their distance from avian vertebrate and intermediate invertebrate hosts. The lower prevalence of parasite taxa in subtidal Mytilus spp. relative to intertidal aquaculture stocks has already been documented (Buck et al., 2005, Brenner et al., 2012). Subtidal Mytilus spp. may not be very susceptible to recorded parasites under current environmental conditions. The effect abiotic environmetnal stressor in the form of nutrient enrichment was inconclusive for the *Mytilus* spp. trials due to the high levels of natural infections.

Data generated by this study contribute novel findings on the factors that influence the distribution and abundance of parasites within the intertidal environment of sites sampled. Chapters 2 and 3 addressed neglected aspects of parasite ecology in relation to marine conservation, EPA quality classifications and the effect of surrounding intertidal biota. At present, few conservation studies consider parasites (Nichols *et al.*, 2011, Gómez *et al.*, 2013). The literature review of Chapter 3 returned 12 studies on the topic of parasites in MPAs. Parasites may be facilitated by conservation and "jump" between hosts (Kermack & McKendrick, 1927, Lebarbenchon *et al.*, 2006).

Chapters 4 & 5 addressed the impact of eutrophication and habitat type on the host susceptibility. Findings from Chapters 2 and 4 suggest that trematode communities, observed, varied in relation to environmental stress. The variation was not consistent between environmental stressors considered or parasite taxa recorded. The magnitude of the effect of environmental stressors on a trematode taxon may vary within one between lifecycle stages and geographic regions (Marcogliese, 2016). Regional variation between has been documented for the trematode *Renicola* spp. utilising *L. littorea*, as a primary intermediate host. In the Barents Sea the optimum temperature range for the release of cercariae from *L. littorea* inhibited the release of cercariae of the same trematode and host taxa in the Wadden Sea (Thieltges *et al.*, 2006, Prokofiev *et al.*, 2016).

Previous parasitological work has illustrated the ubiquity of parasites in intertidal molluscs on the south coast of Ireland and the dominance of trematodes within the parasite communities (Prinz et al., 2010, Fermer et al., 2011). The study highlights the need to adopt a more a nuanced approach to host-parasite ecology. It is frequently reported that environmental stressors can impact water quality, affect the immunological response in aquatic biota and make them more susceptible to parasitic infection and increasing parasite prevalence (Khan and Thulin, 1991). The environmental stressor may also negatively affect the transmission of parasite taxa particularly if the taxa requires multiple hosts to complete its lifecycle (Pietrock & Marcogliese 2003). The response of parasites to environmental stressors is known to be varied and complex (Lafferty 1997, Blanar et al., 2009, Vidal-Martinez et al., 2010). Data from the study agree this the general findings of previous work and suggest that the distribution, prevalence and intensity of trematode taxa varied in response to environmental stressors in the form of nutrient enrichment. The effect of local biotic factors are likely context depenant as the significantly was not consistent between sites or host species sampled. In Chapter 2 the prevalence of Echinostephilla *patllae* positively correlated with the density of its host *Patella vulgata* in the site of good water quality but not in the marine reserve.

This study attempted to provide robust baselines of trematode abundance and distributions in host species screened and sites sampled. The main technique of the study, tissue compression, provided a cost effective and conservative estimate of parasite abundance in invertebrates screened. In the detection of trematode taxa, tissue compression has been demonstrated to be equally effective or even outperform resource-heavy molecular techniques such as quantitative polymerase chain reaction (qPCR) (Markowitz et al., 2016) or fluorescent tagging, respectively (LaFonte et al., 2015). Tissue compression still provides a conservative estimate of trematode abundance as it is not able to detect early primary trematode infections (miracidia), and externally encysting metacercariae. Tissue compression was thus adopted to allow for a more cost effective and accurate estimate of trematode identity and abundance. Parasite identification was based on morphology, so cryptic species, which occur particularly in trematodes relative to other helminths, were grouped in a species complex (Leung et al., 2009, Pérez-Ponce de León et al., 2017). Larval trematodes infect specific host tissues preferentially e.g. palps or abductor muscles (Thieltges 2006, de Montaudouin et al., 2009, Prinz et al., 2010) and may be missed in the 1cm sampled from the of host for histological screening. Trematode taxa are difficult to accurately identify from histological slides without further molecular means.

There are several avenues future work could investigate from the current study. Work could consider the factors that influence the distribution and abundance of parasites within an environment and consider additional hosts other than molluscs when investigating the effect of environmental stressors on trematode prevalences and intensities (Thieltges &Wegner, 2016, Poulin *et al.*, 2016). The potential for protected areas and varying levels of eutrophication to facilitate trematode infection also
warrants further work. It is acknowledged that Lough Hyne may not be representative of the majority of MPAs. Unlike other MPAs, Lough Hyne is an extremely sheltered area and is well established as it was designated 36 years ago. The potential of MPAs to encourage parasite abundance within their boundaries should be evaluated on a larger spatial scale.

The potential of eutrophication to facilitate trematode abundance, should be investigated with manipulative laboratory trials conducted in tandem with field assessments of cercarial abundance in the water column. Manipulative field experiments in which the abundance of trematode is altered, e.g. exclusion from certain areas, are often logistically challenging (Sousa 1999, Kaplan et al., 2009). Field surveys of trematode cercariae are also difficult, as cercariae are soft bodied, short lived and typically missed with traditional plankton survey techniques (Fingerut et al., 2003, Zimmer et al., 2009). However, estimations of cercariae in the water column have been done by screening specific hosts in the field and enumerating the amount of cercariae "shed" during timed intervals. After collection the invertebrate, was placed within a modified a container with several funnels than connect to a pump which water is pulled through, later filtered over filtered through a mesh (Fingerut et al., 2003). Alternatively, a broader approach was undertaken in Californian estuaries where seawater was directly filtered through a mesh with a modified pump. Abundance of cercariae in estuaries was calculated by timed interval samples as a proxy for volume filtered (Zimmer et al., 2009).

To predict the effects of current and potential environmental stressors on parasites, robust estimates of their abundance and response to biologically realistic conditions are needed. Predications on the response of parasite taxa to predicated changes have largely focused on the effect of temperature on a single trematode taxon (Poulin 2006, Macrogliese 2016). Thus, it could be postulated that accurate predictions of parasite

taxa's response to predicated climate change are difficult to make, due to the gaps in the knowledge, the inherent complexity of the host-parasite relationship and the neglect of environmental stressors outside of predicted anthropogenic warming (Harvell *et al.*, 1999, Lafferty & Ward 2004, Poulin, 2006, Marcogliese, 2016).

One such gap is the sublethal effects of parasites on intermediate hosts and the potential effect on surrounding biota. This is need in the case of the commonly parasitised taxa *P. vulgata* and *C. maenas* due to the high intensities of trematode metacercariae recorded and the ecological roles of the invertebrate hosts. Only primary intermediate infections of *E. patellae* in *P. vulgata* have been considered (Rees, 1934). Although, the effects of parasite taxa recorded within *P. vulgata* in Chapter 2 and 3 are largely considered non-pathogenic to the gastropod (Crewe, 1951, Lauckner, 1980) or have yet to be studied, as in the case of the cestode *O. insignis* (Crewe, 1947). The pathology of metacercariae is dependent on the intensity of infection (Fredensborg *et al.*, 2004, Thieltges, 2006).

Findings from the study thesis suggest that while trematode prevalence, intensity and distribution seemed to vary in relation to environmental stressors, further work should investigate the taxa's sensitivity to a range of environmental stressors prior to their use as potential bioindicators (Goodsell *et al.*, 2009).

There may be potential for trematodes as indicators for "Good Environmental Status" (GES) as defined by the Marine Strategy Framework Directive (MSFD). The food web descriptor (D4) in MSFD has been considered the productivity of key species or tropic groups, the proportion of selected species at the top of the food web and the abundance/ distribution of key trophic species (Rombouts *et al.*, 2013, Berg *et al.*, 2015). In particular, digenean trematodes require multiple hosts to complete their lifecycle, can be tropically transmitted and have been used to estimate the abundance

of final hosts (Byers *et al.*, 2011). However, at present, it is recommended that trematodes should be used in tandem with traditional approaches such as traditional biomarkers or communities' surveys. The reasons for the inclusion of trematodes are the taxa's recorded dominance in parasite communities, ubiquity and reported ability to affect host physiology and behaviour of invertebrate hosts (Lauckner, 1980; 1983, Theiltges *et al.*, 2006). This recommendation, is particularly relevant for studies that focus on the physiological responses of invertebrates as indicators of environmental stress or studies aiming to evaluate the effects of predicted climate change (Mouritsen *et al.*, 2002a, Kim *et al.*, 2008, Rodrigues *et al.*, 2014).

REFERENCES

- Berg T, Fürhaupter K, Teixeira H, Uusitalo L, Zampoukas N (2015) The Marine Strategy Framework Directive and the ecosystem-based approach pitfalls and solutions. *Marine Pollution Bulletin*, **96**, 18-28.
- Biguet J, Deblock S, Capron A (1958) Contribution to knowledge of Microphallidae. Annales de Parasitologie Humaine et Comparee, **33**, 396-444.
- Blanar CA, Munkittrick KR, Houlahan J, MacLatchy DL, Marcogliese DJ (2009) Pollution and parasitism in aquatic animals: A meta-analysis of effect size. *Aquatic Toxicology*, **93**, 18-28.
- Brenner M, Broeg K, Frickenhaus S, Buck BH, Koehler A (2014) Multi-biomarker approach using the blue mussel (*Mytilus edulis* L.) to assess the quality of marine environments, Season and habitat-related impacts. *Marine Environmental Research*, **95**, 13-27.
- Buck BH, Thieltges DW, Walter U, Rosenthal H (2005) Inshore–offshore comparison of parasite infestation in *Mytilus edulis*, implications for open ocean aquaculture. *Journal of Applied Ichthyology*, **21**, 107-113.
- Byers JE, Altman I, Grosse AM, Huspeni TC, Maerz JC (2011) Using Parasitic Trematode Larvae to Quantify an Elusive Vertebrate Host. *Conservation Biology*, **25**, 85-93.
- Crewe W, (1951) The occurrence of *Cercaria patellae* Lebour (Trematoda) and its effects on the host; with notes on some other helminth parasites of british limpets. *Parasitology*, **41**, 15-22.
- Cross M, Irwin S, Fitzpatrick S (2001) Effects of heavy metal pollution on swimming and longevity in cercariae of *Cryptocotyle lingua* (Digenea: Heterophyidae). *Parasitology*, **123**, 499-507.
- Cross MA, Irwin SWB, Fitzpatrick SM (2005) Effects of host habitat quality on the viability of *Cryptocotyle lingua* (Trematoda : Digenea) cercariae. *Parasitology*, **130**, 195-201.
- Cross MA, Irwin SWB, Fitzpatrick, S., Manga N (2003) Trematode parasite influence on copper, iron and zinc content of polluted *Littorina littorea*: infection, host

sex and time effects. *Journal of the Marine Biological Association of the United Kingdom*, **83**, 1269-1272.

- de Montaudouin X, Thieltges DW, Gam M, Krakau M, Jensen K and others (2009) Digenean trematode species in the cockle *Cerastoderma edule*:identification key and distribution along the north-east Atlantic shoreline. *Journal of the Marine Biological Association of the United Kingdom*, **89**, 543–556.
- de Montaudouin X, Wegeberg AM, Jensen KT, Sauriau PG (1998) Infection characteristics of *Himasthla elongata* cercariae in cockles as a function of water current. *Diseases of Aquatic Organisms*, **34**, 63-70.
- Evans DW, Irwin SWB, Fitzpatrick SM (1997) Metacercarial encystment and in vivo cultivation of *Cercaria lebouri* Stunkard 1932 (Digenea: Notocotylidae) to adults identified as *Paramonostomum chabaudi* van Strydonck 1965. *International Journal for Parasitology*, **27**, 1299-1304.
- Fermer J, Culloty SC, Kelly TC, O'Riordan RM (2011b) Parasitological survey of the edible cockle *Cerastoderma edule* (Bivalvia) on the south coast of Ireland. *Journal of the Marine Biological Association of the United Kingdom*, **91**, 923-928.
- Fingerut JT, Zimmer CA, Zimmer RK (2003) Patterns and Processes of Larval Emergence in an Estuarine Parasite System. *The Biological Bulletin*, **205**, 110-120.
- Galaktionov KV, Bustnes JO, Bardsen BJ, Wilson JG, Nikolaev KE, Sukhotin AA, Skirnisson K, Saville DH, Ivanov MV, Regel KV (2015) Factors influencing the distribution of trematode larvae in blue mussels *Mytilus edulis* in the North Atlantic and Arctic Oceans. *Marine Biology*, **162**, 193-206.
- Goodsell PJ, Underwood AJ, Chapman MG (2009) Evidence necessary for taxa to be reliable indicators of environmental conditions or impacts. *Marine Pollution Bulletin*, **58**, 323-331.
- Harvell C, Kim K, Burkholder J, Colwell R, Epstein PR, Grimes D, Hofmann E, Lipp E, Osterhaus A, Overstreet RM (1999) Emerging marine diseases-climate links and anthropogenic factors. *Science*, **285**, 1505-1510.
- Huspeni TC, Hechinger RF, Lafferty KD (2005) Trematode parasites as estuarine indicators: Opportunities, applications, and comparisons with conventional community approaches. *Estuarine Indicators*, (ed Bortone, SA) CRC Press, 297-314.

- James B (1968) The distribution and keys of species in the family Littorinidae and of their digenean parasites, in the region of Dale, Pembrokeshire. *Field Studies*, 2, 615-650.
- Kaplan AT, Rebhal S, Lafferty K, Kuris A (2009) Small estuarine fishes feed on large trematode cercariae: lab and field investigations. *Journal of Parasitology*, 95, 477-480.
- Khan RA, Thulin J (1991) Influence of pollution on Parasites of Aquatic Animals. *Advances in Parasitology*, **30**, 201-238.
- Kim Y, Powell EN, Wade TL, Presley BJ (2008) Relationship of parasites and pathologies to contaminant body burden in sentinel bivalves: NOAA Status and Trends 'Mussel Watch' Program. *Marine Pollution Bulletin*, 65, 101–127.
- Kollien A (1996) *Cercaria patellae* Lebour, 1911 developing in *Patella vulgata* is the cercaria of *Echinostephilla patellae* (Lebour, 1911) n. comb. (Digenea, Philophthalmidae). *Systematic Parasitology*, **34**, 11-25.
- Lafferty KD (1997) Environmental parasitology: What can parasites tell us about human impacts on the environment? *Parasitology Today*, **13**, 251-255.
- LaFonte BE, Raffel TR, Monk IN, Johnson (2015) Quantifying larval trematode infections in hosts, A comparison of method validity and implications for infection success. *Experimental Parasitology*, **154**, 155-162.
- Lauckner G (1980) Diseases of Mollusca: Gastropoda In: *Diseases of Marine Animals*. (ed by Kinne O), John Wiley & Sons, New York. **1**, 311–424.
- Lauckner G (1980) Diseases of Mollusca: Gastropoda In: *Diseases of Marine Animals*. (ed by Kinne O), John Wiley & Sons, New York. **1**, 311–424.
- Lauckner G (1983) Introduction, Bivalvia to Scaphopoda In: *Diseases of Marine Animals.* (ed by Kinne O), John Wiley & Sons **2**, 477-897.
- Lauckner G (1983) Introduction, Bivalvia to Scaphopoda In: *Diseases of Marine Animals.* (ed by Kinne O) **2**, 477-897.
- Leung TLF, Donald KM, Keeney DB, Koehler AV, Peoples RC, Poulin R (2009) Trematode parasites of Otago Harbour (New Zealand) soft-sediment intertidal ecosystems, life cycles, ecological roles and DNA barcodes. *New Zealand Journal of Marine and Freshwater Research*, **43**, 857-865.

- Loos-Frank B (1967) Experimentelle Untersuchungen über Bau, Entwicklung und Systematik der Himasthlinae (Trematoda, Echinostomatidae) des Nordseeraumes. *Zeitschrift für Parasitenkunde*, **28**, 299-351.
- Loos-Frank B (1969) Studies on Gymnophallid Trematodes of North Sea. I. Alternative Life-Cycles of *Gymnophallus choledochus* Odhner 1900. *Zeitschrift Fur Parasitenkunde*, **32**, 135-156.
- Marcogliese DJ (2005) Parasites of the superorganism: Are they indicators of ecosystem health? *International Journal for Parasitology*, **35**, 705–716.
- Marcogliese, D J (2016) The Distribution and Abundance of Parasites in Aquatic Ecosystems in a Changing Climate: More than Just Temperature. *Integrative and Comparative Biology*, **56**, 611-619.
- Markowitz K, Williams J, Krause M (2016) Development of quantitative PCR assay for detection of the trematode parasite *Proctoeces maculatus* in the blue mussel *Mytilus edulis*. *Diseases of Aquatic Organisms*, **122**, 125-136.
- Mouritsen KN, Poulin R (2002a) Parasitism, climate oscillations and the structure of natural communities. *Oikos*, **97**, 462-468.
- Mouritsen KN, Poulin R (2002b) Parasitism, community structure and biodiversity in intertidal ecosystems. *Parasitology*, **124**, 101-117.
- Nybakken JW, Bertness MD (2005) Marine biology: an ecological approach. Sixth edition. **1-6**, 1-579.
- Pérez-Ponce de León, G and Poulin, R (2017) An updated look at the uneven distribution of cryptic diversity among parasitic helminths. *Journal of Helminthology*, **6**, 1-6.
- Pietrock M, Marcogliese DJ (2003) Free-living endohelminth stages: at the mercy of environmental conditions. *Trends in Parasitology*, **19**, 293-299.
- Poulin R (2006) Global warming and temperature-mediated increases in cercarial emergence in trematode parasites. *Parasitology*, **132**, 143-151.
- Poulin R, Blasco-Costa I, Randhawa HS (2016) Integrating parasitology and marine ecology: Seven challenges towards greater synergy. *Journal of Sea Research*, 113, 3-10.
- Prinz K, Kelly TC, O'Riordan RM, Culloty SC (2010b) Infection of *Mytilus edulis* by the trematode *Echinostephilla patellae* (Digenea: Philophthalmidae). *Journal of Helminthology*, **84**, 193-198.

- Prokofiev VV, Galaktionov KV, Levakin IA (2016) Patterns of parasite transmission in polar seas: Daily rhythms of cercarial emergence from intertidal snails. *Journal of Sea Research*, **113**, 85-98.
- Rees FG (1934) Cercaria patellae Lebour, 1911, and its effect on the digestive gland and gonads of Patella vulgata. Proceedings of the Zoological Society of London, 104, 45-53.
- Rodrigues ET, Pardal MA (2014) The crab *Carcinus maenas* as a suitable experimental model in ecotoxicology. *Environment International*, **70**, 158-182.
- Rombouts I, Beaugranda G, Fizzala X, Gaill F, Greenstreet SPR, Lamarec S, Le Loc'hf F, McQuatters-Gollop A, Mialet B, Niquil N, Percelay J, Renaudb F, Rossbergj AG, Féral JP (2013) Food web indicators under the Marine Strategy Framework Directive: From complexity to simplicity? *Ecological Indicators*, 29, 246-254.
- Saville DH, Irwin SWB (1991) In ovo cultivation of *Microphallus primas* (Trematoda: Microphallidae) metacercariae to ovigerous adults and the establishment of the life-cycle in the laboratory. *Parasitology*, **103**, 479-484.
- Sousa WP (1994) Patterns and processes in communities of helminth parasites. *Trends in Ecology & Evolution*, **9**, 52-57.
- Stunkard H W (1930) The life history of *Cryptocotyle lingua* (Creplin), with notes on the physiology of the metacercariae. *Journal of Morphology*, **50**, 143-191.
- Stunkard HW, Uzmann JR (1959) The life-cycle of the digenetic trematode, *Proctoeces maculatus* (Looss, 1901) Odhner, 1911 [syn. *P. subtenuis* (Linton, 1907) Hanson, 1950], and description of *Cercaria adranocerca* n. sp. Biological Bulletin (Woods Hole) **116**, 184-193.
- Sures B, Nachev M, Selbach C, Marcogliese DJ (2017) Parasite responses to pollution: what we know and where we go in 'Environmental Parasitology'. *Parasites & Vectors*, **10**, 1-19.
- Thieltges DW, Rick J (2006) Effect of temperature on emergence, survival and infectivity of cercariae of the marine trematode *Renicola roscovita* (Digenea, Renicolidae). *Diseases of Aquatic Organisms*, **73**, 63-68.
- Thieltges DW, Wegner KM (2016) Ecology and evolution of marine parasites and diseases: International NIOZ–AWI symposium, 10-14 March 2014, The Netherlands. *Journal of Sea Research*, **113**, 1-2.

- Thieltges DW. (2006) Effect of infection by the metacercarial trematode Renicola roscovita on growth in intertidal blue mussel Mytilus edulis. Marine Ecology Progress Series, **319**,129-134.
- Thieltges DW, Engelsma MY, Wendling CC, Wegner KM (2013) Parasites in the Wadden Sea food web. *Journal of Sea Research*, 82, 122-133.Thompson RC, Crowe TP, Hawkins SJ (2002) Rocky intertidal communities: past environmental changes, present status and predictions for the next 25 years. *Environmental Conservation*, 29, 168-191.
- Timon-David J, Rebecq J (1959) Les métacercaires parasites de l'annélide *Nereis diversicolor* O.F. Müller et leur développement expérimental. Comptes rendus des séances de la Société de biologie. **152**,1731–1733.
- Vanoverschelde R, Vaes F (1980) Studies on the life-cycle of *Himasthla militaris* (Trematoda: Echinostomatidae). *Parasitology*, **81**, 609-617.
- Vidal-Martinez VM, Pech D, Sures B, Purucker ST, Poulin R (2010) Can parasites really reveal environmental impact? *Trends in Parasitology*, **26**, 44-51.
- Yamaguti S (1971) Synopsis of digenetic trematodes of vertebrates. 2, 1-1074.
- Zimmer RK, Fingerut JT, Zimmer CA (2009) Dispersal pathways, seed rains, and the dynamics of larval behavior. *Ecology*, **90**, 1933-1947.



Fig. 1 Averaged percentage cover of macroalgal taxa recorded per sampling event per site (a) Cusheen, (b) Lough Hyne (c) Sandycove and (d) Whitegate. NS represent not sampled as Lough Hyne was only sampled & surveyed every second month.

224

Appendix



Fig. 2 Averaged density of gastropod host species screened taxa recorded per sampling event per site (a) Cusheen, (b) Lough Hyne, (c) Sandycove and (d) Whitegate. NS represent not sampled as Lough Hyne was only sampled & surveyed every second month. Note the differing y-axes

225

Location	Collection method	Shell height Trematode Polychaete		Cestode	Ciliate	Mite	
		(mm)	(%)	(%)	(%)	(%)	(%)
Whitegate	Hand	22.97 ±3.90	6.67	3.33	NA	3.33	0.00
	QR	22.74±4.39	5.71	8.57	NA	0.00	0.00
	QR	21.28±5.81	4.00	4.00	NA	4.00	0.00
	QT	20.81±5.28	11.76	2.94	NA	5.88	0.00
Sandycove	QR	21.51±8.91	11.11	0.00	NA	0.00	0.00
	QT	21.56±3.79	5.88	0.00	NA	0.00	2.94
Whitegate	QR	24.36±5.74	0.00	0.00	NA	0.00	0.00
	QT	28.13±3.95** ¹	2.78	0.00	NA	0.00	0.00
Tralispean	QR	12.79±2.96* ²	15.63	0.00	15.63	0.00	0.00
	QT	11.23±4.52	33.33	0.00	2.56	0.00	0.00
Sandycove	Hand	15.40±2.99** ³	83.33* ⁴	20.00	15.63	33.33	6.67
	QR	12.97±5.30	53.33	43.33	33.33	56.67	3.33

Table 1 Comparison of collection methods of host species on sample morphometrics and parasite prevalence. Asterisk den Appendix level of significant P-values *<0.05, **<0.001, ***<0.0001.

¹ F=7.86, df=1, P<0.01 ² F=4.67, df=1, P<0.05 ³ F=8.76, df=1, P<0.01

 $^{4}\chi^{2}$ =4.93, df=1, P<0.05

Table 2 Prevalence of trematode taxa in foot tissues of *Mytilus* spp. screened during in the three trials run in Chapter 5. Prevalence of *Echinostephilla patellae* is excluded from the table as in the trials *Mytilus* spp. were exposed to *E. patellae*. – or + indicate whether *Mytilus* spp. was exposed to *Echinostephilla patellae* cercariae. For Trial 1, I represent intertidal and S subtidal *Mytilus* spp. collected in Lough Hyne. For Trial 2, Cu represents Cusheen, LH Lough Hyne, R Ringaskiddy. For Trial 3, C represents the control, L low nutrient enrichment, M medium enrichment, and H High enrichment treatments.

Trial	Treatment	Foot tissue				
		<i>Renicola</i> spp.	Himasthla spp.	Echinostephilla patellae		
1	I-	0.00	0.00	58.33 (1-191)		
1	I+	0.00	0.00	66.67 (2-120)		
1	S-	0.00	0.00	8.33 (22)		
1	S+	0.00	0.00	33.33 (1-5)		
2	Cu-	33.33 (1-2)	88.89 (1-13)	33.33 (1-2)		
2	Cu+	11.11 (1)	77.78 (1-9)	11.11 (1)		
2	LH-	22.22 (1)	44.44 (2-19)	22.22 (1)		
2	LH+	11.11 (1)	66.67 (3-25)	11.11 (1)		
2	R-	0.00	11.11 (2)	0.00		
2	R+	22.22 (1)	33.33 (1-9)	22.22 (1)		
3	C-	29.41 (1-4)	41.18 (1-10)	5.88 (1)		
3	C+	50.00 (1-3)	18.75 (2-4)	6.25 (1)		
3	L-	42.86 (1-2)	57.14 (1-10)	14.29 (1-3)		
3	L+	47.06 (1-5)	70.59 (1-7)	0.00		
3	M-	37.50 (1-4)	43.75 (1-37)	12.50 (1-5)		
3	M+	43.75 (1-6)	62.50 (1-7)	6.25 (1)		
3	H-	33.33 (1-4)	33.33 (1-4)	0.00		
3	H+	41.18 (1-7)	5.88 (1)	11.76 (1-5)		
Grand '	Fotal	26.52 (1-7)	35.65 (1-37)	16.52 (1-191)		

Appendix

Table 3 Overall prevalence and range in intensity of trematode taxa recorded in host taxa in the thesis. The life cycle stage of trematode taxa are listed in order of infection i.e., sporocyst, metacercaria and adult worm. TC or H indicates the method of detection was either tissue compression or histological. Host taxa habitat is classified as rocky shore intertidal (RI), rocky shore subtidal (RS) or as estuarine intertidal (EI). Trematode taxa are spilt relative to their lifecycle stages due to their differing routes of transmission and effects on host taxa. *Mytilus* spp.^{*} represent individuals not included in trials in Chapter 5.

Host taxa	Trematode taxa	Total prevalence (%)	Range of Intensity	Method	Host taxa habitat	Estimated from Chapter
SPOROCYST						*
(Primary intermediate	hosts)					
Littorina littorea	Paramonostomum chabaudi	0.20 (n=1, 572)	NA	TC	RI	2
	Cryptocotyle lingua	1.63	NA	TC	RI	2
	Himasthla spp.	0.27	NA	TC	RI	2
	Podocotyle atomon	1.56	NA	TC	RI	2
	Renicola spp.	0.07	NA	TC	RI	2
Mytilus spp.	Trematode taxa	0.14 (n=731)	NA	Н	RS	5
v 11	(likely Bucephalus spp.)					
Nucella lapillus	Parorchis acanthus	0.10 (n=956)	NA	TC	RI	2
METACERCARIA						
(Second intermediate	hosts)					
Carcinus maenas	Microphallus spp.	86.90 (n=290)	1-2, 436	TC	EI	4
Hediste diversicolor	Echinostomid	15.41 (n=713)	1-13	TC	EI	4
	Gymnophallus spp.	5.44	1-19	TC	EI	4
L. littorea	Renicola spp.	7.13	1-875	TC	RI	2
	Echinostephilla patellae	0.68	1	TC	RI	2
	Himasthla spp.	0.07	1	TC	RI	2

Table 3 Continued

Host taxa	Trematode taxa	Total prevalence	Range of	Method	Host taxa	Estimated
		(%)	Intensity		habitat	from
						Chapter
Mytilus spp.	Trematode taxa	3.53 (n=731)	1-11	Н	RS	5
<i>Mytilus</i> spp.*	E. patellae	15.20 (n=125)	1-16	TC	RI	5
	Himasthla spp.	37.60	1-56	TC	RI	5
	<i>Renicola</i> spp.	77.60	1-123	TC	RI	5
		10.00 (n=10)	1	TC	RS	5
N. lapillus	<i>Renicola</i> spp.	1.15	1-12	TC	RI	2
-	E. patellae	0.31	1-4	TC	RI	2
Patella vulgata	<i>E. patellae</i> $(n=1,745)$	40.90	1-223	TC	RI	2 & 3
Ū.	Gymnophallus spp.	0.23	1-103	TC	RI	2 & 3
	Renicola spp.	0.28	1-8	TC	RI	2
ADULT WORM						
(Final hosts)						
N. lapillus	Proctoeces sp.	0.10	1	TC	RI	2
P. vulgata	Ĩ	0.07	1	TC	RI	2

Family	Genera	First intermediate host	Second intermediate host	Final host	Free living life stages	Reference
Echinostomatidae	<i>Himasthla</i> sp.	Gastropods Littorina littorea	Molluscs & polychaetes L. littorea	Birds	2	James 1968a
	Himasthla militaris	Gastropods	Polychaetes Hediste diversicolor	Larus glaucescens, Arenaria interpres, Calidris maritime, S. mollissima, Chen canagica, Motacilla alba	2	Timon-David & Rebecq 1959, Loos- Frank, 1967, Vanoverschelde & Vaes 1980
Fellodistomidae	Proctoeces sp.	Gastropods	Gastropods	Gastropods Patella vulgata Nucella lapillus	1	Stunkard & Uzmann 1959, Lauckner, 1980
Gymnophallidae	<i>Gymnophallus</i> sp.	Gastropods	Gastropods & Bivalves Patella vulgata	Birds	2	Stunkard & Uzmann 1958, Lauckner, 1980
	Gymnophallus choledochus	Gastropods	Molluscs & polychaetes	Charadriiform, lariform and anatid birds.		Loos-Frank 1969
Heterophyidae	Cryptocotyle lingua	Gastropods Littorina littorea	Bony fish	Birds and mammals	2	Stunkard 1930, James 1968a

Table 4 Summary of lifecycles of trematode taxa recorded during this study. All chapters and sampling events combined.

Table 4 Continued

Family	Genera	First intermediate host	Second intermediate host	Final host	Free living life stages	Reference
Microphallidae	Microphallus primas	Gastropods	Crustaceans Carcinus maenas	Birds Charadriiformes and Anseriformes	2	Biguet <i>et al.</i> , 1958, Saville & Irwin 1991
Notocotylidae	Paramonostomum chabaudi	Gastropods Littorina littorea	Vegetation or substrate	Birds	1	James 1968, Evans <i>et</i> <i>al.</i> , 1997
Opecoelidae	Podocotyle atomon	Gastropods L. littorea	Benthic crustaceans, mainly amphipods	Teleosts	2	James 1968a, Thieltges & Reise 2007
Philophthalmidae	Echinostephilla patellae	Gastropods P. vulgata	Gastropods and Bivalves <i>Mytilus</i> spp. <i>P. vulgata</i>	Wading birds	2	Kollien, 1996, Prinz et al., 2010b
	Parorchis acanthus	Gastropods Nucella lapillus	Gastropods	Seabirds	2	Yamaguti 1971, Lauckner, 1980,
Renicolidae	<i>Renicola</i> spp.	Gastropods L. littorea	Molluscs L. littorea	Fish consuming birds	2	Stunkard & Uzmann 1959, James 1968, Lauckner, 1980

Table 4 Continued

Family	Genera	First intermediate host	Second intermediate host	Final host	Free living life stages	Reference
Echinostomatidae	Himasthla sp.	Gastropods Littorina littorea	Molluscs & polychaetes	Birds	2	James 1968a
	Himasthla militaris	Gastropods	Polychaetes Hediste diversicolor	Larus glaucescens, Arenaria interpres, Calidris maritime, S. mollissima, Chen canagica, Motacilla alba	2	Timon-David & Rebecq 1959, Loos- Frank, 1967, Vanoverschelde & Vaes 1980
Fellodistomidae	Proctoeces sp.	Gastropods	Gastropods	Gastropods Patella vulgata Nucella lapillus	1	Stunkard & Uzmann 1959, Lauckner, 1980
Gymnophallidae	<i>Gymnophallus</i> sp.	Gastropods	Gastropods & Bivalves Patella vulgata	Birds	2	Stunkard & Uzmann 1958, Lauckner, 1980
	Gymnophallus	Gastropods				Loos-Frank 1969
	Cryptocotyle lingua	Gastropods Littorina littorea	Bony fish	Birds and mammals	2	Stunkard 1930, James 1968a

Table 4 Continued

Family	Genera	First intermediate	Second intermediate host	Final host	Free living life stages	Reference
Microphallidae	Microphallus primas	Gastropods	Crustaceans Carcinus maenas	Birds Charadriiformes and Anseriformes	2	Biguet <i>et al.</i> , 1958, Saville & Irwin 1991
Notocotylidae	Paramonostomum chabaudi	Gastropods Littorina littorea	Vegetation or substrate	Birds	1	James 1968, Evans <i>et al.</i> , 1997
Opecoelidae	Podocotyle atomon	Gastropods L. littorea	Benthic crustaceans, mainly amphipods	Teleosts	2	James 1968, Thieltges & Reise 2007
Philophthalmidae	Echinostephilla patellae	Gastropods P. vulgata	Gastropods and Bivalves Mytilus spp. P. vulgata	Wading birds	2	Kollien, 1996, Prinz et al., 2010b
	Parorchis acanthus	Gastropods Nucella lapillus	Gastropods	Seabirds	2	Yamaguti 1971, Lauckner, 1980,
Renicolidae	Renicola roscovita	Gastropods L. littorea	Molluscs L. littorea	Fish consuming birds	2	Stunkard & Uzmann 1959, James 1968, Lauckner, 1980