

Title	Heritability estimation via molecular pedigree reconstruction in a wild fish population reveals substantial evolutionary potential for sea-age at maturity, but not size within age-classes
Authors	Reed, Thomas E.;Prodöhl, Paulo A.;Bradley, Caroline;Gilbey, John;McGinnity, Philip;Primmer, Craig R.;Bacon, Phil J.
Publication date	2018-07-30
Original Citation	Reed, T. E., Prodöhl, P. A., Bradley, C., Gilbey, J., McGinnity, P., Primmer, C. R. and Bacon, P. J. (2019) 'Heritability estimation via molecular pedigree reconstruction in a wild fish population reveals substantial evolutionary potential for sea-age at maturity, but not size within age-classes', Canadian Journal of Fisheries and Aquatic Sciences, 76(5), pp. 790-805. doi: 10.1139/cjfas-2018-0123
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://doi.org/10.1139/cjfas-2018-0123 - 10.1139/cjfas-2018-0123
Rights	© 2018, the Authors. Published by NRC Research Press. All rights reserved.
Download date	2024-05-14 12:25:57
Item downloaded from	https://hdl.handle.net/10468/7876

Heritability estimation via molecular pedigree reconstruction in a wild fish population reveals substantial evolutionary potential for age at maturity, but not size within age-classes.

Can. J. Fish. Aqu. Sci.

Acceptance date: 08-Jul-2018

Thomas E Reed^{1†*}, Paulo Prodöhl^{2†}, Caroline Bradley², John Gilbey³, Philip McGinnity^{1,4},
Craig R. Primmer^{5,6}, and Philip J. Bacon³.

¹ School of Biological, Earth and Environmental Sciences, University of Cork, North Mall, Cork, Ireland.

² Institute for Global Food Security, School of Biological Sciences, Queen's University of Belfast, Medical Biology Centre, 97 Lisburn Road, Belfast, BT9 7BL, UK.

³ Marine Scotland Science, Freshwater Laboratory, Faskally, Pitlochry, Perthshire PH16 5LB, Scotland, UK.

⁴ Marine Institute, Furnace, Newport, Co. Mayo, Ireland.

⁵ Department of Biosciences, University of Helsinki, 00014, Finland

⁶ Institute of Biotechnology, University of Helsinki, 00014, Finland

† Joint first authors.

* Corresponding author: treed@ucc.ie

ABSTRACT

While evolutionary responses require heritable variation, estimates of heritability (h^2) from wild fish populations remain rare. A 20-year molecular pedigree for a wild Scottish population of Atlantic salmon (*Salmo salar*) was used to investigate genetic contributions to (co)variation in two important, correlated, phenotypic traits: “sea-age” (number of winters spent at sea prior to spawning) and size-at-maturity (body length just prior to spawning). Sea-age was strongly heritable ($h^2 = 0.51$) and size exhibited moderate heritability ($h^2 = 0.27$). A very strong genetic correlation ($r_G = 0.96$) between these traits implied the same functional loci must underpin variation in each. Indeed, body size within sea-ages had much lower heritability that did not differ significantly from zero. Thus, within wild *S. salar* populations, temporal changes in sea-age composition could reflect evolutionary responses, whereas rapid changes of body-size within sea-ages are more likely due to phenotypic plasticity. These inheritance patterns will influence the scope of evolutionary responses to factors such as harvest or climate change and, hence, have management implications for salmonid populations comprising a mix of sea ages.

Keywords. salmonids; population dynamics, quantitative genetics, Bayesian, evolvability, adaptation, management, conservation genetics

INTRODUCTION

Life history traits such as age and size at first reproduction are typically expected to be under strong selection, given their close links with Darwinian fitness (Stearns 1992, Roff 2002). All else being equal, genotypes that reproduce as early as possible in life should have higher relative fitness than those that first reproduce at older ages (Brommer et al. 2002). In species with indeterminate growth such as most fishes, however, there may be strong breeding advantages (especially for females) to delaying first reproduction to older ages, and hence larger sizes (Heino and Kaitala 1999). The optimum age and size at first reproduction will then depend on factors such as (seasonal) environmental opportunities for growth, age-specific extrinsic mortality schedules, and the extent to which reproductive success depends on body size (Stearns 1992, Charlesworth 1994, Roff 2002). Human-induced changes to any of these parameters can alter the selective pressures on populations; for example, harvests of wild animals can drive “unnatural selection” on phenotypic traits such as age at maturity and body size (Allendorf and Hard 2009, Heino et al. 2015, Harvey et al. 2017), while climate change can impact selective dynamics in a range of complex ways (Munday et al. 2013, Sydeman et al. 2015). Inferring the potential for evolutionary responses, however, requires information on the extent to which observed phenotypic (co)variation is underpinned by genetic (co)variation (Kuparinen and Hutchings, 2017). Detailed, long-term, studies of naturally regulated populations, in which the life histories, reproductive success and pedigrees of individual animals are measured or inferred can provide this information (Naish and Hard 2008, Clutton-Brock and Sheldon 2010).

In Atlantic salmon (*Salmo salar* L.) and other salmonid fishes, the sea-age of adults returning to spawn for the first time is an important trait from both ecological and economic perspectives, as older fish tend to be larger and therefore more valuable from a commercial and recreational fishing perspective, while also having higher fecundity and therefore

contributing more towards total annual egg numbers. Sea-age variation may also contribute towards ecological and evolutionary portfolio effects via bet-hedging mechanisms (Schindler et al., 2015), where risk is spread across multiple sea-age cohorts. Some individuals, and populations of Atlantic salmon, return after just one winter at sea (1SW), whereas others known as multi-sea winter (MSW) fish return after two or more winters at sea (Klemetsen et al, 2003). The marine feeding destination, sea-age, adult-return size and adult run-timing are all inter-related (Malcolm et al, 2010). The average size of returning adults generally increases with time spent at sea, both across and within sea-age classes; for example, MSW fish tend to be on average much larger than 1SW, and later-running fish of a given sea-age class are typically larger than early-running fish (Hutchings & Jones 1998, Bacon et al. 2009, 2010, Barson et al. 2015). A substantial genetic component underlying sea-age variation has been shown or implied by several studies of Atlantic salmon (Nævdal 1983, Gjerde 1984, Johnston et al. 2014, Ayllon et al. 2015, Barson et al. 2015) and related species such as Chinook salmon (*Oncorhynchus tshawytscha*) (Hankin et al. 1993, Quinn et al. 2001) and steelhead/rainbow trout (*Oncorhynchus mykiss*) (Tipping 1991, Kause et al. 2003).

Body-length at maturity has also been shown to be heritable in salmonids generally, with Carlson & Seamons (2008) reporting a median heritability (the fraction of phenotypic variation explained by genetic variation) estimate of 0.21 across six salmonid studies. Only one of these studies (Dickerson et al. 2005), however, involved a situation where the fish (in that case, pink salmon, *Oncorhynchus gorbuscha*) spawned and reared naturally in a wild river environment; the rest all involved either hatchery/farm-origin broodstock and/or artificial rearing conditions (Carlson and Seamons 2008). Since then a limited number of additional salmonid studies have appeared that estimate heritability of length-at-age in freshwater in fully natural environments (Serbezov et al. 2010, Morrissey and Ferguson 2011,

Reed et al. 2015), but these studies did not focus on heritability of size-at-maturity specifically.

Availing from the (still rare) existence of a multi-generation data set from a wild Atlantic salmon population, the specific aims of this study were to: (1) construct a pedigree based on molecular data, (2) estimate the heritability of both adult body-length and sea-age-at-maturity; (3) test for any genetic correlation between these traits; and (4) test for temporal trends in estimated breeding values for both traits, which would indicate microevolutionary responses over the study period. Significant phenotypic trends in body size and sea-age are apparent in this population during our 1991-2011 study period (Glover, pers comm.) and other populations in the region more generally (Bacon et al. 2009), but it remains unknown whether there is any genetic component to these trends. Answering the latter question could suggest fruitful new avenues for research and interpretation of monitoring data.

METHODS

Study site

The Girnock Burn (Fig.1) is an intensively studied upland catchment of c. 30 km² on the Aberdeenshire Dee (confluence 57 ° 0.6' N, 3° 5.3' W), in north-eastern Scotland, which has been operated by Marine Scotland Science Freshwater Fisheries Laboratory since 1966. Full details of the site, monitoring data and a list of associated publications are available at URL1. In brief, the catchment contains some 8 km of nursery stream for Atlantic salmon ranging in altitude from 230 to 900m. Adult salmon entering the Girnock are caught in an ascending trap near the confluence with the mainstem of the Aberdeenshire Dee, given an individual identification (floy) tag, sexed based on phenotypic characteristics, measured (adult fork

body-length, hereafter body-length) and aged by scale-reading, following international protocols.

Apart from unusual scale- or tissue-sample loss, the DNA samples obtained from adults can be taken as a complete set of breeders (only one adult is known to have got above the traps without being caught in 50 years, during emergency trap-maintenance during an autumn spate). Since the early 1990s tissue samples (adipose-fin clips) of adults were also taken, and stored in molecular grade (99%) ethanol.

The Girnock provided an example of a fully wild Atlantic salmon population from the start of the historic study (1966) until 1999, with adult fish being allowed to spawn naturally above the traps. During the late 1990s, numbers of adult females fell very low (9, 11 & 22 over three years), and from 2000 to 2010 increasing proportions of adults were temporarily held, stripped and their eggs stocked into the river section above the trap (see Bacon et al. 2015; URL2; Glover, *pers. comm.*). The fish were artificially mated with a crossing protocol designed to increase mating variation and promote the maintenance of broad genetic variation (see Bacon et al. 2015 for full details). From 2011 the system was returned to fully natural spawning (but no offspring from that period returned during this study). Most juveniles from the Girnock undergo parr-to-smolt transformation and seaward migration at age 2+ (third year of life) or 3+ (fourth year of life). A minority of adults return after 1SW, the majority after 2SW, typical of upland Scottish salmon sites (Gurney et al. 2015; URL3). 3SW adults are now uncommon, and repeat spawners are regionally relatively rare (less than 2%, as is typical for eastern Scotland, Bacon et al. 2012). The life span of Atlantic salmon from the Girnock is therefore typically 5 to 6 years, but can range from 4 to 8 years. 1SW adults typically suffer only about half the marine mortality of 2SW fish, but, being around half the size (biomass), female 1SW salmon produce about half as many eggs as 2SW females (Gurney et al. 2012, 2015).

DNA samples and microsatellite genotyping

DNA samples were derived from three sources during our study period 1991-2011. In the mid-1990s, aliquots of genomic DNA extracted during earlier studies were available, prepared according to protocols given in Taggart et al. (2001). Most of the other samples from the late 1990s onwards were obtained from adipose fin-clips taken from adults, stored in molecular grade (99%) ethanol, and with genomic DNA extracted using the Promega, Wizard® SV 96 Genomic DNA Purification System (www.promega.com) (see Keenan et al. 2013a). The remaining samples, mainly from the early 1990s, plus some fish that were not adipose-clipped in later years, were obtained from scale samples, archived in paper packets, air-dried and with the genomic DNA extracted as above. Genomic DNA for all samples was checked for quality and concentration through visual comparison with a *Hind*III digested λ DNA size standard on Ethidium Bromide stained 0.8% 0.5X TBE agarose gels.

All samples were examined for a panel consisting of 17 putatively neutral microsatellite marker loci organised in two multiplex PCR. These microsatellites were selected based on information content (polymorphism), consistent co-amplification reliability and non-overlapping size compatibility. Multiplex- 1 comprised ten markers and Multiplex 2 consisted of the seven remaining microsatellites (see Appendix A for details); , in addition to a sex specific marker based on the sdY locus, which is a conserved region for sex determination in salmonids (Yano et al. 2012, 2013). This later marker was used to double-check concordance between the biometric records and the genotype results (to guard against recording errors, including samples potentially omitted from sequences).

PCRs were carried out in 96 well microtitre plates in 3.5 μ l volumes consisting of 1 μ l DNA (~2-5 ng/ μ l), 1.75 μ l of 2x PPP Top-Bio mastermix (Top-Bio) and 0.75 μ l of a cocktail

of ABI fluorescent labelled forward and/or unlabelled reverse ‘pig’ tailed PCR primers (Appendix A.1 for details on labelled and unlabelled primers, and specific primer concentrations per marker loci). Samples were overlain with 10 µl of mineral oil to prevent evaporation. Thermocycling conditions for both multiplex panels were as follows: an initial denaturation step of 15 min at 95°C, followed by 28 cycles of 95°C for 30 sec, 57°C for 90 sec and 72°C for 60 sec. This was followed by a final extension step of 30 min at 60°C. All reactions were carried out using Techne TC-Plus thermal cyclers, with a heated lid at 105°C. Amplified fragments were diluted one-tenth with double-distilled H₂O, and 1µl of this dilution was added to 9 µl of HiDi formamide (Thermo Fisher Scientific) mixed with Gene Scan 600-LIZ size ladder (Thermo Fisher Scientific).

Diluted PCR products were analysed on a 96 capillary ABI 3730XL DNA analyser (Thermo Fisher Scientific), and the fragment size analysis (i.e. allelic calls) for genotypes carried out using GENEMAPPERv4.1 (Thermo Fisher Scientific). Genotypes for each microsatellite locus/specimen were individually checked and manually confirmed prior to their addition to an electronic database. Over 50% of the genotyping was independently repeated to ensure consistent scoring (i.e. to minimise scoring errors). Locus-specific statistics including allele numbers, heterozygosity, allelic richness and deviations from Hardy-Weinberg expectations were estimated both on annual caught samples (i.e. candidate parents) and the pooled data set using the R package *diveRsity* (Keenan et al. 2013b). The power of the markers to correctly assign individuals to unique families was carried out using Family Assignment Program (FAP; Taggart 2007).

Pedigree reconstruction

Parental cohorts were defined as adults returning to spawn at the Girnock in each autumn. For each such spawning cohort, a sub-set of later returning adults that comprised the full span of their sea-going progeny, by all possible river-age (1 to 4) and sea-age (1 to 3) range combinations, was assembled as putative offspring. For instance, returning adults caught in the trap between 1993 and 1998 were analysed as putative offspring for the 1991 spawning cohort. Pedigree reconstruction was carried out using FAP (Taggart 2007), which estimates exclusion-based family assignment probabilities within family mixtures where all parental genotypes are known. In summary, FAP was used to search all potential offspring for the given spawning cohort and to exclude those that could not derive from a given putative pair of male and female adult parents (based on field-observed sexes). The output was a list of ‘family-trio’ identities (potential female parent, male parent, offspring), together with information on the number of loci matching exactly, the number of evidently mismatching loci, and the list of such mismatching offspring loci. This process was repeated, separately, for each adult cohort from 1991 to 2006 (i.e. for all spawning cohorts for which full-sets of putative returning adult offspring were available in the sample database).

The resulting cohort-specific lists of putative family trios were manually examined and any mismatching loci for the putative offspring and its parents were double checked back to the original ABI raw chromatogram data. Most allele mismatches involved single marker loci and resulted from one individual (one parent or the offspring) being incorrectly assigned an adjacent-sized allele, rather than the likely-true parental one. These could be easily rectified following visual inspection. Following this second stage of quality control, the validated family trios were used as the basis for the pedigree. Only complete parent-offspring records were used in this study (a sampled offspring and both its anadromous father and mother), excluding any offspring with only one known adult female parent, presumably sired by an un-sampled precocious parr male. FAP was also used to estimate the ‘parental

exclusion power' of the data (i.e. power of the markers to correctly assign individuals to a given set of known families).

Power analyses for heritability estimation

In order to quantify the statistical power in this dataset to detect true non-zero h^2 , a power analysis (see Appendix B.3 for full details) was conducted in which a range of h^2 values were simulated ($h^2 = 0.05, 0.10, 0.15, 0.20, 0.25, 0.30, 0.40, 0.50, 0.75$) using the *rbv* function in *MCMCglmm* (Hadfield 2010). One thousand replicate simulations were run for each simulated h^2 value, and each time the estimated h^2 and its associated P-value based on likelihood ratio tests were returned and saved. Power was then calculated as the proportion of P-values that were <0.05 .

Estimation of quantitative genetic parameters using animal models

Bayesian animal models were used to estimate quantitative genetic parameters of interest using the R package *MCMCglmm* (Hadfield 2010). Full details on the estimation procedures for the quantitative genetic parameters, as well as more details on the pedigree structure, are given in Appendix B.1

Univariate animal models for the data were first devised (one trait as a single response variable) to estimate the narrow-sense heritability (h^2) of each of the phenotypic traits of interest: adult body-length and sea-age-at-maturity. These univariate models partitioned the total phenotypic variance V_P in each trait into contributions from additive genetic effects (V_A), maternal effects (V_M), year effects (V_Y) and residual effects (V_R), the latter assumed to arise from environmental effects and non-additive genetic effects. Thus $V_P = V_A + V_M + V_Y +$

V_R , and $h^2 = V_A/V_P$. A fixed effect of sex (two level factor: adult males and adult females) was included in the univariate models for both traits, to control for the fact that males may mature at a younger age, on average, and hence also are smaller as adults, than females. Year-to-sea (the year the focal individual migrated to sea as a smolt, known from scale readings) was used to define temporal cohorts (i.e. V_Y , see Appendix B.2).

Adult body size (measured in centimetres and treated as a Gaussian variable) was first natural logarithm (hereafter simply log) transformed before fitting the animal models, as preliminary exploratory analyses indicated that this transformation minimised autocorrelation between MCMC samples and yielded less skewed posterior distributions of the parameters.

In Atlantic salmon, among-individual variation in adult body size is strongly affected by sea-age, as fish that remain at sea an extra year have more time for marine growth. The mean size of 1SW salmon in our 1990-2012 sample, for example, was 55.8 ± 2.1 (S.E.) cm, whereas the mean size of MSW salmon was 68.4 ± 2.1 cm. If sea age itself is heritable (i.e. if the propensity to stay >1 year at sea has a genetic basis), then the h^2 of adult body size as estimated from a univariate animal model will conceptually encompass the effects of genes influencing sea age (which in turn drive size variation among sea-age classes), as well as, genes influencing body size variation within sea-ages. To capture this heritable effect, a new variable “sea-age-corrected body-length”, (denoted $\hat{S}_{i,a}$), was constructed as follows:

$$\hat{S}_{i,a} = S_{i,a} - \bar{S}_a,$$

where $S_{i,a}$ corresponds to the log body-length of individual i of sea-age category a (1SW or MSW) and \bar{S}_a is the mean log-body-length of all individuals of sea-age a . A univariate animal model of $\hat{S}_{i,a}$ was then run (UV model 2) with the same MCMCglmm settings and model structure as UV model 1 to estimate h^2 of body-length conditional on sea-age.

The trait sea-age was treated as a binary variable, with a value of 0 corresponding to maturation after one sea-winter and a value of 1 to maturation after two or more sea-winters (178 of the records in the dataset were 1SW adults; among the MSW adults, only 5 out of 323 spent three winters at sea, and only one fish spent four winters at sea). A probit link function (family = “threshold” in *MCMCglmm*) was used, where the resulting residual variance refers to the variance of the link function, here fixed at 1 in order to render V_A estimable. The h^2 reported from this univariate model of sea-age (UV model 3) thus refers to h^2 on the liability scale, rather than the observed binary scale (see Appendix B.2).

In the next step, bivariate animal models (both traits included as response variables) were used to partition the overall phenotypic (co)variance between body-length and sea-age-at-maturity into additive genetic versus environmental (residual) components. Maternal effects and year effects were not included in these models (see Appendix B.1). In the first bivariate animal model (BV model 1), trait 1 was log adult body-length and trait 2 was binary sea-age. Unstructured covariance matrices for the random effects were specified, in order to provide estimates of COV_A and the residual covariance, COV_R . The additive genetic correlation (r_G) between body-length and sea-age was calculated by dividing COV_A by the product of the square root of V_A for each trait. Similarly, the environmental (residual) correlation (r_E) was calculated by dividing the residual covariance by the product of the square root of V_R for each trait. Statistical support for r_G and r_E was assessed by checking whether the 95% highest posterior density (HPD) interval for each included zero.

A second bivariate animal model (BV model 2) was run where this time the two response variables were sea-age-corrected body length (on the log scale, i.e. $\hat{S}_{i,a}$) and sea-age. This effectively corrected adult body size for variation due to sea-age, and the resulting r_G and r_E estimates from BV model 2 correspond to the genetic and environmental correlations (respectively) between “body size-within-sea-age” and sea age.

All quantitative genetic parameter estimates are reported as posterior means \pm HPD intervals.

Testing for microevolutionary trends

In a quantitative genetic sense, microevolutionary responses are apparent when estimated breeding values (estimates of the additive genetic “merit” of an individual for a given trait) exhibit a temporal trend at the population level. We used the procedure recommended by Hadfield et al. (2010 b) to test for evolutionary change in body-length corrected for sea-age, and in sea-age itself (see Appendix B.4 for full details).

RESULTS

Population composition and phenotypic trends

During our 1991-2011 study, a total of 1,733 anadromous adults came to the Girnock to spawn [annual average 75.3 (range 29 to 144, total adults)], of which the great majority (97%, range 88% to 100%) in every year, were of known sex, sea-age and river-age. The annual proportion of returning adults that had prior-clipped adipose fins, indicating that they were both reared above the Girnock trap and trapped as juvenile emigrants, averaged 47% (range 31% to 76%). The remaining 53% of unclipped returning adults are assumed to have either: 1) passed over the smolt trap and not been marked; or, more likely, (2) to have been reared below the Girnock trap. If reared below the Girnock trap, they were probably reared outside of the Girnock catchment (only 18% of fully-grown parr are estimated to be reared within the Girnock catchment but also below the Girnock trap, a region comprising just 8% of the Girnock’s wetted area (Glover pers comm.)). An annual average of 47% of known-sex

adults were females (range 30% to 60%; male% \approx 100% - female%). During the sub-period of ova-manipulation (2000 to 2010) an average of 80% of females (range 48% to 100%) were artificially spawned. Fluctuations and/or trends in population numbers and composition occurred (see Discussion).

Molecular genetics and pedigree reconstruction

The rate of genotyping success was high, with 92% of all individual samples amplifying for at least 14 or more marker loci (i.e. 82% of screened markers per sample). Full genotype information (17 loci) was obtained for 83% of all samples screened. Loci summary statistics are detailed in Appendix A.2. Across the 17 loci, excluding the sex marker, there were, on average across the 20 cohorts, 11.5 alleles per locus per year (\pm 7.9 S.D., range 2 to 47). Within annual cohorts, observed heterozygosity values averaged 0.748 across loci per year (\pm 0.178 S.D., range 0.150 to 1.000). Allelic richness within loci was similar over all the study years. There was no evidence for departures from Hardy-Weinberg expectations (HWE) or linkage disequilibrium from any cohort.

Power analysis indicates that the marker panel (17 loci) used provides over 99% accuracy to identify putative offspring to family. There is only a very minor decrease in power when 14+ markers are considered with some variation related to the particular marker loci also being noted (i.e. high versus moderately polymorphic marker).

From some 1,600 records of adult genotypes available for paternity assignment during 1991-2011, 273 full parent-offspring trios (mother, father, offspring) were discerned and retained after quality-controlling. 1991 to 1995 and 2007 to 2011 have records of incomplete families (not all eventual 'older' offspring will have returned and been sampled). Inclusion of such records could bias interpretation of the data. The retained complete parentage-records (both

parents and all offspring of varying river- and sea-ages) mainly spanned the years 1995~2006; these years happened also to span a period of low returning adult numbers. Consequently, the estimated annual values of sea-age and body-length are particularly variable.

In order to retain only those parent-offspring trio records that were informative for the quantitative genetic analyses, trios were excluded in which any of the involved individuals were repeat-spawners (i.e. had spawned in previous years); the sex of either parent was unknown; body-length was not measured; or neither sea-age nor river-age were readable from scales (see below). This rigorous filtering reduced the number of useable trios from an initial 336 to 276 (82% retained).

After such filtering, the years of spawning for both putative parents matched in every case. For one of the 276 trios, a missing value for the offspring's sea-age was estimated based on its year of spawning, its known river-age, and the pedigree-determined year of spawning of its parents. Similarly, for 14 of the 276 trio records, the offspring's river-age was estimated from its year of spawning, its known sea-age, and the pedigree-determined year of spawning of its parents. The offspring's total scale-read age closely matched its pedigree-determined total-age in most cases: in 78% of cases (216/276 trios) the match was exact; in 19% the age-comparisons differed by only a single year (53/276); the remaining seven age-discrepancies comprised four records where age-comparisons differed by 2-years; one at 3-years; and two at 5-years. The latter three records involving offspring-ageing discrepancies of >2 years were removed, resulting in 273 rigorously useable parental-trio records.

Further details on the pedigree are given in Table 1 and Appendix B.1.

Animal models

Convergence was good for all animal models (results not shown). The univariate model of body-length (UV model 1) showed that additive genetic effects (V_A) accounted for approximately one-quarter of the total phenotypic variance (Table 2, h^2 of log adult body-length = 0.27). Of the remaining variance (i.e. not including V_A), maternal effects (V_M) accounted for approximately 13% (9% of the total phenotypic variance V_P), year-to-sea effects (V_Y) for 15% (11% of V_P) and residual environmental effects (V_R) for 72% (53% of V_P). Males were on average smaller (61.62 ± 0.58 cm) than females (66.10 ± 0.33 cm). When body-length was corrected for sea-age (UV model 2), V_A then accounted for a considerably smaller fraction of V_P (Table 2, h^2 of body length within sea-ages = 0.14). V_M accounted for approximately 14% of V_P , V_Y for 9%, and V_R accounted for the remaining 64%. The univariate animal model of sea-age (UV model 3) showed that V_A accounted for a little over half of V_P (Table 2, h^2 of sea-age = 0.51). Of the remaining variance (i.e. not including V_A), V_M accounted for approximately 10% (5% of V_P), V_Y for 44% (20% of V_P) and V_R for 46% (21% of V_P).

The results of the bivariate animal model of adult body length and sea-age (BV model 1) mirrored those of the univariate animal models in terms of the estimated heritabilities for the two traits (h^2 of adult body length = 0.26 and h^2 of sea-age = 0.48 in BV model 1; Table 3). The estimated genetic correlation between these traits was very high in the bivariate model where body length was not adjusted for sea age ($r_G = 0.96$; Table 3). The results of the bivariate animal model of sea-age-corrected body length and sea-age (BV model 2) also largely mirrored those of the univariate models in term of the estimated heritabilities for the two traits, although the h^2 of sea-age-corrected body length was somewhat lower (h^2 of sea-age-corrected body length = 0.06 and h^2 of sea-age = 0.53 in BV model 2; Table 3). The estimated genetic correlation between these traits was positive ($r_G = 0.33$) but the HPD

intervals overlapped zero (Table 3), indicating that the r_G was not significantly different from zero.

A power analysis showed that the Girnock data had sufficient power to detect $h^2 > 0.22$ with >80% confidence with no inherent bias in the data-set (Fig.2).

Testing for microevolutionary trends

The temporal trend in estimated breeding values for the trait “sea-age corrected body-length” was estimated at -0.0001 log(cm)/year (95% quantiles: -0.0008 to 0.0004). This trend in estimated breeding values was not statistically significantly negative, as only 67.4% of the posterior distribution of estimated temporal slopes was less than zero. The probability that this very small trend in estimated breeding values was more negative than one would expect based on random genetic drift was 63.3%, again indicating a lack of any evidence for a microevolutionary response to selection. .

The temporal trend in estimated breeding values for the trait sea-age was estimated at -0.017 liability units per year (95% quantiles: -0.053 to 0.009). This trend in estimated breeding values was not statistically significantly negative, as only 81.8% of the posterior distribution of estimated temporal slopes was less than zero. The probability that this trend was more negative than one would expect based on random genetic drift was 74.3%, again indicating a lack of solid evidence for a microevolutionary response to selection.

DISCUSSION

Our data from a well-studied, single, wild population of Atlantic salmon show that sea-age and body-length are both quite strongly heritable and, importantly, that the two traits are strongly genetically correlated. However, when sea-age is alternatively correlated to body-

length *within* each sea-age class (i.e. sea-age corrected body-length), the genetic correlation is then not significantly different from zero, and nor is the h^2 for sea-age-corrected body-length, implying that individual variation in size within sea-ages does not have a heritable basis. Most studies estimate quantitative genetic parameters in laboratory, or in the case of fish, aquaculture settings, which may not reveal much about evolutionary potential under wild natural conditions (Charmantier and Garant 2005). Our results thus add important general information regarding the potential for wild fish populations to respond to natural selection (or “unnatural selection”, c.f. Allendorf & Hard 2009) and complement recent findings showing strong genetic components to age and size at maturity in Atlantic salmon (e.g. Ayllon et al. 2015, Barson et al. 2015, Lepais et al. 2017).

Inheritance patterns

Substantial amounts of additive genetic variance in sea-age were not entirely unexpected, given that recent studies of Atlantic salmon discovered several genetic loci underpinning variation in sea-age, including a single gene (*VGLL3*) that explains almost 40% of the total phenotypic variation (Ayllon et al. 2015, Barson et al. 2015). These studies used a very different approach – genome-wide associations combining data from a large number of populations across a broad geographic region – and homed-in on specific quantitative trait loci (QTL), whereas we followed a classical quantitative genetics approach that treats the underlying genetic architecture as a “black box” and focussed on a single population. The concordant inferences between studies conducted at very different scales using different methodologies – namely, that sea-age variation in wild Atlantic salmon populations has a strong genetic basis – points towards a general finding for the species. Studies of other fish species have also documented heritable differences in age-at-maturity within and among populations (e.g. mosquitofish *Gambusia affini* introduced to Hawaii, (Stearns 1983); guppies *Poecilia reticulata* in Trinidad, (Reznick et al. 1990); Chinook salmon *Oncorhynchus*

tshawytscha introduced to New Zealand, (Quinn et al. 2001). This raises interesting questions regarding the evolutionary forces that maintain genetic variation in this key life-history trait, which may include sex-dependent dominance (Barson et al. 2015), frequency-dependent selection and spatiotemporal variation in environmental selection pressures (Gurney et al. 2012). In this study we ignored potential sex-specific inheritance patterns, which is justified if inter-sex genetic correlations for our traits of interest are positive and high, but this assumption should be tested in future studies, given sufficient statistical power. It is also possible that the period of captive propagation might have had an influence on the results. The crossing protocol, which involved mating multiple males to females and vice versa, may have increased the number of links in the pedigree beyond that which might have occurred in the absence of captive propagation, in which case it could have afforded increased power and accuracy to our animal models. On the other hand, this represented a deviation from natural spawning behaviours and mate choice, which may have affected the expression of genetic variation (Pigliucci 2006), but we have no way of knowing *a priori* the direction and magnitude of such influences.

Despite these strong genetic effects, environmental influences clearly also play a role; in our study, for example, approximately half of the variation in sea-age was attributable to environmental variation, although some of this may have been due to non-additive genetic effects (dominance and epistasis) which we could not separate out with our pedigree structure (indeed, Barson et al. 2015 showed that dominance effects occur at the *VGLL3* locus). To put this in practical terms, we observed over our study period that of the 114 offspring in total produced by matings between two MSW parents, 84 (74%) of the resulting offspring themselves returned as MSW; thus a high heritability “biases” this probability towards “like producing like”. Friedland & Hass (1996) found that the fraction of 1SW adult returns from a hatchery-dependent stock of Atlantic salmon was positively associated with late summer

marine growth, implying that age-at-maturity responds plastically to marine environmental conditions. The literature suggests that freshwater conditions may also affect the sea-age of anadromous Atlantic salmon, as implied by positive correlations between smolt size and the proportion of fish returning after 1SW (O’Connell and Ash 1993, Salminen 1997). Similarly, inverse relationships between freshwater age and ocean age have been found in wild steelhead trout *Oncorhynchus mykiss* (Ward and Slaney 1988). However, if sub-populations are spatially structured, as in the North Esk in Scotland (Gurney et al., 2015; Bacon et al. 2012), with higher fractions of 1SW fish in lower parts of the catchment, where smolts also tend to get larger quicker (e.g. due to better growth at higher temperatures), then such correlations would arise from the spatial structuring alone, even if freshwater conditions have no causal effect on sea-age variation. Thus it remains unclear to what extent such patterns reflect plastic responses, spatial structuring, or genetic correlations among traits expressed at different life stages.

Genetic covariance among suites of growth-related and life-history traits have been demonstrated in rainbow/steelhead trout (*Oncorhynchus mykiss*) (Hecht et al. 2015) and brook charr (*Salvelinus fontinalis*) (Thériault et al. 2007), while some QTL for early male maturation status in Atlantic salmon also collocated with those affecting spring weights of juveniles (Lepais et al. 2017). We lacked individual-level data on freshwater or marine growth rates, and hence could not test explicitly for plastic effects of specific traits/cues on sea-age, nor could we test for genotype-by-environment interactions, which may be important here. Lepais et al. (2017) recently demonstrated that early male maturation in Atlantic salmon emerges as an interaction between individual growth rate (an environmentally-sensitive status trait) and a genetically variable maturation threshold (underpinned by at least three detectable QTL), using a powerful new approach called the ‘latent environmental threshold model’ (LETM). It would be very interesting to apply the LETM approach to future studies of sea-

age variation in Atlantic salmon or other anadromous fish; obtaining individual-level data on relevant status traits, such as post-smolt growth, that may act as cues for sea-age decisions will be challenging in many situations, but circuli spacing on scales could be used, for example (e.g. Friedland & Hass 1996). Furthermore, it would be very interesting to dissect the mechanisms by which genetic variation in sea-age comes about; for example, Scottish-origin Atlantic salmon that eventually become MSW evidently use different oceanic feeding areas (e.g. the waters off Greenland) than those that adopt the 1SW life history (which are not recorded as going to Greenland, e.g. Malcolm et al. 2010), raising the intriguing possibility that genetic variation in migration routes and/or destinations might partially explain variation in sea-age, although the correlation between sea-age and migration patterns may not reflect any causal connection between these traits.

The significant positive genetic correlation we documented between sea-age and overall body-length is also consistent with the results of Barson et al. (2015), who documented pleiotropic effects of specific genomic regions on sea-age and size-at-maturity. Such pleiotropic effects are inevitable to a degree, in that alleles predisposing individuals towards later maturation at older sea-ages should also produce larger fish, given that Atlantic salmon keep growing the longer they remain at sea. A more surprising result was the fact that the genetic correlation between sea-age and body-length was so high ($r_G = 0.96$) as to imply that these are effectively the same trait at a genetic level. That does not mean, however, that body size is completely genetically-determined and indeed, in our case, it appears that individual variation in marine growth – a key determinant of body size at breeding – was predominantly environmental, rather than genetic, in origin. This is because sea-age-corrected body-length exhibited a low h^2 that did not differ significantly from zero (although we only had the statistical power to detect $h^2 > \text{ca. } 0.22$, so a true h^2 of less than this cannot be ruled out). While this trait is not a direct measure of marine growth, Atlantic salmon put on

most of their growth at sea (smolt sizes are very small, see URL4) and thus variation in marine growth rates and/or variation in coastal return-times must explain a large fraction of the variation in size of returning adults within each sea-age class. Smolt size could also indirectly account for some of the variation in adult size if, for example, larger smolts grow faster at sea (e.g. due to feeding advantages of initially larger size within shoals of post-smolts, which would set up a positive feedback).

A lack of (strong) genetic influences on marine growth is somewhat unexpected, given that genetic variation could operate via a range of mechanisms here; e.g. some genotypes could be better foragers, or target different prey types, or have different inherited marine migration pathways or destinations, that expose them to more or less growth opportunity, or be more efficient at converting food into somatic growth, etc. Moreover, return migration timing has been shown to have a heritable basis in Atlantic salmon (Stewart et al. 2002; Cauwelier et al. In Press) and other salmonids (Smoker et al. 1998, Quinn et al. 2000, O'Malley and Banks 2008, Kovach et al. 2012). This alone could produce corresponding genetic variation in body-length within sea-age classes, unless return periods were very short or if late-returning genotypes have lower marine growth rates than early-returning genotypes, such that final size differences among them are minimal. At present we can only speculate on this, as we do not have reliable information on coastal return-dates to estimate the h^2 of that trait and potential genetic correlations with body size, or indeed sea-age. Our tentative conclusion at this point is that environmental drivers of body-length variation within each sea-age class are simply much larger in magnitude than any genetic influences.

Our tests for microevolution during the period 1990~2012 did not reveal any genetically based trends towards reduced average sea-age, nor towards smaller body-size. However, over the longer period 1966~2016, Atlantic salmon from two Deeside streams (the

Girnock and Baddoch) have shown a significant decline in MSW numbers whilst 1SW numbers remained stable, implying a significant decrease in average sea-age ratios, slightly reducing the average sea-age; they also showed concomitant downward trends in body size, which were coherent for both 1SW and MSW fish, (Glover et al. 2018 and *pers comm*). It may simply be that the phenotypic trends documented by Glover et al. are entirely driven by phenotypic plasticity, i.e. environmental influences on average sea-age and body size, rather than microevolutionary responses to any directional selection pressures that might have occurred over this period. Mortality pressures from both marine fisheries and freshwater angling were strong on Scottish Atlantic salmon during the period 1960 to 1990, which conceivably could have led to fisheries-induced selection (FIS) on, and subsequent evolution of, sea-age and/or body size. The data analysed in this study fall mainly into a period when marine fishery pressures were much reduced and freshwater angling pressures decreasing. As evolutionary recovery of phenotypic traits from FIS is expected to be much slower than declines caused by it (Heino et al. 2015), our findings with respect to microevolution could simply reflect a period when any real changes were quite small and too low for our sample sizes to detect. Moreover, the current study was focussed on a shorter time period and involved smaller sample sizes than the Glover et al. study, and thus the power to detect a real but small-magnitude evolutionary response may have been limiting. A further complicating factor is that broad-scale climatic changes also occurred over this period, which may have contributed to the observed phenotypic trends in Girnock salmon (Glover et al. 2018), either via phenotypic plasticity, microevolution, or both. Given the apparently very low heritability of body size within sea-ages, phenotypic plasticity – possibly driven by climate change – represents the most parsimonious explanation for the observed length decreases in 1SW and MSW Girnock salmon. The fact that these trends are parallel between sea-ages, despite differing marine feeding zones, may point towards coherent climate effects acting across a

large geographic scale, but factors other than climate may of course be at play. A clearer understanding of the relative roles of microevolution and plasticity and their environmental/anthropogenic drivers could be achieved by extending this pedigree study over a longer period to match the longer-term phenotypic data..

ACKNOWLEDGEMENTS

The opportunity for such genetic work was first realised by Alan Youngson, John Taggart and Eric Verspoor, whom we thank for initiating the sample-collection. John Taggart kindly provided tissue, and some DNA-extract aliquots, from the earlier years. We thank FFL field staff over two decades, particularly Iain Malcolm, Iain McLaren, Alastair Thorne, Mike Donaghy and Ross Glover, for collecting the data, scales and tissue samples. Alastair Thorne, Iain McLaren and Ross Glover read the scales and Aya Thorne prepared scales for DNA analysis. Iain Malcolm and Ross Glover collated and quality-controlled the biometric data. Iain Malcolm, Ross Glover and Stuart Middlemas made constructive comments on earlier versions of the manuscript. Data collection at the Girnock is funded by the Scottish Government, through Marine Scotland Science Freshwater Fisheries Laboratory under Service Level Agreement FW01t. TR was supported by a European Research Council (ERC) Starting Grant (639192), CRP by the Academy of Finland (grant numbers 302873, 307593). PMcG, PP and CB were supported by the Beaufort Marine Research award in Fish Population Genetics funded by the Irish Government under the Sea Change Programme.

576 Author Contributions

577 The Girnock Atlantic Salmon Pedigree project was envisaged by PJB and organised by him
578 and PMcG, in collaboration with PP, JG and CRP. FFL contributed the tissue-samples and
579 biometric-data. DNA extraction and genotyping were undertaken at QUB by CB and PP. PP
580 and PJB merged the biometric and genotype data, PP undertook the parentage-assignments
581 and he and PJB thereby quality-controlled the original genotype scorings. PJB extracted and
582 quality-controlled the merged sub-set of data for this paper, including preliminary analyses.
583 TR fitted and interpreted the animal models. PP wrote the Methods and Results for parentage
584 analyses. TR and PJB wrote the Methods and Results for the quantitative genetic analyses
585 and an outline for the paper. All authors contributed to the final draft

586

587

REFERENCES

- Allendorf, F.W., and Hard, J.J. 2009. Human-induced evolution caused by unnatural selection through harvest of wild animals. *Proc. Natl. Acad. Sci.* **106**(Supplement 1): 9987–9994.
- Ayllon, F., Kjærner-Semb, E., Furmanek, T., Wennevik, V., Solberg, M.F., Dahle, G., Taranger, G.L., Glover, K.A., Almén, M.S., and Rubin, C.J. 2015. The *vgl13* locus controls age at maturity in wild and domesticated Atlantic salmon (*Salmo salar* L.) males. *PLoS Genet* **11**(11): e1005628.
- Bacon, P.J., Gurney, W.S., McKenzie, E., Whyte, B., Campbell, R., Laughton, R., Smith, G., and MacLean, J. 2010. Objective determination of the sea age of Atlantic salmon from the sizes and dates of capture of individual fish. *ICES J. Mar. Sci.* **68**(1): 130–143.
- Bacon, P.J., MacLean, J.C., Malcolm, I.A., and Gurney, W.S.C. 2012. Ova fecundity in Scottish Atlantic salmon *Salmo salar*: predictions, selective forces and causal mechanisms. *J. Fish Biol.* **81**(3): 921–938.
- Bacon, P.J., Palmer, S.C.F., MacLean, J.C., Smith, G.W., Whyte, B.D.M., Gurney, W.S.C., and Youngson, A.F. 2009. Empirical analyses of the length, weight, and condition of adult Atlantic salmon on return to the Scottish coast between 1963 and 2006. *ICES J. Mar. Sci.* **66**(5): 844–859.
- Barson, N.J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G.H., Fiske, P., Jacq, C., Jensen, A.J., Johnston, S.E., and Karlsson, S. 2015. Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*. **528**(7582): 405–408.
- Brommer, J.E., Merilä, J., and Kokko, H. 2002. Reproductive timing and individual fitness. *Ecol. Lett.* **5**(6): 802–810.

613 Butler, D., Cullis, B.R., Gilmour, A.R., and Gogel, B.J. 2015. Mixed models for S language
614 environments. ASReml-R reference manual. ASReml estimates variance components
615 under a general linear mixed model by residual likelihood (REML). Version 3.0.
616 Queensland, Australia.

617 Cauwelier, E., Gilbey, J., Sampayo, J., Stradmeyer, L. & Middlemas, S. (In press).
618 Identification of a single genomic region associated with seasonal river return timing
619 in adult Scottish Atlantic salmon (*Salmo salar* L.) identified using a genome-wide
620 association study. Can. J. Fish. Aq. Sci.

621 Carlson, S.M., and Seamons, T.R. 2008. A review of quantitative genetic components of
622 fitness in salmonids: implications for adaptation to future change. Evol. Appl. **1**(2):
623 222–238.

624 Charlesworth, B. 1994. Evolution in age-structured populations. Cambridge University Press
625 Cambridge.

626 Charmantier, A., and Garant, D. 2005. Environmental quality and evolutionary potential:
627 lessons from wild populations. Proc. R. Soc. B Biol. Sci. **272**(1571): 1415–1425.

628 Clutton-Brock, T., and Sheldon, B.C. 2010. Individuals and populations: the role of long-
629 term, individual-based studies of animals in ecology and evolutionary biology. Trends
630 Ecol. Evol. **25**(10): 562–573.

631 Dickerson, B.R., Willson, M.F., Bentzen, P., and Quinn, T.P. 2005. Heritability of life history
632 and morphological traits in a wild pink salmon population assessed by DNA
633 parentage analysis. Trans. Am. Fish. Soc. **134**(5): 1323–1328.

634 Enberg, K., Jorgensen, C., Dunlop, E. S., Heino, M. & Dieckmann, U. 2009. Implications of
635 fisheries-induced evolution for stock rebuilding and recovery. Evol. Appl. **2**: 394-414.

636 Gjerde, B. 1984. Response to individual selection for age at sexual maturity in Atlantic
637 salmon. Aquaculture **38**(3): 229–240.

638 Glover, R. S., Fryer, R.J., Bacon, P.J., Soulsby, C., Malcolm, I.A. 2018. Do trends in the
 639 size of wild female Atlantic salmon have a substantial effect on egg deposition? Fish.
 640 Man. and Ecol. **25**: 158-161. <https://doi.org/10.1111/fme.12273>
 641 Gurney, W.S., Bacon, P.J., Speirs, D.C., McGinnity, P., and Verspoor, E. 2012. Sea-age
 642 variation in maiden Atlantic salmon spawners: phenotypic plasticity or genetic
 643 polymorphism? Bull. Math. Biol. **74**(3): 615–640.
 644 Gurney, W.S.C., Bacon, P.J., Malcolm, I.A., MacLean, J.C., and Youngson, A.F. 2015.
 645 Atlantic salmon (*Salmo salar*) population as discerned for an eastern Scottish river.
 646 Scott. Mar. Freshw. Sci. **6**(12): 1–72. doi:10.7489/1662-
 647 Hadfield, J.D. 2010. MCMC methods for multi-response generalized linear mixed models:
 648 the MCMCglmm R package. J. Stat. Softw. **33**(2): 1–22.
 649 Hadfield, J.D., Wilson, A.J., Garant, D., Sheldon, B.C., and Kruuk, L.E. 2010. The misuse of
 650 BLUP in ecology and evolution. Am. Nat. **175**(1): 116–125.
 651 Hankin, D.G., Nicholas, J.W., and Downey, T.W. 1993. Evidence for inheritance of age of
 652 maturity in chinook salmon (*Oncorhynchus tshawytscha*). Can. J. Fish. Aquat. Sci.
 653 **50**(2): 347–358.
 654 Harvey, A. C., Tang, Y., Wennevik, V., Skaala, Ø. & Glover, K. A. 2017. Timing is
 655 everything: Fishing- season placement may represent the most important angling-
 656 induced evolutionary pressure on Atlantic salmon populations. Ecol. and Evol. **7**:
 657 7490-7502.
 658 Hecht, B.C., Hard, J.J., Thrower, F.P., and Nichols, K.M. 2015. Quantitative genetics of
 659 migration-related traits in rainbow and steelhead trout. G3 Genes Genomes Genet.
 660 **5**(5): 873–889.
 661 Heino, M., and Kaitala, V. 1999. Evolution of resource allocation between growth and
 662 reproduction in animals with indeterminate growth. J. Evol. Biol. **12**(3): 423–429.

663 Heino, M., Diaz Pauli, B., and Dieckmann, U. 2015. Fisheries-induced evolution. *Ann. Rev.*
 664 *Ecol. Evol. Syst.*, **46**: 461-480.

665 Hutchings, J.A. and Jones, M.E., 1998. Life history variation and growth rate thresholds for
 666 maturity in Atlantic salmon, *Salmo salar*. *Can. J. Fish. Aq. Sci.*, **55**(S1): 22-47.

667 Johnston, S.E., Orell, P., Pritchard, V.L., Kent, M.P., Lien, S., Niemelä, E., Erkinaro, J., and
 668 Primmer, C.R. 2014. Genome-wide SNP analysis reveals a genetic basis for sea-age
 669 variation in a wild population of Atlantic salmon (*Salmo salar*). *Mol. Ecol.* **23**(14):
 670 3452–3468.

671 Kause, A., Ritola, O., Paananen, T., Mäntysaari, E., and Eskelinen, U. 2003. Selection against
 672 early maturity in large rainbow trout *Oncorhynchus mykiss*: the quantitative genetics
 673 of sexual dimorphism and genotype-by-environment interactions. *Aquaculture* **228**(1):
 674 53–68.

675 Keenan, K., Bradley, C.R., Magee, J.J., Hynes, R.A., Kennedy, R.J., Crozier, W.W., Poole,
 676 R., Cross, T.F., McGinnity, P., and Prodöhl, P.A. 2013a. Beaufort trout MicroPlex: a
 677 high-throughput multiplex platform comprising 38 informative microsatellite loci for
 678 use in resident and anadromous (sea trout) brown trout *Salmo trutta* genetic studies. *J.*
 679 *Fish Biol.* **82**(6): 1789–1804. doi:10.1111/jfb.12095.

680 Keenan, K., McGinnity, P., Cross, T.F., Crozier, W.W., and Prodöhl, P.A. 2013b. diveRsity:
 681 An R package for the estimation and exploration of population genetics parameters
 682 and their associated errors. *Methods Ecol. Evol.* **4**(8): 782–788.

683 Klemetsen, A., P-A. Amundsen, J. B. Dempson, B. Jonsson, N. Jonsson, M. F. O'Connell,
 684 and Mortensen, E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L.
 685 and Arctic charr *Salvelinus alpinus* (L.): a review of aspects of their life histories."
 686 *Ecol. Fresh. Fish* 12(1): 1-59.

687 Kovach, R.P., Gharrett, A.J., and Tallmon, D.A. 2012. Genetic change for earlier migration
688 timing in a pink salmon population. *Proc. R. Soc. Lond. B Biol. Sci.* **279**(1743):
689 3870–3878.

690 Kuparinen, A. & Hutchings, J. A. 2017. Genetic architecture of age at maturity can generate
691 divergent and disruptive harvest-induced evolution. *Phil. Trans. Roy. Soc. B:*
692 *Biological Sciences* **372** (1712): 20160035.

693 Lepais, O., Manicki, A., Glise, S., Buoro, M., and Bardonnnet, A. 2017. Genetic architecture
694 of threshold reaction norms for male alternative reproductive tactics in Atlantic
695 salmon (*Salmo salar* L.). *Sci. Rep.* **7**.

696 Malcolm, I. A., Godfrey, J., and Youngson, A. F. 2010. Review of migratory routes and
697 behaviour of Atlantic salmon, sea trout and European eel in Scotland’s coastal
698 environment: implications for the development of marine renewables. *Scottish Marine*
699 *and Freshwater Science*, **1**(14): 1-72.

700 Morrissey, M.B., and Ferguson, M.M. 2011. A test for the genetic basis of natural selection:
701 an individual-based longitudinal study in a stream-dwelling fish. *Evolution* **65**(4):
702 1037–1047.

703 Munday, P.L., Warner, R.R., Monro, K., Pandolfi, J.M., and Marshall, D.J. 2013. Predicting
704 evolutionary responses to climate change in the sea. *Ecol. Lett.* **16**(12): 1488–1500.

705 Nævdal, G. 1983. Genetic factors in connection with age at maturation. *Aquaculture* **33**(1):
706 97–106.

707 Naish, K.A., and Hard, J.J. 2008. Bridging the gap between the genotype and the phenotype:
708 linking genetic variation, selection and adaptation in fishes. *Fish Fish.* **9**(4): 396–422.

709 O’Connell, M.F., and Ash, E.G.M. 1993. Smolt size in relation to age at first maturity of
710 Atlantic salmon (*Salmo salar*): the role of lacustrine habitat. *J. Fish Biol.* **42**(4): 551–
711 569.

712 Olafsson, K., Hjorleifsdottir, S., Pampoulie, C., Hreggvidsson, G.O., and Gudjonsson, S.
 713 2010. Novel set of multiplex assays (SalPrint15) for efficient analysis of 15
 714 microsatellite loci of contemporary samples of the Atlantic salmon (*Salmo salar*).
 715 Mol. Ecol. Resour. **10**(3): 533–537.

716 O'Malley, K.G., and Banks, M.A. 2008. A latitudinal cline in the Chinook salmon
 717 (*Oncorhynchus tshawytscha*) Clock gene: evidence for selection on PolyQ length
 718 variants. Proc. R. Soc. Lond. B Biol. Sci. **275**(1653): 2813–2821.

719 Pigliucci, M. 2006. Genetic variance–covariance matrices: a critique of the evolutionary
 720 quantitative genetics research program. Biol. and Phil. **21**(1): 1–23.

721 Quinn, T.P., Kinnison, M.T., and Unwin, M.J. 2001. Evolution of chinook salmon
 722 (*Oncorhynchus tshawytscha*) populations in New Zealand: pattern, rate, and process.
 723 Genetica **112**(1): 493–513.

724 Quinn, T.P., Unwin, M.J., and Kinnison, M.T. 2000. Evolution of temporal isolation in the
 725 wild: genetic divergence in timing of migration and breeding by introduced chinook
 726 salmon populations. Evolution **54**(4): 1372–1385.

727 Reed, T.E., Prodöhl, P., Hynes, R., Cross, T., Ferguson, A., and McGinnity, P. 2015.
 728 Quantifying heritable variation in fitness-related traits of wild, farmed and hybrid
 729 Atlantic salmon families in a wild river environment. Heredity. **115**(2): 173–184.

730 Reznick, D.A., Bryga, H., and Endler, J.A. 1990. Experimentally induced life-history
 731 evolution in a natural population. Nature **346**(6282): 357–359.

732 Roff, D.A. 2002. Life History Evolution. 1 edition. Sinauer Associates, Inc., Sunderland,
 733 Mass.

734 Salminen, M. 1997. Relationships between smolt size, postsmolt growth and sea age at
 735 maturity in Atlantic salmon ranches in the Baltic Sea. J. Appl. Ichthyol. **13**(3): 121–
 736 130.

737 Serbezov, D., Bernatchez, L., Olsen, E.M., and Vøllestad, L.A. 2010. Quantitative genetic
 738 parameters for wild stream-living brown trout: heritability and parental effects. J.
 739 Evol. Biol. **23**(8): 1631–1641.

740 Smoker, W.W., Gharrett, A.J., and Stekoll, M.S. 1998. Genetic variation of return date in a
 741 population of pink salmon: a consequence of fluctuating environment and dispersive
 742 selection. Alsk. Fish. Res. Bull. **5**(1): 46–54.

743 Stearns, S.C. 1983. The evolution of life-history traits in mosquitofish since their introduction
 744 to Hawaii in 1905: rates of evolution, heritabilities, and developmental plasticity. Am.
 745 Zool. **23**(1): 65–75.

746 Stearns, S.C. 1992. The evolution of life histories. Oxford University Press Oxford.

747 Stewart, D.C., Smith, G.W., and Youngson, A.F. 2002. Tributary-specific variation in timing
 748 of return of adult Atlantic salmon (*Salmo salar*) to fresh water has a genetic
 749 component. Can. J. Fish. Aquat. Sci. **59**(2): 276–281.

750 Sydeman, W.J., Poloczanska, E., Reed, T.E., and Thompson, S.A. 2015. Climate change and
 751 marine vertebrates. Science **350**(6262): 772–777.

752 Taggart, J.B. 2007. FAP: an exclusion-based parental assignment program with enhanced
 753 predictive functions. Mol. Ecol. Resour. **7**(3): 412–415.

754 Taggart, J.B., McLaren, I.S., Hay, D.W., Webb, J.H., and Youngson, A.F. 2001. Spawning
 755 success in Atlantic salmon (*Salmo salar* L.): a long-term DNA profiling-based study
 756 conducted in a natural stream. Mol. Ecol. **10**(4): 1047–1060.

757 Thériault, V., Garant, D., Bernatchez, L., and Dodson, J.J. 2007. Heritability of life-history
 758 tactics and genetic correlation with body size in a natural population of Brook Charr
 759 (*Salvelinus fontinalis*). J. Evol. Biol. **20**(6): 2266–2277.

760 Tipping, J.M. 1991. Management briefs: heritability of age at maturity in steelhead. North
 761 Am. J. Fish. Manag. **11**(1): 105–108.

762 Ward, B.R., and Slaney, P.A. 1988. Life history and smolt-to-adult survival of Keogh River
 763 steelhead trout (*Salmo gairdneri*) and the relationship to smolt size. Can. J. Fish.
 764 Aquat. Sci. **45**(7): 1110–1122.

765 Yano, A., Guyomard, R., Nicol, B., Jouanno, E., Quillet, E., Klopp, C., Cabau, C., Bouchez,
 766 O., Fostier, A., and Guiguen, Y. 2012. An immune-related gene evolved into the
 767 master sex-determining gene in rainbow trout, *Oncorhynchus mykiss*. Curr. Biol.
 768 **22**(15): 1423–1428.

769 Yano, A., Nicol, B., Jouanno, E., Quillet, E., Fostier, A., Guyomard, R., and Guiguen, Y.
 770 2013. The sexually dimorphic on the Y-chromosome gene (sdY) is a conserved male-
 771 specific Y-chromosome sequence in many salmonids. Evol. Appl. **6**(3): 486–496.

772

773 **Websites cited:**

774 URL1: [http://www.gov.scot/Topics/marine/Salmon-Trout-](http://www.gov.scot/Topics/marine/Salmon-Trout-Coarse/Freshwater/Monitoring/Traps)
 775 [Coarse/Freshwater/Monitoring/Traps](http://www.gov.scot/Topics/marine/Salmon-Trout-Coarse/Freshwater/Monitoring/Traps)

776 URL2: [http://www.gov.scot/Topics/marine/Salmon-Trout-](http://www.gov.scot/Topics/marine/Salmon-Trout-Coarse/Freshwater/Monitoring/Traps/AdultReturnsPopStatus)
 777 [Coarse/Freshwater/Monitoring/Traps/AdultReturnsPopStatus](http://www.gov.scot/Topics/marine/Salmon-Trout-Coarse/Freshwater/Monitoring/Traps/AdultReturnsPopStatus)

778 URL3: <http://data.marine.gov.scot/dataset/girnock-and-baddoch-adult-returns-traps>

779 Citation: Glover, R., Malcolm, I. 2015. Girnock and Baddoch: Adult Returns to the Traps.
 780 DOI: 10.7489/1588-1

781 URL4: [http://www.gov.scot/Topics/marine/Salmon-Trout-](http://www.gov.scot/Topics/marine/Salmon-Trout-Coarse/Freshwater/Monitoring/Traps/JuvenileEmigrants)
 782 [Coarse/Freshwater/Monitoring/Traps/JuvenileEmigrants](http://www.gov.scot/Topics/marine/Salmon-Trout-Coarse/Freshwater/Monitoring/Traps/JuvenileEmigrants)

783 Citation: Glover, R., Malcolm, I. 2015. Girnock and Baddoch: Emigrant Numbers by Year of
 784 Emigration. DOI: 10.7489/1017-1

785

786 **Appendices:**

787 Appendix A:

788 A.1 Microsatellite marker information

789 A.2 Summary sample statistics per locus / cohort

790 Appendix B:

791 B.1 Details on the reconstructed pedigree

792 B.2 Full details on the estimation procedures for the quantitative genetic parameters

793 B.3 Power analyses for heritability estimation

794 B.4 Testing for microevolutionary trends

795

Table 1 Summary of annual sample sizes of adult Atlantic salmon within the reconstructed Girnock pedigree. Column 2 shows the total number of individual fish spawning in that year (Column 1) that appeared either as parents or offspring within “parent-offspring trios” (where a trio represents one offspring and both its genetically-assigned parents). For example, an adult returning in 1996 could be recorded as being in a trio in that year (i) because it mated with another fish in that year and produced at least one surviving offspring that itself was DNA-sampled as a spawning adult (e.g. six years later in 2002); (ii) because it was itself assigned two parents (which might have spawned and been DNA-sampled in 1991, for example), or (iii) for both reasons (i.e. some fish appear in the pedigree as both offspring and parents). Columns 3 and 4 break Column 2 down by sex. Column 5 specifies how many of the fish enumerated in Column 2 were themselves assigned two parents, while Column 6 specifies how many were assigned no parents.

Adult Return Year	Total	Females	Males	Number with both [†] parents known	Number with neither parents known
1991	8	5	3	0 ^{††}	8
1992	9	4	5	0	9
1993	6	3	3	0	6
1994	2	1	1	0	2
1995	13	7	6	0	13
1996	12	8	4	7	5
1997	6	3	3	3	3
1998	10	6	4	6	4
1999	15	6	9	5	10
2000	28	11	17	8	20
2001	10	6	4	5	5
2002	11	3	8	4	7
2003	13	5	8	5	8
2004	51	28	23	9	42
2005	40	25	15	14	26
2006	46	28	18	9	37
2007	23	13	10	5	18
2008	19	7	12	14	5
2009	37	14	23	37	0
2010	60	25	35	60	0
2011	65	42	23	65	0
Grand totals:	484	250	234	256	228

[†] Note that this pedigree study focussed only on “full trios”, i.e. cases where both parents of a given offspring were assigned. An unknown fraction of offspring each year are sired by precociously maturing males, which are not DNA-sampled and therefore cannot be assigned as true fathers, but the numbers here refer only to full parent-offspring trios.

^{††} Note that prior to 1996, the parents of sampled spawning adults were not discernible as these parents would have themselves spawned prior to 1991, when DNA samples were not available.

Table 2 Results of the univariate animal models. The values for adult body-length are on the natural logarithm scale, while the values for sea-age-at-maturity are on the underlying liability scale from the probit model (note the residual variance was fixed at 1). The intercept refers to the fixed effect of the reference sex, females. HPD = highest posterior density.

Model	Parameter	Mean	Lower HPD interval	Upper HPD interval
<i>(UV model 1)</i> Univariate for adult body- length	Intercept (females)	4.197	4.173	4.2223
	Sex: males	-0.072	-0.090	-0.049
	Additive genetic variance V_A	0.0039	0.0016	0.0062
	Maternal variance V_M	0.0014	0.0002	0.0029
	Year variance V_M	0.0016	0.0004	0.0033
	Residual variance V_R	0.0078	0.0058	0.0101
	Heritability h^2	0.265	0.125	0.414
<i>(UV model 2)</i> Univariate for sea-age- corrected body length	Intercept (females)	0.000	-0.0148	0.0136
	Sex: males	0.0179	-0.0048	0.0307
	Additive genetic variance V_A	0.0008	0.0003	0.0015
	Maternal variance V_M	0.0008	0.0003	0.0015
	Year variance V_M	0.0005	0.0001	0.0010
	Residual variance V_R	0.0037	0.0030	0.0045
	Heritability h^2	0.140	0.044	0.261
<i>(UV model 3)</i> Univariate for sea-age-at- maturity	Intercept (females)	2.207	1.299	3.363
	Sex: males	-2.511	-3.647	-1.539
	Additive genetic variance V_A	2.580	0.114	6.143
	Maternal variance V_M	0.212	0.000	0.809
	Year variance V_M	0.937	0.081	2.158
	Residual variance V_R	1.000	1.000	1.000
	Heritability h^2	0.511	0.267	0.734

Table 3 Results of the bivariate animal models of (1) adult body-length and sea-age (binary trait), and (2) sea-age-corrected body length and sea-age. The values for adult body-length and sea-age-corrected body length are on the natural logarithm scale, while the values for sea-age are on the underlying liability scale from the probit model (note that V_R for sea-age was fixed at 1). The fixed effects are not reported as they mirror the findings of the univariate models.

Model	Parameter	Mean	Lower HPD interval	Upper HPD interval
(BV model 1) Bivariate for adult body- length and sea-age	V_A for body length	0.0042	0.0019	0.0065
	V_A for sea-age	1.0914	0.1436	2.2986
	Additive genetic covariance COV_A	0.0633	0.0171	0.1157
	V_R for body length	0.0116	0.0095	0.0138
	V_R for sea-age	1.0000	1.0000	1.0000
	Residual covariance COV_R	0.0635	0.0471	0.0791
	Heritability h^2 for body length	0.2637	0.1261	0.3878
	Heritability h^2 for sea-age	0.4823	0.2174	0.7284
	Genetic correlation r_G	0.9551	0.9096	0.9879
	Environmental correlation r_E	0.5893	0.4586	0.6834
(BV model 2) Bivariate for adult body- length within sea-ages, and sea-age	V_A for body length within sea-ages	0.0004	0.0002	0.0007
	V_A for sea-age	1.5715	0.1436	2.2986
	Additive genetic covariance COV_A	0.0079	-0.0077	0.0256
	V_R for body length within sea-ages	0.0069	0.0061	0.0078
	V_R for sea-age	1.0000	1.0000	1.0000
	Heritability h^2 for body length within sea-ages	0.0560	0.0295	0.0976
	Heritability h^2 for sea-age	0.5347	0.2316	0.8495
	Genetic correlation r_G	0.3334	-0.2604	0.7205





Fig.1 Map of the Girnock Burn study system, a tributary of the River Dee, Aberdeenshire, Scotland.



876

877

878 **Fig.2** Power analysis for heritability estimation for an arbitrary trait given the observed
 879 pedigree and phenotypic data structure, with statistical power (A) and estimated heritability
 880 (mean plus 95% confidence intervals, B) plotted against simulated heritability. Dotted line in
 881 A corresponds to a power of 80%; dotted line in B shows the 1:1 line.