

Title	Advanced glycation end product intake during pregnancy and offspring allergy outcomes: prospective cohort study
Authors	Venter, Carina;Pickett, Kaci;Starling, Anne;Maslin, Kate;Smith, Pete K.;Palumbo, Michaela P.;O'Mahony, Liam;Ben Abdallah, Miriam;Dabelea, Dana
Publication date	2021-10-05
Original Citation	Venter, C., Pickett , K., Starling, A., Maslin, K., Smith, P. K., Palumbo, M. P., O'Mahony, L., Ben Abdallah, M. and Dabelea, D. (2021) 'Advanced glycation end product intake during pregnancy and offspring allergy outcomes: prospective cohort study', Clinical and Experimental Allergy. doi: 10.1111/cea.14027
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1111/cea.14027
Rights	© 2021, John Wiley & Sons, Inc. This is the peer reviewed version of the following item: Venter, C., Pickett , K., Starling, A., Maslin, K., Smith, P. K., Palumbo, M. P., O'Mahony, L., Ben Abdallah, M. and Dabelea, D. (2021) 'Advanced glycation end product intake during pregnancy and offspring allergy outcomes: prospective cohort study', Clinical and Experimental Allergy, doi: 10.1111/cea.14027, which has been published in final form at https://doi.org/10.1111/cea.14027 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Download date	2025-08-03 18:18:05
Item downloaded from	https://hdl.handle.net/10468/12081



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

DR CARINA VENTER (Orcid ID : 0000-0002-7473-5355)

DR KATE MASLIN (Orcid ID : 0000-0002-9768-7245)

Article type : Original Article

Corresponding author mail id : carina.venter@childrenscolorado.org

**Advanced glycation end product intake during pregnancy and offspring allergy outcomes:
prospective cohort study**

Running title: Maternal AGEs intake and offspring allergies

Authors: Carina Venter PhD RD,^{a,b} Kaci Pickett MS,^b Anne Starling PhD^{c,d}, Kate Maslin, PhD RD,^e Pete K. Smith MD^{f,g}, Michaela P. Palumbo MS^c, Liam O'Mahony PhD^h, Miriam Ben Abdallah MD^b, Dana Dabelea MD PhD^{b,c,d}

Affiliations:

- a. Section of Allergy & Immunology, Department of Pediatrics, Children's Hospital Colorado, University of Colorado School of Medicine, Aurora, Colorado
- b. Department of Pediatrics, University of Colorado School of Medicine, University of Colorado Denver, Aurora, Colorado
- c. Lifecourse Epidemiology of Adiposity and Diabetes Center, University of Colorado Anschutz Medical Campus, University of Colorado Denver, Aurora, Colorado
- d. Department of Epidemiology, Colorado School of Public Health, University of Colorado Denver, Aurora, Colorado
- e. University of Plymouth, Plymouth, United Kingdom
- f. Qld Allergy Services, 17/123 Nerang ST Southport Qld 4215 Australia
- g. Griffith University Dept. of Clinical Medicine, Southport Qld 4215 Australia
- h. Departments of Medicine and Microbiology, APC Microbiome Ireland, University College

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/CEA.14027](https://doi.org/10.1111/CEA.14027)

This article is protected by copyright. All rights reserved

Cork, Cork, Ireland

Corresponding author

Carina Venter PhD RD
Associate Professor of Pediatrics, Section of Allergy & Immunology
University of Colorado Denver School of Medicine
Children's Hospital Colorado
13123 East 16th Avenue, Box B518
Anschutz Medical Campus
Aurora, CO 80045

Declarations of interest

Dr. Venter reports grants from Reckitt Benckiser, grants from Food Allergy Research and Education, grants from National Peanut Board, during the conduct of the study; personal fees from Reckitt Benckiser, personal fees from Nestle Nutrition Institute, personal fees from Danone, personal fees from Abbott Nutrition, personal fees from Else Nutrition, and personal fees from Before Brands, outside the submitted work. Dr Smith reports personal fees from the Nestle Nutrition Institute and speaker fee from Danone and Bayer outside of the submitted work.

Dr. O'Mahony reports personal fees from Alimentary Health, grants from GSK, outside the submitted work. The other authors declare no interests.

Author contributions:

CV initiated the paper and drafted the first version of the paper. KP performed the data analysis with guidance and review by MP. AS guided on the epidemiological aspects of the study. PS and LOM advised on interpretation of AGEs intake and cytokine data. MAB extracted the allergy data from the electronic medical records. KM assisted with interpretation of the dietary and AGEs intake. DD is the principal investigator and assessed and guided the data analysis plan. All authors reviewed and commented on various drafts of the paper.

Informed consent

Informed consent was obtained from all subjects involved in the study.

Institutional review board

The Healthy Start study protocol was approved by the Colorado Multiple Institutional Review Board. (IRB number: 09-0563; Healthy Start 1; 2009-2014 and Healthy Start 2; 2015-present).

Data availability statement

Data available on request from the authors from <https://coloradosph.cuanschutz.edu/research-and-practice/centers-programs/lead>.

Funding/ Acknowledgements:

This work was supported by the of Health, grant numbers: R01 DK076648/DK/NIDDK NIH HHS/United States, R01 GM121081/GM/NIGMS NIH HHS/United States, UG3 OD023248/OD/NIH HHS/United States, UH3 OD023248/OD/NIH HHS/United States, R25GM111901-S1, R25GM11190, NIH grant R00ES025817

1 **Abstract**

2 **Background**

3 Associations have been shown between concurrent assessment of dietary intake of AGEs and
4 childhood allergic outcomes. We examined the association between maternal AGEs intake and
5 development of offspring asthma, wheeze, atopic dermatitis, allergic rhinitis, and food allergies, and
6 sought to determine whether intake of AGEs was associated with cord sera cytokines/chemokines.

7 **Methods**

8 Pregnant women ≥ 16 years were recruited in the Healthy Start study, a prospective pre-birth cohort
9 from Colorado (N =1410). The analysis included 962 dyads with adequate diet (≥ 2 recalls) and
10 allergy outcome details. AGEs intake was estimated for each mother by matching intakes reported
11 using 24-hour dietary recalls during pregnancy to a reference database of commonly consumed
12 foods' AGEs values. Child diagnoses of asthma and allergies up to 8 years were obtained from
13 electronic medical records. Cord sera cytokines and chemokines were analyzed in a subset (N =
14 462) of children.

15 **Results**

16 The median [IQR] AGEs intake for the overall sample was 11919 kU/day [8293, 16573]. Unadjusted
17 analysis showed a positive association between maternal AGEs intake in pregnancy and rhinitis up
18 to 8 years of age (HR = 1.03; 95% CI: 1.01, 1.06), but the association was attenuated and no longer
19 significant in adjusted models (HR = 1.01; 95% CI: 0.98, 1.04). Both adjusted and unadjusted models
20 showed no associations between AGEs intake in pregnancy and any of the other outcomes ($p>0.05$).
21 There were no significant associations between any cytokine or chemokine measured and AGEs
22 intake or any of the outcomes studied ($p>0.05$).

23 **Conclusion**

24 The study showed that maternal AGEs intake was not associated with offspring asthma and allergy
25 outcomes. AGEs exposure during pregnancy may not have the same impact on child development to
26 postnatal exposure.

27

28

29 **Key messages**

- 30
- 31
- 32
- 33
- 34
- AGEs exposure during postnatal life has been associated with multiple adverse health outcomes.
 - Maternal intake of AGEs during pregnancy was not associated with offspring allergies.
 - Maternal intake of AGEs during pregnancy was not associated with cord blood cytokine or chemokine levels.

35

36

37 Introduction

38 Allergic diseases are an increasing public health concern.^{1,2} The four major presentations of allergic
39 diseases include asthma, atopic dermatitis, allergic rhinitis, and food allergies. Atopic dermatitis is
40 usually the first manifestation of allergic diseases, followed by food allergies, asthma and allergic
41 rhinitis, a process of allergic disease development referred to as the atopic march.³ Although allergic
42 diseases share a common immunological profile involving T-helper 2 (Th2) and inflammatory cells
43 and their cytokines and chemokines, each condition may also appear separately with its own
44 individual gene/environment interactions.⁴

46 Early life factors, including maternal diet during pregnancy, have been studied as a possible risk
47 factor for offspring allergic diseases. However, despite the large number of studies focusing on
48 maternal dietary intake in pregnancy and offspring allergic outcomes, the results do not give clear
49 guidance on which dietary factors to address.^{5,6} The European Academy of Allergy and Clinical
50 Immunology also concludes that the role of the maternal diet in the development of offspring allergy
51 outcomes is unclear.⁷ Much more progress has been made in terms of the infant diet and allergy
52 prevention, with most international allergy prevention guidelines advising against delaying
53 introduction of food allergens, particularly peanut and egg.^{8,7,9,10}

55 Observational data indicates that increased intake in childhood of foods typical of the Western diet,
56 such as burgers, sugar and high fructose corn syrup,¹¹ and food preparations, such as frying, may be
57 associated with the increase in allergic diseases.¹² Advanced glycation end products (AGEs) are
58 compounds formed when sugar binds to protein or it is formed via lipoxidation.¹¹ It is postulated that
59 the Western diet is high in AGEs and via its effect on both Th2 and inflammatory cytokines, may
60 contribute to the increase in allergic outcomes. This is also referred to as the false alarm
61 hypothesis.¹²

62 Inflammatory cytokines and chemokines include, but are not limited to IL-1, IL-6, IL-8, IL-4, IL-5, IL-
63 13, alarmins (IL-25, IL-33 and TSLP), and TNF- α .^{13,14} The Receptor for Advanced Glycation End
64 Products (RAGE) is activated by ligands, including endogenous pathogen associated molecular
65 patterns (PAMPs), and also dietary derived AGEs, which are high in the western diet. The AGEs
66 receptor (RAGE) is linked to up-regulation of TNF- α , IL-1, IL-6 and IL-8.¹³ Induction of alarmins (IL25,
67 IL33, TSLP) and the importance of this is reinforced by models of atopic dermatitis and allergic

68 asthma that show RAGE and its activation ligands are central to the development of sensitization
69 and allergic responses.^{12,15,16} In addition, higher levels of soluble RAGE (a decoy ligand) is protective
70 against asthma.¹⁷ Further inflammatory mechanisms related to AGEs products are: 1) induction
71 of glycation of intracellular proteins including transcription factors, 2) alteration of the pericellular
72 matrix, resulting in signaling changes and cellular dysfunction, 3) oxidative stress and mitochondrial
73 dysfunction, 4) methylglyoxal – an archetypal AGE binds to lysine on DNA and induces
74 oxidative/nitrogen induced damage and DNA cleavage. Each of the fore-mentioned mechanisms
75 may have more impact on a developing fetus than a child/adult and is worthy of investigation in the
76 context of development of atopic disease. In animal models of food allergy, resveratrol, a well-
77 defined anti-oxidant, reduces development of ovalbumin allergy in a cholera-toxin adjuvant model
78 with both reduced sensitization of ovalbumin and a reduction in dendritic cell activation.¹⁸ Soy
79 isoflavones, daidzien and genistein, have been shown to suppress allergic reactions in a murine
80 model of peanut allergy.¹⁹ Furthermore, in-vitro studies of resveratrol have shown a reduction of
81 AGEs induced dendritic cell (DC) maturation, decreased dendritic cell RAGE receptor activation in
82 response to AGE-albumin, and reduced dendritic cell activation to AGE-albumin stimulated DC
83 (cytokines, co-stimulatory cells, mitogen activated protein kinases and NF- κ B).²⁰

84 Based on the underlying inflammatory processes in allergic conditions, particularly asthma and
85 wheeze, we previously studied the association between the maternal diet and asthma/wheeze in the
86 child. We concluded that the inflammatory profile of the maternal diet was not associated with
87 cytokine and chemokine levels at birth. However, the results suggested that a maternal diet that
88 scored higher on the dietary inflammatory index (DII) was associated with increased odds of
89 offspring asthma and/or wheeze by age 4 years.²¹ The DII is a complex index that provides an
90 estimate of the inflammatory potential of the diet based on the inflammatory potential of foods and
91 nutrients in single index.²² AGEs are proteins or lipids that become glycated after exposure to sugars
92 during heating and is considered to be one aspect of an inflammatory diet.¹¹

94 In our pursuit to find the specific dietary factors in pregnancy that are associated with offspring
95 outcomes, we investigated the role of maternal dietary AGEs intake and offspring asthma, wheeze,
96 atopic dermatitis, allergic rhinitis and food allergy. The primary aim of this study was to assess the
97 associations between maternal AGEs intake during pregnancy and offspring diagnosis of asthma,
98 wheeze, allergic rhinitis, atopic dermatitis, and food allergy up to 8 years. We hypothesized that

99 increased maternal AGEs intake during pregnancy would be associated with an increased rate of
100 development of these outcomes in offspring. The secondary aim of this study was to examine
101 associations between maternal AGEs intake and cord sera levels of cytokines and chemokines. We
102 hypothesized that maternal AGEs intake would be associated with cord sera cytokines and
103 chemokines, and that the effect might be exacerbated by maternal obesity, as both AGEs and
104 obesity are related to inflammatory processes.^{12,23}

105

106 **Methods**

107 ***Study sample***

108 This analysis included data from a longitudinal pre-birth cohort of 1410 mother-child dyads. Pregnant
109 women aged 16 years or older with singleton pregnancies were recruited from obstetrics clinics at
110 the local hospital from 2009 to 2014. The Healthy Start study protocol was approved by the Colorado
111 Multiple Institutional Review Board (IRB number: 09-0563; Healthy Start 1; 2009-2014 and Healthy
112 Start 2; 2015-present) and was registered as an observational study
113 at clinicaltrials.gov as NCT02273297. Further details regarding the study have been published
114 elsewhere.²³

115

116 At recruitment, the cohort included 1410 mother-child dyads. Following written consent, pregnant
117 women completed questionnaires on medical history at enrollment in early pregnancy and were
118 asked to give consent for the review of offspring electronic medical records up to age 4 years.
119 Additional consent for review of offspring electronic medical records up to age 8 was requested when
120 mothers and children came in for a follow-up visit after age 4.

121

122 Participants were excluded from this analysis if they did not provide consent for child medical record
123 review from birth up to 4 years of age (n=9), had offspring who died prior to birth (n=6), or who had
124 insufficient data available to search for the child in the electronic medical records system (n=66).
125 Among the 1329 participants eligible for inclusion in the electronic medical record search, 68 children
126 had no records in the electronic medical record system, resulting in a total of 1261 participants with
127 allergy outcome information. Mothers were asked to complete 1 dietary recall per month during
128 pregnancy. The analytic cohort included 962 mother-offspring dyads that completed ≥ 2 dietary
129 recalls over the course of pregnancy and had valid offspring allergy outcome information. For the

immunological analysis, data from N=462 mother-offspring dyads with data on cord sera cytokine and chemokine levels were used (Supplemental Figure 1).

Maternal daily AGEs intake

Maternal dietary intake was measured 2-8 times throughout pregnancy, from the second trimester onward, using the Automated Self-Administered 24-hour dietary recall (ASA24).²⁴ To estimate AGEs intake from the reported dietary intake, a reference database of commonly consumed foods' AGEs values (in kU/100g) built by Uribarri et al.¹¹ was used. Food descriptions from the ASA24 diaries were matched to the AGEs database by name and cooking process where possible. Foods without simple matches as well as complex multi-ingredient foods were decomposed into gram equivalents for 23 food components specified in My Pyramid,²⁴³ e.g. 1 cup of broccoli = 1 dark-green vegetable and 1 English muffin = 1 grain equivalent. The gram equivalents for each code were estimated from My Pyramid by decomposing food codes using the USDA food and nutrient database for dietary studies 3.0 ingredient list.²⁵⁶ Average AGEs (kU/g) values were then created from the Uribarri et al.¹¹ AGEs list for each of the 23 components based on general cooking type (fried, baked, stewed, or fresh), as cooking method is the main determinant of AGEs scores. Components for discretionary fats and sugars were excluded from this step due to difficulties with calculating accurate AGEs scores. Finally, daily dietary AGEs intake was calculated (in kU) based on quantity of the food/food component consumed. Median daily AGEs intake (kU/day) was calculated for each individual based on total number of dietary recalls available due to the right skewed nature of the AGEs distribution. Median daily AGEs were also standardized to amount per daily caloric intake (kU/1000kcal/day) to compare to previous literature.

Child allergy outcomes

The outcomes of interest for the present paper were child diagnosis of allergic diseases (asthma, wheeze, allergic rhinitis, atopic dermatitis, food allergy) up to 8 years of age based on data obtained from the electronic medical records. "Any outcome" was defined as diagnosis of one or more of the following diseases: asthma, wheeze, allergic rhinitis, atopic dermatitis, food allergy. Children's medical record data was abstracted for participants who consented to child medical record review and whose records were available in the Epic medical records system, as described previously.²³ In short, the following search terms were used: 1) allergic rhinitis: "allergic rhinitis", "allergic rhinoconjunctivitis", "hay fever", "rhinitis", "seasonal allergies"; 2) asthma: "asthma"; 3) atopic

dermatitis/eczema: “atopic dermatitis”, “eczema”; 4) wheeze: “wheeze.” Search terms related to IgE-mediated food allergies and food allergens included: “food allergy”, “almond”, “cashew”, “clam”, “crab”, “egg”, “fin fish”, “fish”, “milk”, “pecan”, “peanut”, “salmon”, “sesame seed”, “scallop”, “shellfish”, “shrimp”, “soy”, “sunflower seed”, “tree nut”, “tuna”, “wheat”. The electronic medical records were reviewed by two clinician researchers, who assigned diagnoses for asthma, wheeze, allergic rhinitis, atopic dermatitis/eczema, and food allergies, after extensive review of medical notes.

For any participant with suspected food allergy, we recorded whether the child had IgE-mediated food allergy, the age of diagnosis, and the age at development of tolerance to the food if tolerance had been developed. Children with Eosinophilic Esophagitis (EoE) or Food Protein induced enterocolitis (FPIES) were not considered to have IgE-mediated food allergies unless they also had co-existing IgE-mediated food allergies.

173

174 ***Cord blood collection***

Cord blood samples were obtained at delivery, stored on ice for up to 20 minutes, and processed by centrifugation. Serum aliquots were stored at 4°C for up to 24 hours before being transported (on ice) to an 80°C freezer for long-term storage.²⁶⁷

178

179 ***Cord sera cytokines and chemokines***

Under a separate ancillary study (NIH: R00ES025817) focusing on sera analysis from maternal-offspring dyads with available sera and cord blood, stored frozen umbilical cord sera extracted at birth was analyzed for a range of cytokines and chemokines. Cord blood inflammatory biomarkers were processed by the University of Colorado Cancer Center Flow Cytometry Shared Resource. Plasma cytokine/chemokine concentrations were determined by multiplex panel immunoassay according to manufacturer’s instructions (EMD Millipore Corporation, Billerica, MA 01821). Cytokines including IL-1 β , IL-4, IL-6, IL-10, and TNF- α ; and chemokines IL-8 and MCP-1, were measured in units of pg/mL. For each analysis, samples were run in duplicate, and the percent coefficient of variation was computed as a quality control measure. If both replicates were out of range, the value was designated as being below (or above) the limit of detection. Values that were below the lower limit of detection were marked as “out-of-range low” and values that were above the upper limit of detection were marked as “out-of-range high”.²¹ The cytokines and chemokines studied were previously selected to be analyzed for an NIH grant (R00ES025817) and included those that were related to either air pollution exposure or pregnancy/birth outcomes.

194

195 **Covariate data**

196 Data regarding maternal race/ethnicity, parity, maternal history of allergy (asthma and/or allergic
197 rhinitis), and age of introduction of solid foods were obtained through self-reported questionnaires.

198 Maternal history of asthma was assessed using the following question, *“Has a health professional*
199 *such as a doctor, physician assistant, or nurse practitioner ever told you that you have asthma?”*

200 Maternal history of allergic rhinitis was assessed using the following question, *“Has a health*
201 *professional such as a doctor, physician assistant, or nurse practitioner ever told you that you have*

202 *hay fever, seasonal allergies or allergic rhinitis?”* Mothers were considered to have a history of
203 allergy if they answered yes to one or both of the preceding questions. Pre-pregnancy weight was

204 obtained from either medical records or self-reported early in pregnancy. Maternal height was
205 measured at the first research visit via stadiometer. Pre-pregnancy body mass index (BMI) was

206 calculated using pre-pregnancy weight (kg) divided by height (m) squared. Pre-pregnancy BMI was
207 categorized as follows: lean (BMI <25 kg/m²), overweight (BMI 25-29.99 kg/m²), and obese (BMI ≥30

208 kg/m²).²⁴ Observed gestational weight gain was calculated as the difference between the last
209 available weight recorded during pregnancy and the pre-pregnancy weight.²⁷⁸ Gestational weight

210 gain was categorized as less than recommended, within the recommended range, or more than
211 recommended (excessive weight gain) based on pre-pregnancy BMI categories, as described by the

212 2009 Institute of Medicine (IOM) guidelines.²⁸⁹ Information on total caloric intake during pregnancy
213 (kcal/day) was obtained using repeated 24-hour recalls, as described above. Breastfeeding duration

214 was computed as breastmilk months, a product of breastfeeding duration and intensity, using feeding
215 information reported by mothers at the 18 months postnatal interview. For exclusively breastfed

216 infants, breastmilk months is equivalent to the duration of breastfeeding (e.g., 8 months of
217 breastfeeding = 8 breastmilk months). For infants fed both breastmilk and formula, breastmilk months

218 is the duration of exclusive breastfeeding plus the weighted duration of mixed feeding (e.g., 4 months
219 of exclusive breastfeeding + 2 months of 50% breastmilk and 50% formula = 5 breastmilk months.)

220 For infants fed formula exclusively, breastmilk months is 0.³⁰

221

222 **Statistical Analysis**

223 Descriptive statistics were calculated for maternal and offspring characteristics, including means and
224 standard deviations or median and interquartile range (IQR) for continuous variables dependent on

225 distribution. Categorical variables were presented as frequencies and percentages. Demographic

226 variables were compared between those with or without available cytokine data using statistical
227 tests, including t-tests for normally distributed continuous variables, Wilcoxon rank sum tests for non-
228 normally distributed continuous variables, and chi-square tests for categorical variables. We
229 computed cumulative incidence of the medical record verified respiratory and allergy outcomes up to
230 1 year, 2 years, 3 years, and 4 years of age to describe the burden of these diseases in the cohort.
231 Hypothesis testing was conducted at an alpha level of 0.05. In the final models, we reported, where
232 appropriate, beta estimates, hazard ratios, 95% confidence intervals, and p-values and for the
233 associations.

234

235 *Associations between maternal AGEs intake during pregnancy and offspring diagnosis of asthma,* 236 *wheeze, allergic rhinitis, atopic dermatitis, and food allergy*

237 Cox proportional hazards models were fit to examine the associations between maternal daily AGEs
238 intake (kU) and development of child allergic diseases (asthma, atopic dermatitis/eczema, food
239 allergy, allergic rhinitis, wheeze, and any allergy) up to age 8 years. A literature review²⁹ identified a
240 set of covariates used in previous publications seeking to identify associations between maternal diet
241 during pregnancy and offspring allergy and respiratory outcomes. Two models were performed for
242 the outcome: a base adjusted model 1, and an expanded adjusted model 2. Adjusted Model 1 was
243 the base model with factors most commonly associated with allergy outcomes^{2,30} and included child
244 race/ethnicity, child sex, nulliparity, gestational smoking, and energy intake during pregnancy.
245 Adjusted Model 2 was based on factors previously included when studying maternal diet and
246 childhood allergy outcomes.²⁹ These included all variables in model 1, and additionally: mode of
247 delivery, maternal history of allergy, breastfeeding duration, age of introduction of solid foods,
248 maternal pre-pregnancy BMI category (Supplemental Figure 2). We constructed a directed acyclic
249 graph to represent hypothesized causal relationships and confirmed that none of the covariates were
250 either intermediates or colliders. To test the hypothesis that the effect of maternal AGEs intake on
251 allergy onset may be exacerbated by maternal obesity and excessive gestational weight gain, we
252 included interaction terms between maternal AGEs intake with pre-pregnancy BMI category and IOM
253 gestational weight gain category into Adjusted model 2. Non-significant interactions ($p > 0.05$) were
254 removed from the final adjusted models. The follow-up age was different for each child. This
255 occurred for one of two reasons: 1) consent was only given to review child medical records from birth
256 up to age 4 years, but not from age 4 to 8 years; or 2) consent was given to review child medical
257 records from birth up to age 8 years, but the child had not yet reached 8 years of age at the time of

the electronic medical record search (e.g. the child was only 6 years old when the search was conducted). The Cox proportional hazards modeling approach allowed us to censor participants at the latest follow-up age for which they had available electronic medical record data, due to one of the two reasons described. For children with multiple recorded diagnoses of a disease outcome, their age at their first diagnosis was used. We checked that the assumption of proportional hazards was met for the “any outcome” prior to interpreting the results of these models. A sensitivity analysis was also performed dividing AGEs intake into quartiles to test for non-linear associations.

Associations between maternal AGEs intake and cord sera levels of cytokines and chemokines.

To examine the associations between maternal intake of AGEs and cytokines, two separate modeling strategies were used depending on the detection rate of the values. For each of the cytokines and chemokines, we determined the number and percentage of values that were outside the detection range. In addition, we calculated the median and IQR for each of the cytokines and chemokines. Cord sera cytokines and chemokines with <20% of values outside the detection range were treated as continuous variables (IL-6, IL-8, TNF- α , MCP-1).¹ When the cytokines or chemokines were treated as continuous variables, values below the limit of detection were assigned values equal to half the lowest value observed on the standard curve^{28 29} and values above the limit of detection were assigned values equal to 1.5 times the highest value on the standard curve. Cord sera cytokines and chemokines with \geq 20% of values outside the detection range were treated as categorical variables and dichotomized as detectable or not detectable (IL-1 β , IL-4).

For chemokines with <20% of values outside of the detection range, separate general linear univariate models were fit. Prior to fitting models for each of the cytokines and chemokines, the values of these cytokines and chemokines (IL-6, IL-8, TNF- α , MCP-1) were natural log transformed to account for the positively skewed distributions. For cytokines with \geq 20% values outside of the detection range (IL-1 β , IL-4), separate logistic regression models were fit to estimate the odds that the value of the cytokine or chemokine was detectable. For both cytokines and chemokines with <20% or \geq 20% of values outside the detection range, unadjusted models included maternal intake of AGEs as the only predictor. Adjusted models were fit for each cytokine/chemokine, with the covariates and hypothesized interactions tested determined a priori. The adjusted models for the inflammatory cytokines (IL-1 β , IL-6, TNF- α) and chemokines (IL-8, MPC-1) included nulliparity, child race/ethnicity, child sex, pre-pregnancy BMI category, IOM gestational weight gain category, and total caloric intake (kcal/day) as covariates (Supplemental Figure 3). The adjusted models for the T-regulatory cytokine (IL-10) and Th2 cytokine (IL-4) included the same covariates as were included for

the inflammatory cytokines, in addition to maternal history of allergy. To test the hypothesis that the effect of maternal AGEs intake on inflammatory cytokines and chemokines may be exacerbated by maternal obesity and excessive gestational weight gain, we included interaction terms between maternal AGEs intake with pre-pregnancy BMI category and with IOM gestational weight gain category. Non-significant interactions ($p > 0.05$) were removed from the final adjusted models. A sensitivity analysis was again performed dividing AGEs intake into quartiles to test for non-linear associations.

297

298 **Results**

299 *Descriptive statistics*

Table 1 reports descriptive statistics for maternal and offspring characteristics of the overall sample of Healthy Start participants who had both maternal AGEs intake and offspring electronic medical record data. Table 1 also compares maternal and offspring characteristics between the subset of participants with cytokine/chemokine data and the subset without cytokine/chemokine data. There were no significant demographic differences in the cohort included in analysis ($N = 962$, Supplemental Figure 1) and those excluded (data not shown). The median [IQR] AGEs intake for the overall sample was 11919 kU/day [8293, 16573]. Accounting for daily caloric intake, the median [IQR] AGEs intake was 6485.0 kU/1000kcal [4669.6, 8546.6]. Birthweight (g) was statistically, but not clinically, significantly higher in participants with cytokine and chemokine data compared to participants without cytokine and chemokine data (3290 ± 433 vs. 3196 ± 541 , $p=0.002$), as was gestational age at birth (weeks) (39.48 ± 1.24 vs. 39.21 ± 1.95 , $p=0.01$). Maternal age (years) at delivery was also statistically, while not substantially, different at 28.25 ± 6.03 for those with cytokine and chemokine data compared to 29.17 ± 5.84 for those without ($p=0.02$). AGEs intake, total caloric intake, breastfeeding duration, gestational age, age of introduction of solids, nulliparity, gestational smoking, ethnicity, pre-pregnancy BMI, IOM gestational weight gain, maternal history of allergy, and child sex were not significantly different between those with and without cord sera cytokine and chemokine data. Supplemental Table 1 shows cumulative incidence of allergy outcomes in the offspring. There were no statistically significant differences between those participants with or without cytokine data for any of the outcomes studied: any outcome, wheeze, asthma, atopic dermatitis, allergic rhinitis, and food allergy ($p > 0.05$, data not shown).

320

321 *Associations between maternal AGEs intake and offspring diagnosis of allergic outcomes*

Unadjusted analysis showed a positive association between a 1000kU/day increase in maternal AGEs intake in pregnancy and development of allergic rhinitis up to 8 years of age (HR = 1.034; 95% CI: 1.012, 1.057), but the association was attenuated and no longer significant in adjusted model 1 (HR = 1.01 (95% CI: 0.986, 1.043). In the base adjusted analysis for any outcome, asthma, atopic dermatitis, food allergy, and wheeze, there were no statistically significant associations between AGEs scores and childhood allergy outcomes up to 8 years of age, with all Hazard Ratio values nearing 1 ($p > 0.05$, Table 2). For these allergic outcomes, fully adjusted regression models gave similar results (Figure 1). There was also no evidence of a non-linear relationship between AGEs intake and any outcome ($p > 0.05$, Supplemental Table 4). There were no significant interactions between AGEs and BMI category ($p > 0.05$) for any of outcomes except for allergic rhinitis ($p = 0.01$) and wheeze ($p = 0.02$), and thus interactions are not reported in the adjusted model 2 in Table 2. For allergic rhinitis, women with a pre-pregnancy BMI categorized as “lean” were the only group with a significant AGEs effect with a hazard ratio of 1.054 for a 1000kU/day increase in AGEs (95% CI: 1.014, 1.095; $p = 0.008$). For wheeze, only women with a pre-pregnancy BMI categorized as “obese” had a significant AGEs effect with a hazard ratio of 0.934 for a 1000kU/day increase in AGEs (95% CI: 0.883, 0.989; $p = 0.02$) (Supplemental Table 3).

Associations between maternal AGEs intake and cord sera cytokine/chemokine levels

Supplemental Table 2 reports the frequency and percentage of values outside of the detection range for each cytokine/chemokine, and records whether the out-of-range values were low (below the lower limit of detection) or high (above the upper limit of detection). The median and IQR for levels of each cytokine/chemokine are also reported.

The hypothesized interaction between AGEs intake and pre-pregnancy BMI category for each of the cytokines/chemokines examined was non-significant (all $p > 0.05$). After removing all non-significant interaction terms, the associations between continuous maternal AGEs scores and cord sera levels of the cytokines/chemokines remained non-significant in the final adjusted models (all $p > 0.05$; Tables 3 and 4). There was also no evidence of a non-linear relationship between AGEs intake and any cord sera levels of the cytokines/chemokines ($p > 0.05$, Supplemental Table 5)

Discussion

353 Previous studies have indicated that AGEs intake may be implicated in non-communicable diseases
354 such as diabetes,^{31,32} chronic kidney disease,³³ non-alcoholic fatty liver disease,³⁴ coronary heart
355 disease,³² obstructive airway diseases,³⁵ cancer,³⁶ dementia and aging³⁷, arthritis,³⁸
356 asthma/wheeze,^{39,40} chronic bronchitis.^{41 37,42,43} Studies focusing on allergy outcomes, showed an
357 association between the childhood diet, studied at a similar time as allergic outcomes.^{12,44,45} In this
358 study, we examined the association between maternal AGEs intake during pregnancy and offspring
359 allergy outcomes. The unadjusted analysis showed a positive association between maternal AGEs
360 intake in pregnancy and offspring allergic rhinitis up to 8 years of age, but the association was no
361 longer significant in adjusted models. For all other offspring allergy outcomes, we did not find a
362 significant association with maternal AGEs intake, before or after adjusting for potential confounders.
363 For allergic rhinitis, a significant interaction between maternal pre-pregnancy BMI and AGEs intake
364 was observed, showing that the association between increased AGEs intake and “lean” based on
365 their pre-pregnancy BMI. A significant interaction between AGEs intake and pre-pregnancy BMI was
366 also observed for wheeze, indicating increased intake of AGEs was associated with reduced risk of
367 offspring wheeze only among women categorized as obese based on their pre-pregnancy BMI.
368 Immunological data from previous studies^{12,15,16} indicates that intake of AGEs may be associated with
369 development of allergic disease via its effect on RAGE receptors, leading to a cascade of
370 inflammatory processes. In the study presented here, no significant associations between maternal
371 AGEs intake and cord sera levels any of the cytokines or chemokines studied were observed.
372
373 Studies examining the relationship between AGEs intake and allergy outcomes, have been
374 conducted in children, focusing on the child’s diet and the child’s concurrent allergic symptoms. The
375 International Study of Asthma and Allergies in Children (ISAAC) study reported that eating fast foods
376 ≥ 3 times a week by adolescents was associated with asthma, rhino-conjunctivitis and eczema in
377 these study participants.⁴⁵ The authors hypothesized this this effect may be due to the AGEs content
378 of the food. Another prospective study reported on intake of AGEs in children and found that higher
379 fast food consumption rates in urban children in South Africa was associated with atopic dermatitis
380 than those with lower consumption rates. Urban children with high fried/microwaved meat
381 consumption also had higher rates of any allergy compared to those with lower intakes.⁴⁴ These
382 foods were once again used as a proxy for AGEs intake, rather than using a robust measurement of
383 AGEs content of the whole diet, as was done for the Healthy Start cohort in the study presented
384 here. More recently, a study from the US indicated that increased AGEs intake was significantly

385 associated with increased odds of wheezing, wheeze-disrupted sleep, and wheezing requiring
386 prescription medication.⁴⁰

387

388 Following on from the results of these studies,^{40,44,45} we sought to test whether maternal AGEs intake
389 during pregnancy was associated with offspring allergic outcomes. In both the unadjusted and
390 adjusted models for any outcome, asthma, atopic dermatitis, and food allergy up to 8 years, we
391 found no statistically significant findings for the association between maternal AGEs intake and
392 offspring allergy outcomes. The primary reason we suspect may explain the discrepancy between
393 our study findings and the findings of the previous studies listed, is that previous studies examined
394 concurrent AGEs intake of the child in which allergic disease was also studied, whereas we
395 prospectively studied maternal intake of AGEs during pregnancy and examined allergy in offspring.
396 This may indicate that AGEs intake in pregnancy may not affect child allergy outcomes. The results
397 of our study indicated that among women with a pre-pregnancy BMI categorized as lean weight
398 women, increased AGEs intake during pregnancy was associated with increased risk of offspring
399 allergic rhinitis; and among women with a pre-pregnancy BMI categorized as obese, increased AGEs
400 intake was associated reduced risk of offspring wheeze. These findings are difficult to interpret and
401 further studies may be able to clarify these results.

402

403 The discrepancy in findings between our study and previous studies may also be explained by
404 different measures and methods used to determine AGEs intake, or different methods for
405 assessment of allergy outcomes. Previous studies investigating the association between AGEs and
406 disease outcomes, used serum levels of AGEs,^{31,46} or activity of the age receptor (RAGE)^{37,42,43}
407 rather than dietary intake of AGEs as in our study. Some studies measured reported high-fructose
408 corn syrup sweetened soft drinks, fruit drinks³⁸ or apple juice, apple juice, fruit drinks and soda
409 intake.^{39,41} It is postulated that intake high fructose corn syrup, leads to *in situ* formation of AGEs,
410 hence making these foods a suitable proxy for studying AGEs intake.⁴⁷

411

412 Only three studies used dietary intake of AGEs measured by food frequency questionnaires to define
413 AGEs intake; one study defined AGEs intake by food group³⁵ and two studies quantified AGEs intake
414 based on FFQs measuring AGEs intake of commonly eaten foods.^{33,37} Our study is the first study to
415 our knowledge that investigated AGEs intake in pregnancy and its association with child health
416 outcomes. We have used detailed 24 hour recalls completed 2-8 times over the course of pregnancy

417 to calculate average AGEs intake throughout pregnancy. The median AGEs intake of the women in
418 the Healthy Start study was 6485.0 kU/1000 kcal. In our cohort, the median [IQR] AGEs intake for
419 the overall sample was 11919 kU/day [8293, 16573]. There are no studies reporting AGEs intake in
420 pregnant women, but a review paper by Nowotny et al.⁴⁸ indicates that AGEs intakes in adults can
421 range from 6 000 – 27 000 kU/day, which indicates that our calculated AGEs intake falls within this
422 range.

423
424 No previous studies have explored the relationship between maternal AGEs intake during pregnancy
425 and cord blood cytokines and chemokines. We had expected to see that higher maternal AGEs
426 intake would be positively associated with proinflammatory cytokines and chemokines (IL-1 β , IL-6,
427 TNF- α , MCP-1, IL-8)⁴⁹ and negatively associated with anti-inflammatory cytokines such as IL-10.⁴ In
428 particular, we expected to see some association between AGEs intake and TNF- α and MCP-1 levels.
429 TNF- α is a known pro-inflammatory cytokine and higher levels of TNF- α has been associated with
430 increased expression of RAGE.⁵⁰ MCP-1 is thought to play a role in cell-surface expression of
431 adhesion molecules⁵¹ and RAGE is closely related to other genes coding for cell adhesion
432 molecules. However, we saw no significant associations between maternal AGEs intake and cord
433 sera cytokine or chemokine levels.

434 A strength of this study is that we present information from a well-characterized cohort with a large
435 sample size, which enabled us to take various relevant covariates into account. We were able to
436 study the association between a comprehensive and detailed measure of dietary AGEs intake during
437 pregnancy, physician diagnosed recorded outcomes, and a range of cytokines and chemokines in
438 cord sera. We are also the first research group to study the association between maternal AGEs and
439 cord sera cytokine levels as a possible underlying mechanism for the development of offspring
440 allergies.

441 We have used detailed 24 hour recalls completed 2-8 times over the course of pregnancy to
442 calculate average AGEs intake throughout pregnancy. Approximately 76% of the participants
443 completed ≥ 2 dietary recalls over the pregnancy, with a median of 3 recalls. Two recalls can be
444 representative of the entire pregnancy, given that dietary intake is relatively stable across
445 pregnancy⁵² and two or more recalls are sufficient to estimate usual dietary intake, per the National
446 Cancer Institute Dietary Assessment Primer.⁵³ In addition, we also compared the estimated daily
447 AGEs values for those with below the median number of intakes (<3) to those with 4 or more entries.

448 The distributions were not statistically different with those with 3 or less entries (N = 556) having a
449 median estimated daily AGEs intake of 12104 (IQR: 7400-17148) and those with 4+ entries (N =
450 406) with a median of 11824 (IQR: 8978-15577) (p = 0.49). We have used a robust method to
451 calculate AGEs intake, but the AGEs content of many foods is unknown and calculation of the AGEs
452 scores of composite foods is difficult to standardize. One particular limitation of the dietary AGEs
453 calculation may be that discretionary fats and sugars were excluded due to difficulties with
454 calculating accurate AGEs scores. These issues may partially explain the null associations.

455
456 Future studies may benefit from using an AGEs food frequency questionnaire, validated against
457 reliable biomarkers, such as AGEs serum levels, and other forms of dietary intake to measure
458 AGEs instead. This information will enable us to assess if current levels of foods are indeed correct,
459 especially composite foods and provide us with validated measures of dietary intake. There are
460 currently no studies reporting on validating measurement of AGEs intake by using biological samples
461 such as blood¹² or skin.^{33,54} Some covariate information was collected through self-report rather than
462 objective measures, which might have influenced data quality.

463
464 Our results are limited as only one Th2 and one T-regulatory cytokine were measured. Measuring
465 cytokines in cord blood mononuclear cells, after sufficient stimulation with antigen, is preferable to
466 using frozen cord sera. Future studies should focus on measuring a wider range of cord sera
467 cytokines and chemokines, using antigen stimulation of cord blood mononuclear cells.

468
469 Other limitations included the potential bias associated with the self-report of dietary intake.^{55,56} In
470 addition, the sensitivity and specificity of using electronic medical record data for allergy outcomes is
471 not known, but this approach has been used in European cohort studies to report and validate on
472 asthma outcomes.⁵⁷ We also acknowledge that the mechanistic consequences of the AGE/RAGE
473 interaction could either impact on immune responses increasing a tendency to allergic sensitization
474 and/or enhancing susceptibility to inflammation independent of allergy. The latter may have been
475 impacted by pregnancy intake of AGEs and future studies should focus on separating “allergic” from
476 “inflammatory” outcomes. In particular, wheezing in infancy may not indicate an allergic phenotype,
477 though its inflammatory characteristics⁵⁸ have been well described. Most of the findings have wide
478 confidence intervals, highlighting the fact that further studies are required.

479

480 **Conclusion**

481 In this study, we did not observe significant associations between maternal AGEs intake during
482 pregnancy, and offspring allergy outcomes or cord blood cytokines and chemokines. These findings
483 suggest that maternal AGEs intake during pregnancy may not impact development of allergic
484 diseases in offspring. Exposure to AGEs during pregnancy may not have the same impact on child
485 development as postnatal AGEs exposure.

486

487

488 **References**

- 489 1. Institute of Medicine. Food Allergies: Global Burden, Causes, Treatment, Prevention and
490 Public Policy. National Academy of Sciences.
491 <http://www.nationalacademies.org/hmd/Activities/Nutrition/FoodAllergies.aspx>. Accessed
492 September 2021.
- 493 2. European Academy of Allergy and Clinical Immunology. Global atlas of Allergy.
494 <http://www.eaaci.org/globalatlas/GlobalAtlasAllergy.pdf>. Accessed September 2021.
- 495 3. Dharmage SC, Lowe AJ, Matheson MC, Burgess JA, Allen KJ, Abramson MJ. Atopic
496 dermatitis and the atopic march revisited. *Allergy*. 2014;69(1):17-27.
- 497 4. Akdis M, Aab A, Altunbulakli C, et al. Interleukins (from IL-1 to IL-38), interferons,
498 transforming growth factor beta, and TNF-alpha: Receptors, functions, and roles in diseases.
499 *J Allergy Clin Immunol*. 2016;138(4):984-1010.
- 500 5. Garcia-Larsen V, Ierodiakonou D, Jarrold K, et al. Diet during pregnancy and infancy and risk
501 of allergic or autoimmune disease: A systematic review and meta-analysis. *PLoS Med*.
502 2018;15(2):e1002507.
- 503 6. Venter C, Greenhawt M, Meyer RW, et al. EAACI position paper on diet diversity in
504 pregnancy, infancy and childhood: Novel concepts and implications for studies in allergy and
505 asthma. *Allergy*. 2020;75(3):497-523.
- 506 7. Halken S, Muraro A, de Silva D, et al. EAACI guideline: preventing the development of food
507 allergy in infants and young children (2020 update). *Pediatr Allergy Immunol*. 2021,
508 32(5):843-858
- 509 8. Greer FR, Sicherer SH, Burks AW, Committee On N, Section On A, Immunology. The Effects
510 of Early Nutritional Interventions on the Development of Atopic Disease in Infants and
511 Children: The Role of Maternal Dietary Restriction, Breastfeeding, Hydrolyzed Formulas, and
512 Timing of Introduction of Allergenic Complementary Foods. *Pediatrics*. 2019;143(4).
- 513 9. Turner PJ, Feeney M, Meyer R, Perkin MR, Fox AT. Implementing primary prevention of food
514 allergy in infants: New BSACI guidance published. *Clin Exp Allergy*. 2018;48(8):912-915.
- 515 10. Fleischer DM, Chan ES, Venter C, et al. A Consensus Approach to the Primary Prevention of
516 Food Allergy Through Nutrition: Guidance from the American Academy of Allergy, Asthma,
517 and Immunology; American College of Allergy, Asthma, and Immunology; and the Canadian
518 Society for Allergy and Clinical Immunology. *J Allergy Clin Immunol Pract*. 2021, 9(1):22-43

- 519 11. Uribarri J, Woodruff S, Goodman S, et al. Advanced glycation end products in foods and a
520 practical guide to their reduction in the diet. *J Am Diet Assoc.* 2010;110(6):911-916 e912.
- 521 12. Smith PK, Masilamani, M., Li, X., Sampson, H.A. "The False Alarm" hypothesis: Food allergy
522 is associated with high dietary advanced glycation end products and pro-glycating dietary
523 sugars that mimic alarmins. *Journal of Allergy and Clinical Immunology.*
524 2016;<http://dx.doi.org/10.1016/j.jaci.2016.05.040>.
- 525 13. Guarneri F, Custurone P, Papaiani V, Gangemi S. Involvement of RAGE and Oxidative
526 Stress in Inflammatory and Infectious Skin Diseases. *Antioxidants (Basel).* 2021;10(1).
- 527 14. Han X, Krempski JW, Nadeau K. Advances and novel developments in mechanisms of
528 allergic inflammation. *Allergy.* 2020;75(12):3100-3111.
- 529 15. Hammad H, Lambrecht BN. Barrier Epithelial Cells and the Control of Type 2 Immunity.
530 *Immunity.* 2015;43(1):29-40.
- 531 16. Ullah MA, Loh Z, Gan WJ, et al. Receptor for advanced glycation end products and its ligand
532 high-mobility group box-1 mediate allergic airway sensitization and airway inflammation. *J*
533 *Allergy Clin Immunol.* 2014;134(2):440-450.
- 534 17. Lyu Y, Zhao H, Ye Y, et al. Decreased soluble RAGE in neutrophilic asthma is correlated with
535 disease severity and RAGE G82S variants. *Mol Med Rep.* 2018;17(3):4131-4137.
- 536 18. Okada Y, Oh-oka K, Nakamura Y, et al. Dietary resveratrol prevents the development of food
537 allergy in mice. *PLoS One.* 2012;7(9):e44338.
- 538 19. Masilamani M, Wei J, Bhatt S, Paul M, Yakir S, Sampson HA. Soybean isoflavones regulate
539 dendritic cell function and suppress allergic sensitization to peanut. *J Allergy Clin Immunol.*
540 2011;128(6):1242-1250.e1241.
- 541 20. Buttari B, Profumo E, Capozzi A, et al. Advanced glycation end products of human β_2
542 glycoprotein I modulate the maturation and function of DCs. *Blood.* 2011;117(23):6152-6161.
- 543 21. Venter C, Palumbo MP, Sauder KA, et al. Examining Associations Between Dietary
544 Inflammatory Index in Pregnancy, Pro-inflammatory Cytokine and Chemokine Levels at Birth,
545 and Offspring Asthma and/or Wheeze by Age 4 Years. *J Acad Nutr Diet.* 2021, doi:
546 10.1016/j.jand.2021.02.015.
- 547 22. Shivappa N, Steck SE, Hurley TG, Hussey JR, Hebert JR. Designing and developing a
548 literature-derived, population-based dietary inflammatory index. *Public Health Nutr.*
549 2014;17(8):1689-1696.

- 550 23. Ahluwalia N, Andreeva VA, Kesse-Guyot E, Hercberg S. Dietary patterns, inflammation and
551 the metabolic syndrome. *Diabetes Metab.* 2013;39(2):99-110.
- 552 24. U.S. Department of Agriculture Human Nutrition Information Service. Food Guide Pyramid: A
553 guide to daily food choices. Home and Garden Development of the Food Guide Pyramid.
554 Nutrition. *Nutrition Today.* 1992;27:12.
- 555 25. US Department of Agriculture. The USDA food and nutrient database for dietary studies, 3.0-
556 Documentation and User guide.
557 https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/fndds/fndds3_doc.pdf. Accessed
558 September 2021.
- 559 26. Lemas DJ, Brinton JT, Shapiro AL, Glueck DH, Friedman JE, Dabelea D. Associations of
560 maternal weight status prior and during pregnancy with neonatal cardiometabolic markers at
561 birth: the Healthy Start study. *Int J Obes (Lond).* 2015;39(10):1437-1442.
- 562 27. Starling AP, Brinton JT, Glueck DH, et al. Associations of maternal BMI and gestational
563 weight gain with neonatal adiposity in the Healthy Start study. *Am J Clin Nutr.*
564 2015;101(2):302-309.
- 565 28. In: Rasmussen KM, Yaktine AL, eds. *Weight Gain During Pregnancy: Reexamining the*
566 *Guidelines.* Washington (DC)2009.
- 567 29. Venter C, Palumbo MP, Glueck DH, et al. The Maternal Diet Index in Pregnancy is
568 associated with offspring allergic diseases: The Healthy Start Study. *Allergy.* 2021.
- 569 30. World Health Organization. World Allergy Organization White Book on Allergy.
570 2011:http://www.worldallergy.org/UserFiles/file/WAO-White-Book-on-Allergy_web.pdf.
571 Accessed September 2021.
- 572 31. de la Cruz-Ares S, Cardelo MP, Gutiérrez-Mariscal FM, et al. Endothelial Dysfunction and
573 Advanced Glycation End Products in Patients with Newly Diagnosed Versus Established
574 Diabetes: From the CORDIOPREV Study. *Nutrients.* 2020;12(1).
- 575 32. Poirier P, Eckel RH. Advanced glycation end products and coronary heart disease in type 2
576 diabetes. *Diabetes Care.* 2000;23(9):1441-1442.
- 577 33. Chen J, Waqas K, Tan RC, et al. The association between dietary and skin advanced
578 glycation end products: the Rotterdam Study. *Am J Clin Nutr.* 2020;112(1):129-137.
- 579 34. Fernando DH, Forbes JM, Angus PW, Herath CB. Development and Progression of Non-
580 Alcoholic Fatty Liver Disease: The Role of Advanced Glycation End Products. *Int J Mol Sci.*
581 2019;20(20).

- 582 35. Lam R, Kwon S, Riggs J, et al. Dietary phenotype and advanced glycation end-products
583 predict WTC-obstructive airways disease: a longitudinal observational study. *Respir Res.*
584 2021;22(1):19.
- 585 36. Peterson LL, Park S, Park Y, Colditz GA, Anbardar N, Turner DP. Dietary advanced glycation
586 end products and the risk of postmenopausal breast cancer in the National Institutes of
587 Health-AARP Diet and Health Study. *Cancer.* 2020;126(11):2648-2657.
- 588 37. Chen J, Mooldijk SS, Licher S, et al. Assessment of Advanced Glycation End Products and
589 Receptors and the Risk of Dementia. *JAMA Netw Open.* 2021;4(1):e2033012.
- 590 38. DeChristopher LR, Uribarri J, Tucker KL. Intake of high-fructose corn syrup sweetened soft
591 drinks, fruit drinks and apple juice is associated with prevalent arthritis in US adults, aged 20-
592 30 years. *Nutr Diabetes.* 2016;6:e199.
- 593 39. DeChristopher LR, Uribarri J, Tucker KL. Intakes of apple juice, fruit drinks and soda are
594 associated with prevalent asthma in US children aged 2-9 years. *Public Health Nutr.*
595 2016;19(1):123-130.
- 596 40. Wang JG, Liu B, Kroll F, et al. Increased advanced glycation end product and meat
597 consumption is associated with childhood wheeze: analysis of the National Health and
598 Nutrition Examination Survey. *Thorax.* 2021;76(3):292-294.
- 599 41. DeChristopher LR, Uribarri J, Tucker KL. Intake of high fructose corn syrup sweetened soft
600 drinks is associated with prevalent chronic bronchitis in U.S. Adults, ages 20-55 y. *Nutr J.*
601 2015;14:107.
- 602 42. Brinkley TE, Semba RD, Kritchevsky SB, Houston DK. Dietary protein intake and circulating
603 advanced glycation end product/receptor for advanced glycation end product concentrations
604 in the Health, Aging, and Body Composition Study. *Am J Clin Nutr.* 2020;112(6):1558-1565.
- 605 43. Zha Z, Han Q, Huo S. The protective effects of bexarotene against advanced glycation end-
606 product (AGE)-induced degradation of articular extracellular matrix (ECM). *Artif Cells*
607 *Nanomed Biotechnol.* 2020;48(1):1-7.
- 608 44. Levin ME, Botha M, Basera W, et al. Environmental factors associated with allergy in urban
609 and rural children from the South African Food Allergy (SAFFA) cohort. *J Allergy Clin*
610 *Immunol.* 2020;145(1):415-426.
- 611 45. Ellwood P, Asher MI, Garcia-Marcos L, et al. Do fast foods cause asthma, rhinoconjunctivitis
612 and eczema? Global findings from the International Study of Asthma and Allergies in
613 Childhood (ISAAC) phase three. *Thorax.* 2013;68(4):351-360.

- 614 46. Kim Y, Keogh JB, Deo P, Clifton PM. Differential Effects of Dietary Patterns on Advanced
615 Glycation end Products: A Randomized Crossover Study. *Nutrients*. 2020;12(6).
- 616 47. DeChristopher LR, Uribarri J, Tucker KL. The link between soda intake and asthma: science
617 points to the high-fructose corn syrup, not the preservatives: a commentary. *Nutr Diabetes*.
618 2016;6(11):e234.
- 619 48. Nowotny K, Schröter D, Schreiner M, Grune T. Dietary advanced glycation end products and
620 their relevance for human health. *Ageing Res Rev*. 2018;47:55-66.
- 621 49. Shivappa N, Hebert JR, Marcos A, et al. Association between dietary inflammatory index and
622 inflammatory markers in the HELENA study. *Molecular Nutrition & Food Research*.
623 2017;61(6):06.
- 624 50. Tanaka N, Yonekura H, Yamagishi S, Fujimori H, Yamamoto Y, Yamamoto H. The receptor
625 for advanced glycation end products is induced by the glycation products themselves and
626 tumor necrosis factor-alpha through nuclear factor-kappa B, and by 17beta-estradiol through
627 Sp-1 in human vascular endothelial cells. *J Biol Chem*. 2000;275(33):25781-25790.
- 628 51. Jiang Y, Beller DI, Frendl G, Graves DT. Monocyte chemoattractant protein-1 regulates
629 adhesion molecule expression and cytokine production in human monocytes. *The Journal of*
630 *Immunology*. 1992;148(8):2423-2428.
- 631 52. Crozier SR, Robinson SM, Godfrey KM, Cooper C, Inskip HM. Women's dietary patterns
632 change little from before to during pregnancy. *J Nutr*. 2009;139(10):1956-1963.
- 633 53. National Cancer Institute 24-hour Dietary Recall (24HR) At a Glance.
634 <https://dietassessmentprimer.cancer.gov/profiles/recall/index.html>. Accessed September
635 2021.
- 636 54. Birukov A, Cuadrat R, Polemiti E, Eichelmann F, Schulze MB. Advanced glycation end-
637 products, measured as skin autofluorescence, associate with vascular stiffness in diabetic,
638 pre-diabetic and normoglycemic individuals: a cross-sectional study. *Cardiovasc Diabetol*.
639 2021;20(1):110.
- 640 55. Lovell A, Bulloch R, Wall CR, Grant CC. Quality of food-frequency questionnaire validation
641 studies in the dietary assessment of children aged 12 to 36 months: a systematic literature
642 review. *J Nutr Sci*. 2017;6:e16.
- 643 56. Ishihara J. Challenges in Dietary Exposure Assessment in Epidemiology: Research Trends. *J*
644 *Nutr Sci Vitaminol (Tokyo)*. 2015;61 Supl:S33-35.

- 645 57. Moth G, Vedsted P, Schiotz PO. National registry diagnoses agree with medical records on
646 hospitalized asthmatic children. *Acta Paediatr.* 2007;96(10):1470-1473.
- 647 58. Tsuang A, Chan ES, Wang J. Food-Induced Anaphylaxis in Infants: Can New Evidence
648 Assist with Implementation of Food Allergy Prevention and Treatment? *J Allergy Clin*
649 *Immunol Pract.* 2021;9(1):57-69.
- 650

651 **Figure Legends/Text:**

652 Figure 1. Hazard ratios with confidence intervals for the associations between continuous maternal
653 AGE (kU/day) intake and Offspring Allergy Outcomes up to 8 years; adjusted for maternal
654 race/ethnicity, child sex, nulliparity, maternal smoking prior to pregnancy, maternal history of allergy,
655 mode of delivery, breastfeeding duration, age of introduction of solid foods, maternal pre-pregnancy
656 BMI category, and energy intake during pregnancy (all models exclude interactions).

657

Table 1. Maternal and offspring characteristics in the Healthy Start cohort

	Overall sample	Subset with cytokine/chemokine data	Subset without cytokine/chemokine data	
Sample size (N)	962	462	500	
Continuous variables	Mean* \pm SD^a	Mean* \pm SD^a	Mean* \pm SD^a	p-value^b
Maternal characteristics				
Advanced Glycation End Product Intake (AGEs) (kU/day)(Median [IQR])	11919 [8293, 16573]	11921.2 [7875.5, 15998.7]	11918.7 [8648.2, 16890.9]	0.21
AGEs (kU/1000kcal) (Median [IQR])	6485.0 [4669.6, 8546.6]	6434.0 [4510.2, 8448.9]	6434.0 [4510.2, 8448.9]	0.17
Age at delivery (years)	28.7 \pm 5.9	28.25 \pm 6.03	29.17 \pm 5.84	0.02
Total caloric intake (kcal/day)	2070 \pm 390	2055.58 \pm 378.81	2083.66 \pm 399.48	0.26
Breastfeeding duration (breastmilk months)³⁴	8.7 \pm 6.7	8.28 \pm 6.64	9.11 \pm 6.71	0.08
Nulliparous	475 (49%)	239 (52%)	236 (47%)	0.18
Smoking in pregnancy	64 (7%)	32 (7%)	32 (6%)	0.84
Pre-pregnancy body mass index³¹				0.28

Underweight (<18.5 kg/m²)	24 (2%)	14 (3%)	10 (2%)	
Lean (25 kg/m²)	510 (53%)	232 (50%)	278 (56%)	
Overweight (25-29.99 kg/m²)	245 (25%)	120 (26%)	125 (25%)	
Obese (≥30 kg/m²)	183 (19%)	96 (21%)	87 (17%)	
IOM^d gestational weight gain³³				0.25
Less than recommended	213 (22%)	92 (20%)	121 (24%)	
Within recommended range	286 (30%)	143 (31%)	143 (29%)	
More than recommended	460 (48%)	227 (49%)	233 (47%)	
Maternal history of allergy	334 (35%)	160 (35%)	174 (35%)	1
Offspring characteristics				
Race/ethnicity				0.94
Non-Hispanic white	548 (57%)	262 (57%)	285 (57%)	
Non-Hispanic black	98 (10%)	50 (11%)	48 (10%)	
Hispanic	213 (22%)	101 (22%)	112 (22%)	
Other^c	104 (11%)	49 (11%)	55 (11%)	
Sex – female	503 (52%)	247 (53%)	256 (51%)	0.52
Birthweight (grams)	3242 ± 595	3290.4 ± 432.8	3196.3 ± 541.12	0.002
Gestational age at birth (weeks)	39.3 ± 1.7	39.48 ± 1.24	39.21 ± 1.95	0.01

Age solid foods introduced (months)	6.1 ± 2.0	6.1 ± 1.9	6.2 ± 2.0	0.12
--	------------------	------------------	------------------	-------------

a) SD: Standard deviation

b) p-value for hypothesis test comparing demographic variables between those with or without any cytokine data. Statistical tests included t-tests for normally distributed continuous variables, Wilcoxon rank sum tests for non-normally distributed continuous variables, and chi-square tests for categorical variables.

c) Other race/ethnicity includes non-Hispanic Asian, American Indian/Alaska Native, Hawaiian/Pacific Islander, and multi-racial

d) IOM: Institute of Medicine

***AGE intake was expressed as median [IQR] as a non-normally distributed variable.**

Table 2. Results of Cox proportional hazard models examining the association between maternal daily AGE intake and child allergy diagnosis

	Unadjusted Model ((N = 912))		Adjusted Model 1 [±] ((N = 912))		Adjusted Model 2 [#] (N = 782)	
Outcome	HR*(95% CI)	p-value	HR*(95% CI)	p-value	HR*(95% CI)	p-value
Any Allergy	1.007 (0.993, 1.02)	0.32	0.995 (0.978, 1.01)	0.56	0.997 (0.978, 1.01)	0.74
Asthma	1.014 (0.993, 1.03)	0.19	1.009 (0.984, 1.03)	0.48	1.008 (0.98, 1.03)	0.57
Atopic Dermatitis	1.003 (0.986, 1.02)	0.70	0.993 (0.972, 1.01)	0.49	0.999 (0.976, 1.02)	0.93
Food Allergy	1.012 (0.971, 1.05)	0.56	0.978 (0.928, 1.03)	0.41	0.975 (0.915, 1.03)	0.44
Rhinitis	1.034 (1.012, 1.057)	0.002	1.014 (0.986, 1.04)	0.34	1.012 (0.98, 1.04)	0.58 [#]
Wheeze	0.99 (0.969, 1.01)	0.35	0.975 (0.951, 1.00)	0.05	0.981 (0.955, 1.00)	0.17 [#]

* Hazard ratio's (HR) given for 1000kU increase in AGE value.

[±]adjusted for maternal race/ethnicity, child sex, nulliparity, maternal smoking prior to pregnancy and energy intake during pregnancy.

[#]adjusted for maternal race/ethnicity, child sex, nulliparity, maternal smoking prior to pregnancy, maternal history of allergy, mode of delivery, breastfeeding duration, age of introduction of solid foods, maternal pre-pregnancy BMI category, and energy intake during pregnancy

[#]Significant interaction between AGE and BMI category observed. Results from models including interactions shown in Supplemental Table 3.

Table 3. Associations between maternal AGE intake (kU) and natural log levels of cord sera cytokines and chemokines

Outcome	Unadjusted (N= 462)			Adjusted* (N= 462)		
	β^*	SE	p-value	β^*	SE	p-value
IL-6	-0.001	0.01	0.95	0.009	0.011	0.43
IL-8	-0.01	0.008	0.18	-0.005	0.009	0.57
TNF- α	-0.0003	0.003	0.91	0.004	0.004	0.27
MCP-1	0.004	0.004	0.23	0.007	0.004	0.09
IL-10	-0.007	0.006	0.22	-0.001	0.013	0.92

*The beta estimate represents the change in log-pg/mL of each outcome per each 1000kU increase in maternal AGE intake.

+ Adjusted for child race/ethnicity, child sex, nulliparity, energy intake during pregnancy, IOM gestational weight gain category, BMI Category. The model for IL-10 additionally adjusts for maternal history of allergy.

Table 4. Associations between maternal AGE intake (kU) and the odds of the cord sera cytokine and chemokine value being detectable

Outcome	Unadjusted Model (N= 462)			Adjusted Model* (N= 462)		
	OR*	95% CI	p-value	OR*	95% CI	p-value
IL-1 β	0.992	(0.966, 1.019)	0.57	0.981	(0.949, 1.014)	0.26
IL-4	0.998	(0.97, 1.026)	0.87	1.015	(0.98, 1.051)	0.41

* The odds ratio (OR) represents the change in the odds that the outcome is detectable per each 1000kU increase in maternal AGE intake.

+ Adjusted for child race/ethnicity, child sex, nulliparity, energy intake during pregnancy, IOM gestational weight gain category, BMI Category. The model for IL-4 additionally adjusts for maternal history of allergy.

