

Title	eHealth interventions to support self-management in people with musculoskeletal disorders, "eHealth: It's TIME"—A scoping review
Authors	Kelly, Marie;Fullen, Brona M.;Martin, Denis;McMahon, Sinéad;McVeigh, Joseph G.
Publication date	2022-01-13
Original Citation	Kelly, M., Fullen, B., Martin, D., McMahon, S. and McVeigh, J. G (2022) 'eHealth interventions to support self-management in people with musculoskeletal disorders, "eHealth: It's TIME"—A scoping review', Physical Therapy, 102(4), pzab307 (11pp). https://doi.org/10.1093/ptj/pzab307
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1093/ptj/pzab307
Rights	© 2022, the Author(s). Published by Oxford University Press on behalf of the American Physical Therapy Association. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited. - https://creativecommons.org/licenses/by/4.0/
Download date	2026-08-21 07:11:37
Item downloaded from	https://hdl.handle.net/10468/16579



UCC

University College Cork, Ireland
 Coláiste na hOllscoile Corcaigh

eHealth Interventions to Support Self-Management in People With Musculoskeletal Disorders, “eHealth: It’s TIME”—A Scoping Review

Marie Kelly , MSc^{1,2,*}, Brona Fullen, PhD³, Denis Martin, PhD^{4,5}, Sinéad McMahon, PhD³, Joseph G. McVeigh , PhD¹

¹Discipline of Physiotherapy, School of Clinical Therapies, College of Medicine and Health, University College Cork, Cork, Ireland

²Department of Physiotherapy, Mercy University Hospital, Cork, Ireland

³School of Public Health, Physiotherapy and Sports Science, University College Dublin, Dublin, Ireland

⁴School of Health and Life Sciences, Teesside University, Middlesbrough, United Kingdom

⁵NIHR Applied Research Collaborative, North East and North Cumbria, United Kingdom

*Address all correspondence to Marie Kelly at: 119227522@umail.ucc.ie

Abstract

Objective. eHealth-mediated interventions have been proposed as one option to support self-management in those with musculoskeletal disorders (MSDs). This scoping review aimed to chart the evidence regarding eHealth modalities, musculoskeletal diagnosis, and outcomes of eHealth-mediated self-management support interventions in persons with MSDs and identify any gaps within the literature.

Methods. Six electronic databases (MEDLINE, CINAHL, PsycINFO, Embase, Scopus, and the Cochrane Database of Systematic Reviews), 7 grey literature sources (eg, OpenGrey), and reference and citation lists of included studies were searched from database inception to July 2020. Published studies of adult participants with a MSD utilizing an eHealth intervention to support self-management were included. Studies were limited to those published in English. Two reviewers independently screened all studies. Data were extracted by 1 reviewer and reviewed by another reviewer.

Results. After screening 3377 titles and abstracts followed by 176 full texts, 87 studies fulfilled the eligibility criteria. The majority were published in the last 5 years ($n = 48$; 55%), with almost one-third originating in the United States ($n = 28$; 32%). The most common eHealth modality type was internet based ($n = 22$; 35%), with almost one-half ($n = 41$; 47%) of the included studies involving participants with widespread musculoskeletal symptoms. The most commonly reported outcomes were related to body functions (ie, pain intensity) ($n = 67$; 45%), closely followed by activities and participation (ie, function) ($n = 65$; 44%), with environmental factors (ie, health care utilization) the least commonly reported ($n = 17$; 20%).

Conclusions. There is considerable variation within the eHealth-mediated self-management support intervention literature. Research is needed on the role of eHealth-mediated self-management support interventions across a broad range of MSDs to guide clinical practice.

Impact. This scoping review has identified gaps in the literature relating to specific eHealth modalities, musculoskeletal diagnoses, and health care utilization data, which should guide future research.

Keywords: Digital Health, eHealth, Musculoskeletal Pain, Review, Self-Management

Received: June 9, 2021. **Revised:** October 27, 2021. **Accepted:** November 29, 2021

© The Author(s) 2022. Published by Oxford University Press on behalf of the American Physical Therapy Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

The profound burden of disease associated with musculoskeletal disorders (MSDs) is well established. Accounting for one-fifth of the world's total "years lived with disability,"¹ MSDs such as knee osteoarthritis and back pain also represent a significant economic cost.² This global burden is expected to rise in the coming decades due to an increase in related factors such as obesity, population growth, and aging.³ Health care systems worldwide face the challenge of how to manage the rising demand with limited resources.

Against this background, eHealth has attracted considerable interest as a possible solution. Definitions of eHealth are many and varied⁴; in this paper, an eHealth modality is defined as any specific technology applied in the context of health care.⁴ Examples include internet-based interventions, telephone support (interventions with telephone support from health practitioners), interactive voice response (the use of a phone's touch-tone keypad to provide responses to automated scripts), virtual reality (a 3-dimensional computer-generated environment the individual can explore, interact with, and manipulate), and mobile phone applications (mobile-based or mobile-enhanced programs) that deliver health-related services. Telehealth is an example of eHealth,⁵ and with respect to telehealth, this scoping review focuses on "real-time" telehealth (videoconferencing and telephone-based interventions).⁶ More detailed definitions are available in [Supplementary File](#).

eHealth modalities have the potential to improve the availability of health care services, helping individuals to access services and support from their own home. Furthermore, eHealth modalities may provide more cost-effective treatment options, reducing the requirement for travel and direct health care professional involvement, although actual savings are unknown at present.⁷ Such modalities also have the potential to improve treatment durability, because individuals can obtain support and reinforcement of skills after formal treatment has ceased.⁷ The COVID-19 pandemic, viewed as a "black swan" moment,^{8–10} appears to have been a significant catalyst for the temporary implementation of eHealth modalities into routine practice. However, much of the literature on eHealth predates the COVID-19 pandemic and is very fragmented, with existing systematic reviews mainly involving those with a chronic MSD^{11,12} and often limited to a specific eHealth subgroup such as internet-based¹² or mobile phone applications.¹¹

Self-management strategies for people with MSDs are complex interventions that involve patient education and behavior modification and are designed to encourage people to take an active self-management role to improve health outcomes.¹³ Effective self-management interventions have been identified as a priority implementation area for health care services.¹⁴ Supported self-management aims to empower people to take an active role in managing their condition and health behaviors collaboratively with their health care professional.¹⁵ Supported self-management is an important step toward better health care provision, which results in positive outcomes including reduced pain and improved self-efficacy.¹⁶ eHealth may be used to promote and enable self-management support on a large scale; however, uptake may be limited to those with knowledge and access to appropriate technology and hence may exacerbate health care inequality.¹⁷ Another concern is that eHealth modalities may increase dependence on health care professionals.¹⁸

Self-management and its support comprise a wide range of potential activities and interventions, with no single component identified as being more important than another.¹⁹ Taylor et al¹⁹ suggest that the detail and quality of reporting of complex self-management interventions was a barrier to understanding their effective components and wider implementation,¹⁹ which contributed to the development of the Practical Review in Self-Management Support (PRISMS) taxonomy.²⁰ This taxonomy identified 14 separate elements, which are considered important for self-management support by individuals and caregivers²¹ (eg, information provision and patient education; remote monitoring with feedback and action plans).²² eHealth has the potential to contribute to many of these components.

The PRISMS taxonomy has previously been used in evidence syntheses for other conditions^{22,23}; however, to our knowledge, there has not been a review in the MSD literature that outlines components of self-management support with reference to the PRISMS taxonomy.

Given the variety of eHealth options and the varied nature of self-management interventions, this review aimed to provide a broad overview of the evidence for eHealth-mediated self-management support interventions using scoping review methodology. This methodology is recommended when mapping literature predicted to be large and heterogeneous.²⁴ To characterize the available literature and identify any gaps or limitations, this scoping review had the following objectives: (1) to map the available evidence base on eHealth modalities, musculoskeletal diagnosis, and outcomes; and (2) to chart intervention characteristics, such as intervention provider, duration, and frequency, together with information on outcomes described as improved, unchanged, or worse.

Methods

Scoping Review

This scoping review followed the Joanna Briggs Institute methodology for scoping reviews²⁴ and reported findings utilizing the elements provided in the Preferred Reporting Items for Systematic reviews and meta-analyses extension for scoping reviews.²⁵ Our protocol was registered with the Open Science Framework (<https://osf.io/29rd6>) and published.²⁶ Because our full methods are available in our protocol, they are outlined briefly below.

Data Sources and Searches

For MEDLINE (via EBSCO), a 3-step search strategy was utilized as outlined in the protocol.²⁶ One reviewer (M.K.) developed the search strategy, with assistance from 2 professional librarians. A comprehensive electronic search of MEDLINE (EBSCO), CINAHL (EBSCO), PsycINFO (EBSCO), Embase, Scopus, and the Cochrane Database of Systematic Reviews databases was conducted from inception to July 2020. Studies were limited to those published in English. The main literature search was supplemented by utilizing the Canadian Agency for Drugs and Technology Gray Matters approach²⁷ with Google Scholar, Health Technology Assessment agencies (Canada, Australia, Ireland, UK, and USA) along with Open-Grey searched for grey literature. Additionally, screening of citation and reference lists of included studies was conducted to identify other potential studies that met the inclusion criteria.

Table 1. Characteristics of the Studies Included in Scoping Review^a

Characteristics	Number of Studies With Characteristic (N = 87)
Study design	
Randomized controlled trial	60
Randomized non-inferiority trial	2
Observational	10
Qualitative	8
Mixed methods	7
Origin of study	
United States	28
Australia	16
Sweden	11
United Kingdom	6
Norway	5
Germany	5
Netherlands	4
China	3
Nigeria	2
Switzerland	2
Austria	1
Canada	1
India	1
Italy	1
South Korea	1
Year of publication	
2016–2020	49
2011–2015	26
2006–2010	5
Prior to 2006	7
Musculoskeletal diagnosis	
Widespread musculoskeletal conditions ^a	41
Back pain	27
Osteoarthritis	12
Knee pain	5
Neck pain	2
Duration of symptoms	
Chronic (>12 wk)	79
Acute (<12 wk)	4
Unspecified duration	3

^aIndicates pain at multiple sites.

Study Selection

Literature search results were screened using Covidence online software (<http://www.covidence.org>). Titles and abstracts were screened by 2 review authors (M.K. and S.M.) independently against the eligibility criteria (Suppl. Table 1). Any disagreements were resolved by consensus or by decision of a third reviewer (J.M.). Studies that met the inclusion criteria were then retrieved in full and independently assessed in detail against the eligibility criteria by 2 review authors (M.K. and S.M.). Any conflicts were resolved by consensus or a third reviewer (J.M.).

Data Extraction

Data extraction was conducted using a standardized data extraction form that was developed a priori based on the Joanna Briggs Institute data extraction tool²⁴ and the template for Intervention Description and Replication checklist.²⁸ Data charted are synthesized in Tables 1 to 4 and in Supplementary Table 2. Data were charted for all studies by the lead author (M.K.) and checked by another author for accuracy, with any discrepancies resolved by consultation.

Data Synthesis and Analysis

Quantitative and qualitative data are presented in tabular format and synthesized narratively, prioritizing information reporting on distribution of studies by eHealth modality, musculoskeletal diagnosis, and quantitative and qualitative outcomes. We mapped the interventions in included studies to the components of the PRISMS taxonomy of self-management support.²⁰ Quantitative outcomes are presented in line with the International Classification of Functioning, Disability and Health framework,²⁹ with overall study results categorized descriptively (rather than analytically) as “improved,” “unchanged,” “worse,” or “mixed.” This categorization was based on the results presented by the study authors without further interpretation of the results due to the nature of the review being undertaken. Due to updated scoping review methodology recommendations,²⁴ a qualitative thematic analysis was not performed, which was a deviation from the scoping review protocol.²⁶

Role of the Funding Source

The Irish Society of Chartered Physiotherapists (ISCP) played no role in the design, conduct, or reporting of this review.

Results

As presented in the PRISMA flow diagram (Fig. 1),²⁵ from 3377 unique citations, 87 full-text studies were included.

Description of Studies

A summary of the 87 studies included in this review is provided in Table 1 and Supplementary Table 2 (categorized by eHealth modality).

The most common study designs were randomized controlled trials (n = 60; 69%). Nonrandomized controlled trials consisted of observational (n = 10; 12%), qualitative (n = 8; 9%), mixed methods (n = 7; 8%), and randomized non-inferiority trials (n = 2; 2%). The majority of qualitative and mixed methods studies (n = 12; 80%) utilized interviews for data collection.

The included studies were conducted in 15 countries, including the United States (n = 28; 32%), Australia (n = 16; 18%), and Sweden (n = 11; 13%). Most studies were published in the last 5 years (n = 48; 55%). The mean age range of participants reported across studies was 27.3 to 75 years. Only 3 studies had participants with a mean age greater than 65 years.^{30–32}

eHealth Modality Type

Twenty-four studies reported the same eHealth modality: hence, 63 interventions were outlined (Tab. 2). The most common eHealth modality type was internet based (n = 22; 35%), with the least common single modality being virtual reality (n = 1; 2%). Almost one-fifth of interventions identified (n = 12; 19%) utilizing more than 1 eHealth modality, with the most common combination being internet based with telephone support (n = 8; 13%). Most interventions utilized a combination of face-to-face (ie, in-person) and eHealth modality format (n = 35; 56%).

Musculoskeletal Diagnosis

There were 5 groups of musculoskeletal diagnoses identified. The most frequent diagnosis was widespread musculoskeletal




Figure 1. PRISMA flow diagram.

conditions ($n=41$; 47%) (Tab. 1) followed by back pain ($n=27$; 31%), with neck pain ($n=2$; 2%) as the least common diagnosis. Most studies related to chronic or persistent presentations ($n=79$; 91%).

Study Outcomes and Results

Characteristics of the quantitative study outcomes are presented in Table 3. Sixty-seven studies (45%) assessed outcomes relating to body functions (ie, pain, anxiety, depression, etc), and 65 studies (44%) assessed outcomes relating to activities and participation (ie, function). The least common outcomes reported related to environmental factors such as medication and health care utilization ($n=17$; 20%). The overall study outcomes were reported solely based on the findings reported by the study authors' findings and were not subjected to any form of critical analysis. These results were categorized descriptively as "improved," "unchanged," or "worse." When studies reported a combination of improved, unchanged, and/or worse outcomes, they were categorized as "mixed" results. Based on this method of categorization, 149 outcomes were reported, of which 50% ($n=75$) improved,

1% ($n=2$) worsened, 21% ($n=30$) had no change, and 28% ($n=42$) had mixed results. Of the 75 outcome categories that were improved, 30 (40%) were related to body functions, 38 (51%) were related to activities and participation, and 7 (9%) were related to environmental factors.

Supplementary Table 2 includes the findings from 8 qualitative and 7 mixed methods studies. The majority of these studies ($n=12$; 80%) utilized interviews for data collection. It is difficult to summarize findings into meaningful categories given the variety of eHealth-mediated self-management support interventions. Two studies reported good satisfaction,^{33,34} 3 studies reported good acceptability,³⁵⁻³⁷ and 2 studies reported that participants with MSDs found the eHealth-mediated self-management support interventions to be beneficial.^{38,39}

eHealth-Mediated Self-Management Support Intervention Characteristics

eHealth modalities were overseen by a variety of providers (Tab. 2). The most common category was "multiple providers" ($n=17$; 27%). The most common identifiable group of

Table 2. Overall Characteristics of eHealth-mediated Self-management Support Interventions^a

Variable	Intervention Count (n = 63)
eHealth-modality type	
Internet based	22
Telephone based	15
Mobile phone application	8
Internet based with telephone support	8
Interactive voice response	3
Video teleconferencing	2
Mobile phone application with telephone support	1
Interactive voice response with telephone support	1
Internet based with video conferencing	1
Interactive voice response with telephone support and internet based	1
Virtual reality	1
Format	
Mixture of face-to-face and eHealth	35
eHealth only	28
Components of the interventions mapped to PRISMS taxonomy	
Adherence support and lifestyle interventions	59
Information provision and patient education	58
Remote monitoring with feedback and action plans	53
Training and rehearsal of psychological strategies	41
eHealth-facilitated clinical review	42
Interventions that contained all 5 PRISMS taxonomy components	23
Interventions contained ≤ 2 PRISMS taxonomy components	2
Type of provider	
Multiple	17
Psychologist/graduate psychology students	12
N/A, eHealth modality fully automated	12
Other ^b	10
Physical therapist	9
Nurse	2
Health coach	1
Location of intervention	
Multiple	34
Individual's home	28
Outpatient	1
Frequency	
Mixed	29
Weekly	18
More than once per week	5
Daily	5
Unstructured, degree of frequency not specified	3
Monthly	2
More than once per day	1
Duration	
6–8 wk	18
9–20 wk	16
6 mo–1 y	14
3 wk–5 wk	7
Mixed	5
2 y	1
One-time access	1
Unspecified	1
Follow-up	
None	29
1–5 mo	13
11–12 mo	11
6–9 mo	10
Intervention development	
Use of framework/theory	48
Co-designed with individual	14

^aN/A = not applicable; PRISMS = Practical Review in Self-Management Support. ^bIndicates other providers, that is, health counselor, facilitator, psychiatrist, physical therapist, research staff, exercise physiologist, orthopedic specialist, and/or physician.

Table 3. Summary of Quantitative Study Outcomes^a

Common Outcomes	Description	No. of Studies	No. of Studies in Which Outcome Improved
Body functions	Pain intensity, pain interference, anxiety, depression, sleep, self-efficacy	67	30
Activities and participation	Stress management, walking, function, employment	65	38
Environmental factors	Medication, health care utilization	17	7

^aBased on the International Classification of Functioning, Disability and Health framework.

providers was psychologists or graduate psychology students ($n = 12$; 19%), with the same number of interventions fully automated with no provider involvement ($n = 12$; 19%). The least common was health coach ($n = 1$; 2%). The majority of interventions involved mixed frequencies ranging from 3 times per day to monthly ($n = 29$; 46%), with the least common frequency reported was more than once per day ($n = 1$; 2%). The majority of interventions lasted between 6 and 8 weeks ($n = 18$; 29%), whereas the least common durations reported included one-time access, 2 years, and unspecified ($n = 1$; 2% each). Almost one-half of the included studies ($n = 29$; 46%) reported posttreatment data only. The majority of eHealth-mediated self-management support interventions involved multiple locations such as a combination of outpatient with persons' home ($n = 34$; 54%), with the outpatient setting the least common location reported ($n = 1$; 2%).

Adherence support and lifestyle interventions were the most common components of the PRISMS taxonomy identified ($n = 59$; 94%), whereas training and rehearsal of psychological strategies was the least common ($n = 41$; 65%). Most of the eHealth-mediated self-management support interventions were complex interventions with multiple components; only 2 studies (3%) contained 2 or less PRISMS taxonomy components.^{40,41}

With respect to intervention development, 76% ($n = 48$) indicated the use of theories or frameworks. These included cognitive behavioral theory, acceptance and commitment theory, social cognitive theory, and the transtheoretical model of behavior change. Almost one-quarter of interventions ($n = 14$; 22%) involved individuals in the design of the intervention.

An overview of identified eHealth modality types with corresponding musculoskeletal diagnosis, outcomes, and intervention characteristics is provided in Table 4. The majority of studies evaluating internet-based interventions have been conducted in those with widespread musculoskeletal symptoms ($n = 19$; 63%), whereas the majority of telephone-based interventions have been assessed in populations with back pain ($n = 9$; 45%) and osteoarthritis ($n = 8$; 40%). Of the 10 studies included in this review relating to mobile phone applications, 6 (60%) were evaluated in those with back pain.

Discussion

This scoping review mapped the available evidence on eHealth-mediated self-management support interventions for those with MSDs with regards to eHealth modalities, musculoskeletal diagnosis, and outcomes. To our knowledge, this is the first review comprehensively charting 7 different eHealth modalities and their various combinations with respect to self-management support. In total, 87 studies of considerable variation in eHealth modality, outcomes

assessed, and results obtained fulfilled the inclusion criteria. eHealth-mediated self-management support interventions are an emerging area of practice in musculoskeletal disease, with most of the studies published in the last 5 years. COVID-19 may accelerate this further,⁸ with reported increases in eHealth use in Switzerland and Australia as a result of the pandemic ranging from 44.6% to 60%.^{42,43} However, these figures contrast sharply with findings reported by Werneke et al,⁴⁴ indicating an overall eHealth usage rate of 6%. Furthermore, despite increased use of eHealth reported in certain settings, physical therapists may not plan to continue to provide eHealth modalities after the pandemic.⁴³

Internet-based interventions were the most common eHealth modality. In keeping with previous reviews,^{7,45} the most common intervention delivery format was a combination of eHealth and face-to-face. It should be noted that during the COVID-19 pandemic, allied health clinicians managing those with MSDs rapidly introduced video teleconferencing and telephone into their clinical practice,^{42,43} not internet-based interventions. This is likely related to the fact that more people have access to smartphones than desktop personal computers,⁴⁶ and not all internet-based interventions are accessible via smartphones.⁴⁷ Other reasons may be the cost of developing and implementing internet-based self-management support interventions from an organizational perspective, along with issues around reimbursement that health care services across many countries receive for eHealth interventions.⁴⁸ In response to the COVID-19 pandemic, advocacy from professional bodies supported the reimbursement of telephone and video teleconferencing services by numerous funding bodies,^{49,50} which paved the way for the recent rapid introduction of these eHealth modalities into practice in some settings.^{42,43} Although this move was necessary during the pandemic for many health care providers because it was the only option for individuals to receive musculoskeletal health care,⁶ it is worth noting that video teleconferencing was identified in only 2 included studies^{32,51} and hence requires further investigation. The first-hand experience of video teleconferencing and telephone that many clinicians and individuals with MSDs now have will certainly shift views, either positively or negatively.⁶ Given the lack of user (ie, individuals with MSDs) involvement in eHealth intervention design noted within this review and criticized repeatedly in the literature,^{52,53} it is of paramount importance that future research adopts a co-design model involving various stakeholders (eg, individuals with MSDs, clinicians, software developers, researchers) to ensure the success and long-term sustainability of eHealth as a method for service delivery.⁵⁴

Almost one-half of the included studies ($n = 41$; 47%) involved participants with widespread musculoskeletal

Table 4. Tabular Presentation of Scoping Review Outcomes^a

eHealth Modality Type (Intervention Count = 63)	Internet Based (n = 22)	Telephone Based (n = 15)	Modality + Telephone Support (n = 11)	Mobile Phone Application (n = 8)	IVR (n = 3)	Video Teleconferencing (n = 2)	VR (n = 1)	Internet Based With Video Conferencing (n = 1)
MSK diagnosis (n = 87)	MSK conditions (n = 19) Back pain (n = 7) OA (n = 4)	MSK conditions (n = 1) Back pain (n = 9) OA (n = 8) Knee pain (n = 1) Neck pain (n = 1)	MSK conditions (n = 13) Back pain (n = 5)	MSK conditions (n = 2) Back pain (n = 6)	MSK conditions (n = 3) Back pain (n = 1)	MSK conditions (n = 1)	MSK conditions (n = 1)	
Quantitative study outcomes (n = 149)	Body functions and participation (n = 23) Activities and participation (n = 22) Environmental factors (n = 4)	Body functions (n = 13) Activities and participation (n = 15) Environmental factors (n = 5)	Body functions (n = 15) Activities and participation (n = 13) Environmental factors (n = 4)	Knee pain (n = 1) Neck pain (n = 1) Body functions (n = 9) Activities and participation (n = 8) Environmental factors (n = 2)	Body functions (n = 3) Activities and participation (n = 3) Environmental factors (n = 1)	Knee pain (n = 1) Body functions (n = 2) Activities and participation (n = 2)	Knee pain (n = 2) Body functions (n = 1) Activities and participation (n = 1) Environmental factors (n = 1)	
Intervention provider (n = 63)	N/A, eHealth modality fully automated (n = 9) Multiple (n = 5) Psychologist/graduate psychology students (5) Nurse (n = 1) Physical therapist (n = 1) Other (n = 1)	Multiple (n = 7) Nurse (n = 1) Physical therapist (n = 3) Other (n = 4)	Multiple (n = 2) Psychologist/graduate psychology students (6) Physical therapist (n = 1) Other (n = 1); health coach (n = 1)	N/A, eHealth modality fully automated (n = 2) Multiple (n = 2) Physical therapist (n = 2) Other (n = 2)	Multiple (n = 1) Other (n = 2)	Psychologist/graduate psychology students (1) Physical therapist (n = 1)	N/A, eHealth modality fully automated (n = 1)	
Duration (n = 63)	3–5 wk (n = 5) 6–8 wk (n = 6) 9–20 wk (n = 6) 6 mo–1 y (n = 2) Mixed (n = 2) One time access, unspecified (n = 1) Daily (n = 1) Daily (n = 1) Weekly (n = 9) More than once per day (n = 1) More than once/wk (n = 2)	6–8 wk (n = 2) 9–20 wk (n = 1) 6 mo–1 y (n = 7) Mixed (n = 3) Unspecified (n = 1) 2 y (n = 1) Mixed (n = 8)	6–8 wk (n = 6) 9–20 wk (n = 2) 6 mo–1 y (n = 3)	3 wk–5 wk (n = 2) 6–8 wk (n = 2) 9–20 wk (n = 3) Unspecified (n = 1) Mixed (n = 8) Daily (n = 2)	9–20 wk (n = 1) Mixed (n = 2) Mixed (n = 2)	6–8 wk (n = 1) 9–20 wk (n = 1) Mixed (n = 3) Weekly (n = 2)	3 wk–5 wk (n = 1) 9–20 wk (n = 1) Mixed (n = 1)	
Frequency	Daily (n = 1) Daily (n = 1) Weekly (n = 9) More than once per day (n = 1) More than once/wk (n = 2)	Mixed (n = 8) 2 y (n = 1) Mixed (n = 8) Daily (n = 1) Weekly (n = 9) More than once per day (n = 1) More than once/wk (n = 2)	Mixed (n = 11) Weekly (n = 3)	Mixed (n = 8) Daily (n = 2)	Mixed (n = 2)	Mixed (n = 3) Weekly (n = 2)	Daily (n = 1)	
Outcomes improved, worse, unchanged or mixed	Unstructured (n = 1) Body functions: Mixed (n = 15) Improved (n = 5) Unchanged (n = 3) Activities and participation (n = 9) Improved (n = 9) Unchanged (n = 9) Mixed (n = 3) Worse (n = 1) Environmental factors (n = 2) Improved (n = 2) Unchanged (n = 1) Mixed (n = 1)	Body functions: Mixed (n = 5) Improved (n = 5) Unchanged (n = 3) Activities and participation (n = 9) Improved (n = 9) Unchanged (n = 2) Mixed (n = 4) Environmental factors (n = 1) Improved (n = 1) Unchanged (n = 3) Worse (n = 1)	Body functions: Mixed (n = 6) Improved (n = 8) Unchanged (n = 1) Activities and participation (n = 8) Improved (n = 8) Unchanged (n = 4) Mixed (n = 1) Environmental factors (n = 2) Improved (n = 2) Mixed (n = 1) Unchanged (n = 1)	Unstructured (n = 2) Body functions: Mixed (n = 2) Improved (n = 7) Activities and participation (n = 6) Improved (n = 6) Unchanged (n = 1) Mixed (n = 1) Environmental factors (n = 2) Improved (n = 2) Mixed (n = 1) Unchanged (n = 1)	Body functions: Improved (n = 3) Activities and participation (n = 3) Improved (n = 3) Environmental factors (n = 1)	Body functions: Improved (n = 1) Mixed (n = 1) Activities and participation (n = 1) Improved (n = 1) Mixed (n = 1)	Body functions: Improved (n = 1) Activities and participation (n = 1) Improved (n = 1) Environmental factors (n = 1)	

TVR = interactive voice response; MSK = musculoskeletal; N/A = not applicable; OA = osteoarthritis; VR = virtual reality.

symptoms such as neck and back pain⁵⁵ or fibromyalgia^{56–58} or chronic pain.^{51,59–82} This is not surprising given that research indicates that most people reporting musculoskeletal pain describe multisite pain, with single-site pain relatively rare.⁸³ Furthermore, the more widespread the symptoms, the greater the impact on health and functioning.⁸³ An important point to highlight is that telephone and video teleconferencing consultations were being utilized during the pandemic to manage a myriad of MSDs such as tendinopathy and joint and muscle strains despite very limited evidence of efficacy to support eHealth-mediated self-management support for these presentations⁴²; the majority of the telephone-based studies identified within this review pertained to those with back pain and osteoarthritis. More recently, a retrospective observational design study published by Werneke et al⁴⁴ reported that telehealth was administered equally across numerous musculoskeletal regions, with unadjusted physical function change similar for those using and no using telehealth. However, the authors urge caution in the interpretation of physical function change due to the study design. This, together with the gap identified in this scoping review, highlights the urgent need for high-quality research to develop and evaluate eHealth-mediated self-management support interventions for the broad range of MSDs managed by health care systems to provide better guidance for clinical practice.⁴² Only 5% of included studies involved individuals with an acute presentation (<12 weeks), which is similar to other reviews^{84,85} on self-management interventions. As a result, the effectiveness of self-management support interventions in this cohort remains unknown despite the importance of early self-management in those with MSDs.⁸⁶ This gap within the literature may be partly related to the nature of MSDs, with recent-onset low back pain, for example, resolving quickly,⁸⁷ often with minimal intervention.⁸⁸ However, approximately 26% do not recover within 3 to 6 months,⁸⁹ with research indicating that treatments are rarely effective at returning them to a pain-free life.⁹⁰ It is likely that any eHealth-mediated self-management support intervention for those with acute MSDs would require different advice and support from that offered to those with chronic MSDs, in line with clinical guidelines.⁸⁶

Although research indicates that self-management support has a positive impact on health care utilization in conditions such as asthma²³ and cardiovascular disorders,⁹¹ there is a paucity of literature available on this outcome regarding MSDs. The eHealth-mediated self-management support intervention duration was relatively short despite improved treatment durability considered 1 potential advantage of eHealth modalities. Another limitation of the eHealth-mediated self-management support interventions identified within this scoping review is the lack of follow-up, with almost one-half of studies reporting posttreatment data only, a finding replicated in a review by Garg et al.⁹²

The outcomes of the interventions used in the studies included in the review were descriptively categorized as improved, unchanged, or worse based solely on the study authors' conclusions. The strength of the evidence underpinning the authors' findings was not subjected to critical analysis, and, as a result, conclusions regarding the effectiveness of eHealth-mediated self-management support interventions for those with MSDs cannot be drawn. The aim of this scoping review was to review the breadth of the

current literature and present the gaps that exist with scoping review methodology, not the most valid approach to evaluate effectiveness.⁹³

Study Limitations

Several limitations pertaining to this scoping review need to be acknowledged. This review includes English language papers only, and hence, potentially relevant evidence from other language may have been overlooked. Materials from sources such as guidelines, websites, or book chapters were not included. This may have introduced potential bias into the results. Finally, scoping reviews are inherently limited because the focus is to provide breadth rather than depth of evidence on a particular topic.²⁴ Hence, conclusions cannot be drawn regarding the most effective eHealth-mediated self-management support intervention for those with MSDs. However, this method was appropriate given the variety of options that eHealth includes and the varied nature of self-management interventions.

Future Implications

Accessibility and sustainability are key challenges facing musculoskeletal health care services, challenges that have been further exacerbated by COVID-19. Robust evidence to support eHealth-mediated self-management is critical if these interventions are to be successfully adopted post COVID-19. At present, it is challenging to synthesize and interpret eHealth-mediated self-management support interventions given the different methodological approaches and outcome measurements utilized among the included studies. To ameliorate this, researchers should adopt standard eHealth reporting guidelines.⁹⁴ This scoping review identified 2 studies evaluating the role of video teleconferencing in supporting self-management, an eHealth modality that was rapidly introduced into routine clinical practice in response to the COVID-19 pandemic. It is imperative that future research adopts a co-design model when evaluating the role of video teleconferencing and other eHealth modalities in supporting self-management. The majority of eHealth modalities were evaluated in populations with chronic pain, back pain, and osteoarthritis. To provide better guidance for clinical practice, high-quality research is also required for the broad range of MSDs managed by health care systems. Lastly, future studies should also evaluate the impact of these interventions on health care utilization⁹² and increase the length of follow-up.

Author Contributions

Concept/idea/research design: M. Kelly, B. Fullen, D. Martin, J. McVeigh
Writing: M. Kelly, B. Fullen, D. Martin, J. McVeigh
Data collection: M. Kelly, S. McMahon, J. McVeigh
Data analysis: M. Kelly, S. McMahon, J. McVeigh
Project management: M. Kelly, J. McVeigh
Providing institutional liaisons: J. McVeigh
Consultation (including review of manuscript before submitting):
B. Fullen, D. Martin, J. McVeigh

Funding

This study was funded by the Irish Society of Chartered Physiotherapists (ISCP) COVID-19 Bursary.

Systematic Review Registration

The protocol was registered with the Open Science Framework (<https://osf.io/29rd6>) and published.²⁶

Disclosures and Presentations

The authors completed the ICMJE Form for Disclosure of Potential Conflicts of Interest and reported no conflicts of interest.

Part of this study was presented at the National Irish Society of Chartered Physiotherapists Conference (virtual), October 15, 2021.

This work was registered in Open Science Framework: <https://osf.io/29rd6>.

References

- James SL, Abate D, Abate KH, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease study 2017. *Lancet (London, England)* 2018;392:1789–1858.
- Briggs AM, Woolf AD, Dreinhöfer K, et al. Reducing the global burden of musculoskeletal conditions. *Bull World Health Organ* 2018;96:366.
- Blyth FM, Briggs AM, Schneider CH, Hoy DG, March LM. The global burden of musculoskeletal pain—where to from here? *Am J Public Health* 2019;109:35–40.
- Showell C, Nohr C. How should we define eHealth, and does the definition matter? *Stud Health Technol Inform*. 2012;180:881–884.
- Gee PM, Greenwood DA, Paterniti DA, Ward D, Miller LMS. The eHealth enhanced chronic care model: a theory derivation approach. *J Med Internet Res* 2015;17:e4067.
- Cottrell MA, Russell TG. Telehealth for musculoskeletal physiotherapy. *Musculoskelet Sci Pract*. 2020;48:102193.
- Heapy AA, Higgins DM, Cervone D, Wandner L, Fenton BT, Kerns RD. A systematic review of technology-assisted self-management interventions for chronic pain: looking across treatment modalities. *Clin J Pain* 2015;31:470–492.
- Wind TR, Rijkeboer M, Andersson G, Riper H. The COVID-19 pandemic: the ‘black swan’ for mental health care and a turning point for e-health. *Internet Interv*. 2020;20:100317.
- Tack C, Grodon J, Shorthouse F, Spahr N. “Physio anywhere”: digitally-enhanced outpatient care as a legacy of coronavirus 2020. *Physiotherapy*. 2021;110:A26–A28.
- Telehealth is here to stay. *Nat Med* 2021;27:1121–1121.
- Devan H, Farmery D, Peebles L, Grainger R. Evaluation of self-management support functions in apps for people with persistent pain: systematic review. *JMIR mHealth and uHealth* 2019;7:e13080.
- Eccleston C, Fisher E, Brown R et al. Psychological therapies (internet-delivered) for the management of chronic pain in adults. *Cochrane Database Syst Rev*. 2014;2:CD010152.
- McGowan PT. Self-management education and support in chronic disease management. *Prim Care* 2012;39:307–325.
- Naylor C, Imison C, Addicott R et al. *Transforming Our Health Care System: Ten Priorities For Commissioners*. London, UK: The Kings Fund; 2015.
- da Silva D. Helping people help themselves: a review of the evidence considering whether it is worthwhile to support self-management. London, UK: Health Foundation; 2011. Accessed March 31, 2022. <https://www.health.org.uk/sites/default/files/HelpingPeopleHelpThemselves.pdf>.
- Mann EG, LeFort S, VanDenKerkhof EG. Self-management interventions for chronic pain. *Pain Manag* 2013;3:211–222.
- Janjua S, Banchoff E, Threapleton CJ, Prigmore S, Fletcher J, Disler RT. Digital interventions for the management of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*. 2021;4:CD013246.
- Fairbrother P, Pinnock H, Hanley J, et al. Exploring telemonitoring and self-management by patients with chronic obstructive pulmonary disease: a qualitative study embedded in a randomized controlled trial. *Patient Educ Couns* 2013;93:403–410.
- Taylor SJC, Pinnock H, Epiphaniou E et al. *A Rapid Synthesis of The Evidence on Interventions Supporting Self-Management for People With Long-Term Conditions: Prisms—Practical Systematic Review of Self-Management Support for Long-Term Conditions*. Southampton, UK: NIHR Journals Library; 2014.
- Pearce G, Parke HL, Pinnock H, et al. The PRISMS taxonomy of self-management support: derivation of a novel taxonomy and initial testing of its utility. *J Health Serv Res Policy* 2016;21:73–82.
- Sheridan N, Kenealy T, Kuluski K, McKillop A, Parsons J, Wong-Cornall C. Are patient and carer experiences mirrored in the practice reviews of self-management support (PRISMS) provider taxonomy? *Int J Integr Care* 2017;17:8.
- Hanlon P, Daines L, Campbell C, McKinstry B, Weller D, Pinnock H. Telehealth interventions to support self-management of long-term conditions: a systematic metareview of diabetes, heart failure, asthma, chronic obstructive pulmonary disease, and cancer. *J Med Internet Res* 2017;19:e172.
- Pinnock H, Parke HL, Panagioti M, et al. Systematic meta-review of supported self-management for asthma: a healthcare perspective. *BMC Med* 2017;15:64.
- Peters M, Godfrey C, McInerney P, Munn ZT, AC, Khalil H. Chapter 11: Scoping reviews (2020 version). In: Aromataris E, Munn Z (Eds). *JBIManual for Evidence Synthesis*; 2020. Accessed July 1, 2021. <https://reviewersmanual.joannabriggs.org/>.
- Page MJ, Moher D, Bossuyt PM et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160.
- Kelly M, Fullen B, Martin D, McMahon S, McVeigh JG. eHealth interventions to support self-management in people with musculoskeletal disorders: a scoping review protocol. *JBIM Evid Synth* 2021;19:709–720.
- CADTH. *Grey Matters: A Practical Tool for Searching Health-Related Grey Literature*. Ottawa, Canada: Canadian Agency for Drugs and Technologies in Health; 2019.
- Hoffmann TC, Glasziou PP, Boutron I et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ*. 2014;348:g1687.
- WHO. *International Classification of Functioning, Disability and Health: ICF*. Geneva, Switzerland: World Health Organization; 2001.
- Rini C, Porter LS, Somers TJ, et al. Automated, internet-based pain coping skills training to manage osteoarthritis pain: a randomized controlled trial. *Pain* 2015;156:837.
- Goode AP, Taylor SS, Hastings SN, Stanwyck C, Coffman CJ, Allen KD. Effects of a home-based telephone-supported physical activity program for older adult veterans with chronic low back pain. *Phys Ther* 2018;98:369–380.
- Wong YK, Hui E, Woo J. A community-based exercise programme for older persons with knee pain using telemedicine. *J Telemed Telecare* 2005;11:310–315.
- Guilkey RE, Draucker CB, Wu J, Yu Z, Kroenke K. Acceptability of a telecare intervention for persistent musculoskeletal pain. *J Telemed Telecare* 2016;24:44–50.
- Higgins DM, Buta E, Williams DA et al. Internet-based pain self-management for veterans: feasibility and preliminary efficacy of the pain EASE program. *Pain Pract*. 2020;20:357–370.
- Hinman R, Nelligan R, Bennell K, Delany C. “Sounds a bit crazy, but it was almost more personal.” a qualitative study of patient and clinician experiences of physical therapist-prescribed exercise for knee osteoarthritis via skype. *Arthritis Care Res (Hoboken)* 2017;69:1834–1844.
- Salisbury C, Foster NE, Hopper C, et al. A pragmatic randomised controlled trial of the effectiveness and cost-effectiveness of

- 'PhysioDirect' telephone assessment and advice services for physiotherapy. *Health Technol Assess* 2013;17:1–157.
37. Hasenöhl T, Windschnurer T, Dorotka R, Ambroz C, Crevenna R. Prescription of individual therapeutic exercises via smartphone app for patients suffering from non-specific back pain. *Wien Klin Wochenschr.* 2020;135:115–123.
 38. Hinman RS, Delany CM, Campbell PK, Gale J, Bennell KL. Physical therapists, telephone coaches, and patients with knee osteoarthritis: qualitative study about working together to promote exercise adherence. *Phys Ther* 2016;96:479–493.
 39. Kristjánsdóttir ÓB, Fors EA, Eide E et al. Written online situational feedback via mobile phone to support self-management of chronic widespread pain: a usability study of a web-based intervention. *BMC Musculoskelet Disord.* 2011;12:51–51.
 40. Petrozzi MJ, Leaver A, Ferreira PH, Rubinstein SM, Jones MK, Mackey MG. Addition of MoodGYM to physical treatments for chronic low back pain: a randomized controlled trial. *Chiropr Man Therap* 2019;27:54.
 41. Chhabra H, Sharma S, Verma S. Smartphone app in self-management of chronic low back pain: a randomized controlled trial. *Eur Spine J* 2018;27:2862–2874.
 42. Malliaras P, Merolli M, Williams C, Caneiro J, Haines T, Barton C. 'It's not hands-on therapy, so it's very limited': telehealth use and views among allied health clinicians during the coronavirus pandemic. *Musculoskelet Sci Pract.* 2021;52:102340.
 43. Rausch A-K, Baur H, Reicherzer L, et al. Physiotherapists' use and perceptions of digital remote physiotherapy during COVID-19 lockdown in Switzerland: an online cross-sectional survey. *Archives of physiotherapy* 2021;11:1–10.
 44. Werneke MW, Deutscher D, Grigsby D, Tucker CA, Mioduski JE, Hayes D. Telerehabilitation during the Covid-19 pandemic in outpatient rehabilitation settings: a descriptive study. *Phys Ther* 2021;101:pzab110.
 45. Slattery BW, Haugh S, O'Connor L, et al. An evaluation of the effectiveness of the modalities used to deliver electronic health interventions for chronic pain: systematic review with network meta-analysis. *J Med Internet Res* 2019;21:e11086.
 46. Du S, Liu W, Cai S, Hu Y, Dong J. The efficacy of e-health in the self-management of chronic low back pain: a meta analysis. *Int J Nurs Stud.* 2020;106:103507.
 47. Dario AB, Cabral AM, Almeida L, et al. Effectiveness of telehealth-based interventions in the management of non-specific low back pain: a systematic review with meta-analysis. *Spine J* 2017;17:1342–1351.
 48. Kruse C, Karem P, Shifflett K, Vegi L, Ravi K, Brooks M. Evaluating barriers to adopting telemedicine worldwide: a systematic review. *J Telemed Telecare* 2018;24:4–12.
 49. Tanne JH, Hayasaki E, Zastrow M, Pulla P, Smith P, Rada AG. Covid-19: how doctors and healthcare systems are tackling coronavirus worldwide. *BMJ.* 2020;368:m1090.
 50. McDonald, K. Telehealth in the time of coronavirus. *Pulse+IT.* 2020. Accessed July 21, 2021. <https://www.pulseitmagazine.com.au/blog/5399-telehealth-in-the-time-of-coronavirus>.
 51. Herbert MS, Afari N, Liu L, et al. Telehealth versus in-person acceptance and commitment therapy for chronic pain: a randomized noninferiority trial. *J Pain* 2017;18:200–211.
 52. Kress H-G, Aldington D, Alon E, et al. A holistic approach to chronic pain management that involves all stakeholders: change is needed. *Curr Med Res Opin* 2015;31:1743–1754.
 53. Cranen K, Groothuis-Oudshoorn CG, Vollenbroek-Hutten MM, IJzerman MJ. Toward patient-centered telerehabilitation design: understanding chronic pain patients' preferences for web-based exercise telerehabilitation using a discrete choice experiment. *J Med Internet Res* 2017;19:e26.
 54. Michie S, Yardley L, West R, Patrick K, Greaves F. Developing and evaluating digital interventions to promote behavior change in health and health care: recommendations resulting from an international workshop. *J Med Internet Res* 2017;19:e7126.
 55. Lo WLA, Lei D, Li L, Huang DF, Tong K-F. The perceived benefits of an artificial intelligence-embedded mobile app implementing evidence-based guidelines for the self-management of chronic neck and back pain: observational study. *JMIR mHealth and uHealth* 2018;6:e198.
 56. Ljótsson B, Atterlöv E, Lagerlöf M, et al. Internet-delivered acceptance and values-based exposure treatment for fibromyalgia: a pilot study. *Cogn Behav Ther* 2014;43:93–104.
 57. Williams DA, Kuper D, Segar M, Mohan N, Sheth M, Clauw DJ. Internet-enhanced management of fibromyalgia: a randomized controlled trial. *Pain* 2010;151:694–702.
 58. Friesen LN, Hadjistavropoulos HD, Schneider LH, Alberts NM, Titov N, Dear BF. Examination of an internet-delivered cognitive behavioural pain management course for adults with fibromyalgia: a randomized controlled trial. *Pain* 2017;158:593–604.
 59. Naylor MR, Helzer JE, Naud S, Keefe FJ. Automated telephone as an adjunct for the treatment of chronic pain: a pilot study. *J Pain* 2002;3:429–438.
 60. Hauser-Ulrich S, Künzli H, Meier-Peterhans D, Kowatsch T. A smartphone-based health care chatbot to promote self-management of chronic pain (SELMA): pilot randomized controlled trial. *JMIR Mhealth Uhealth* 2020;8:e15806.
 61. Naylor MR, Keefe FJ, Brigidi B, Naud S, Helzer JE. Therapeutic interactive voice response for chronic pain reduction and relapse prevention. *Pain* 2008;134:335–345.
 62. Naylor MR, Naud S, Keefe FJ, Helzer JE. Therapeutic interactive voice response (TIVR) to reduce analgesic medication use for chronic pain management. *J Pain* 2010;11:1410–1419.
 63. Ruehlman LS, Karoly P, Enders C. A randomized controlled evaluation of an online chronic pain self management program. *Pain* 2012;153:319–330.
 64. Buhrman M, Nilsson-Ihrfeldt E, Jannert M, Ström L, Andersson G. Guided internet-based cognitive behavioural treatment for chronic back pain reduces pain catastrophizing: a randomized controlled trial. *J Rehabil Med* 2011;43:500–505.
 65. Buhrman M, Skoglund A, Husell J, et al. Guided internet-delivered acceptance and commitment therapy for chronic pain patients: a randomized controlled trial. *Behav Res Ther* 2013;51:307–315.
 66. Dear BF, Courtney C, Khor KE et al. The pain course: exploring the feasibility of an internet-delivered pain management program when offered by a tertiary pain management service. *Clin J Pain.* 2018;34:505–514.
 67. Dear BF, Gandy M, Karin E, et al. The pain course: 12-and 24-month outcomes from a randomized controlled trial of an internet-delivered pain management program provided with different levels of clinician support. *J Pain* 2018;19:1491–1503.
 68. Dear BF, Gandy M, Karin E, et al. The pain course: a randomised controlled trial examining an internet-delivered pain management program when provided with different levels of clinician support. *Pain* 2015;156:1920.
 69. Dear BF, Titov N, Perry KN, et al. The pain course: a randomised controlled trial of a clinician-guided internet-delivered cognitive behaviour therapy program for managing chronic pain and emotional well-being. *Pain* 2013;154:942–950.
 70. Kristjánsdóttir ÓB, Fors EA, Eide E, et al. A smartphone-based intervention with diaries and therapist-feedback to reduce catastrophizing and increase functioning in women with chronic widespread pain: randomized controlled trial. *J Med Internet Res* 2013;15:e5.
 71. Kristjánsdóttir ÓB, Fors EA, Eide E, et al. A smartphone-based intervention with diaries and therapist feedback to reduce catastrophizing and increase functioning in women with chronic widespread pain. Part 2: 11-month follow-up results of a randomized trial. *J Med Internet Res* 2013;15:e72.
 72. Kroenke K, Baye F, Lourens SG et al. Automated self-management (ASM) vs ASM-enhanced collaborative Care for Chronic Pain and Mood Symptoms: the CAMMPS randomized clinical trial. *J Gen Intern Med.* 2019;34:1806–1814.

73. Kroenke K, Krebs EE, Wu J, Yu Z, Chumbler NR, Bair MJ. Telecare collaborative management of chronic pain in primary care: a randomized clinical trial. *JAMA* 2014;312:240–248.
74. Buhrman M, Syk M, Burvall O, Hartig T, Gordh T, Andersson G. Individualized guided internet-delivered cognitive-behavior therapy for chronic pain patients with comorbid depression and anxiety: a randomized controlled trial. *Clin J Pain* 2015;31:504–516.
75. Trompetter HR, Bohlmeijer ET, Veehof MM, Schreurs KM. Internet-based guided self-help intervention for chronic pain based on acceptance and commitment therapy: a randomized controlled trial. *J Behav Med* 2015;38:66–80.
76. Wilson M, Roll JM, Corbett C, Barbosa-Leiker C. Empowering patients with persistent pain using an internet-based self-management program. *Pain Manag Nurs* 2015;16:503–514.
77. Nordin CA, Michaelson P, Gard G, Eriksson MK. Effects of the web behavior change program for activity and multimodal pain rehabilitation: randomized controlled trial. *J Med Internet Res* 2016;18:e265.
78. Calner T, Nordin C, Eriksson MK, Nyberg L, Gard G, Michaelson P. Effects of a self-guided, web-based activity programme for patients with persistent musculoskeletal pain in primary healthcare: a randomized controlled trial. *Eur J Pain* 2017;21:1110–1120.
79. Lin J, Paganini S, Sander L, et al. An internet-based intervention for chronic pain: a three-arm randomized controlled study of the effectiveness of guided and unguided acceptance and commitment therapy. *Dtsch Arztebl Int* 2017;114:681.
80. Nes AAG, van Dulmen S, Wicksell R, Fors EA, Eide H. Analyzing change processes resulting from a smartphone maintenance intervention based on acceptance and commitment therapy for women with chronic widespread pain. *Int J Behav Med* 2017;24:215–229.
81. Peters ML, Smeets E, Feijge M, et al. Happy despite pain: a randomized controlled trial of an 8-week internet-delivered positive psychology intervention for enhancing well-being in patients with chronic pain. *Clin J Pain* 2017;33:962.
82. Johnson SS, Levesque DA, Broderick LE, Bailey DG, Kerns RD. Pain self management for veterans: development and pilot test of a stage-based mobile optimized intervention. *JMIR Med Inform* 2017;5:e40.
83. Kamaleri Y, Natvig B, Ihlebaek CM, Bruusgaard D. Localized or widespread musculoskeletal pain: does it matter? *Pain* 2008;138:41–46.
84. Nicholl BI, Sandal LF, Stochkendahl MJ, et al. Digital support interventions for the self-management of low back pain: a systematic review. *J Med Internet Res* 2017;19:e179.
85. Oliveira VC, Ferreira PH, Maher CG, Pinto RZ, Refshauge KM, Ferreira ML. Effectiveness of self-management of low back pain: systematic review with meta-analysis. *Arthritis Care Res (Hoboken)* 2012;64:1739–1748.
86. National Institute for Health and Care Excellence. NICE guideline (NG59): low back pain and sciatica in over 16s: assessment and management. 2016. Accessed Jan 16, 2020. <https://www.nice.org.uk/guidance/ng59>.
87. Henschke N, Maher CG, Refshauge KM et al. Prognosis in patients with recent onset low back pain in Australian primary care: inception cohort study. *BMJ*. 2008;337:a171.
88. Hancock MJ, Maher CG, Latimer J, Herbert RD, McAuley JH. Can rate of recovery be predicted in patients with acute low back pain? Development of a clinical prediction rule. *Eur J Pain* 2009;13:51–55.
89. Chou R, Shekelle P. Will this patient develop persistent disabling low back pain? *Jama* 2010;303:1295–1302.
90. Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet* 2017;389:736–747.
91. Panagioti M, Richardson G, Small N, et al. Self-management support interventions to reduce health care utilisation without compromising outcomes: a systematic review and meta-analysis. *BMC Health Serv Res* 2014;14:356.
92. Garg S, Garg D, Turin TC, Chowdhury MFU. Web-based interventions for chronic back pain: a systematic review. *J Med Internet Res* 2016;18:52–64.
93. Munn Z, Peters MD, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol* 2018;18:1–7.
94. Eysenbach G, Group C-E. CONSORT-EHEALTH: improving and standardizing evaluation reports of web-based and mobile health interventions. *J Med Internet Res* 2011;13:e126.