CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Assessment of plasma B7-H3 levels in pediatric patients with different degrees of surgical stress
Authors	Li, Yan;Yuan, Qing;Huang, Jie;Li, Yi Ping;Pan, Jian;Feng, Xing;Zhang, Xue Guang;Wang, Jiang Huai;Wang, Jian
Publication date	2016-07-26
Original Citation	Li, Y., Yuan, Q., Huang, J., Li, Y. P., Pan, J., Feng, X., Zhang, X. G., Wang, J. H. and Wang, J. (2016) 'Assessment of plasma B7-H3 levels in pediatric patients with different degrees of surgical stress', BMC Pediatrics, 16,110 (6pp). doi: 10.1186/s12887-016-0655-1
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://bmcpediatr.biomedcentral.com/articles/10.1186/s12887-016-0655-1 - 10.1186/s12887-016-0655-1
Rights	© 2016, Li et al. Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated. - https://creativecommons.org/licenses/by/4.0/
Download date	2026-11-16 12:03:13
Item downloaded from	https://hdl.handle.net/10468/4115



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Li et al. BMC Pediatrics (2016) 16:110
DOI 10.1186/s12887-016-0655-1

BMC Pediatrics

RESEARCH ARTICLE Open Access

Assessment of plasma B7-H3 levels in pediatric patients with different degrees of surgical stress

Yan Li[1†], Qing Yuan[2†], Jie Huang[3], Yi Ping Li[4], Jian Pan[4], Xing Feng[5], Xue Guang Zhang[6], Jiang Huai Wang[7*] and Jian Wang[1*]

Abstract

Background: Surgical stress initiates a series of host hormone, metabolism and immune responses, which predominantly affect the homeostatic mechanism of patients with major surgery. B7-H3 is a co-stimulatory molecule and has been shown to participate in both adaptive and innate immune responses. In this study we evaluated the clinical significance of plasma B7-H3 levels in pediatric patients with different types of operation and degrees of surgical stress.

Methods: A total of 48 children received pediatric general and cardiac surgery were recruited into this study. Based on the surgical stress scoring, children were divided into moderate stress ($n = 14$) and severe stress ($n = 34$) groups. Plasma B7-H3 levels were assessed at selected time points: before surgery, immediately after surgery, at day 1, day 3, and day 7 after surgery. Correlations between plasma B7-H3 levels and surgical stress scores were also examined.

Results: Plasma B7-H3 levels were significantly decreased in all 48 pediatric patients after surgery compared to the B7-H3 level before surgery ($p < 0.01$). Children with general surgery showed significant decreases in plasma B7-H3 immediately after surgery, and at day 3 and day 7 after surgery ($p < 0.05$, $p < 0.01$), whereas children with cardiac surgery showed reduced plasma B7-H3 immediately after surgery and at day 3 after surgery ($p < 0.05$). Plasma B7-H3 in cardiac surgery group was dropped much lower than that in general surgery group at day 1 ($p < 0.05$) and day 3 ($p < 0.01$) after surgery. Significantly reduced plasma B7-H3 was observed in the severe stress group, but not in the moderate stress group, immediately after surgery and at day 3 after surgery ($p < 0.05$), and severe stress group had significantly lower plasma B7-H3 levels than moderate stress group at day 1, day 3, and day 7 after surgery ($p < 0.05$). Furthermore, plasma B7-H3 levels at day 1 ($p = 0.01$) and day 3 ($p = 0.025$) after surgery correlated negatively with surgical stress scores.

Conclusions: Plasma B7-H3 levels were decreased significantly in children subjected to pediatric general and cardiac surgery, which is closely associated with the severity of surgical stress. The negative correlation of plasma B7-H3 levels at day 1 and day 3 after surgery with surgical stress scoring implicates that the plasma B7-H3 level might be a useful biomarker for monitoring stress intensity during pediatric surgery.

Keywords: Pediatric surgery, B7-H3, Surgical stress, Children

* Correspondence: jh.wang@ucc.ie; wj196312@gmail.com
†Equal contributors
[7]Department of Academic Surgery, University College Cork, Cork University Hospital, Cork, Ireland
[1]Department of Pediatric Surgery, Affiliated Children's Hospital, Soochow University, Suzhou 215003, China
Full list of author information is available at the end of the article

BioMed Central

© 2016 Li et al. **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated.

Background

Surgical stress initiates a series of host hormone, metabolism and immune responses, which predominantly affect the homeostatic mechanism of patients with major surgery. It has long been demonstrated for the past several decades that surgical procedures can activate the hypothalamic–-pituitary–adrenal (HPA) axis, thereby causing an elevated systemic glucocorticoid level [1]. Recent studies further show markedly altered levels of multiple cytokines after surgery, and these cytokines are mainly involved in initiation of the acute inflammatory reaction and modulation of immune responses [2]. Among them, IL-6 functions as a representative proinflammatory mediator to upregulate the acute inflammatory reaction, whereas IL-10 serves as a typical anti-inflammatory cytokine to inhibit the proinflammatory response. Upregulation of these cytokines in response to surgical stress for a new balance between proinflammatory and anti-inflammatory cytokines is crucial for maintaining the homeostatic stability of the host. Nevertheless, the imbalance between these cytokines could lead to systemic inflammatory response syndrome (SIRS) or compensatory anti-inflammatory response syndrome (CARS), and even multiple-organ dysfunction syndrome (MODS) [2].

In addition, surgical procedures may bring in exogenous antigens. These exogenous antigens interact with antigen-specific T cells, thereby participating in the process of major histocompatibility complex-associated antigen presentation and T cell-mediated activation of the adaptive immune response. Importantly, this process also requires the participation of co-stimulatory molecules. B7-H3 is a co-stimulatory molecule and belongs to the B7 superfamily. B7-H3 possesses a contrasting role in regulating T cell-mediated immune responses by functioning as both a T cell co-stimulator and co-inhibitor [3]. In addition to a well-documented role of B7-H3 in regulation of T cell-mediated immune responses, it has been recently shown that B7-H3 is also involved in the innate immunity-associated inflammatory response by functioning as a co-stimulator to promote proinflammatory cytokine production [4]. Moreover, significantly enhanced soluble B7-H3 in the circulation of septic patients could predict the poor outcome of these patients [4].

In this study, peripheral blood samples were collected from children subjected to pediatric general and cardiac surgery, and plasma B7-H3 levels before surgery, immediately after surgery, and at day 1, day 3, and day 7 after surgery were measured. We further evaluated alterations in plasma B7-H3 levels before and after surgery with types of surgery and severities of surgical stress, and the related clinical significance.

Methods

Patients

A total of 48 children who were admitted into the surgical ward of Affiliated Children's Hospital, Soochow University, Suzhou, China to undergo pediatric general surgery and cardiac surgery between June 2009 and December 2009 were recruited into this study. Among 48 cases, 20 cases underwent general surgery and 28 cases underwent cardiac surgery. Based on the Anand surgical stress scoring system [5], all children were further sub-grouped into moderate stress group ($n = 14$, score 6–10) and severe stress group ($n = 34$, score 11–20). The exclusion criteria included diagnosis of a genetic syndrome or malignant diseases, documented immunodeficiency or taken immunosuppressive drugs in the last 3 months, and refusal of consent. This study was approved by the Institutional Research Ethics Committee of Affiliated Children's Hospital and Soochow University for clinical investigation, and the written informed consent was obtained from parents or guardians of the recruited children prior to enrollment. All experiments and procedures followed were conducted in accordance with the principles of the Declaration of Helsinki involving human subjects.

Surgical stress scoring

The severity of surgical stress was assessed by the Anand scoring method, which is based on the amount of blood loss (score 0–3), the site and duration of surgery (score 0–2 and 0–4), the amount of tissue trauma (score 1–3), extent of visceral trauma (score 2–4), and associated stress factors including localized or generalized infection (score 1–2) and cardiopulmonary bypass (score 4) [5]. The total scores were used to classify the degree of surgical stress as minor (score 0–5), moderate (score 6–10), and severe (score 11–20) as described previously [5].

Blood sampling

Blood samples (2.0 ml) were obtained at the following time points: before surgery (Tc), immediately after surgery (T0), at day 1 (T1), day 3 (T3), and day 7 (T7) after surgery. At each time point, freshly heparinized blood samples were collected and centrifuged. Plasma samples were harvested and stored at −80 °C until analysis.

Measurement of plasma soluble B7-H3

An ELISA kit for soluble B7-H3 detection, with anti-B7-H3 mAbs 4H7 as a capture antibody and biolinylated 21D4 as a detecting antibody, was previously developed in our laboratory (6). In the current study, plasma concentrations of soluble B7-H3 was assessed using a modified ELISA kit in which the biolinylated 21D4 mAb was replaced by the biolinylated 2E6 mAb as a detecting antibody, with an improved sensitivity of soluble B7-H3 detecting limit from a previous 27 pg/ml to 3.3 pg/ml (4).

Prognosis of patients

Postoperative complications including hemorrhage, fever, hypothermia, infection, and incision rupture were monitored and recorded. In addition, the length of hospital stay was also recorded.

Statistical analysis

Statistical analysis was performed using SPSS17.0. All data were tested for normal distribution and expressed as mean ± standard deviation (SD) or otherwise presented as median (inter-quartile range). The Friedman test was used to compare the medians in multiple groups and the Spearman rank correlation coefficient test was used to analysis correlations between plasma soluble B7-H3 levels and surgical stress scores. A p-value of less than 0.05 was considered to be statistically significant.

Results

Patient information

Patient's data and clinical characteristics are summarized in Table 1. Among 48 cases, there were 23 males and 25 females with an average age of 12.74 ± 17.69 months. The length of hospital stay was 16.48 ± 6.75 days, and white blood cell counts, neutrophil counts, and C-reactive protein concentrations before surgery were $(8.79 \pm 4.44) \times 10^9$/L, 61.0 ± 8.82 %, and 3.26 ± 4.17 mg/L, respectively. There were more cases with cardiac surgery but less cases with general surgery in severe stress group compared to moderate stress group ($p < 0.01$). Moreover, children in the severe stress group had much longer hospital stay than children in the moderate stress group ($p < 0.05$).

Significantly reduced plasma B7-H3 levels after surgery

As shown in Fig. 1, soluble B7-H3 concentrations in the circulation were significantly decreased in all 48 pediatric patients after general and cardiac surgery compared to the B7-H3 level before surgery ($p < 0.05$, $p < 0.01$). The plasma B7-H3 level before surgery was 6.58 (4.72–11.32) ng/ml; however, it dropped to 4.75 (3.73–8.34) ng/ml immediately after surgery ($p = 0.000$), 5.63 (4.68–9.29) ng/ml at day 1 ($p = 0.009$), 5.61 (4.12–8.81) ng/ml at day 3 ($p = 0.000$), and 5.61 (4.12–8.81) ng/ml at day 7 ($p = 0.030$) after surgery.




Fig. 1 Plasma B7-H3 levels were significantly decreased after surgery. Heparinized whole blood samples were collected from 48 children subjected to pediatric general and cardiac surgery at following time points: before surgery (Tc), immediately after surgery (T0), at day 1 (T1), day 3 (T3), and day 7 (T7) after surgery. Plasma soluble B7-H3 was assessed by ELISA. Data are presented as median (inter-quartile range). *$p < 0.05$, **$p < 0.01$ compared with plasma B7-H3 before surgery (Tc)

We further assessed plasma B7-H3 levels in children subjected to either general surgery or cardiac surgery. When compared to the B7-H3 level before surgery at 8.73 (5.44–15.56) ng/ml, children with general surgery had significantly decreased plasma B7-H3 immediately after surgery at 5.44 (4.40–10.54) ng/ml ($p = 0.002$), and at day 3 at 7.28 (5.53–10.39) ng/ml ($p = 0.006$) and day 7 at 6.17 (4.53–11.91) ng/ml ($p = 0.028$) after surgery, whereas children with cardiac surgery showed significantly reduced plasma B7-H3 immediately after surgery at 4.32 (3.72–7.86) ng/ml ($p = 0.015$) and at day 3 at 5.43 (3.35–8.12) ng/ml ($p = 0.013$) after surgery (Fig. 2). Furthermore, plasma B7-H3 in cardiac surgery group was dropped much lower than that in general surgery group at day 1 ($p < 0.05$) and day 3 after surgery ($p < 0.01$) (Fig. 2).

Table 1 Patient's data and characteristics

	Total patients	Moderate stress	Severe stress	*P* values
Male / female	23/25	9/5	14/20	0.111
Age (months)	12.7 ± 17.7	8.87 ± 19.4	13.9 ± 17.3	0.004
General surgery (cases %)	20 (41.7)	14 (100)	6 (17.6)	0.000
Cardiac surgery (cases %)	28 (58.3)	0 (0)	28 (82.4)	0.000
Total white cell counts before surgery (10^9/L)	8.79 ± 4.44	8.87 ± 5.70	8.74 ± 3.54	0.513
PMN counts before surgery (%)	61.0 ± 8.82	60.6 ± 7.31	61.6 ± 10.8	0.124
CRP (mg/L)	3.26 ± 4.17	2.78 ± 4.25	3.26 ± 4.21	0.725
Hospital stay (days)	16.5 ± 6.75	13.4 ± 6.28	19.5 ± 7.22	0.031

PMN polymorphonuclear neutrophil, *CRP* C-reactive protein
All data are expressed as mean ± SD. *P* values: comparisons between moderate and severe stress groups




Fig. 2 The effect of pediatric general and cardiac surgery on plasma B7-H3 levels. Heparinized whole blood samples were collected from children with general surgery ($n = 20$) and children with cardiac surgery ($n = 28$) at following time points: before surgery (Tc), immediately after surgery (T0), at day 1 (T1), day 3 (T3), and day 7 (T7) after surgery. Plasma soluble B7-H3 was assessed by ELISA. Data are presented as median (inter-quartile range). $^{*}p < 0.05$, $^{**}p < 0.01$ compared with plasma B7-H3 before surgery (Tc); $^{\neq}p < 0.05$ compared with cardiac surgery group

Correlation between plasma B7-H3 and surgical stress scoring

We next analyze plasma B7-H3 levels between moderate stress group and severe stress group before and after surgery. As shown in Fig. 3, there were no significant differences in plasma B7-H3 levels observed in moderate stress group after surgery compared to the B7-H3 level before surgery; however, significantly reduced plasma B7-H3 was seen in severe stress group immediately after surgery at 4.47 (3.65–8.01) ng/ml ($p = 0.014$) and at day 3 after surgery at 5.15 (3.97–7.84) ng/ml ($p = 0.017$) compared to the plasma B7-H3 level at 5.62 (3.78–10.48) ng/ml before surgery. In addition, children in the severe stress group had significantly lower plasma B7-H3 than children in the moderate stress group at day 1, day 3, and day 7 after surgery ($p < 0.05$) (Fig. 3).




Fig. 3 Moderate and severe surgical stresses differentially affect plasma B7-H3 levels. Heparinized whole blood samples were collected from children with moderate surgical stress ($n = 14$) and children with severe surgical stress ($n = 34$) at following time points: before surgery (Tc), immediately after surgery (T0), at day 1 (T1), day 3 (T3), and day 7 (T7) after surgery. Plasma soluble B7-H3 was assessed by ELISA. Data are presented as median (inter-quartile range). $^{*}p < 0.05$ compared with plasma B7-H3 before surgery (Tc); $^{\neq}p < 0.05$ compared with severe stress group

Plasma B7-H3 levels in children subjected to pediatric general and cardiac surgery correlated negatively with the Anand surgical stress scores at day 1 after surgery (R-square = −0.367, $p = 0.010$) and at day 3 after surgery (R-square = −0.324, $p = 0.025$), respectively (Table 2).

Correlations of surgical stress scoring with prognosis of patients

As shown in Fig. 4, children in the severe stress group showed much longer hospital stay than children in the moderate stress group ($p < 0.05$). Moreover, the Anand surgical stress scores correlated positively with postoperative complications (R-square = 0.288, $p = 0.047$) (Table 3). However, plasma B7-H3 levels after surgery in all measured time points did not correlate with either the length of hospital stay or postoperative complications (data not shown).

Discussions

Host immune system is affected and regulated by multiple mediators and factors including stress, infection, trauma, malignant diseases, and nutrient conditions. Nevertheless, major surgery also affects and modulates the host homeostasis and defense mechanisms. In the present study, we measured plasma B7-H3 levels in 48 children subjected to pediatric general and cardiac surgery at different time points before and after surgery. We found that children with general surgery showed significantly reduced plasma B7-H3 immediately after surgery, and at day 3 and day 7 after surgery, whereas children with cardiac surgery showed significantly reduced plasma B7-H3 immediately after surgery and at day 3 after surgery. Significantly reduced plasma B7-H3 immediately after surgery and at day 3 after surgery was also observed in children with severe surgical stress, but not in children with moderate surgical stress. Moreover,

Table 2 Correlations of plasma B7-H3 levels with the Anand surgical stress scores at different time points before and after surgery

Plasma B7-H3	Anand surgical stress scores	
	R-square	*p* value
Before surgery	−0.218	0.136
Immediately after surgery	−0.194	0.188
At day 1 after surgery	−0.367	0.010
At day 3 after surgery	−0.324	0.025
At day 7 after surgery	−0.218	0.137

R2 values were calculated using the Spearman rank correlation coefficient test




Fig. 4 Children with severe surgical stress show significantly longer hospital stay than children with moderate surgical stress. The length of hospital stay between children in the moderate stress group ($n = 14$) and in the severe stress group ($n = 34$) was recorded and compared. $p = 0.031$ compared with the moderate stress group

plasma B7-H3 levels at day 1 and day 3 after surgery correlated negatively with the surgical stress scoring. These results indicate for the first time that the altered plasma B7-H3 level after surgery is associated with the degree of surgical stress in children who undergo major surgery.

Soluble B7-H3 in the circulation is shed from membrane B7-H3 expressed on the cell surface of human monocyte, dendritic cell, and active T cells [6], where matrix metalloproteinases (MMPs) play a key role to cleave the membrane B7-H3 into soluble B7-H3 via the enzymatic digestion [7]. Circulating soluble B7-H3 is detectable but usually at very low level in normal humans; however, patients with liver cancer [8] or colon cancer [9], and children with mycoplasma pneumonia [10] presented substantially enhanced B7-H3 levels in their circulation. Wei et al. [11] reported that significantly lower B7-H3 levels in the prostatic fluid were observed in patients with chronic prostatitis compared to healthy volunteers, which correlates negatively with the patients' disease status. In the present study, we also found that plasma soluble B7-H3 levels decreased significantly after surgery in children subjected to pediatric general and cardiac surgery. It has been well-documented that B7-H3, as a co-stimulatory molecule, participates in both adaptive and innate immune responses [3, 4]. Thus, the reduced plasma B7-H3 level after surgery may reflect the decreased and/or depressed host immune response in children with major surgery.

Table 3 Correlation between postoperative complications and the Anand surgical stress scores

	Anand surgical stress scores	
	R-square	*p* value
Postoperative complications	0.288	0.047

The R2 value was calculated using the Spearman rank correlation coefficient test

It is generally believed that surgery could suppress host immune responses, at least for a short time period [12–14]. However, other studies have shown that surgery does not alter or even enhance immune responses [15–17]. An early study by Mollitt et al. [18] revealed that neutrophil phagocytosis, migration, chemotaxis, and killing were not affected by surgery in children who underwent minor surgery. Interestingly, Romeo et al. [19] reported an enhanced activity of monocytes observed in children who underwent minor surgery, whereas Merry et al. [20] found no alterations in neutrophil chemotaxis and actin polymerization seen in children who underwent major surgery. These results implicate that different components of host immune system such as monocytes and neutrophils may respond differentially under different surgical stress conditions. Nevertheless, our results indicate the negative impact of major surgery on host immune responses in children.

As plasma B7-H3 levels were dropped substantially after surgery, we next examine whether the reduced plasma B7-H3 correlates with the severity of surgical stress. All 48 children subjected to pediatric general and cardiac surgery were scored using the Anand scoring method to evaluate the severity of surgical stress and sub-grouped into moderate stress group and severe stress group based on their surgical stress scores. Notably, children in the severe stress group showed significantly lower plasma B7-H3 than children in the moderate stress group at day 1, day 3, and day 7 after surgery, indicating that the plasma B7-H3 level is associated with the severity of surgical stress. Furthermore, a negative correlation was revealed between plasma B7-H3 levels at day 1 and day 3 after surgery, but not immediately after surgery and at day 7 after surgery, with surgical stress scores, indicating that surgical stress may reach peak point during the period of day 1 and day 3 after surgery. Zhao et al. [21] selectively inhibited histone deaccetylase-6 to attenuate the stress response in a murine cecal ligation and puncture (CLP)-induced sepsis model and found that the maximized stress response was at day 3 after CLP, which is in consistence with our findings. Our results also revealed that children in the severe stress group had much longer stay in the hospital than children in the moderate stress group, and the Anand surgical stress scores correlated positively with postoperative complications. However, neither the length of hospital stay nor postoperative complications correlated with the plasma B7-H3 levels before surgery, immediately after surgery, at day 1, day 3, and day 7 after surgery.

The major limitation of the current study is the relative small numbers of recruited pediatric patients, in particular when these patients were further sub-grouped into moderate and severe stress groups. Consequently, although reduced plasma B7-H3 levels after surgery correlated negatively with the surgical stress scoring and were closely associated with the severity of surgical stress, they failed to correlate with the clinical outcomes including postoperative complications and the length of hospital stay.

Conclusions

In this study we demonstrated that plasma soluble B7-H3 in children subjected to pediatric general and cardiac surgery were significantly decreased in response to surgical stress, which correlates negatively with the Anand surgical stress scores. These results suggest the clinical significance for detection of plasma B7-H3 in the clinical setting as a biomarker to monitor the stress intensity during pediatric surgery, though large scale of studies will be needed to confirm this.

Abbreviations

CARS, compensatory anti-inflammatory response syndrome; CLP, cecal ligation and puncture; HPA, hypothalamic–pituitary–adrenal; MMPs, matrix metalloproteinases; MODS, multiple-organ dysfunction syndrome; SIRS, systemic inflammatory response syndrome

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant 81272143 and 81420108022), Key Laboratory of Suzhou (Grant SZS201307), Jiangsu Innovation Team (Grant LJ201141), and Jiangsu Province Program of Innovative and Entrepreneurial Talents (2011–2014).

Authors' contributions

YL, JHW, and JW conceptualized and designed the study. YL, QY, JH, YPL, and JP performed the study, YL, XF, XGZ, JHW, and JW analyzed the data, YL, QY, and JHW draft the initial manuscript, JHW and JW approved the final version of manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Author details

[1]Department of Pediatric Surgery, Affiliated Children's Hospital, Soochow University, Suzhou 215003, China. [2]Department of Anesthesiology, Affiliated Children's Hospital, Soochow University, Suzhou 215003, China. [3]Department of Pediatric Cardiology, Affiliated Children's Hospital, Soochow University, Suzhou 215003, China. [4]Institute of Pediatric Research, Affiliated Children's Hospital, Soochow University, Suzhou 215003, China. [5]Department of Neonatology, Affiliated Children's Hospital, Soochow University, Suzhou 215003, China. [6]Institute of Medical Biotechnology, Soochow University, Suzhou 215006, China. [7]Department of Academic Surgery, University College Cork, Cork University Hospital, Cork, Ireland.

Received: 22 January 2015 Accepted: 21 July 2016
Published online: 26 July 2016

References

1. Nakamura M, Suita S, Yamanouchi T, Masumoto K, Ogita K, Taguchi S, Uesugi T. Cortisol and cytokine responses after surgery in different age groups of pediatric patients. Pediatr Surg Int. 2003;19:194–9.
2. Gentile LF, Moldawer LL. HMGB1 as a therapeutic target for sepsis: It's all in the timing. Expert Opin Ther Targets. 2014;18:243–5.
3. Suh WK, Gajewska BU, Okada H, Gronski MA, Bertram EM, Dawicki W, Duncan GS, Bukczynski J, Plyte S, Elia A, Wakeham A, Itie A, Chung S, Da Costa J, Arya S, Horan T, Campbell P, Gaida K, Ohashi PS, Watts TH, Yoshinaga SK, Bray MR, Jordana M, Mak TW. The B7 family member B7-H3 preferentially down-regulates T helper type 1-mediated immune responses. Nat Immunol. 2003;4:899–906.
4. Zhang G, Wang J, Kelly J, Gu G, Hou J, Zhou Y, Redmond HP, Wang JH, Zhang X. B7-H3 augments the inflammatory response and is associated with human sepsis. J Immunol. 2010;185:3677–84.
5. Anand KJ, Brown MJ, Causon RC, Christofides ND, Bloom SR, Aynsley-Green A. Can the human neonate mount an endocrine and metabolic response to surgery? J Pediatr Surg. 1985;20:41–8.
6. Zhang G, Hou J, Shi J, Yu G, Lu B, Zhang X. Soluble CD276 (B7-H3) is released from monocytes, dendritic cells and activated T cells and is detectable in normal human serum. Immunology. 2008;123:538–46.
7. Chen X, Bai Y, Cui W, Wang Z, Zhang G, Xu Y, Zhu X, Li Y, Wang JH. Effects of B7-H3 on the inflammatory response and expression of MMP-9 in mice with pneumococcal meningitis. J Mol Neurosci. 2013;50:146–53.
8. Xu W, Li F, Yu L, Liu J, He X, Fan R, Chen W. Plasma sB7-H3 serves as an assistant diagnostic marker for hepatitis B virus-associated hepatocellular carcinoma. Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi. 2014;30:744–7.
9. Zhou B, Zhang G, Yan G, Zhou H, Lv D, Zhang X. Overexpression of B7-H3 in $CD133^+$ colorectal cancer cells is associated with cancer progression and survival in human patients. J Surg Res. 2014;188:396–403.
10. Chen ZR, Zhang GB, Wang YQ, Yan YD, Zhou WF, Zhu CH, Wang J, Ji W. Soluble B7-H3 elevations in hospitalized children with mycoplasma pneumoniae. Diagn Microbiol Infect Dis. 2013;77:362–6.
11. Wei X, Zhang G, Yuan H, Ding X, Li S, Zhang X, Hou J. Detection and quantitation of soluble B7-H3 in expressed prostatic secretions: a novel marker in patients with chronic prostatitis. J Urol. 2011;185:532–7.
12. Klava A, Windsor A, Boylston AW, Reynolds JV, Ramsden CW, Guillou PJ. Monocyte activation after open and laparoscopic surgery. Br J Surg. 1997;84:1152–6.
13. Lennard TW, Shenton BK, Borzotta A, Donnelly PK, White M, Gerrie LM, Proud G, Taylor RM. The influence of surgical operations on components of the human immune system. Br J Surg. 1985;72:771–6.
14. Inoue T, Obata M, Mishima Y. Polymorphonuclear leukocyte function and serum opsonic activity in surgical patients. Surg Today. 1992;22:233–43.
15. van Dijk WC, Verbrugh HA, van Rijswijk RE, Vos A, Verhoef J. Neutrophil function, serum opsonic activity, and delayed hypersensitivity in surgical patients. Surgery. 1982;92:21–9.
16. Stanley TH, Hill GE, Portas MR, Hogan NA, Hill HR. Neutrophil chemotaxis during and after general anesthesia and operation. Anesth Analg. 1976;55:668–73.
17. Klava A, Windsor AC, Ramsden CW, Guillou PJ. Enhanced polymorphonuclear leucocyte adhesion after surgical injury. Eur J Surg. 1997;163:747–52.
18. Mollitt DL, Steele RW, Marmer DJ, Golladay ES, Costas S. Surgically induced immunologic alterations in the child. J Pediatr Surg. 1984;19:818–22.
19. Romeo C, Cruccetti A, Turiaco A, Impellizzeri P, Turiaco N, Di Bella C, Merlino MV, Cifala S, Basile M, Gentile C, Salpietro DC. Monocyte and neutrophil activity after minor surgical stress. J Pediatr Surg. 2002;37:741–4.
20. Merry C, Puri P, Reen DJ. Effect of major surgery on neutrophil chemotaxis and actin polymerization in neonates and children. J Pediatr Surg. 1997;32:813–7.
21. Zhao T, Li Y, Bronson RT, Liu B, Velmahos GC, Alam HB. Selective histone deacetylase-6 inhibition attenuates stress responses and prevents immune organ atrophy in a lethal septic model. Surgery. 2014;156:235–42.