

Title	Dairy Foods: A Matrix for Human Health and Precision Nutrition— Effect of processing infant milk formula on protein digestion and gut barrier health (in vitro and preclinical)
Authors	Dold, Cathal A.;Sahin, Aylin W.;Giblin, Linda
Publication date	2025
Original Citation	Dold, C. A., Sahin, A. W. and Giblin, L. (2024) 'Effect of processing infant milk formula on protein digestion and gut barrier health (in vitro and preclinical)', Journal of Dairy Science, 108(4), pp. 3088-3108. DOI: 10.3168/jds.2024-25356
Type of publication	Article (peer reviewed)
Link to publisher's version	10.3168/jds.2024-25356
Rights	© 2025, the Authors. Published by Elsevier Inc. on behalf of the American Dairy Science Association®. This is an open access article under the CC BY license (https://creativecommons.org/ licenses/by/4.0/) - https://creativecommons.org/licenses/by/4.0/
Download date	2026-09-21 22:28:31
Item downloaded from	https://hdl.handle.net/10468/17317



Dairy Foods: A Matrix for Human Health and Precision Nutrition— Effect of processing infant milk formula on protein digestion and gut barrier health (in vitro and preclinical)

Cathal A. Dold,^{1,2} Aylin W. Sahin,² and Linda Giblin^{1*}

¹Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork P61 C996, Ireland

²School of Food and Nutritional Sciences, University College Cork, Cork T12 CY82, Ireland

ABSTRACT

The infant gut is immature and permeable with high gastric pH, low protease activities, and underdeveloped intestinal architecture. Protein digestion in the upper gastrointestinal tract of infants is slow and incomplete. During manufacture, infant milk formula (IMF) is typically heat-treated so it is safe for human consumption. This heat treatment causes denaturation and aggregation of milk proteins, and formation of undesirable Maillard reaction products. The aim of this review is to critically summarize the in vitro and preclinical data available on the effect of IMF thermal processing on protein digestion and gut barrier physiology in the immature infant gut. Recent research efforts have focused on reducing thermal loads during IMF manufacturing by sourcing ingredients with low thermal loads, by reducing temperatures during IMF processing itself, and by seeking alternative processing technologies. This review also aims to evaluate whether these thermal reductions have a knock-on effect on protein digestion and gut barrier health in the infant. The ultimate aim is to create a safe next-generation IMF product that more closely mimics human breast milk in its protein digestion kinetics and its ability to promote gut barrier maturity in the infant.

Key words: infant milk formula, processing, protein digestion, gut barrier health, infant

INTRODUCTION

When babies are born their digestive tract is underdeveloped (Jiang et al., 2022). This immaturity of the infant gastrointestinal tract is reflected by higher gastric pH (Bourlieu et al., 2014), lower levels and activities of digestive enzymes (Nguyen et al., 2015; Ménard et al., 2018), immature intestinal architecture (Gleeson et

al., 2021), lower numbers of enterocytes, goblet cells, and Paneth cells (Demers-Mathieu, 2022), a more permeable mucus layer (Macierzanka et al., 2014), higher intestinal permeability (Kosek et al., 2017; Musa et al., 2019; Gleeson et al., 2021), and lower diversity of gut microbiota (Yao et al., 2021), compared with the adult gastrointestinal tract.

All of these characteristics combine to result in differences in nutrient digestion and absorption between infant and adult (Gan et al., 2018). At birth, preterm and term infants have a gastric pH ≥ 7 , which is heavily influenced by fed or fasted state and the infant's ability to secrete gastric acid (Bourlieu et al., 2014; Ménard et al., 2018). This is in sharp contrast to pH 1.5 to 2 in the adult stomach (Fujimori, 2020). The optimum pH for activity of the gastric enzyme, pepsin, is pH 2 (Nguyen et al., 2015; Gan et al., 2018). At pH ≥ 6.5 , pepsin activity is reduced, with complete denaturation and inactivation at pH ≥ 8 (Johnston et al., 2007). Therefore, in the infant stomach, pepsin activity is $<10\%$ of the activity reported in adults (Nguyen et al., 2015). In addition, the activities of the small intestine enzymes, trypsin and chymotrypsin, are reported in the infant at 10% to 60% of the adult activity levels (Nguyen et al., 2015). Whether age differences exist in brush border enzyme activities is less well characterized.

The small intestine surface is a mass of villi to increase surface area and therefore maximize nutrient absorptive capacity (Walton et al., 2016). Nutrients must cross the villi epithelium to be absorbed. The epithelium barrier is a single layer of cells composed primarily of absorptive enterocytes, goblet cells, Paneth cells, stem cells, and neuroendocrine cells. The enterocytes, goblet cells, and neuroendocrine cells begin life in the Paneth controlled crypts before maturing and migrating toward the villus tip (Kong et al., 2018). In the small intestine, villi height is lower in infants compared with adults (Gleeson et al., 2021). Mucus-secreting goblet cells prevent adhesion and attack by pathogenic bacteria (Johansson et al., 2013). In the mature gut, goblet cells account for 4% to 12% of

Received June 28, 2024.

Accepted September 24, 2024.

*Corresponding author: linda.giblin@teagasc.ie

The list of standard abbreviations for JDS is available at adsa.org/jds-abbreviations-25. Nonstandard abbreviations are available in the Notes.

intestinal epithelial cells, but it is widely accepted that the percentage of goblet cells is much lower in the infant intestine (Demers-Mathieu, 2022). Intestinal permeability is also higher in infants than in adults, with this permeability decreasing as the distance between neighboring cells narrows while the gut matures (Chelakkot et al., 2018; Gleeson et al., 2021). In 72 healthy infants the permeability test of lactulose:mannitol ratio was highest at birth, decreasing significantly during the first month of life (mean $\pm$ SD; 1.27 ± 0.73 at d 1 and 0.22 ± 0.21 at 1 mo; $P < 0.05$; Catassi et al., 1995). Neighboring barrier cells connect via tight junctions. The mRNA transcript levels of various tight junction biomarkers (*occludin*, *claudin-3*, *zonula occludens-1*, and *junctional adhesion molecule-A*) were significantly lower in the proximal and distal small intestine of infant mice pups compared with older adult mice ($P < 0.05$; Gleeson et al., 2021). In the infant gut, tight junction protein localization patterns differ from those in adults, with immunofluorescence staining revealing claudin-3 protein localized in the crypts in infant mice compared with in the midvillus and villus tip in adult mice (Gleeson et al., 2021). The infant gut microbiota also has lower microbial diversity compared with the adult (Odamaki et al., 2016; Lugli et al., 2023), and is heavily influenced by environmental factors such as mode of baby delivery, type of feeding, and health status (Hill et al., 2017; Yao et al., 2021).

Up to 6 mo of life, babies will consume either human breast milk or infant formula as a sole food. Breast milk is the optimum source of food, providing not only nutrients but also immunological factors, probiotics, and bioactive components (Henderson et al., 2022; Phosanam et al., 2021). The composition of breast milk is dynamic, changing not only over stage of lactation but also within a feed from foremilk to hindmilk (Martin et al., 2016). From a nutrient perspective, mature breast milk contains approximately 9 to 12 g/L protein, 67 to 78 g/L lactose, 12 to 14 g/L oligosaccharides, and 32 to 40 g/L fat (Ballard and Morrow, 2013; Kim and Yi, 2020). Mature breast milk also contains minerals such as iron (0.3–0.7 mg/L), calcium (200–250 mg/L), phosphorus (120–140 mg/L), magnesium (30–35 mg/L), sodium (150–250 mg/L), chloride (400–450 mg/L), potassium (400–550 mg/L), manganese (3–4 μ g/L), iodine (140–150 μ g/L), selenium (10–25 μ g/L), copper (0.1–0.3 μ g/L), and zinc (1–3 μ g/L; Kim and Yi, 2020). Whey proteins account for 70% to 80% of total milk protein in early to mid-lactation (Martin et al., 2016), with α -LA accounting for 40% and lactoferrin at 26% of whey protein (Guo and Hendricks, 2008). The CN protein content in breast milk increases from 20% in early lactation to 50% in mature breast milk (Lönnerdal, 2003). The CN in breast milk consists primarily of β -CN and κ -CN but also has trace concentrations of α_{s1} -CN (Basdeki et al., 2021).

Breast milk has a digestible indispensable AA score (**DIAAS**) of 101, determined in 25 d old Yucatan piglets ($n = 9$) who received breast milk for 6 d (Charton et al., 2023). Complete gastric emptying of breast milk in infants ($n = 26$, age 4.7 ± 1.7 mo) takes 2.12 ± 1.07 h, as defined by a cross-sectional area of ≤ 3.07 cm², determined by ultrasound (Das et al., 2024). Using simulated infant conditions, digestibility of breast milk was reported at 48.89% at the end of the gastric phase (60 min; Xiao et al., 2023). Under in vitro infant gastric conditions, breast milk samples formed large curds or aggregates in the stomach with particles sizes ranging between 1 and 10 μ m and 10 and 1,000 μ m (He et al., 2022). Surprisingly, a semidynamic in vitro infant gastrointestinal system observed only 6% protein hydrolysis of breast milk after 120 min gastric digestion but $>38\%$ following intestinal digestion (Abrahamse et al., 2022).

It is widely accepted that infants who receive breast milk have a lower incidence of diarrhea and lower risk of developing necrotizing enterocolitis than bottle-fed babies (Lyons et al., 2020). Preterm infants ($n = 62$, ≤ 32 wk of gestation) receiving breast milk also had significantly lower intestinal permeability (lactulose:mannitol ratio) compared with preterm infants fed infant formula at 1 wk postpartum (0.076 vs. 0.205, $P = 0.04$; Taylor et al., 2009). Rasmussen et al. (2016) also observed significantly higher brush border membrane aminopeptidase N, aminopeptidase A, lactase, and dipeptidyl peptidase IV activities in the proximal small intestine of preterm pigs ($n = 47$, 105–106 d gestation) fed breast milk compared with those fed commercial infant formula ($P < 0.05$; Rasmussen et al., 2016). In human enteroid monolayers, exposure to breast milk (for 24 and 72 h) has been shown to increase goblet cell and Paneth cell number and expression of occludin protein levels (Noel et al., 2021). With increases in digestive enzyme capacity, goblet cell and Paneth cell numbers, tight junction markers, and a reduction in permeability, breast milk is beneficial to the intestinal epithelium, helping to mature the infant gut.

In circumstances where breastfeeding is not possible, the World Health Organization recommends the use of infant formula (World Health Organization, 2013). Under Commission Regulation (EU) No. 609/2013 infant formula is described as a “food intended for use by infants during the first months of life and satisfying by itself the nutritional requirements of such infants until the introduction of appropriate complementary feeding” (Commission Regulation, 2013). Before, during, and after processing infant formula is heat-treated. These high thermal treatment steps ensure a microbiologically safe product with an extended shelf life (Wada and Lönnerdal, 2015b). Bovine milk-based infant formulas are the most common, commercially available infant formula on the market (Halabi et al., 2020b; He et al., 2022). Heat treat-

ment of infant milk formula (IMF) will induce denaturation and aggregation of milk proteins (Pellegrino et al., 2011), and the formation of undesirable Maillard reaction products (Lund et al., 2022). This differs substantially from breast milk. The objective of this narrative literature review is to evaluate the consequences of IMF heat treatment on protein digestion and gut barrier health in the infant gut, using data generated solely from in vitro and preclinical studies. For comparison purposes, in vitro and preclinical data generated with mildly heat-treated IMF will also be summarized. The hypothesis is that mildly heat-treated IMF may be more similar to breast milk in the context of protein digestion and gut physiology in the infant.

INFANT FORMULA COMPOSITION

When producing first-stage infant formula, manufacturers aim to produce a product with a macronutrient and micronutrient composition that mimics breast milk (He et al., 2022). In addition, and in an attempt to achieve the health promoting attributes of breast milk, manufacturers will formulate first-stage infant formulas with additional ingredients, such as omega-3 fatty acids and omega-6 fatty acids, oligosaccharides, probiotics, prebiotics, taurine, lactoferrin, folic acid, L-carnitine, α -LA, osteopontin, and milk fat globule membrane (Almeida et al., 2021, Bakshi et al., 2023).

Skim milk is the starting ingredient for IMF. To achieve a whey-to-CN ratio of 60:40 (similar to mature breast milk), skim milk powder (SMP), whey protein isolate (WPI), or whey protein concentrate (WPC) or combinations are added (Phosanam et al., 2021; Bakshi et al., 2023). The base formula will also contain lactose, vegetable oil, and vitamins and minerals (Bakshi et al., 2023). Protein composition of bovine milk differs from mature breast milk with a whey-to-CN ratio of 20:80 (Navis et al., 2020b) and the presence of β -LG, which is absent in breast milk (Ballard and Morrow, 2013). By correcting the whey-to-CN ratio with the addition of WPI or WPC, the protein content of IMF (1.8–2.5 g/100 kcal) is typically higher than breast milk (EFSA Panel on Dietetic Products and Allergies, 2014).

IMF PROCESSING

Ingredient suppliers rarely disclose the processing histories of their dairy ingredients but traditionally SMP, WPI, and WPC are routinely subjected to thermal processing before IMF manufacturing (Lin et al., 2018). Then, during IMF processing, these same ingredients undergo a series of heat-processing steps including pasteurization (72°C–75°C for 15 s) and UHT (135°C–150°C for 1–10 s; Dash et al., 2022). Finally, for ease of transport and

to extend shelf life, liquid IMF is typically spray dried (inlet temperature 180°C–200°C and outlet temperature 80°C–100°C) to produce an IMF powder (Masum et al., 2020). High thermal loads ensure a microbiologically safe product. Routine monitoring and testing is required in manufacturing facilities for the indicator bacteria, *Enterobacteriaceae* (Commission Regulation [EC] No. 2073/2005; Commission Regulation, 2005). Dried infant formula must not contain *Cronobacter sakazakii* or *Salmonella* spp. (Commission Regulation, 2005).

PROTEIN DENATURATION AND AGGREGATION IN THERMALLY PROCESSED IMF

The effects of heat on milk and milk products have been reviewed previously by Pellegrino et al. (2011). Table 1 summarizes the main effects of heat treatment of IMF ingredients (SMP, WPI, and WPC) and IMF final products on (a) milk protein structure, (b) protein interactions, and (c) Maillard reaction products.

For IMF protein ingredients, Lin et al. (2018) observed that 80.75% of the total whey protein content in skim milk was denatured when this milk was heated at 120°C for 120 s and spray dried at inlet and outlet temperatures of 180°C and 85°C, respectively, to produce SMP (Lin et al., 2018; Table 1). In addition, 86.7% of this denatured whey protein had associated with the CN micelle (Lin et al., 2018). This high-heat-treated SMP had a particle size of 213 nm (Lin et al., 2018). In skim milk heated at 75°C to 100°C, the CN micelle size increased by 30 to 35 nm (Anema and Li, 2003). For WPC, 70% of whey protein was denatured when this ingredient was heated to 73°C for 30 s followed by 80°C for 6 min (Navis et al., 2020b). Only 21% of the protein was soluble (Navis et al., 2020b). Where WPI was preheated to either 72°C for 2 min or 85°C for 2 min, whey protein denaturation was 25.5% and 74.8%, respectively (Joyce et al., 2017). Li et al. (2013) observed that a WPC produced from acid whey that was double pasteurized and spray dried, contained 0.05 g/L native lactoferrin and 0.2 g/L native IgG (Li et al., 2013; Table 1).

Once protein ingredients are sourced and IMF processing begins, it is important to note that protein denaturation and aggregation in heated IMF powders is dependent on β -LG concentration (Crowley et al., 2016). β -lactoglobulin is more heat labile than α -LA because the compact globular structure of α -LA results in higher heat tolerance and a resistance to forming covalent bonds with other proteins (Buggy et al., 2017). Joyce et al. (2017) observed that IMF processed at 85°C for 2 min resulted in $94.7 \pm 0.66\%$ denaturation of β -LG (Joyce et al., 2017). This rose to $98 \pm 0.22\%$ denaturation when a heat-treated WPI ingredient was used (Joyce et al., 2017). Denaturation of α -LA was lower at $78 \pm 1.07\%$, which rose to

Table 1. Effect of processing on IMF protein ingredients and proteins in IMF¹

Ingredient or IMF product	Processing parameter	Effect on protein structure, protein interactions, and AGE	Reference
2 SMP	Ingredient High-heat SMP: 120°C for 120 s Pasteurized SMP: 72°C for 15 s	High-heat SMP ↑ whey protein denaturation, ↑ interactions with CN micelle and ↑ particle size vs. pasteurized SMP	Lin et al., 2018
2 WPC from whole milk produced by cold filtration	Extensively heated WPC: 73°C for 30 s and 80°C for 6 min	Extensively heated WPC ↑ whey protein denaturation and ↑ CML levels vs. pasteurized WPC	Navis et al., 2020b
3 WPC	Pasteurized WPC: 73°C for 30 s Double-pasteurized skim milk WPC: 72°C for 15 s and 80°C for 30 s Stored (37°C for 6 wk) double-pasteurized skim milk WPC	Pasteurized WPC had 100% native whey protein Stored double-pasteurized WPC ↑ CML and furosine levels vs. pasteurized and double-pasteurized skim milk WPC	Aasmul-Olsen et al., 2024
3 WPC	Pasteurized skim milk WPC (from sweet whey): 72°C for 15 s Double-pasteurized WPC (from acid whey), membrane filtered and standard spray dried Pasteurized WPC (from acid whey), membrane filtered and low-temperature spray dried Membrane-filtered WPC (from sweet whey), freeze-dried and γ-irradiated WPI heated at 85°C for 2 min WPI heated at 72°C for 2 min IMF final product IMF heated at 85°C for 2 min IMF heated at 72°C for 2 min	Double-pasteurized WPC: 0.05 g/L native lactoferrin and 0.2 g/L native IgG Pasteurized WPC: 0.66 g/L native lactoferrin and 3.8 g/L native IgG Membrane-filtered WPC: 0.49 g/L native lactoferrin and 4.6 g/L native IgG WPI heated at 85°C for 2 min ↑ whey protein denaturation vs. WPI heated at 72°C for 2 min	Li et al., 2013
2 WPI	IMF heated at 85°C for 2 min	IMF heated at 85°C for 2 min ↑ denaturation of β-LG and α-LA and ↑ protein particle size in (with preheated WPI) vs. IMF heated at 72°C for 2 min	Joyce et al., 2017
2 liquid IMF	IMF heated at 72°C for 2 min	IMF heated at 85°C for 2 min ↑ denaturation of β-LG and α-LA and ↑ protein particle size in (with preheated WPI) vs. IMF heated at 72°C for 2 min	Joyce et al., 2017
2 IMF powders produced at pilot-plant scale (100 kg)	Extensively heated IMF: double pasteurization at 72°C for 30 s and 90°C for 2–3 s, evaporated and heated at 85°C for 2 min and spray dried Membrane-filtered IMF: 0.8- and 0.1-μm membranes and spray dried	Extensively heated IMF ↑ whey protein denaturation vs. membrane-filtered IMF (58 ± 0% vs. 6 ± 4%) Extensively heated IMF ↓ bacterial load No difference in CML or furosine	Yu et al., 2021
2 IMF powders produced at laboratory scale	High-temperature IMF: 125°C for 5 s and spray dried Membrane-filtered IMF: 1.4- and 0.2-μm membranes and spray dried	High-temperature IMF ↓ native whey protein vs. membrane-filtered IMF (2.5% vs. 18.3%) High-temperature IMF ↓ total bacterial count vs. membrane-filtered IMF (5.6 × 10 ¹ vs. 1.3 × 10 ² cfu/g of powder) No difference in CML between high-temperature IMF and membrane-filtered IMF (3.97 ± 0.7 and 3.65 ± 0.74 μg/g of powder)	Chen et al., 2021
2 IMF powders produced at pilot-plant scale (250 kg)	High-temperature IMF: 125°C for 5 s and spray dried Membrane-filtered IMF: 1.4 and 0.2 μm membranes and spray dried	High-temperature IMF ↓ native whey protein vs. membrane-filtered IMF (4.5% vs. 59.9%) High-temperature IMF ↓ total bacterial count vs. membrane-filtered IMF (8.3 × 10 ² vs. 1.8 × 10 ⁷ cfu/g of powder)	Chen et al., 2023; Dold et al., 2024
2 liquid IMF produced at laboratory scale	IMF heated at 90°C for 15 s IMF heated at 75°C for 15 s	IMF heated at 90°C for 15 s ↓ native whey protein vs. IMF heated at 75°C for 15 s	Halabi et al., 2020b

Continued

Table 1 (Continued), Effect of processing on IMF protein ingredients and proteins in IMF¹

Ingredient or IMF product	Processing parameter	Effect on protein structure, protein interactions, and AGE	Reference
2 liquid IMF produced at laboratory scale	IMF with commercial sodium caseinate and WPI in batch sterilized at 110°C for 38 min IMF with sodium caseinate and WPC (acid whey) from pasteurized skim milk in batch sterilized at 110°C for 38 min	In-batch sterilized IMF (with commercial ingredients) ↑ furosine and lysinoalanine levels vs. in-batch sterilized IMF (with sodium caseinate and WPC from pasteurized skim milk) No difference in pyrraline levels	Cattaneo et al., 2017
3 liquid IMF produced at manufacturing scale	UHT IMF: 143°C for 6 s Stored UHT IMF (40°C for 60 d) Pasteurized IMF: 72°C for 10 s	UHT IMF and stored UHT IMF ↑ CML and furosine levels and ↑ protein aggregation vs. pasteurized IMF	Sun et al., 2022

¹SMP = skim milk powder; WPC = whey protein concentrate; WPI = whey protein isolate; IMF = infant milk formula; AGE = advanced glycation end product; CML = N(6)-carboxymethyllysine; CEL = N(6)-carboxyethyllysine. Up and down arrows signify increases and decreases, respectively.

94.2 ± 0.78% with the heat-treated WPI ingredient (Joyce et al., 2017). Not surprisingly, the use of preheated WPI also resulted in increased protein particle size from 176 ± 4 nm to 278 ± 15 nm (Joyce et al., 2017). Fourier-transform infrared spectroscopy data demonstrated that as IMF powders (11 g of protein/100g IMF powder) are heated beyond 50°C, whey protein structures will switch from β-sheets to β-turns, with no changes in α-helices (Ye et al., 2017). In an extensively heat processed IMF powder (double pasteurization of milk at 72°C for 30 s and 90°C for 2–3 s before evaporation and 85°C for 2 min before spray drying), 58% of whey proteins were denatured (Yu et al., 2021). An IMF powder heated at 125°C for 5 s followed by single-stage spray drying contained 97.5% denatured whey protein (Chen et al., 2021). Interestingly, the WPI ingredient used in this formulation already contained 8.5% denatured protein (Chen et al., 2021). Only 5.19% of the total β-LG content in the IMF was in its native, monomeric form compared with 16.57% of the total α-LA (Chen et al., 2021; Table 1).

In the quest to mimic breast milk, IMF manufacturers have toyed with various α-LA:β-LG ratios. Crowley et al. (2016) observed that unheated IMF, regardless of α-LA:β-LG ratio, had protein particle size similar to CN micelles, whereas IMF heated to 140°C had a protein particle size ranging from 319 ± 80.2 nm for IMF (0.1 α-LA:1 β-LG) to 165 ± 2 nm for IMF (4.6 α-LA:1 β-LG; Crowley et al., 2016). Similarly, after heat treatment at 90°C for 15 s, 81% of whey proteins remained in their native state in a liquid IMF formulated with 0.06 g β-LG/100 g IMF in contrast to 48% in an IMF containing 0.51 g β-LG/100 g IMF (Halabi et al., 2020b; Table 1).

Where IMF ingredients are heat-treated or liquid IMF is thermally processed, spray drying has only a minor effect on whey protein denaturation and aggregation (Oldfield et al., 2005; Yu et al., 2021).

ADVANCED GLYCATION END PRODUCTS IN THERMALLY PROCESSED IMF

The Maillard reaction is a nonenzymatic, complex series of reactions between reducing sugars and AA that occurs at high temperatures (Kumar et al., 2017). The final result is “browning,” that is, the formation of advanced glycation end products (AGE) such as N(6)-carboxymethyllysine (CML), N(6)-carboxyethyllysine (CEL), and pyrraline. During thermal processing of IMF, the early-stage Maillard reaction occurs between the carbonyl group of the reducing sugar lactose and an available free amino group, which results in the creation of a Schiff base (Van Boekel, 1998). Rearrangements in the unstable Schiff base leads to the formation of the stable early Maillard reaction products known as Amadori products (Horvat and Jakas, 2004). Detection of furosine

is commonly used as an indicator of early-stage Maillard reaction (Chen et al., 2019; Li et al., 2021b). Furfural compounds such as 5-hydroxymethylfurfural, 5-methyl-2-furaldehyde, 2-furyl-methyl ketone, and 2-furaldehyde are indicators of intermediate Maillard reaction products (Chávez-Servín et al., 2015). Finally, AGE such as CML, CEL, and pyrraline are signals of advanced-stage Maillard reaction (Chen et al., 2019; van der Lugt et al., 2020; Li et al., 2021b). Several studies have reported furosine and CML concentrations in commercially available liquid IMF and IMF powders (Fenaille et al., 2006; Chen et al., 2019). In addition, AGE are commonly quantified during IMF processing and in the final product as an indicator of the extent of Maillard reaction (Chen et al., 2019; Lee et al., 2019; Chen et al., 2021; Yu et al., 2021; Sun et al., 2022; Aasmul-Olsen et al., 2024). Importantly, micronutrients such as vitamin C and iron, added to IMF formulations, can promote the Maillard reaction by acting as precursors (Leclère et al., 2002; Pischetsrieder et al., 2005). However, it must be noted that it is difficult to compare reported levels of Maillard reaction products, including furosine, in final IMF products. Concentrations of these compounds will vary, depending on detection methods used, ingredient quality, protein concentration, and lactose content in the formulation, different IMF processing strategies, and storage conditions.

Table 1 lists the effect of heat treatment of IMF ingredients and IMF final products on Maillard reaction products. Before IMF processing, thermal processing of dairy ingredients will result in the presence of Maillard reaction products where lactose is present. Aasmul-Olsen et al. (2024) reported furosine and CML concentrations of $2,893 \pm 105$ $\mu\text{g/g}$ of protein and 160 ± 4 $\mu\text{g/g}$ of protein, respectively, in a double-pasteurized WPC (72°C for 15 s and 80°C for 30 s) before incorporation into an IMF suitable for piglets (Aasmul-Olsen et al., 2024). The ingredient contained 13.2% lactose (Aasmul-Olsen et al., 2024). However, CEL levels were below the limits of detection of liquid chromatography (LC)-MS/MS (Aasmul-Olsen et al., 2024). Using ultra-fast LC/fluorescence, Navis et al. (2020b) reported approximately 19 $\mu\text{g/mL}$ CML in an extensively heated WPC (73°C for 30 s and 80°C for 6 min; Navis et al., 2020b; Table 1).

Cattaneo et al. (2017), using in-batch sterilization produced liquid IMF by sourcing commercial sodium caseinate and WPI (Cattaneo et al., 2017). This IMF had $6,378 \pm 225$ μg furosine/g of protein, 55 ± 2 μg pyrraline/g of protein and 999 ± 95 μg lysinoalanine/g of protein (Cattaneo et al., 2017; Table 1).

In UHT-treated liquid IMF, quantified by LC-MS/MS, Sun et al. (2022) reported furosine, CML, and CEL concentrations of $2,798 \pm 136$ $\mu\text{g/g}$ of protein, 242 ± 9 $\mu\text{g/g}$ of protein and 71 ± 1 $\mu\text{g/g}$ of protein, respectively (Sun et al., 2022). After spray drying, a heat-treated IMF powder

contained 3.97 ± 0.7 μg CML/g of powder (Chen et al., 2021). In an extensively heat processed IMF powder, furosine concentration was approximately 5,000 $\mu\text{g/g}$ of protein and CML concentrations was 106 ± 0 $\mu\text{g/g}$ of protein (Yu et al., 2021; Table 1).

Lee et al. (2019) reported a contribution of the spray-drying process to CML content in IMF powder, with an increase from approximately 30 $\mu\text{g/g}$ of protein in the heat-treated liquid IMF to 60 $\mu\text{g/g}$ of protein in the final powder, albeit no processing temperatures before spray drying were given (Lee et al., 2019). Yu et al. (2021) have previously shown no contribution by spray drying to furosine or CML content in IMF with high thermal loads. However, it is well established that spray drying and storage of dairy powders will promote the Maillard reaction (van Lieshout et al., 2020). The level of AGE in the final IMF powder will be determined by the level of protein denaturation during processing, the inlet and outlet temperatures of spray drying, storage temperature, and relative humidity (Zhu et al., 2018; van Lieshout et al., 2020).

INFLUENCE OF THERMAL PROCESSING OF IMF ON PROTEIN DIGESTION IN THE INFANT

Thermally processed IMF will deliver aggregated, denatured proteins and AGE to the infant gut. The protein digestibility (DIAAS) of heat-treated IMF in the infant gut cannot be generalized because it is governed by the extent of these factors in the specific IMF product and the level they interfere with digestive enzymes. Recently, van Lieshout et al. (2020) and Li et al. (2021a) reviewed the effect of these components, within a variety of dairy products, on protein digestion in the adult stomach and small intestine (van Lieshout et al., 2020; Li et al., 2021a). Specifically, Charton et al. (2023) recently reported a DIAAS of 83 for an extensively heat processed IMF powder, which is considerably lower than breast milk at a DIAAS of 101, as determined in young pigs. Table 2 summarizes the effect of IMF processing on protein digestion in the infant gut using data generated from in vitro and preclinical models.

Chen et al. (2022) produced a high-temperature (125°C for 5 s) IMF powder, which was subjected to an in vitro semidynamic digestion mimicking the physiological conditions of the infant stomach (Chen et al., 2022). Once IMF was added to the gastric chamber, the pH rose immediately from 2.5 to 7.57 ± 0.13 (Chen et al., 2022). A large compact curd was formed and remained to the end of the gastric phase (156 min, pH 3.2; Chen et al., 2022). Confocal laser scanning microscopy revealed microstructures of protein aggregates remaining on fat droplet surfaces at 156 min of gastric digestion (Chen et al., 2022). The SDS-PAGE analysis showed

Product	Processing parameter	Method	Measure of digestibility	Effect on protein digestibility	Reference
2 IMF powders produced at laboratory scale	High-temperature IMF: 125°C for 5 s and spray dried Membrane-filtered IMF: 1.4- and 0.2-μm membranes and spray dried	INFOGEST in vitro semidynamic infant gastric digestion and INFOGEST in vitro static infant gastrointestinal digestion	SDS-PAGE and degree of protein hydrolysis	High-temperature IMF ↑ gastric pH at 0 and 31 min, ↓ protein hydrolysis and ↑ β-LG digestion vs. membrane-filtered IMF High-temperature IMF formed large aggregates in stomach High-temperature IMF ↓ water-soluble material in gastric phase at 0, 31, 62, 94, and 125 min vs. membrane-filtered IMF High-temperature IMF and membrane-filtered IMF had similar protein hydrolysis and no intact proteins visible at 60 min of intestinal digestion	Chen et al., 2022
2 IMF powders produced at pilot-plant scale (250 kg)	High-temperature IMF: 125°C for 5 s and spray dried Membrane-filtered IMF: 1.4- and 0.2-μm membranes and spray dried	28-d-old pigs (n = 20) fed IMF diets (35% inclusion) for 28 d	Degree of protein hydrolysis and AA analysis	High-temperature IMF ↓ protein hydrolysis in stomach and duodenum vs. membrane-filtered IMF (stomach: 509 ± 17 vs. 599 ± 45 μmol free amines/g of protein and duodenum: 1,174 ± 124 vs. 1,530 ± 136 μmol free amines/g of protein) High-temperature IMF ↓ protein/peptide content in lumen, ↓ release of Pro in duodenum and ↓ taurine and Trp in jejunum vs. membrane-filtered IMF Similar DIAAS for extensively heated IMF and membrane-filtered IMF (0.82 and 0.8)	Chen et al., 2023
2 IMF powders produced at pilot-plant scale (100 kg)	Extensively heated IMF: double pasteurization at 72°C for 30 s, 90°C for 2–3 s before evaporation and 85°C for 2 min before spray drying Membrane-filtered IMF: 0.8- and 0.1-μm membranes and spray dried	28 d Wistar Han rats (n = 60) fed IMF diets (40% inclusion) for 14 d	Total nitrogen digestibility, DIAAS, and plasma AA concentration	Extensively heated IMF ↓ plasma indispensable and total AA concentrations 1 h after ingestion vs. membrane-filtered IMF No change in dietary nitrogen in stomach and SI in extensively heated IMF vs. membrane-filtered IMF (stomach: 0.14 ± 0.34% vs. 0.04 ± 0.09% and SI: 1 ± 1.66% vs. undetectable)	Calvez et al., 2024
2 liquid IMF produced at laboratory scale	IMF heated at 67.5°C for 200 min or 80°C for 7 min Unheated IMF	INFOGEST in vitro static infant gastrointestinal digestion	SDS-PAGE and AA analysis	Heated IMF ↓ protein hydrolysis vs. unheated IMF at 5 min of intestinal digestion	Halabi et al., 2020a
2 IMF powders produced at manufacturing scale	IMF dry heated at 70°C for 335 h Unheated IMF	INFOGEST in vitro static infant gastrointestinal digestion	SDS-PAGE and degree of protein hydrolysis	No difference in bioaccessibility of AA Dry-heated IMF ↓ protein hydrolysis vs. unheated IMF at 60 min of gastric digestion and 120 min of intestinal digestion Dry-heated IMF ↑ peptide length in intestinal digesta vs. unheated IMF	Zenker et al., 2020

Journal of Dairy Science Vol. 108 No. 4, 2025

Table 2 (Continued). Effect of IMF processing on protein digestibility¹

Product	Processing parameter	Method	Measure of digestibility	Effect on protein digestibility	Reference
2 liquid IMF produced at manufacturing scale	UHT IMF: 143°C for 6 s (direct or indirect) Pasteurized IMF: 72°C for 10 s	In vitro static infant gastrointestinal digestion	SDS-PAGE and LC-MS/MS	β -LG undigested in pasteurized IMF UHT IMF $\uparrow$ protein aggregates in digesta vs. pasteurized IMF	Ye et al., 2022
2 liquid IMF produced at laboratory scale with commercial sodium caseinate and WPI or sodium caseinate and WPC (acid whey) from pasteurized skim milk	IMF in batch sterilized at 110°C for 38 min Unheated IMF	INFOGEST in vitro static gastrointestinal digestion	Degree of proteolysis calculated based on nitrogen content of a 3 kDa fraction of digesta	IMF with commercial sodium caseinate and WPI: in-batch sterilized IMF $\uparrow$ proteolysis vs. unheated IMF (40.3 $\pm$ 1.2% vs. 35.6 $\pm$ 0.6%) IMF with sodium caseinate and WPC (acid whey) from pasteurized skim milk: in-batch sterilized IMF $\downarrow$ proteolysis vs. unheated IMF (54 $\pm$ 0.9% vs. 35% $\pm$ 0.9%)	Cattaneo et al., 2017

¹IMF = infant milk formula; DIAAS = digestible indispensable amino acid score; SI = small intestine.

that at 125 min intact CN were still present (Chen et al., 2022). By the end of the gastric phase, β -LG bands had weakened considerably but α -LA was still present (Chen et al., 2022). The degree of protein hydrolysis, determined by the o-phthalaldehyde method, at the end of the gastric phase was 845 ± 78 μ mol free amines/g of protein (Chen et al., 2022). Not surprisingly, the release of peptides (<10 kDa) from heated IMF digesta increased significantly from the beginning to the end of gastric digestion ($P < 0.05$; Chen et al., 2022). At the end of the intestinal phase, using an in vitro static protocol for infant digestion (Ménard et al., 2018), no intact proteins were visible by SDS-PAGE with this high-temperature IMF (Bavaro et al., 2021). In a follow-on preclinical study, young pigs were fed a diet composed of this high-temperature IMF and lumen samples were collected from 4 different gut locations 180 min after the final meal (Chen et al., 2023). Pigs fed the heated IMF had a degree of protein hydrolysis of 509 ± 17 μ mol free amines/g of protein in the stomach and $1,174 \pm 124$ μ mol free amines/g of protein in the duodenum, with an overall value for the stomach, duodenum, jejunum, and ileum of $1,022 \pm 43$ μ mol free amines/g of protein (Chen et al., 2023). The concentration of free AA released was 59.4 ± 3.05 μ mol/g of protein in the gastric phase, 171.9 ± 29.7 μ mol/g of protein in the duodenum, 211.4 ± 22.75 μ mol/g of protein in the jejunum, and 285 ± 28.51 μ mol/g of protein in the ileum (Chen et al., 2023). In contrast, Calvez et al. (2024) fed Wistar Han rats (n = 30, 28 d old) with extensively heat-treated IMF at 40% inclusion (Calvez et al., 2024). Levels of dietary nitrogen recovered in the stomach ($0.14 \pm 0.34\%$) and the small intestine ($1 \pm 1.66\%$) 6 h after ingestion were not influenced by heating (Calvez et al., 2024).

The results of Halabi et al. (2020a) showed that in vitro infant gastric digestion of heat-treated IMF (80°C) formed small curd particles of 10 μ m. Thermal treatment of IMF at either 67.5°C or 80°C supported gastric CN hydrolysis with only $6 \pm 1\%$ of residual intact CN still present after 30 min of gastric digestion (Halabi et al., 2020a). However, the degree of protein hydrolysis was low in IMF heated at 80°C, quantified at only $2 \pm 1\%$ after 60 min of gastric digestion, rising to $14 \pm 2\%$ within 5 min of the intestinal phase and finishing at $35 \pm 2\%$ at the end of the intestinal phase (Halabi et al., 2020a). The bioaccessibility of AA as a percentage of total AA was lower than expected, with $35.5 \pm 1.2\%$ for EAA and $10.3 \pm 0.6\%$ for NEAA in IMF heated at 80°C (Halabi et al., 2020a). Lysine, Phe, Tyr, Leu, and Arg were higher than all other AA, and free Pro concentration was negligible (Halabi et al., 2020a). Gómez-Gallego et al. (2016) produced an IMF powder with heat treatment at 70°C and then 105°C before spray drying. An in vitro infant gastrointestinal digestion protocol was performed (Gómez-

Gallego et al., 2016). No intact proteins were visible at the end of the intestinal phase by SDS-PAGE and levels of NPN rose from 0.19 ± 0.02 mg/g before digestion to 0.95 ± 0.2 mg/g after digestion yielding an estimated in vitro digestibility of $45.19 \pm 11.29\%$ (Gómez-Gallego et al., 2016). Liquid chromatography-MS/MS identified 87 unique peptides from in vitro digestion of the final IMF product at the end of the intestinal phase (Gómez-Gallego et al., 2016). Cattaneo et al. (2017) reported that the degree of proteolysis, during in vitro infant gastrointestinal of in-batch sterilized liquid IMF, was $40.3 \pm 1.2\%$ (Cattaneo et al., 2017). This was calculated based on the nitrogen content of a 3 kDa fraction of digesta (Cattaneo et al., 2017). A standard IMF produced by pasteurization, evaporation, and spray drying underwent an in vitro semidynamic gastric digestion simulating infant digestion (Lambers et al., 2023). The degree of protein hydrolysis ranged from approximately 0.11 to 0.18 μmol free amines/mg of protein from the early stage of gastric digestion at pH 6 to the end of digestion (final pH of 3.5; Lambers et al., 2023). The molecular weight distribution analysis revealed that approximately 53% of peptides were >50 kDa at pH 3.5, indicating low levels of digestion in the gastric phase (Lambers et al., 2023). Ye et al. (2022) produced a liquid IMF by direct or indirect UHT (143°C for 6 s) processing (Ye et al., 2022). This study performed an in vitro static infant gastrointestinal digestion with gastric digestion at pH 4 for 2 h followed by 6 h intestinal digestion at pH 6.5 (Ye et al., 2022). At the end of the gastric phase, aggregates (>260 kDa) in both UHT IMF remained resistant to pepsin (Ye et al., 2022). Following intestinal digestion all protein bands, including aggregates, were digested (Ye et al., 2022). The LC-MS/MS analysis revealed the total number of peptides released during digestion was similar in both direct and indirect UHT treatments (Ye et al., 2022). Interestingly, there were significantly more β -CN peptides in the intestinal digesta from indirect UHT IMF compared with the direct UHT IMF ($P < 0.05$; Ye et al., 2022). Direct UHT processing had a lower heat load indicated by significantly lower levels of lactosylation in released β -LG peptides from direct UHT IMF compared with indirect UHT IMF ($P < 0.05$; Ye et al., 2022).

Previous studies have shown that structural changes to dairy proteins induced by the Maillard reaction can influence protein digestion by exposing or blocking enzymatic cleavage sites (Joubran et al., 2015; Joubran et al., 2017). Zenker et al. (2020) induced a Maillard reaction in a commercially-sourced IMF powder by dry heating to 70°C for 335 h (Zenker et al., 2020). This generated a powder with 2,022 mg furosine/100 g of protein (Zenker et al., 2020). At the end of in vitro infant gastric digestion, the degree of protein hydrolysis was only $4.3 \pm 0.2\%$ rising to $29.8 \pm 0.8\%$ at the end of the intestinal

phase (Zenker et al., 2020). A rapid disappearance in CN bands was observed in the gastric phase whereas intact β -LG was still present at 60 min of intestinal digestion (Zenker et al., 2020). This corresponds to results from peptide identification experiments where fewer whey protein-derived peptides ($3.9 \pm 0.19\%$) were identified compared with CN-derived peptides ($96.1 \pm 0.19\%$) at the end of the intestinal phase (Zenker et al., 2020). Levels of furosine in the IMF did not influence peptide size distribution at the end of the gastric phase but at the end of the intestinal phase there were significantly more peptides with 15–19 AA, 25–29 AA, and 30–34 AA, and significantly fewer peptides with 5–9 AA compared with the control IMF (308 mg furosine/100 g of protein; $P < 0.05$; Zenker et al., 2020). Overall, the average peptide length was significantly higher in dry-heated IMF compared with the control IMF at the end of the intestinal phase ($P < 0.05$; Zenker et al., 2020). Zenker et al. (2020) calculated that 44.5% of the AA Lys was blocked from enzymatic hydrolysis due to its reaction with lactose (Zenker et al., 2020). This blockage impaired the breakdown of larger peptides (Zenker et al., 2020).

IMPACT OF THERMAL PROCESSING OF IMF ON GUT BARRIER HEALTH

Recently, researchers have used both in vitro and pre-clinical approaches to examine the relationship between thermally processed IMF and gut barrier health (Li et al., 2013; Navis et al., 2020a; Navis et al., 2020b; Bavaro et al., 2021; Sun et al., 2022; Aasmul-Olsen et al., 2024; Dold et al., 2024). Table 3 summarizes the results of these studies. In many cases, differences observed in gut barrier response have been attributed to higher levels of Maillard reaction products in IMF with high thermal loads (Sun et al., 2022; Aasmul-Olsen et al., 2024). Individual AGE compounds can interact with the gut barrier directly to mediate an adverse effect (Guibourdenche et al., 2021; Wu et al., 2022). However, Maillard reaction products in thermally processed IMF may not be the sole contributor to barrier response, with protein structure also playing a role (Li et al., 2013; Navis et al., 2020a; Navis et al., 2020b; Bavaro et al., 2021; Dold et al., 2024). Indeed it is likely that both protein structure and AGE levels will modulate digestive enzyme accessibility to milk protein cleavage sites, resulting in differences in bioactive peptides released during digestion. This difference in bioactive capacity may also drive gut response.

Sun et al. (2022) fed preterm pigs UHT liquid IMF for 5 d by an orogastric feeding tube. A wide range of Maillard reaction products and cross-linked AA were quantified in the UHT IMF (Sun et al., 2022). To further increase these levels, the UHT IMF was stored at 40°C for 60 d before the intervention trial (Sun et al., 2022). The ileal mucosa

Table 3. Impact of IMF processing on gut barrier health¹

Product	Processing parameter	Method	Gut barrier health assessment	Impact on gut barrier health	Reference
3 liquid IMF produced at manufacturing scale	UHT IMF: 143°C for 6 s Stored UHT IMF (40°C for 60 d) Pasteurized IMF: 72°C for 10 s	Preterm pigs (n = 56, 106-d gestation) received IMF by orogastric feeding tube for 5 d	Histology, intestinal permeability (lactulose:mannitol ratio), and brush border enzyme activity	UHT IMF ↓ SI villus height and ↑ distal SI crypt depth vs. pasteurized IMF Stored UHT IMF ↓ SI villus height, ↑ SI crypt depth and ↓ proximal and middle SI lactase activity vs. pasteurized IMF No difference in lactulose:mannitol ratio IMF with stored double-pasteurized WPC ↑ CML and furosin in distal SI tissue vs. IMF with pasteurized WPC and double-pasteurized WPC No change in villus height, crypt depth, and brush border enzymes	Sun et al., 2022
3 IMF powders produced with different WPC	Double-pasteurized skim milk WPC: 72°C for 15 s and 80°C for 30 s Stored (37°C for 6 wk) double-pasteurized skim milk WPC Pasteurized skim milk WPC (from sweet whey): 72°C for 15 s	Preterm piglets (n = 72, 105-d gestation) received IMF by orogastric feeding tube for 5 d	Histology, AGE quantified in distal SI tissue by LC-MS/MS, and brush border enzyme activity		Aasmul-Olsen et al., 2024
2 IMF powders produced at pilot-plant scale (250 kg)	High-temperature IMF: 125°C for 5 s and spray dried Membrane-filtered IMF: 1.4- and 0.2-µm membranes and spray dried	28-d-old pigs (n = 20) fed IMF diets (35% inclusion) for 28 d	Histology, RT-qPCR, Western blot, acidic mucus levels, and brush border enzyme activity	In duodenum high-temperature IMF ↓ brush border lactase activity and ↓ <i>claudin-1</i> , <i>mucin-1</i> , and <i>mucin-2</i> mRNA transcripts vs. membrane-filtered IMF In jejunum high-temperature IMF ↓ goblet cell number, ↓ acidic mucus levels, ↓ mucin-2 protein, and ↑ <i>RAGE</i> mRNA transcripts vs. membrane-filtered IMF No change in villus height or crypt depth	Dold et al., 2024
2 IMF powders produced at laboratory scale	High-temperature IMF: 125°C for 5 s and spray dried Membrane-filtered IMF: 1.4- and 0.2-µm membranes and spray dried	INFOGEST in vitro static infant gastrointestinal digestion and 21-d-old Caco-2 monolayers treated with 200 µg of protein/cm ²	TEER and Western blot	High-temperature IMF ↓ TEER and claudin-1 protein in Caco-2 monolayers vs. membrane-filtered IMF	Bavaro et al., 2021
3 liquid IMF produced with different WPC	IMF with double-pasteurized WPC (from acid whey) IMF with pasteurized WPC (from acid whey) IMF with membrane-filtered WPC (from sweet whey)	Preterm pigs (n = 55, 105-d gestation) received IMF by orogastric feeding tube for 5 d	Histology, Western blot, intestinal permeability (lactulose:mannitol ratio), inflammatory biomarkers, and brush border enzyme activity	IMF with double-pasteurized WPC ↓ proximal SI villus height vs. IMF with pasteurized WPC and IMF with membrane-filtered WPC IMF with double-pasteurized WPC ↑ lactulose:mannitol ratio vs. IMF with membrane filtered WPC No change in inflammatory biomarkers	Li et al., 2013
2 liquid IMF produced with different WPC	IMF with extensively heated WPC: 73°C for 30 s and 80°C for 6 min IMF with pasteurized WPC: 73°C for 30 s	Preterm (n = 34, 106-d gestation) and near-term (n = 18, 113-d gestation) piglets received IMF by orogastric feeding tube for 5 d	Histology, brush border enzyme activity, intestinal permeability (lactulose:mannitol ratio), and inflammatory biomarkers	IMF with extensively heated WPC ↑ crypt depth, ↓ colonic goblet cells, ↓ intestinal alkaline phosphatase activity, ↑ lactulose:mannitol ratio and ↑ IL-8, IL-1β, and TNF-α vs. IMF with pasteurized WPC	Navis et al., 2020a,b

¹IMF = infant milk formula; WPC = whey protein concentrate; AGE = advanced glycation end product; SI = small intestine; RT-qPCR = reverse transcription quantitative real-time PCR; TEER = transepithelial electrical resistance; CML = N(6)-carboxymethyllysine; RAGE = receptor for advanced glycation end products; IL-8 = interleukin-8; IL-1β = interleukin-1β; TNF-α = tumor necrosis factor α.

of pigs who received stored UHT IMF had furosine levels $>450 \mu\text{g/g}$ of dry weight and CML levels $>130 \mu\text{g/g}$ of dry weight (Sun et al., 2022). Preterm pigs fed the stored UHT IMF had significantly lower villus height at the proximal small intestine compared with those fed the UHT IMF ($P < 0.05$), but similar villus height in the middle and distal small intestine (Sun et al., 2022). Pigs who received either UHT IMF or stored UHT IMF had similar crypt depth and brush border lactase activity in the small intestine (Sun et al., 2022). Gut permeability (lactulose:mannitol ratio) did not decrease by increasing AGE levels in the IMF (Sun et al., 2022). Also, no changes occurred in intestinal mRNA transcript levels of pathogen recognition receptors, acute inflammatory response, and oxidative stress markers with increasing levels of AGE in IMF (Sun et al., 2022). However, dietary intervention with stored UHT IMF resulted in significant increases in mRNA transcript levels of AGE receptors compared with UHT IMF ($P < 0.05$; Sun et al., 2022).

Preterm piglets were fed for 5 d, by orogastric feeding tube, an IMF produced with a double-pasteurized WPC (Aasmul-Olsen et al., 2024). Again, in this study storage of the WPC ingredient was used to increase levels of furosine and CML in the final IMF product (Aasmul-Olsen et al., 2024). In contrast to the results of Sun et al. (2022), there was no significant effect of processing on villus height and crypt depth ($P > 0.05$; Aasmul-Olsen et al., 2024). Brush border aminopeptidase N, aminopeptidase A, lactase, sucrase, maltase, and dipeptidyl peptidase IV activities were all similar in the small intestine of piglets fed the IMF with double-pasteurized WPC or stored double-pasteurized WPC (Aasmul-Olsen et al., 2024). In the distal small intestine, tissue furosine and CML levels were significantly higher in piglets fed IMF with stored double-pasteurized WPC ($2,894 \pm 427 \mu\text{g/g}$ of protein and $639 \pm 67 \mu\text{g/g}$ of protein) compared with those piglets fed the IMF with double-pasteurized WPC ($1,301 \pm 188 \mu\text{g/g}$ of protein and $331 \pm 38 \mu\text{g/g}$ of protein; $P < 0.01$; Aasmul-Olsen et al., 2024).

We have recently fed young pigs a high-temperature IMF ($3.97 \mu\text{g CML/g}$; 35% inclusion in the diet) for 28 d (Chen et al., 2023). Both the duodenum and jejunum were assessed for gut physiology and digestive enzyme capacity (Dold et al., 2024). For example, in the duodenum of pigs fed high-temperature IMF diets, villus height was $449.60 \pm 24.72 \mu\text{m}$, crypt depth was $366.78 \pm 23.63 \mu\text{m}$, and goblet cell number was 30.81 ± 1.67 (Dold et al., 2024). In the duodenal lumen, trypsin activity was $0.91 \pm 0.2 \text{ U/mg}$ of protein and in the duodenal brush border membrane vesicles aminopeptidases N activity was $7.44 \pm 0.69 \text{ U/mg}$ of protein, aminopeptidase A activity was $155.17 \pm 22.83 \text{ mU/mg}$ of protein, intestinal alkaline phosphatase levels were $167.44 \pm 30.49 \text{ U/mg}$ of protein and lactase activity was $110.75 \pm 7.21 \text{ nmol glucose/}$

mg of protein (Dold et al., 2024). Previously, treatment of polarized human epithelial cell monolayers (Caco-2) with in vitro gastrointestinal digested high-temperature IMF resulted in a significant reduction in monolayer integrity (transepithelial electrical resistance [TEER]) compared with untreated control cell monolayers ($P < 0.05$; Bavaro et al., 2021). However, this barrier disruption did not alter levels of the tight junction proteins occludin, claudin-4, and zonula occludens-1 or junctional adhesion molecule-A (Bavaro et al., 2021).

Li et al. (2013) described a study where preterm pigs received an IMF formulated with double-pasteurized, spray-dried WPC for 5 d by orogastric feeding tube (Li et al., 2013). Maillard reaction products were not determined in the final formulation (Li et al., 2013). In the proximal small intestine these preterm pigs had a villus height of $440 \pm 47 \mu\text{m}$ (Li et al., 2013). Brush border enzyme activities were determined in small intestinal tissue of these preterm pigs with an aminopeptidase N activity of $5.9 \pm 1.4 \text{ U/g}$ of tissue, aminopeptidase A activity of $2.5 \pm 0.6 \text{ U/g}$ of tissue, lactase activity of $9.7 \pm 0.8 \text{ U/g}$ of tissue, sucrase activity of $0.15 \pm 0.01 \text{ U/g}$ of tissue, maltase activity of $3.1 \pm 0.5 \text{ U/g}$ of tissue, and dipeptidyl peptidase IV activity of $2.3 \pm 0.7 \text{ U/g}$ of tissue (Li et al., 2013). The inflammatory biomarker, IL-8 in the distal small intestinal tissue of preterm pigs was approximately 58 pg/mg of tissue (Li et al., 2013). The urinary lactulose:mannitol ratio was >0.5 (Li et al., 2013).

In a 5-d intervention trial, preterm and near-term piglets received increasing doses, by orogastric feeding tube, of an IMF produced using an extensively heated WPC (Navis et al., 2020a,b). This WPC had approximately $19 \mu\text{g CML/mL}$ (Navis et al., 2020b). The colon crypt depth in the piglets from both age groups fed this IMF formulated with extensively heated WPC was $>200 \mu\text{m}$ (Navis et al., 2020a).

ALTERNATIVES TO THERMAL PROCESSING FOR IMF PRODUCTION

To preserve native proteins and reduce AGE levels in IMF, recent research efforts have focused on reducing thermal loads by (a) sourcing mildly processed protein ingredients with lower thermal loads (i.e., sweet whey), (b) by decreasing IMF processing temperatures or looking to alternative processing technologies (membrane filtration and UV irradiation), or both, and (c) by eliminating spray-drying steps (i.e., liquid IMF).

Food authorities are likely to limit AGE concentrations in IMF in the near future. Certainly, the EU Commission Regulation (EU) No. 2017/2158 has already set the benchmark level for the presence of acrylamide in “baby foods” at $40 \mu\text{g/kg}$ of BW (Commission Regulation, 2017). Acrylamide is a toxic, potentially carcinogenic,

byproduct of the Maillard reaction (Augustine and Bent, 2022). Sourcing ingredients manufactured with lower thermal loads and lowering the temperature during IMF processing will result in lower levels of AGE (Navis et al., 2020b; Aasmul-Olsen et al., 2024; Table 1). From an ingredient perspective, Navis et al. (2020b) noted that pasteurized WPC (73°C for 30 s) had significantly lower levels of CML compared with an extensively heated WPC (73°C for 30 s and 80°C for 6 min; $P < 0.05$; Navis et al., 2020b). The same pasteurized WPC had close to 100% native whey protein compared with only 30% in the extensively heated WPC (Navis et al., 2020b). In this pasteurized WPC, 72% soluble protein remained following processing compared with only 21% in the extensively heated WPC (Navis et al., 2020b). In a pasteurized WPC (72°C for 15 s) there was significantly lower furosine and CML levels compared with a double-pasteurized WPC (72°C for 15 s and 80°C for 30 s) stored at 37°C for 6 wk ($3,172 \pm 231$ and 182 ± 13 µg/g of protein vs. $6,166 \pm 115$ and 399 ± 14 µg/g of protein; $P < 0.05$; Aasmul-Olsen et al., 2024). No distinctive protein aggregates were observed by SDS-PAGE in the pasteurized WPC compared with large disulfide-linked protein aggregates seen in the double-pasteurized WPC (Aasmul-Olsen et al., 2024). Lin et al. (2018) observed significantly lower levels of whey protein denaturation in a pasteurized SMP versus a high-heat-treated SMP (5.44% vs. 80.75%; $P < 0.05$; Lin et al., 2018). A significant reduction in particle size to 179 nm was seen in pasteurized SMP versus 213 nm in the SMP with high-heat ($P < 0.05$; Lin et al., 2018). Where WPC is produced using sweet whey rather than acid whey, lower temperatures are often employed. The study by Cattaneo et al. (2017) produced sodium caseinate and WPC by skim milk acidification before IMF processing (Cattaneo et al., 2017). Furosine ($3,492 \pm 110$ µg/g of protein) and lysinoalanine (385 ± 43 µg/g of protein) were significantly reduced compared with IMF formulated with commercial ingredients ($P < 0.05$; Cattaneo et al., 2017). There was no change in pyrroline levels (Cattaneo et al., 2017; Table 1). A WPC, produced using membrane filtration (sweet whey) and sterilized by γ -irradiation, had higher amounts of native lactoferrin (0.49 g/L) and native IgG (4.6 g/L) compared with a double-pasteurized WPC from acid whey (Li et al., 2013; Table 1).

Lowering the temperature during IMF processing is also an option (Lund et al., 2022; Sun et al., 2022). Reducing the thermal load from 90°C for 15 s to 72°C for 15 s increased retention of native whey proteins in liquid IMF to 98% and 99%, respectively (Halabi et al., 2020b; Table 1). Pasteurized liquid IMF (72°C for 10 s) had lower furosine and CML concentrations ($1,797 \pm 59$ µg/g of protein and 38 ± 2 µg/g of protein) compared with the UHT liquid IMF (143°C for 6 s; $2,798 \pm 136$ µg/g of protein and 242 ± 9 µg/g of protein; Sun et al.,

2022). The CEL concentrations were similar between pasteurized and UHT IMF (Sun et al., 2022). Not surprisingly, storage of the same UHT liquid at 40°C for 60 d increased furosine, CML, and CEL concentrations (Sun et al., 2022). Sodium dodecyl sulfate-PAGE revealed less protein aggregation in pasteurized liquid IMF compared with UHT-treated liquid IMF and stored UHT liquid IMF (Sun et al., 2022). An IMF powder produced with pasteurized skim milk and WPC (72°C for 15 s) had a CML concentration of 143 ± 15 µg/g of protein, a CEL concentration of 8.99 ± 1.93 µg/g of protein, and pyrroline levels of 36.2 ± 2.9 µg/g of protein (Lund et al., 2022). However, in these studies the total bacterial load of these low thermally processed IMF was not reported, which is critical to determine whether a product is safe for human consumption.

Membrane filtration technology using filters to trap bacteria has been used recently to produce IMF at pilot-plant scale (Chen et al., 2021; Yu et al., 2021). Yu et al. (2021) produced 100 kg of IMF powder (whey:CN ratio of 60:40) from fresh bovine milk using 0.8- and 0.1-µm pore size membranes (Yu et al., 2021). Reverse-phase HPLC revealed significantly lower levels of whey protein denaturation in the membrane-filtered IMF powder ($6 \pm 4\%$ denatured whey protein) compared with the IMF powder produced using extensive heat processing (58% denatured whey protein; $P < 0.05$; Yu et al., 2021; Table 1). Surprisingly, using LC-MS/MS, Yu et al. (2021) found no significant differences in AGE (furosine and CML) concentrations between IMF at different stages of production, from mixing to final powder ($P > 0.05$; Yu et al., 2021). In the final IMF powder collected after spray drying, furosine levels were comparable between the membrane-filtered and extensively heat processed IMF powders (approximately 4,800 and 5,000 µg furosine/g of protein; Yu et al., 2021). The CML concentrations were 88 ± 2 µg/g of protein in the membrane-filtered IMF and 106 ± 0 µg/g of protein in the extensively heat processed IMF (Yu et al., 2021). From a safety perspective, membrane-filtered IMF contained 3.63×10^2 cfu/g of powder compared with only 0.95×10^2 cfu/g of powder in the extensively heated IMF powder (Yu et al., 2021), albeit absent of *C. sakazakii* and *Salmonella* spp., adhering to the specifications outlined under Commission Regulation (EC) No. 2073/2005 on microbiological criteria for foodstuffs (Commission Regulation, 2005).

In our group, a ceramic 1.4-µm microfiltration membrane followed by a polyvinylidene difluoride polymeric 0.2-µm membrane was used to produce an IMF powder (whey:CN ratio of 60:40) at laboratory scale (Chen et al., 2021). A control, high-temperature IMF was also produced by heating at 125°C for 5 s (Chen et al., 2021). Both homogenized IMF were spray dried at 185°C (inlet temperature) and 80°C (outlet temperature; Chen et al.,

2021). There was no significant difference in CML concentrations in both spray-dried IMF powders (3.65 ± 0.74 µg/g of membrane-filtered IMF powder vs. 3.97 ± 0.7 µg/g of high-temperature IMF powder; $P > 0.05$; Chen et al., 2021; Table 1). Taken together with the results of Yu et al. (2021), this may indicate that where thermal load is low before entering the dryer, spray drying then becomes a major contributor to CML levels in IMF.

At pilot-plant scale (250 kg), this membrane-filtered IMF had a significant increase in native whey protein content (59.9%, 4.02 g native whey protein) compared with high-temperature IMF (4.5%, 0.3 g native whey protein; $P < 0.001$; Chen et al., 2021; Chen et al., 2023). There was no loss in protein by membrane filtration with no differences in total protein content (10.9 g total protein/100 g of membrane-filtered IMF vs. 11.4 g total protein/100 g of high-temperature IMF; Chen et al., 2021; Chen et al., 2023). In agreement with Yu et al. (2021), pathogenic bacteria were absent in the final powders, but the total bacterial load was significantly higher in membrane-filtered IMF (1.8×10^7 cfu/g of powder) compared with high-temperature IMF (8.3×10^2 cfu/g of powder; $P < 0.05$; Dold et al., 2024).

Proof of principle research has demonstrated that UV irradiation can reduce the bacteria load from commercial IMF powder spiked with *C. sakazakii* (Liu et al., 2012). However, manufacturing IMF at pilot-plant scale using UV irradiation as an alternative to thermal processing is still under development.

WILL REDUCTION IN THERMAL PROCESSING OF IMF ALTER PROTEIN DIGESTIBILITY?

Limiting the exposure of ingredients and IMF to high thermal loads during processing may influence protein digestion in the upper infant gut (Cattaneo et al., 2017; Halabi et al., 2020a; Navis et al., 2020b; Zenker et al., 2020; Bavaro et al., 2021; Ye et al., 2022; Chen et al., 2022; Chen et al., 2023; Calvez et al., 2024; Table 2).

Producing IMF with a low-heat WPC ingredient supported faster gastric emptying of IMF compared with IMF produced with extensively heat-treated WPC (Navis et al., 2020b; Table 2). Both preterm and near-term piglets received an IMF formulated with pasteurized WPC for 5 d (Navis et al., 2020b). At 60 min post feeding, these piglets had significantly lower gastric content (approximately 15 g/kg and 30 g/kg) compared with piglets that received an IMF with extensively heated WPC (approximately 35 g/kg and 45 g/kg; $P < 0.01$; Navis et al., 2020b). In another study, the rate of protein hydrolysis differed during in vitro intestinal digestion between unheated IMF and heated IMF (Halabi et al.,

2020a). At 5 min intestinal digestion, unheated IMF had a significantly higher degree of protein hydrolysis versus IMF heated at 67.5°C or 80°C ($24 \pm 2\%$ vs. $14 \pm 2\%$; $P < 0.05$; Halabi et al., 2020a). However, by the end of intestinal digestion, the degree of protein hydrolysis was similar ($35 \pm 2\%$; Halabi et al., 2020a). Unheated and IMF heat-treated at 80°C had comparable bioaccessibility of EAA ($34.1 \pm 1.5\%$ vs. $35.5 \pm 1.2\%$) and NEAA ($10.2 \pm 0.7\%$ vs. $10.3 \pm 0.6\%$) at the end of the intestinal phase (Halabi et al., 2020a; Table 2). Zenker et al. (2020) produced an unheated IMF powder with relatively low furosine levels (308 mg furosine/100 g of protein). This IMF powder had a significantly higher degree of protein hydrolysis following gastric ($5.9 \pm 0.4\%$) and intestinal ($46.8 \pm 3.9\%$) digestion compared with the dry-heated IMF powder with 2,022 mg furosine/100 g of protein ($P < 0.001$; Zenker et al., 2020).

Using a semidynamic infant in vitro gastric digestion model, protein digestion of membrane-filtered IMF was tracked (Chen et al., 2022). Once membrane-filtered IMF was added to the gastric phase, an increase in pH from 2.5 to 7.38 ± 0.02 was observed (Chen et al., 2022). By the end of gastric digestion (156 min, pH 3.2), a significantly higher degree of protein hydrolysis was recorded in the membrane-filtered IMF ($1,338 \pm 253$ µmol free amines/g of protein) versus the high-temperature IMF (845 ± 78 µmol free amines/g of protein; $P < 0.05$; Chen et al., 2022). The membrane-filtered IMF had a fragmented curd throughout in vitro gastric digestion (Chen et al., 2022). From 94 min gastric digestion, a clear phase separation could be observed as fat droplets began to separate from protein aggregates, indicating that membrane-filtered IMF had a less dense curd than high-temperature IMF (Chen et al., 2022). Still, SDS-PAGE analysis showed intact β-LG was still present in membrane-filtered IMF at the end of gastric digestion (Chen et al., 2022). No significant difference was observed in the release of peptides <10 kDa from either membrane-filtered IMF or high-temperature IMF at the end of the gastric phase ($P > 0.05$; Chen et al., 2022). In the follow-up pig trial, 180 min after a final feed of a membrane-filtered IMF-based diet, young pigs had a significantly higher degree of protein hydrolysis in the lumen collected from the stomach compared with stomach contents of pigs fed high-temperature IMF diets (599 ± 45 µmol free amines/g of protein vs. 509 ± 17 µmol free amines/g of protein; $P < 0.05$; Chen et al., 2023). In the stomach of these pigs there was a significantly higher concentrations of water-soluble proteins and peptides compared with high-temperature IMF fed pigs ($P < 0.05$; Chen et al., 2023). The total concentration of free AA released from the stomach of pigs fed diets with

membrane-filtered IMF or high-temperature IMF were comparable (56.5 ± 4.6 $\mu\text{mol/g}$ of protein vs. 59.4 ± 3.05 $\mu\text{mol/g}$ of protein; Chen et al., 2023).

In the duodenum, there was a significant increase in the degree of protein hydrolysis from pigs fed membrane-filtered IMF diets ($1,530 \pm 136$ μmol free amines/g of protein) compared with pigs fed diets with high-temperature IMF ($1,174 \pm 124$ μmol free amines/g of protein; $P < 0.05$; Chen et al., 2023). The total concentration of free AA released was similar in the duodenum of pigs in both IMF treatment groups (Chen et al., 2023). However, membrane-filtered IMF fed pigs had a significantly higher Pro concentration in the duodenum compared with high-temperature IMF fed pigs (7.8 ± 2.61 $\mu\text{mol/g}$ of protein vs. 2.2 ± 0.9 $\mu\text{mol/g}$ of protein; $P < 0.05$; Chen et al., 2023). There was a similar level of protein hydrolysis recorded in the jejunum and ileum of pigs in both IMF treatment groups (Chen et al., 2023). In the jejunum, taurine and Trp concentrations were both significantly higher in pigs fed the membrane-filtered IMF diets compared with pigs fed high-temperature IMF diets (4.7 ± 1.55 and 4.9 ± 0.37 $\mu\text{mol/g}$ of protein, respectively, vs. 1.5 ± 0.37 and 3.5 ± 0.48 $\mu\text{mol/g}$ of protein; $P < 0.05$; Chen et al., 2023). Similar amounts of total free AA were released from the ileum of membrane-filtered IMF fed pigs (251.5 ± 33.1 $\mu\text{mol/g}$ of protein) and pigs fed high-temperature IMF (285 ± 28.51 $\mu\text{mol/g}$ of protein; Chen et al., 2023). Pigs fed membrane-filtered IMF did have lower concentrations of Tyr in the ileum compared with high-temperature IMF fed pigs (8.8 ± 0.98 $\mu\text{mol/g}$ of protein vs. 11.5 ± 0.86 $\mu\text{mol/g}$ of protein; $P < 0.05$; Chen et al., 2023). Overall, membrane-filtered IMF fed pigs had a higher total degree of protein hydrolysis in the gastrointestinal tract compared with high-temperature IMF fed pigs ($1,173 \pm 46$ μmol free amines/g of protein vs. $1,022 \pm 43$ μmol free amines/g of protein; $P < 0.05$) and higher concentrations of water-soluble proteins and peptides ($P < 0.001$; Chen et al., 2023). However, some discrepancies between in vitro and preclinical experiments did exist, with an in vitro static infant gastrointestinal digestion reporting similar levels of proteolysis at the end of intestinal digestion for membrane-filtered and high-temperature IMF (471 ± 36.5 μmol free amines/mg of protein vs. 481 ± 7.7 μmol free amines/mg of protein; Bavaro et al., 2021). In vitro, at the end of the intestinal phase, Lys, Leu, and Tyr were the most abundant AA released from membrane-filtered IMF compared with Lys, Leu, and Phe from high-temperature IMF (Bavaro et al., 2021).

From a peptide-profiling perspective, there is evidence that processing will affect individual peptides released during digestion. Although processing pa-

rameters were not disclosed, following in vitro gastrointestinal digestion, Wada and Lönnerdal (2015a) identified 39 bioactive peptides in intestinal digesta from a standard IMF powder compared with only 9 in the intestinal digesta from an extensively hydrolyzed IMF powder (Wada and Lönnerdal, 2015a). Bavaro et al. (2021) treated Caco-2 cell monolayers with IMF digesta for 4 h. Analysis by LC-MS/MS identified 488 dairy peptides in the apical chambers common to both membrane-filtered and high-temperature IMF, with 112 peptides unique to membrane-filtered IMF and 97 peptides unique to high-temperature IMF (Bavaro et al., 2021). Bavaro et al. (2021) did not determine if this resulted in differences in peptide bioactive capacity. Interestingly, the bioactive β -casomorphin-7 opioid peptide (Claustre et al., 2002; Trompette et al., 2003; Zoghbi et al., 2006), which is released from β -CN following gastrointestinal digestion, was identified in both IMF (Bavaro et al., 2021).

Calvez et al. (2024) calculated a DIAAS of 82 for a membrane-filtered IMF powder, which was similar to 80 for an extensively, heat-treated IMF powder. In a preclinical study by Calvez et al. (2024), Wistar Han rats ($n = 30$, 28-d old) were fed a diet containing this membrane-filtered IMF at 40% inclusion for 14 d. There was no significant effect of IMF processing on dietary nitrogen levels in the stomach ($P = 0.5$) or the small intestine ($P = 0.17$) 6 h after final meal ingestion (Calvez et al., 2024). However, there were differences observed in AA bioavailability, with Calvez et al. (2024) observing significant increases in postprandial indispensable and total AA concentrations in rats fed membrane-filtered IMF within 1 h, compared with rats fed thermally processed IMF ($P < 0.001$). No significant effect of processing on AA bioavailability was observed at any other time point ($P > 0.05$; Calvez et al., 2024).

However, lowering thermal processing may not always speed up digestion. An in vitro infant digestion of a pasteurized liquid IMF (72°C for 10 s) resulted in visible β -LG and α -LA protein bands on SDS-PAGE at the end of the gastric phase, while β -LG was the only whey protein still present at the end of intestinal digestion (Ye et al., 2022). In contrast, all proteins in the intestinal digesta of UHT-treated liquid IMF were completely degraded (Ye et al., 2022). A similar number of peptides were released from the intestinal digesta of pasteurized liquid IMF and UHT IMF (Ye et al., 2022; Table 2). In another study, Cattaneo et al. (2017) reported that the degree of proteolysis was significantly lower in an unheated IMF ($35.6 \pm 0.6\%$) compared with an IMF produced by in-batch sterilization at 110°C for 38 min ($40.3 \pm 1.2\%$; $P < 0.05$) albeit this was reversed when ingredients (sodium

caseinate and WPC) were processed in-house rather than purchased commercially (sodium caseinate and WPI; Cattaneo et al., 2017).

EFFECTS OF LOWER HEAT TREATMENT DURING IMF PRODUCTION ON GUT BARRIER

Several studies have focused on whether or not reduced thermal processing of IMF will benefit gut physiology and gut barrier health (Li et al., 2013; Navis et al., 2020a,b; Bavaro et al., 2021; Sun et al., 2022; Dold et al., 2024; Table 3).

In a study by Sun et al. (2022), preterm pigs received pasteurized IMF for 5 d by orogastric feeding tube. These pigs had significantly higher villus height and lower crypt depth in the proximal, middle, and distal small intestine compared with pigs receiving 60 d stored UHT liquid IMF ($P < 0.05$; Sun et al., 2022). Intestinal permeability was unaffected by IMF processing (Sun et al., 2022).

In another study by Li et al. (2013), preterm pigs receiving an IMF formulated with membrane-filtered WPC for 5 d by orogastric feeding tube had significantly increased villus height in the proximal small intestine compared with pigs receiving IMF produced with double-pasteurized WPC ($667 \pm 46 \mu\text{m}$ vs. $440 \pm 47 \mu\text{m}$; $P < 0.05$; Li et al., 2013). Interestingly, Western blot analysis revealed that claudin-4 protein levels were significantly lower in pigs receiving IMF with membrane-filtered WPC compared with the pigs that received IMF formulated with double-pasteurized WPC ($P < 0.05$; Li et al., 2013). The urinary lactulose:mannitol ratio was significantly lower in pigs that received the IMF with membrane-filtered WPC (>0.1) compared with pigs receiving the IMF produced with double-pasteurized WPC (>0.5 ; $P < 0.05$; Li et al., 2013). Inflammatory biomarkers IL-8, tumor necrosis factor α ($0.6 \pm 0.7 \text{ pg/mg}$ of tissue), and IL-1 β ($3.7 \pm 0.5 \text{ pg/mg}$ of tissue) in the distal small intestine of preterm pigs were unaffected by processing type (Li et al., 2013).

Preterm and near-term piglets received an IMF produced with pasteurized WPC by orogastric feeding (Navis et al., 2020a,b). Crypt depth was significantly lower in the colon of preterm ($P < 0.05$) and near-term ($P < 0.01$) piglets receiving IMF with pasteurized WPC compared with piglets receiving an IMF with extensively heated WPC (Navis et al., 2020a). Using periodic acid-Schiff staining, Navis et al. (2020a) reported an observational increase in colonic goblet cell number in piglets receiving the IMF made with pasteurized WPC compared with those piglets that received an IMF with extensively heated WPC, although no difference was observed in colonic mRNA transcript levels of *mucin-1* and *mucin-2*. *Mucin-2* mRNA transcripts were also unaffected by processing type in the distal small intestine (Navis et al.,

2020a). Intestinal permeability was influenced by processing type, with near-term piglets receiving IMF with pasteurized WPC having a significantly lower urinary lactulose:mannitol ratio compared with piglets receiving IMF with extensively heated WPC (>0.07 vs. > 0.2 ; $P < 0.05$; Navis et al., 2020b). Interleukin-8 protein levels and mRNA transcripts were significantly reduced in the colon of preterm piglets that received IMF with pasteurized WPC compared with piglets that received IMF with extensively heated WPC ($P < 0.05$; Navis et al., 2020a). Near-term piglets receiving IMF with pasteurized WPC also had significantly lower mRNA transcript levels of *IL-8* ($P < 0.05$; Navis et al., 2020a). Interleukin-1 β mRNA transcript levels were significantly lower in the distal small intestine of preterm piglets and colon of near-term piglets receiving IMF with pasteurized WPC ($P < 0.05$; Navis et al., 2020a). Protein concentrations and mRNA transcript levels of tumor necrosis factor α were significantly lower in near-term piglets that received IMF with pasteurized WPC compared with piglets that received IMF with extensively heated WPC ($P < 0.05$; Navis et al., 2020a).

In preterm piglets fed for 5 d by orogastric feeding tube an IMF produced with a pasteurized WPC, there was no difference in villus height or crypt depth compared with piglets fed an IMF produced with a stored double-pasteurized WPC (Aasmul-Olsen et al., 2024).

For our preclinical study with 28-d-old pigs, there were also no significant differences observed in villus height, crypt depth, or mucosa thickness in the upper small intestine (duodenum and jejunum) of pigs fed membrane-filtered or high-temperature IMF ($P > 0.05$; Dold et al., 2024). However, there was a significant increase in goblet cell number in the jejunum of pigs fed membrane-filtered IMF diets compared with high-temperature IMF diets (17.39 ± 1.43 vs. 11.7 ± 1.27 ; $P < 0.05$; Dold et al., 2024). In addition, acidic mucus levels and mucin-2 protein concentrations in the jejunum were significantly increased in membrane-filtered IMF fed pigs compared with pigs fed high-temperature IMF diets ($P < 0.05$; Dold et al., 2024). A significant increase in mRNA transcript levels of *claudin-1* ($P < 0.001$), *mucin-1* ($P < 0.01$), and *mucin-2* ($P < 0.001$) was measured in the duodenal mucosal scrapings of pigs fed membrane-filtered IMF diets versus pigs fed high-temperature IMF diets (Dold et al., 2024). In parallel, in in vitro experiments where Caco-2 polarized cell monolayers were treated with intestinal digesta from membrane-filtered IMF, TEER values and claudin-1 protein concentrations were significantly increased compared with high-temperature IMF treatment ($P < 0.05$; Bavaro et al., 2021). Other biomarkers were unaffected by treatment type, in the pig study mRNA transcripts of *occludin*, *claudin-2*, *claudin-4*, *zonula occludens-1*, and

Table 4. Sensitivity of gut biomarkers to high-temperature processing

Biomarker (no. of studies)	Sensitive	Occasionally sensitive	Insensitive or resistant	Reference
Villus height (n = 4)		↓ in villus height in 2 of 4 studies		Li et al., 2013; Sun et al., 2022; Aasmul-Olsen et al., 2024; Dold et al., 2024
Crypt depth (n = 4)		↑ in crypt depth in 2 of 4 studies		Navis et al., 2020b; Sun et al., 2022; Aasmul-Olsen et al., 2024; Dold et al., 2024
Tight junctions (n = 3)		↓ in tight junction mRNA transcripts and protein levels in 2 of 3 studies		Li et al., 2013; Bavaro et al., 2021; Dold et al., 2024
Goblet cell number (n = 2)	↓ in goblet cell number in 2 of 2 studies			Navis et al., 2020a; Dold et al., 2024
Mucins (n = 2)		↓ in mucin mRNA transcripts and protein levels in 1 of 2 studies		Navis et al., 2020a; Dold et al., 2024
Intestinal permeability (n = 3)		↑ in intestinal permeability (lactulose:mannitol ratio) in 2 of 3 studies		Li et al., 2013; Navis et al., 2020b; Sun et al., 2022
Aminopeptidase activity (n = 5)			↑ in aminopeptidase N activity in 1 of 5 studies (no change in aminopeptidase A activity in all 5 studies)	Li et al., 2013; Navis et al., 2020b; Sun et al., 2022; Aasmul-Olsen et al., 2024; Dold et al., 2024
Intestinal alkaline phosphatase activity (n = 2)		↓ in intestinal alkaline phosphatase activity in 1 of 2 studies		Navis et al., 2020a; Dold et al., 2024
Lactase activity (n = 5)		↓ in lactase activity in 2 of 5 studies		Li et al., 2013; Navis et al., 2020b; Sun et al., 2022; Aasmul-Olsen et al., 2024; Dold et al., 2024

junctional adhesion molecule-A were similar (Dold et al., 2024) and in vitro occludin, claudin-4, zonula occludens-1, and junctional adhesion molecule-A protein levels were unaffected by processing (Bavaro et al., 2021). Interestingly, in the pig study, a significant reduction in mRNA transcript levels for the receptor for AGE was also detected in the jejunum of membrane-filtered IMF fed pigs compared with high-temperature IMF fed pigs ($P < 0.05$; Dold et al., 2024), even though both IMF had similar CML concentrations (Chen et al., 2021).

From a digestive capacity perspective in the gut, Sun et al. (2022) reported a significant increase in brush border lactase activity in the proximal and middle small intestine of pigs fed the pasteurized liquid IMF compared with stored UHT liquid IMF ($P < 0.05$; Sun et al., 2022). No differences were reported in activities of the other brush border enzymes analyzed (aminopeptidase N, aminopeptidase A, sucrase, maltase, and dipeptidyl peptidase IV; Sun et al., 2022). In a study by Li et al. (2013) processing of WPC used in the formulation of IMF had no significant effect on brush border aminopeptidase N, aminopeptidase A, lactase, sucrase, maltase, and dipeptidyl peptidase IV activities ($P > 0.05$). For a study by Navis et al. (2020a) brush border intestinal alkaline phosphatase activity was significantly higher in the colon of preterm piglets receiving IMF with pasteurized WPC compared with piglets receiving the IMF with extensively heated WPC ($P < 0.05$), with similar levels recorded in the colon of near-term piglets (Navis et al., 2020a). For preterm and near-term piglets, similar levels of intestinal alkaline phosphatase activity were recorded in the distal small intestine, regardless of IMF diet (Navis et al., 2020a). Aminopeptidase N, aminopeptidase A, and lactase in the middle small intestine were also unaffected by IMF processing (Navis et al., 2020b). In the trial of Aasmul-Olsen et al. (2024) brush border aminopeptidase N, aminopeptidase A, lactase, sucrase, maltase, and dipeptidyl peptidase IV activities were similar between treatment types. Consistent with the findings of Sun et al. (2022), we have also recently measured a significant increase in lactase activity in isolated duodenal brush border membrane vesicles in pigs fed membrane-filtered IMF diets compared with pigs fed high-temperature IMF diets (166.59 ± 22.09 nmol glucose/mg of protein vs. 110.75 ± 7.21 nmol glucose/mg of protein; $P < 0.05$; Dold et al., 2024). In addition, duodenal trypsin activities were significantly higher in membrane-filtered IMF fed pigs compared with pigs fed high-temperature IMF diets (1.36 ± 0.19 mU/mg of protein vs. 0.91 ± 0.2 mU/mg of protein; $P < 0.05$; Dold et al., 2024). However, in the duodenum, brush border aminopeptidase N activity was significantly lower in pigs fed membrane-filtered IMF compared with pigs on high-temperature IMF diets (5.23 ± 0.26 U/mg of protein vs. 7.44 ± 0.69 U/mg of protein;

$P < 0.05$; Dold et al., 2024). Processing type had no significant effect on duodenal brush border aminopeptidase A or intestinal alkaline phosphatase activities and no effect on the digestive enzyme capacity of the jejunum ($P > 0.05$; Dold et al., 2024).

By comparing gut responses to IMF produced with different thermal loads, it is apparent that several gut barrier biomarkers are sensitive to IMF thermal processing (Table 4). These include villus height, crypt depth, tight junction proteins, goblet cell number, mucus levels, brush border enzyme activities, and intestinal permeability. Further work is needed to determine whether the main driver of gut responses to processing is IMF protein structure or IMF AGE levels or a contribution from both. Mechanistic studies are required to investigate the response of gut cells to individual AGE compounds and individual bioactive peptides.

CONCLUSIONS

In general, in vitro and preclinical studies have revealed that thermal processing of IMF reduces protein digestibility and alters gut physiology in the young gut. A major limitation to date is that all studies have been performed in vitro or preclinical. How this data correlates to the human infant requires follow-up intervention studies in babies, which is challenging. Research progress is also hampered by the failure of IMF manufacturers to disclose processing temperatures. A full profile of AGE, quantified by standardized methods, should be given with each IMF product to allow accurate comparisons. Processing improvements, at pilot-plant scale, are also required to develop alternative technologies for IMF processing (i.e., UV irradiation and membrane filtration), whilst ensuring the production of a safe product. Ultimately, the impact of any IMF product on protein digestion and gut barrier physiology in the infant gut must be compared with breast milk.

NOTES

This work was funded by Science Foundation Ireland (SFI; Dublin, Ireland) and the Department of Agriculture, Food, and the Marine, Agriculture House (Dublin, Ireland), on behalf of the Government of Ireland under Grant No. 16/RC/3835 VistaMilk. Cathal A. Dold is in receipt of a Teagasc Walsh Scholarship (No. 2019220) from the Teagasc Food Research Centre (Moorepark, Fermoy, Co. Cork, Ireland). Author contributions are as follows: Cathal A. Dold, conceptualization, writing (original draft, review and editing); Aylin W. Sahin, supervision, writing (review and editing); Linda Giblin, conceptualization, project administration, funding acquisition, supervision, writing (review and editing).

No human or animal subjects were used, so this analysis did not require approval by an Institutional Animal Care and Use Committee or Institutional Review Board. The authors have not stated any conflicts of interest.

Nonstandard abbreviations used: AGE = advanced glycation end product; CEL = N(6)-carboxyethyllysine; CML = N(6)-carboxymethyllysine; DIAAS = digestible indispensable amino acid score; IMF = infant milk formula; LC = liquid chromatography; RAGE = receptor for advanced glycation end products; RT-qPCR = reverse transcription quantitative real-time PCR; SI = small intestine; SMP = skim milk powder; TEER = transepithelial electrical resistance; WPC = whey protein concentrate; WPI = whey protein isolate.

REFERENCES

- Aasmul-Olsen, K., H. G. Akıllıoğlu, L. I. Christiansen, K. Engholm-Keller, A. Brunse, D. V. Stefanova, A. Bjørnshave, M. R. Bechshøft, K. Skovgaard, T. Thymann, P. T. Sangild, M. N. Lund, and S. B. Bering. 2024. A gently processed skim milk-derived whey protein concentrate for infant formula: Effects on gut development and immunity in preterm pigs. *Mol. Nutr. Food Res.* 68:2300458. <https://doi.org/10.1002/mnfr.202300458>.
- Abrahamse, E., G. G. M. Thomassen, I. B. Renes, P. A. Wierenga, and K. A. Hettinga. 2022. Gastrointestinal protein hydrolysis kinetics: opportunities for further infant formula improvement. *Nutrients* 14:1512. <https://doi.org/10.3390/nu14071512>.
- Almeida, C. C., B. F. Mendonça Pereira, K. C. Leandro, M. P. Costa, B. F. Spisso, and C. A. Conte-Junior. 2021. Bioactive compounds in infant formula and their effects on infant nutrition and health: A systematic literature review. *Int. J. Food Sci.* 2021:1–31. <https://doi.org/10.1155/2021/8850080>.
- Anema, S. G., and Y. Li. 2003. Association of denatured whey proteins with casein micelles in heated reconstituted skim milk and its effect on casein micelle size. *J. Dairy Res.* 70:73–83. <https://doi.org/10.1017/S0022029902005903>.
- Augustine, D. A., and G.-A. Bent. 2022. Acrylamide, a toxic maillard by-product and its inhibition by sulfur-containing compounds: A mini review. *Front. Food Sci. Technol.* 2:1072675. <https://doi.org/10.3389/ffst.2022.1072675>.
- Bakshi, S., V. K. Paswan, S. P. Yadav, B. K. Bhinchhar, S. Kharkwal, H. Rose, P. Kanetkar, V. Kumar, Z. A. S. Al-Zamani, and D. S. Bunkar. 2023. A comprehensive review on infant formula: nutritional and functional constituents, recent trends in processing and its impact on infants' gut microbiota. *Front. Nutr.* 10:1194679. <https://doi.org/10.3389/fnut.2023.1194679>.
- Ballard, O., and A. L. Morrow. 2013. Human milk composition: Nutrients and bioactive factors. *Pediatr. Clin. North Am.* 60:49–74. <https://doi.org/10.1016/j.pcl.2012.10.002>.
- Basdeki, A.-M., D. G. Fatouros, C. G. Biliaderis, and T. Moschakis. 2021. Physicochemical properties of human breast milk during the second year of lactation. *Curr. Res. Food Sci.* 4:565–576. <https://doi.org/10.1016/j.crfs.2021.08.001>.
- Bavaro, S. L., G. Mamone, G. Picariello, M. J. Callanan, Y. Chen, A. Brodkorb, and L. Giblin. 2021. Thermal or membrane processing for infant milk formula: Effects on protein digestion and integrity of the intestinal barrier. *Food Chem.* 347:129019. <https://doi.org/10.1016/j.foodchem.2021.129019>.
- Bourlieu, C., O. Ménard, K. Bouzerzour, G. Mandalari, A. Macierzanka, A. R. Mackie, and D. Dupont. 2014. Specificity of infant digestive conditions: Some clues for developing relevant in vitro models. *Crit. Rev. Food Sci. Nutr.* 54:1427–1457. <https://doi.org/10.1080/10408398.2011.640757>.
- Buggy, A. K., J. J. McManus, A. Brodkorb, N. M. Carthy, and M. A. Fenelon. 2017. Stabilising effect of α -lactalbumin on concentrated infant milk formula emulsions heat treated pre- or post-homogenisation. *Dairy Sci. Technol.* 96:845–859. <https://doi.org/10.1007/s13594-016-0306-1>.
- Calvez, J., A. Blais, A. Deglaire, C. Gaudichon, F. Blachier, and A. M. Davila. 2024. Minimal processed infant formula vs. conventional shows comparable protein quality and increased postprandial plasma amino acid kinetics in rats. *Br. J. Nutr.* 131:1115–1124. <https://doi.org/10.1017/S0007114523002696>.
- Catassi, C., A. Bonucci, G. V. Coppa, A. Carlucci, and P. L. Giorgi. 1995. Intestinal permeability changes during the first month: Effect of natural versus artificial feeding. *J. Pediatr. Gastroenterol. Nutr.* 21:383–386. <https://doi.org/10.1002/j.1536-4801.1995.tb11955.x>.
- Cattaneo, S., M. Stuknytė, F. Masotti, and I. De Noni. 2017. Protein breakdown and release of β -casomorphins during in vitro gastrointestinal digestion of sterilised model systems of liquid infant formula. *Food Chem.* 217:476–482. <https://doi.org/10.1016/j.foodchem.2016.08.128>.
- Charton, E., G. Henry, A. Cahu, Y. Le Gouar, P. Dahirel, P. J. Moughan, C. A. Montoya, A. Bellanger, D. Dupont, I. Le Huërou-Luron, and A. Deglaire. 2023. Ileal digestibility of nitrogen and amino acids in human milk and an infant formula as determined in neonatal minipiglets. *J. Nutr.* 153:1063–1074. <https://doi.org/10.1016/j.tjnut.2023.02.025>.
- Chávez-Servín, J. L., K. de la Torre Carbot, T. García-Gasca, A. I. Castellote, and M. C. López-Sabater. 2015. Content and evolution of potential furfural compounds in commercial milk-based infant formula powder after opening the packet. *Food Chem.* 166:486–491. <https://doi.org/10.1016/j.foodchem.2014.06.050>.
- Chelakkot, C., J. Ghim, and S. H. Ryu. 2018. Mechanisms regulating intestinal barrier integrity and its pathological implications. *Exp. Mol. Med.* 50:1–9. <https://doi.org/10.1038/s12276-018-0126-x>.
- Chen, Y., M. Callanan, L. Giblin, J. Tobin, and A. Brodkorb. 2022. Comparison of conventional heat-treated and membrane filtered infant formulas using an in vitro semi-dynamic digestion method. *Food Funct.* 13:8158–8167. <https://doi.org/10.1039/D2FO00342B>.
- Chen, Y., M. Callanan, C. Shanahan, J. Tobin, L. F. Gamon, M. J. Davies, L. Giblin, and A. Brodkorb. 2021. The use of membrane filtration to increase native whey proteins in infant formula. *Dairy* 2:515–529. <https://doi.org/10.3390/dairy2040041>.
- Chen, Z., A. Kondrashina, I. Greco, L. F. Gamon, M. N. Lund, L. Giblin, and M. J. Davies. 2019. Effects of protein-derived amino acid modification products present in infant formula on metabolic function, oxidative stress, and intestinal permeability in cell models. *J. Agric. Food Chem.* 67:5634–5646. <https://doi.org/10.1021/acs.jafc.9b01324>.
- Chen, Y., H. Rooney, C. Dold, S. Bavaro, J. Tobin, M. J. Callanan, A. Brodkorb, P. G. Lawlor, and L. Giblin. 2023. Membrane filtration processing of infant milk formula alters protein digestion in young pigs. *Food Res. Int.* 166:112577. <https://doi.org/10.1016/j.foodres.2023.112577>.
- Claustre, J., F. Toumi, A. Trompette, G. Jourdan, H. Guignard, J. A. Chayvialle, and P. Plaisancié. 2002. Effects of peptides derived from dietary proteins on mucus secretion in rat jejunum. *Am. J. Physiol. Gastrointest. Liver Physiol.* 283:G521–G528. <https://doi.org/10.1152/ajpgi.00535.2001>.
- Commission Regulation. 2005. Commission Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs. Accessed Mar. 26, 2024. <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC061603/>.
- Commission Regulation. 2017. Commission Regulation (EU) 2017/2158 establishing mitigation measures and benchmark levels for the reduction of the presence of acrylamide in food. Accessed May 17, 2024. <https://www.fao.org/faolex/results/details/en/c/LEX-FAOC171340/>.
- Commission Regulation. 2013. Commission Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive

- 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009. Accessed Jun. 24, 2024. <https://eur-lex.europa.eu/eli/reg/2013/609/oj>.
- Crowley, S. V., A. P. Dowling, V. Caldeo, A. L. Kelly, and J. A. O'Mahony. 2016. Impact of α -lactalbumin: β -lactoglobulin ratio on the heat stability of model infant milk formula protein systems. *Food Chem.* 194:184–190. <https://doi.org/10.1016/j.foodchem.2015.07.077>.
- Das, D., A. Trikha, M. Kaur, T. Muthiah, R. K. Pandey, B. Ankalji, K. S. Madhusudhan, D. N. Srivastava, and P. M. Singh. 2024. Comparison of gastric emptying time between breast-fed and formula milk-fed infants less than 6 months old, using gastric ultrasound—An observational study. *Pediatr. Anaesth.* 34:430–437. <https://doi.org/10.1111/pan.14841>.
- Dash, K. K., U. Fayaz, A. H. Dar, R. Shams, S. Manzoor, A. Sundarsingh, P. Deka, and S. A. Khan. 2022. A comprehensive review on heat treatments and related impact on the quality and microbial safety of milk and milk-based products. *Food Chem. Adv.* 1:100041. <https://doi.org/10.1016/j.focha.2022.100041>.
- Demers-Mathieu, V. 2022. The immature intestinal epithelial cells in preterm infants play a role in the necrotizing enterocolitis pathogenesis: A review. *Health Sci. Rev. (Oxf.)* 4:100033. <https://doi.org/10.1016/j.hsr.2022.100033>.
- Dold, C. A., S. L. Bavaro, Y. Chen, M. J. Callanan, D. Kennedy, J. Cassidy, J. Tobin, A. W. Sahin, P. G. Lawlor, A. Brodtkorb, and L. Giblin. 2024. Infant milk formula produced by membrane filtration promotes mucus production in the upper small intestine of young pigs. *Food Res. Int.* 187:114343. <https://doi.org/10.1016/j.foodres.2024.114343>.
- EFSA Panel on Dietetic Products and Allergies. 2014. Scientific opinion on the essential composition of infant and follow-on formulae. *EFSA J.* 12:3760. <https://doi.org/10.2903/j.efsa.2014.3760>.
- Fenaille, F., V. Parisod, P. Visani, S. Populaire, J.-C. Tabet, and P. A. Guy. 2006. Modifications of milk constituents during processing: A preliminary benchmarking study. *Int. Dairy J.* 16:728–739. <https://doi.org/10.1016/j.idairyj.2005.08.003>.
- Fujimori, S. 2020. Gastric acid level of humans must decrease in the future. *World J. Gastroenterol.* 26:6706–6709. <https://doi.org/10.3748/wjg.v26.i43.6706>.
- Gan, J., G. Bornhorst, B. Henrick, and B. German. 2018. Protein digestion of baby foods: Study approaches and implications for infant health. *Mol. Nutr. Food Res.* 62:1700231. <https://doi.org/10.1002/mnfr.201700231>.
- Gleeson, J., K. Fein, N. Chaudhary, R. Doerfler, A. Newby, and K. Whitehead. 2021. The enhanced intestinal permeability of infant mice enables oral protein and macromolecular absorption without delivery technology. *Int. J. Pharm.* 593:120120. <https://doi.org/10.1016/j.ijpharm.2020.120120>.
- Gómez-Gallego, C., I. Recio, V. Gómez-Gómez, I. Ortuño, M. J. Bernal, G. Ros, and M. J. Periago. 2016. Effect of processing on polyamine content and bioactive peptides released after in vitro gastrointestinal digestion of infant formulas. *J. Dairy Sci.* 99:924–932. <https://doi.org/10.3168/jds.2015-10030>.
- Guibourdenche, M., J. Haug, N. Chevalier, M. Spatz, N. Barbezier, J. Gay-Quéheillard, and P. M. Anton. 2021. Food contaminants effects on an in vitro model of human intestinal epithelium. *Toxics* 9:135. <https://doi.org/10.3390/toxics9060135>.
- Guo, M., and G. Hendricks. 2008. Chemistry and biological properties of human milk. *Curr. Nutr. Food Sci.* 4:305–320. <https://doi.org/10.2174/157340108786263667>.
- Halabi, A., T. Croguennec, S. Bouhallab, D. Dupont, and A. Deglaire. 2020a. Modification of protein structures by altering the whey protein profile and heat treatment affects in vitro static digestion of model infant milk formulas. *Food Funct.* 11:6933–6945. <https://doi.org/10.1039/D0FO01362E>.
- Halabi, A., A. Deglaire, P. Hamon, S. Bouhallab, D. Dupont, and T. Croguennec. 2020b. Kinetics of heat-induced denaturation of proteins in model infant milk formulas as a function of whey protein composition. *Food Chem.* 302:125296. <https://doi.org/10.1016/j.foodchem.2019.125296>.
- He, T., W. Rombouts, A. W. C. Einerhand, N. Hotrum, and F. van de Velde. 2022. Gastric protein digestion of goat and cow milk infant formula and human milk under simulated infant conditions. *Int. J. Food Sci. Nutr.* 73:28–38. <https://doi.org/10.1080/09637486.2021.1921705>.
- Henderson, D., C. A. Murphy, A.-C. Glynn, M. A. Boyle, and N. McCallion. 2022. Feeding practices and the prevalence of cow's milk protein allergy in Irish preterm infants. *J. Hum. Nutr. Diet.* 35:535–541. <https://doi.org/10.1111/jhn.12971>.
- Hill, C. J., D. B. Lynch, K. Murphy, M. Ulaszewska, I. B. Jeffery, C. A. O'Shea, C. Watkins, E. Dempsey, F. Mattivi, K. Tuohy, R. P. Ross, C. A. Ryan, P. W. O'Toole, and C. Stanton. 2017. Evolution of gut microbiota composition from birth to 24 weeks in the INFANTMET Cohort. *Microbiome* 5:4. <https://doi.org/10.1186/s40168-016-0213-y>.
- Horvat, S., and A. Jakas. 2004. Peptide and amino acid glycation: New insights into the Maillard reaction. *J. Pept. Sci.* 10:119–137. <https://doi.org/10.1002/psc.519>.
- Jiang, H., S. Gallier, L. Feng, J. Han, and W. Liu. 2022. Development of the digestive system in early infancy and nutritional management of digestive problems in breastfed and formula-fed infants. *Food Funct.* 13:1062–1077. <https://doi.org/10.1039/D1FO03223B>.
- Johansson, M. E. V., H. Sjövall, and G. C. Hansson. 2013. The gastrointestinal mucus system in health and disease. *Nat. Rev. Gastroenterol. Hepatol.* 10:352–361. <https://doi.org/10.1038/nrgastro.2013.35>.
- Johnston, N., P. W. Dettmar, B. Bishwokarma, M. O. Lively, and J. A. Koufman. 2007. Activity/stability of human pepsin: Implications for Reflux attributed laryngeal disease. *Laryngoscope* 117:1036–1039. <https://doi.org/10.1097/MLG.0b013e31804154c3>.
- Jouban, Y., A. Moscovici, and U. Lesmes. 2015. Antioxidant activity of bovine alpha lactalbumin Maillard products and evaluation of their in vitro gastro-duodenal digestive proteolysis. *Food Funct.* 6:1229–1240. <https://doi.org/10.1039/C4FO01165A>.
- Jouban, Y., A. Moscovici, R. Portmann, and U. Lesmes. 2017. Implications of the Maillard reaction on bovine alpha-lactalbumin and its proteolysis during in vitro infant digestion. *Food Funct.* 8:2295–2308. <https://doi.org/10.1039/C7FO00588A>.
- Joyce, A. M., A. Brodtkorb, A. L. Kelly, and J. A. O'Mahony. 2017. Separation of the effects of denaturation and aggregation on whey-casein protein interactions during the manufacture of a model infant formula. *Dairy Sci. Technol.* 96:787–806. <https://doi.org/10.1007/s13594-016-0303-4>.
- Kim, S. Y., and D. Y. Yi. 2020. Components of human breast milk: From macronutrient to microbiome and microRNA. *Clin. Exp. Pediatr.* 63:301–309. <https://doi.org/10.3345/cep.2020.00059>.
- Kong, S., Y. H. Zhang, and W. Zhang. 2018. Regulation of Intestinal Epithelial Cells Properties and Functions by Amino Acids. *BioMed Res. Int.* 2018:2819154. <https://doi.org/10.1155/2018/2819154>.
- Kosek, M. N., G. O. Lee, R. L. Guerrant, R. Haque, G. Kang, T. Ahmed, P. Bessong, A. Ali, E. Mduma, P. Peñataro Yori, W. A. Faubion, A. A. M. Lima, M. Paredes Olortegui, C. Mason, S. Babji, R. Singh, S. Qureshi, P. S. Kosek, A. Samie, J. Pascal, S. Shrestha, B. J. J. McCormick, J. C. Seidman, D. R. Lang, A. Zaidi, L. E. Caulfield, and M. Gottlieb. 2017. Age and sex normalization of intestinal permeability measures for the improved assessment of enteropathy in infancy and early childhood: Results from the MAL-ED study. *J. Pediatr. Gastroenterol. Nutr.* 65:31–39. <https://doi.org/10.1097/MPG.0000000000001610>.
- Kumar, N., M. Raghavendra, J. Tokas, and H. R. Singal. 2017. Chapter 10: Flavor addition in dairy products: Health benefits and risks. Pages 123–135 in *Nutrients in Dairy and Their Implications on Health and Disease*. R. R. Watson, R. J. Collier, and V. R. Preedy, ed. Academic Press.
- Labbers, T. T., J. Wissing, and J. Roggekamp. 2023. In vitro gastric digestion of an experimental infant formula containing both intact and hydrolyzed milk proteins. *J. Dairy Sci.* 106:4524–4532. <https://doi.org/10.3168/jds.2022-22667>.
- Leclère, J., I. Birlouez-Aragon, and M. Meli. 2002. Fortification of milk with iron-ascorbate promotes lysine glycation and tryptophan oxidation. *Food Chem.* 76:491–499. [https://doi.org/10.1016/S0308-8146\(01\)00369-7](https://doi.org/10.1016/S0308-8146(01)00369-7).

- Lee, H.-M., S.-Y. Yang, J. Han, Y. K. Kim, Y. J. Kim, M. S. Rhee, and K.-W. Lee. 2019. Optimization of spray drying parameters and food additives to reduce glycation using response surface methodology in powdered infant formulas. *Food Sci. Biotechnol.* 28:769–777. <https://doi.org/10.1007/s10068-018-0524-9>.
- Li, S., A. Ye, and H. Singh. 2021a. Impacts of heat-induced changes on milk protein digestibility: A review. *Int. Dairy J.* 123:105160. <https://doi.org/10.1016/j.idairyj.2021.105160>.
- Li, Y., M. V. Østergaard, P. Jiang, D. E. W. Chatterton, T. Thymann, A. S. Kvistgaard, and P. T. Sangild. 2013. Whey protein processing influences formula-induced gut maturation in preterm pigs. *J. Nutr.* 143:1934–1942. <https://doi.org/10.3945/jn.113.182931>.
- Li, Y., W. Quan, X. Jia, Z. He, Z. Wang, M. Zeng, and J. Chen. 2021b. Profiles of initial, intermediate, and advanced stages of harmful Maillard reaction products in whole-milk powders pre-treated with different heat loads during 18 months of storage. *Food Chem.* 351:129361. <https://doi.org/10.1016/j.foodchem.2021.129361>.
- Lin, Y., A. L. Kelly, J. A. O'Mahony, and T. P. Guinee. 2018. Effect of heat treatment, evaporation and spray drying during skim milk powder manufacture on the compositional and processing characteristics of reconstituted skim milk and concentrate. *Int. Dairy J.* 78:53–64. <https://doi.org/10.1016/j.idairyj.2017.10.007>.
- Liu, Q., X. Lu, B. G. Swanson, B. A. Rasco, and D. H. Kang. 2012. Monitoring ultraviolet (UV) radiation inactivation of *Cronobacter sakazakii* in dry infant formula using Fourier transform infrared spectroscopy. *J. Food Sci.* 77:M86–M93. <https://doi.org/10.1111/j.1750-3841.2011.02503.x>.
- Lönnerdal, B. 2003. Nutritional and physiologic significance of human milk proteins. *Am. J. Clin. Nutr.* 77:1537S–1543S. <https://doi.org/10.1093/ajcn/77.6.1537S>.
- Lugli, G. A., L. Mancabelli, C. Milani, F. Fontana, C. Tarracchini, G. Alessandri, D. van Sinderen, F. Turroni, and M. Ventura. 2023. Comprehensive insights from composition to functional microbe-based biodiversity of the infant human gut microbiota. *NPJ Biofilms Microbiomes* 9:25. <https://doi.org/10.1038/s41522-023-00392-6>.
- Lund, P., M. R. Bechshøft, C. A. Ray, and M. N. Lund. 2022. Effect of processing of whey protein ingredient on Maillard reactions and protein structural changes in powdered infant formula. *J. Agric. Food Chem.* 70:319–332. <https://doi.org/10.1021/acs.jafc.1c05612>.
- Lyons, K. E., C. A. Ryan, E. M. Dempsey, R. P. Ross, and C. Stanton. 2020. Breast milk, a source of beneficial microbes and associated benefits for infant health. *Nutrients* 12:1039. <https://doi.org/10.3390/nu12041039>.
- Macierzanka, A., A. R. Mackie, B. H. Bajka, N. M. Rigby, F. Nau, and D. Dupont. 2014. Transport of particles in intestinal mucus under simulated infant and adult physiological conditions: Impact of mucus structure and extracellular DNA. *PLoS One* 9:e95274. <https://doi.org/10.1371/journal.pone.0095274>.
- Martin, C. R., P. R. Ling, and G. L. Blackburn. 2016. Review of infant feeding: Key features of breast milk and infant formula. *Nutrients* 8:279. <https://doi.org/10.3390/nu8050279>.
- Masum, A. K. M., J. Chandrapala, T. Huppertz, B. Adhikari, and B. Zisu. 2020. Influence of drying temperatures and storage parameters on the physicochemical properties of spray-dried infant milk formula powders. *Int. Dairy J.* 105:104696. <https://doi.org/10.1016/j.idairyj.2020.104696>.
- Ménard, O., C. Bourlieu, S. C. De Oliveira, N. Dellarosa, L. Laghi, F. Carrière, F. Capozzi, D. Dupont, and A. Deglaire. 2018. A first step towards a consensus static in vitro model for simulating full-term infant digestion. *Food Chem.* 240:338–345. <https://doi.org/10.1016/j.foodchem.2017.07.145>.
- Musa, M. A., M. Kabir, M. I. Hossain, E. Ahmed, A. Siddique, H. Rashid, M. Mahfuz, D. Mondal, T. Ahmed, W. A. Petri, and R. Haque. 2019. Measurement of intestinal permeability using lactulose and mannitol with conventional five hours and shortened two hours urine collection by two different methods: HPAE-PAD and LC-MSMS. *PLoS One* 14:e0220397. <https://doi.org/10.1371/journal.pone.0220397>.
- Navis, M., V. Muncan, P. T. Sangild, L. Møller Willumsen, P. J. Koelink, M. E. Wildenberg, E. Abrahamse, T. Thymann, R. M. van Elburg, and I. B. Renes. 2020a. Beneficial effect of mildly pasteurized whey protein on intestinal integrity and innate defense in preterm and near-term piglets. *Nutrients* 12:1125. <https://doi.org/10.3390/nu12041125>.
- Navis, M., L. Schwebel, S. Søndergaard Kappel, V. Muncan, P. T. Sangild, E. Abrahamse, L. Aunsholt, T. Thymann, R. M. van Elburg, and I. B. Renes. 2020b. Mildly pasteurized whey protein promotes gut tolerance in immature piglets compared with extensively heated whey protein. *Nutrients* 12:3391. <https://doi.org/10.3390/nu12113391>.
- Nguyen, T. T. P., B. Bhandari, J. Cichero, and S. Prakash. 2015. A comprehensive review on in vitro digestion of infant formula. *Food Res. Int.* 76:373–386. <https://doi.org/10.1016/j.foodres.2015.07.016>.
- Noel, G., J. G. In, J. M. Lemme-Dumit, L. R. DeVine, R. N. Cole, A. L. Guerrero, J. D. Campbell, O. Kovbasnjuk, and M. F. Pasetti. 2021. Human breast milk enhances intestinal mucosal barrier function and innate immunity in a healthy pediatric human enteroid model. *Front. Cell Dev. Biol.* 9:685171. <https://doi.org/10.3389/fcell.2021.685171>.
- Odamaki, T., K. Kato, H. Sugahara, N. Hashikura, S. Takahashi, J. Z. Xiao, F. Abe, and R. Osawa. 2016. Age-related changes in gut microbiota composition from newborn to centenarian: A cross-sectional study. *BMC Microbiol.* 16:90. <https://doi.org/10.1186/s12866-016-0708-5>.
- Oldfield, D. J., M. W. Taylor, and H. Singh. 2005. Effect of preheating and other process parameters on whey protein reactions during skim milk powder manufacture. *Int. Dairy J.* 15:501–511. <https://doi.org/10.1016/j.idairyj.2004.09.004>.
- Pellegrino, L., S. Cattaneo, and I. De Noni. 2011. Nutrition and health: Effects of processing on protein quality of milk and milk products. Pages 1067–1074 in *Encyclopedia of Dairy Sciences*. 2nd ed. J. W. Fuquay, ed. Academic Press, San Diego, CA.
- Phosanam, A., J. Chandrapala, T. Huppertz, B. Adhikari, and B. Zisu. 2021. In vitro digestion of infant formula model systems: Influence of casein to whey protein ratio. *Int. Dairy J.* 117:105008. <https://doi.org/10.1016/j.idairyj.2021.105008>.
- Pischetsrieder, M., B. Larisch, and T. Severin. 2005. The Maillard reaction of ascorbic acid with amino acids and proteins—Identification of products. Pages 107–112 in *The Maillard Reaction in Foods and Medicine*. J. O'Brien, H. E. Nursten, M. J. C. Crabbe, and J. M. Ames, ed. Woodhead Publishing.
- Rasmussen, S. O., L. Martin, M. V. Østergaard, S. Rudloff, Y. Li, M. Roggenbuck, S. B. Bering, and P. T. Sangild. 2016. Bovine colostrum improves neonatal growth, digestive function, and gut immunity relative to donor human milk and infant formula in preterm pigs. *Am. J. Physiol. Gastrointest. Liver Physiol.* 311:G480–G491. <https://doi.org/10.1152/ajpgi.00139.2016>.
- Sun, J., H. G. Akilloğlu, K. Aasmul-Olsen, Y. Ye, P. Lund, X. Zhao, A. Brunse, C. F. Nielsen, D. E. W. Chatterton, P. T. Sangild, M. N. Lund, and S. B. Bering. 2022. Ultra-high temperature treatment and storage of infant formula induces dietary protein modifications, gut dysfunction and inflammation in preterm pigs. *Mol. Nutr. Food Res.* 66:2200132. <https://doi.org/10.1002/mnfr.202200132>.
- Taylor, S. N., L. A. Basile, M. Ebeling, and C. L. Wagner. 2009. Intestinal permeability in preterm infants by feeding type: Mother's milk versus formula. *Breastfeed. Med.* 4:11–15. <https://doi.org/10.1089/bfm.2008.0114>.
- Trompette, A., J. Claustre, F. Caillon, G. Jourdan, J. A. Chayvialle, and P. Plaisancié. 2003. Milk bioactive peptides and beta-casomorphins induce mucus release in rat jejunum. *J. Nutr.* 133:3499–3503. <https://doi.org/10.1093/jn/133.11.3499>.
- Van Boekel, M. A. J. S. 1998. Effect of heating on Maillard reactions in milk. *Food Chem.* 62:403–414. [https://doi.org/10.1016/S0308-8146\(98\)00075-2](https://doi.org/10.1016/S0308-8146(98)00075-2).
- van der Lugt, T., K. Venema, S. van Leeuwen, M. F. Vrolijk, A. Opperhuizen, and A. Bast. 2020. Gastrointestinal digestion of dietary advanced glycation endproducts using an in vitro model of the gastrointestinal tract (TIM-1). *Food Funct.* 11:6297–6307. <https://doi.org/10.1039/D0FO00450B>.
- van Lieshout, G. A. A., T. T. Lambers, M. C. E. Bragt, and K. A. Hettinga. 2020. How processing may affect milk protein digestion and overall physiological outcomes: A systematic review. *Crit. Rev. Food Sci. Nutr.* 60:2422–2445. <https://doi.org/10.1080/10408398.2019.1646703>.

- Wada, Y., and B. Lönnerdal. 2015a. Bioactive peptides released by in vitro digestion of standard and hydrolyzed infant formulas. *Peptides* 73:101–105. <https://doi.org/10.1016/j.peptides.2015.09.005>.
- Wada, Y., and B. Lönnerdal. 2015b. Effects of industrial heating processes of milk-based enteral formulas on site-specific protein modifications and their relationship to in vitro and in vivo protein digestibility. *J. Agric. Food Chem.* 63:6787–6798. <https://doi.org/10.1021/acs.jafc.5b02189>.
- Walton, K. D., A. M. Freddo, S. Wang, and D. L. Gumucio. 2016. Generation of intestinal surface: An absorbing tale. *Development* 143:2261–2272. <https://doi.org/10.1242/dev.135400>.
- World Health Organization. 2013. Essential nutrition actions: Improving maternal, newborn, infant and young child health and nutrition. World Health Organization, Geneva, Switzerland.
- Wu, Y., M. Zong, Z. Zhang, Y. Wu, L. Li, X. Zhang, H. Wu, and B. Li. 2022. Selective transportation and energy homeostasis regulation of dietary advanced glycation end-products in human intestinal Caco-2 cells. *Food Chem.* 391:133284. <https://doi.org/10.1016/j.foodchem.2022.133284>.
- Xiao, T., J. Zeng, C. Zhao, Y. Hou, T. Wu, Z. Deng, and L. Zheng. 2023. Comparative analysis of protein digestion characteristics in human, cow, goat, sheep, mare, and camel milk under simulated infant condition. *J. Agric. Food Chem.* 71:15035–15047. <https://doi.org/10.1021/acs.jafc.3c03123>.
- Yao, Y., X. Cai, Y. Ye, F. Wang, F. Chen, and C. Zheng. 2021. The role of microbiota in infant health: From early life to adulthood. *Front. Immunol.* 12:708472. <https://doi.org/10.3389/fimmu.2021.708472>.
- Ye, M. P., R. Zhou, Y. R. Shi, H. C. Chen, and Y. Du. 2017. Effects of heating on the secondary structure of proteins in milk powders using mid-infrared spectroscopy. *J. Dairy Sci.* 100:89–95. <https://doi.org/10.3168/jds.2016-11443>.
- Ye, Y., K. Engholm-Keller, Y. Fang, C. F. Nielsen, A. Jordà, M. N. Lund, and D. E. W. Chatterton. 2022. UHT treatment and storage of liquid infant formula affects protein digestion and release of bioactive peptides. *Food Funct.* 13:344–355. <https://doi.org/10.1039/D1FO02619D>.
- Yu, X., N. Leconte, S. Méjean, G. Garric, S. Even, G. Henry, F. J. Tessier, M. Howsam, T. Croguennec, G. Gésan-Guizieu, D. Dupont, R. Jeantet, and A. Deglaire. 2021. Semi-industrial production of a minimally processed infant formula powder using membrane filtration. *J. Dairy Sci.* 104:5265–5278. <https://doi.org/10.3168/jds.2020-19529>.
- Zenker, H. E., G. A. A. van Lieshout, M. P. van Gool, M. C. E. Bragt, and K. A. Hettinga. 2020. Lysine blockage of milk proteins in infant formula impairs overall protein digestibility and peptide release. *Food Funct.* 11:358–369. <https://doi.org/10.1039/C9FO02097G>.
- Zhu, R., H. Cheng, L. Li, H. R. Erichsen, M. A. Petersen, J. Soerensen, and L. H. Skibsted. 2018. Temperature effect on formation of advanced glycation end products in infant formula milk powder. *Int. Dairy J.* 77:1–9. <https://doi.org/10.1016/j.idairyj.2017.09.005>.
- Zoghbi, S., A. Trompette, J. Claustre, M. El Homsí, J. Garzón, G. Jourdan, J.-Y. Scoazec, and P. Plaisancié. 2006. β -Casomorphin-7 regulates the secretion and expression of gastrointestinal mucins through a μ -opioid pathway. *Am. J. Physiol. Gastrointest. Liver Physiol.* 290:G1105–G1113. <https://doi.org/10.1152/ajpgi.00455.2005>.

ORCIDS

Cathal A. Dold, <https://orcid.org/0000-0001-9307-2705>

Aylin W. Sahin, <https://orcid.org/0000-0002-1138-6063>

Linda Giblin <https://orcid.org/0000-0002-9354-3121>