

Title	Anthropogenic pressures within the breeding range of the Hen Harrier <i>Circus cyaneus</i> in Ireland
Authors	Caravaggi, Anthony;Irwin, Sandra;Lusby, John;Ruddock, Marc;Mee, Allan;Nagle, Tony;O'Toole, Lorcán;O'Neill, Shane;O'Halloran, John
Publication date	2020-03-27
Original Citation	Caravaggi, A., Irwin, S., Lusby, J., Ruddock, M., Mee, A., Nagle, T., O'Toole, L., O'Neill, S. and O'Halloran, J. (2020) 'Anthropogenic pressures within the breeding range of the Hen Harrier <i>Circus cyaneus</i> in Ireland', <i>Bird Study</i> , 66(4), pp. 461-470. doi: 10.1080/00063657.2020.1725420
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1080/00063657.2020.1725420
Rights	© 2020, British Trust for Ornithology. All rights reserved. This is an Accepted Manuscript of an item published by Taylor & Francis in <i>Bird Study</i> on 27 March, 2020, available online: https://doi.org/10.1080/00063657.2020.1725420
Download date	2024-05-14 00:44:27
Item downloaded from	https://hdl.handle.net/10468/9856

Original Research Paper

Anthropogenic pressures within the breeding range of the Hen Harrier (*Circus cyaneus*) in Ireland

Anthony Caravaggi^{1,2}, Sandra Irwin¹, John Lusby³, Marc Ruddock⁴, Allan Mee⁵, Tony

Nagle⁶, John O'Halloran^{1*}

¹ School of Biological, Earth and Environmental Sciences, University College Cork,
Distillery Field, North Mall, Cork, Ireland, T23 XA50.

² University of South Wales, 9 Graig Fach, Pontypridd, UK CF37 4BB.

³ BirdWatch Ireland, Unit 20, Block D, Bullford Business Campus, Kilcoole, Co. Wicklow,
Ireland, A63 RW83.

⁴ Golden Eagle Trust Ltd, 12 Ely Place, Dublin 2, Ireland, D02 T651.

⁵ Ardpatrick, Kilmallock, Co. Limerick, Ireland.

⁶ The Rookery, Ballyfeard, Co. Cork, Ireland.

*Corresponding author: j.ohalloran@ucc.ie

Keywords: raptor; human impacts; upland conservation; bird of prey; land management; policy

Word count (excluding references, table and figure): XXXX

Short title: Impacts on Hen Harrier breeding habitat

Summary

Capsule

Patterns in the frequency and co-occurrence of anthropogenic pressures associated with suitable breeding habitat for Hen Harriers demonstrates the need for specific, focussed management and policy options aimed at mitigating impacts on this threatened population.

Aims

To describe anthropogenic pressures and threats in the upland breeding range of Hen Harriers and to explore their potential impacts on the declining Hen Harrier population.

Methods

We used text mining, mixed effects models, Principal Component Analysis and clustering methods to explore anthropogenic pressures on suitable breeding and foraging habitats for Hen Harriers in Ireland, based on the 2015 national breeding Hen Harrier survey data.

Results

Mixed-effects models described a strong influence of agriculture, forestry, predator activity and recreational activities on survey areas that contained Hen Harrier territories. Cluster analyses described three discrete pressure clusters and showed consistent co-occurrence of independent pressures.

Conclusions

Areas of suitable habitat for Hen Harriers in the uplands overlap with areas that experience anthropogenic pressures known to negatively impact on this vulnerable bird species. Combined with clear evidence for the co-occurrence of multiple pressures at regional scale, this demonstrates a clear need for statutory agencies to consider the potential cumulative impacts of individual pressures when developing conservation strategies for Hen Harriers.

48

49 **Introduction**

50 Many species, worldwide, are threatened by anthropogenic pressures that require intervention
51 to mitigate or eliminate their negative impacts (Wilcove et al. 1998; Carroll et al. 2015; Di
52 Minin et al. 2016). Such pressures can result in stress responses or reduced fitness in wildlife
53 that, in some cases, has severe impacts on individuals or populations (Wilcove et al. 1998;
54 Taylor & Knight 2003; Johnson et al. 2005; Ciuti et al. 2012; Coetzee & Chown 2016).
55 Conservation processes typically aim to prevent species population declines and extinctions
56 (Soule 1985). However, conservation policy must also be cognisant of the sustainable
57 management of environmental resources and other activities of economic and social
58 importance including commercial forestry, agriculture and recreation (Young et al. 2005;
59 Kareiva & Marvier 2012; Kennedy et al. 2016; Vangansbeke et al. 2017)

60 Human activities in the vicinity of breeding birds can lead to increased rates of nest
61 desertion (White & Thurow 1985), and reduced rates of site occupancy (Webber et al. 2013),
62 territory establishment (Bötsch et al. 2017), breeding success (Balotari-Chiebao et al. 2016)
63 and survival (Ruhlen et al. 2003; including illegal killing, e.g. Smart et al. 2010).
64 Quantifying the extent and ecological relevance of each of these impacts informs our
65 understanding of human-wildlife interactions and underpins conservation and resource
66 management processes. It is essential, therefore, that human activities that have the potential
67 to affect wildlife, particularly vulnerable species of conservation concern, are properly
68 assessed and understood, so that appropriate measures can be developed to facilitate
69 conservation and sustainable land and resource use.

70 Hen Harriers (*Circus cyaneus*) are medium-sized raptors that nest largely upland
71 areas, typically heather moorland in Britain (Redpath et al. 1998; Amar et al. 2008; Watson
72 2017), during the summer breeding season. Upland habitats in Ireland have been subjected to

degradation and land-use change and, in the absence of their preferred open heath and blanket bog nesting habitat. As a result of a large-scale afforestation programme in the Republic of Ireland from the 1950s and the conversion of ‘traditional’ open habitats to forest, Hen Harriers in Ireland are frequently associated with young (i.e. pre-thicket) conifer plantations that provide them with areas for nesting and foraging (Wilson et al. 2009; Irwin et al. 2012; Wilson et al. 2012; Ruddock et al. 2016). Anthropogenic impacts such as afforestation and forest management (NPWS 2015), landscape degradation and land-use change (Wilson et al. 2009; Wilson et al. 2012), livestock grazing (O’Rourke & Kramm 2009), illegal burning (Renou-Wilson et al. 2011), peat extraction (O’Riordan et al. 2015), recreation (Hynes & Buckley 2007) and wind energy development (Wilson et al. 2017) could have important implications for breeding Hen Harriers. It should be noted that the level of persecution observed in Britain (e.g. Redpath et al. 2010; Murgatroyd et al. 2019) is not observed in Ireland as there are no areas that are managed solely for driven grouse shooting. However, raptors are known to migrate within the British Isles (Mead, 1973) and persecution of Hen Harriers in Britain could have hitherto undescribed impacts on the Irish population. The Hen Harrier population in Ireland is of national conservation concern (Colhoun & Cummins 2013), with a population of between 108 and 157 breeding pairs recorded in the most recent national survey (Ruddock et al. 2016). The species is listed under Annex I of the European Commission Birds Directive (2009/147/EC) that requires Member States to designate Special Protection Areas (SPAs) for their survival and reproduction. Six Hen Harrier SPAs containing important breeding areas for the species were designated in Ireland in 2007.

Hen Harrier conservation research in Ireland to date has focussed on the impacts of afforestation (Irwin et al., 2012; Wilson et al., 2009, 2012) and wind farm development (Fernández-Bellon et al., 2015; Wilson et al., 2017) on their populations, as required to inform conservation management. Due to the targeted nature of previous research, very little

information is available in the published literature regarding the broader range of anthropogenic pressures that might impact breeding Hen Harriers and associated foraging and breeding habitat. Furthermore, previous research has considered how individual pressures impact separately and in specific contexts while consideration of the synergies between pressures is lacking. To address these gaps, we explored data on anthropogenic pressures affecting Hen Harriers within their breeding range in Ireland, with the aim of deriving information that would inform conservation and management processes for this threatened species.

Materials and methods

The 2015 National Survey of Breeding Hen Harrier in Ireland was conducted between April and August 2015 in suitable Hen Harrier habitat in upland areas, largely, but not exclusively, between 200m and 600m above sea level (asl) and within the Hen Harrier breeding range (Ruddock et al. 2016). Survey squares of 10 km² (n = 268) were defined using the Irish National Grid (Fig. 1a). Anthropogenic activities that could potentially impact on breeding Hen Harriers ('pressures' from hereon) were recorded from vantage points within each survey square during each of 4-6 dedicated watches per square, during the breeding season. Where sites were occupied, vantage points were a minimum of 500 m from nests sites. Vantage points were identified *a-priori* based on habitat suitability, topographical constraints and the potential for observers to cause disturbance to breeding birds (Ruddock and Whitfield 2007; Philip Whitfield et al. 2008). Hen Harrier territories (n = 100, across 54 survey squares) were recorded where identified; occupancy was based on observations of Hen Harrier breeding behaviour and the repeated presence of birds (Ruddock et al., 2016). Data were collected by staff, members and volunteers from the National Parks & Wildlife Service

(NPWS), BirdWatch Ireland (BWI), Irish Raptor Study Group (IRSG), Golden Eagle Trust (GET), university researchers, and independent commercial and voluntary ornithological surveyors

Pressures were divided into 47 discrete categories (Appendix I) aligned with the EU Birds Directive (2009/147/EC) reporting matrix. The frequency of occurrence of each pressure within 2 km of vantage point locations was recorded within each survey square. Initial exploration of the data revealed extreme outliers, therefore we adopted a precautionary approach and applied consistent thresholds throughout. Values for individual pressures that occurred beyond two standard deviations (SDs) from the mean were replaced with the maximum value as defined by the aforementioned threshold, rounded to the nearest whole integer. This allowed us to capture the prevalence of each pressure at each location while mitigating over-inflation. The sum frequency of each pressure was calculated (i) across all survey squares where the total number of recorded pressures was >0 ($n=146$; Appendix II); (ii) across squares located within SPA boundaries only ($n=24$); and (iii) across squares where confirmed Hen Harrier territories were present ($n=54$). It was necessary to account for variation in survey effort as the number of visits made to vantage points varied between observers. Therefore, a Pressure Index (PI) was created, where the total number of pressures was divided by the total number of visits (Ruddock et al. 2016). PI scores were normalised between 0 and 1 to facilitate comparisons between sites. General Linear Models (GLMs) were used to investigate differences between PI scores – with zero counts removed and remaining data log transformed to meet model assumptions – where PI was the dependent variable and the location of vantage points relative to SPA boundaries (inside/outside) and confirmed Hen Harrier territories (present/absent) were explanatory variables. Models explored each category (SPA boundaries and Hen Harrier territories) independently as well as part of a fully-factorial model that included an interaction term.

Principal Component Analysis (PCA; Jolliffe & Cadima, 2016) and linear mixed-effects models were used to investigate relationships between the presence/absence of Hen Harrier territories and pressure categories. Data were Box Cox transformed to remove skewness, centred and standardised to have a $\bar{x}=0$ and $\sigma = 1$ prior to analysis. Principal Components (PCs) that cumulatively accounted for >50% of the variance were retained for inclusion in models. The presence/absence of Hen Harrier territories (Fig. 1b) was entered as a binary dependent variable, retained PCs were included as explanatory variables and surveyor identity was included as a random variable. Model permutations were ranked using the Akaike Information Criterion (AIC); the top subset of models was found within $\Delta AIC \leq 2$ units (Burnham & Anderson 2002).

Cluster analysis was used to quantify associations between individual pressure categories across all survey squares. The various methods that comprise cluster analyses provide a means of classifying multivariate data into subgroups according to the similarity of their attributes, thus revealing the underlying structure (Everitt et al. 2009). We calculated the distance of each recorded pressure from the cluster's mean using a Euclidean distance index and applied the Ward error sum of squares hierarchical clustering method (Ward 1963) to the resultant data. The optimal number of clusters (k_t) was identified using average silhouettes (Kaufman & Rousseeuw 1990) and Approximately Unbiased (AU) p -values with multiscale bootstrap resampling ($B = 10,000$) where clusters with $p \geq 0.95$ were strongly supported (Suzuki and Shimodaira 2006). All data analyses and plotting were carried out using the statistical programme R (R Core Team 2017), specifically the packages *cluster* (Maechler et al. 2018) and *pvclust* (Suzuki and Shimodaira 2015), *dendextend* (Galili 2015), *nlme* (Pinheiro et al. 2017) and *caret* (Kuhn 2017). Data are subject to data-sharing agreements and, therefore, cannot be redistributed. However, R code used for data exploration and analyses are available at <http://doi.org/10.5281/zenodo.3549584>.

172

173 **Results**

174 A total of 2,873 individual pressure occurrences were recorded during this study. There were
175 no anthropogenic pressures recorded in 45% of survey squares. The most frequently recorded
176 pressures across all survey squares were *forest management and use* (13% of occurrences),
177 *paths, tracks, forest roads* (11%), *uncontrolled burning* (6%) and *wind energy production*
178 (6%). Similar pressures were recorded inside and outside of SPA boundaries: *forest*
179 *management and use* (14% and 11% of occurrences, respectively), *paths, tracks, forest roads*
180 (10%, 0%), *forest planting on open ground* (0%, 8%), *uncontrolled burning* (6.6%, 7%) and
181 *wind energy production* (9.5%, 7%). The most frequently recorded pressures associated with
182 confirmed Hen Harrier territories were *loss of habitat features* (13.7%), *dispersed habitation*
183 (10.5%), *paths, tracks, forest roads* (9.2%) and *forest management* (8.1%). In contrast,
184 pressures at vantage points not associated with Hen Harrier territories were *forest*
185 *management and use* (16%), off-road motorised driving (12%), *forest planting on open*
186 *ground* (11%), and *mechanical removal of peat* (11%).

187 Pressure Indices varied between survey squares (Fig. 1c) and only one survey square
188 had a $PI > 0.5$. Survey squares where vantage points occurred within SPAs had a maximum
189 PI of 0.22 ($\bar{x} = 0.08 \pm 0.07$), which was significantly higher than those outside SPAs ($t =$
190 0.028 ; $\beta = -0.44 \pm 0.20$; $P = 0.03$). Survey squares where vantage points were associated with
191 Hen Harrier territories had a maximum PI of 0.42 ($\bar{x} = 0.10 \pm 0.09$; Appendix I), which was
192 significantly higher than those that were not associated with territories ($t = 0.038$; $\beta = -0.39 \pm$
193 0.19 ; $P = 0.04$; Table 1).

194 Both silhouette and AU clustering methods supported three discrete clusters ($P \geq$
195 0.05). The largest cluster (ii) consisted of 25 pressure categories while the smallest (iii) was

the most distinct and consisted of five pressure categories. One sub-cluster was statistically supported (*iv*) and was comprised of 17 pressure categories (Fig. 2).

A total of seven Principal Component axes, accounting for >50% of the total variance, were retained for inclusion in mixed-effects models investigating the relationship between the presence/absence of Hen Harrier territories and associated pressures. The top subset of models ($\Delta AIC \leq 2$) included PC1, PC2 and PC3. PC1 accounted for the greatest proportion of total variance (20%); loadings were most strongly weighted towards aspects of agricultural and forestry activity and predators; PC2 (8%) was weighted towards forest management and site access; and PC3 was weighted towards forest clearance and recreational activities (Table 2). The best approximating model was positively influenced by PC1 and PC3, and negatively influenced by PC2 (Table 2). It should be noted that PC1 includes *nest destruction*, *predation by birds* and *predation by mammals*. These pressures can only occur where Hen Harriers nest, hence the observed positive association is to be expected.

Discussion

Our results show that suitable Hen Harrier breeding habitats in Ireland are subjected to a wide range of anthropogenic pressures that could have significant implications for this vulnerable species. The number and variety of pressures recorded demonstrates the potential of disturbance to prospecting Hen Harriers early in the breeding season and/or to foraging Hen Harriers once territories have been established (e.g. González et al. 2006). Furthermore, the co-occurrence of pressures as described by cluster analyses demonstrates the considerable potential for cumulative effects. Anthropogenic impacts are not homogenous in their severity or extent. This is certainly true in the current study, where some pressures will have more severe consequences for Hen Harriers or will act at different spatial scales. However, there is

a dearth of quantitative data on the impacts of described pressures on Hen Harriers. Our results highlight the importance of managing pressures in an integrated manner rather than on an individual basis. This provides support for the effective management of suitable breeding areas to minimise the potential impact of anthropogenic pressures on vulnerable Hen Harrier populations.

Planted forests and the presence of tracks or roads were recorded at high frequencies in all survey squares across Ireland. Large areas of Irish upland habitat have been afforested in recent decades and total forest cover is expected to continue to increase from the current 11% to as much as 18% in the next 30 years (NPWS 2015). In the absence of their traditional open heath and blanket bog habitat, Hen Harriers in Ireland are frequently associated with young (i.e. pre-thicket) conifer plantations that provide areas that Hen Harriers use for nesting and foraging (Wilson et al. 2009; Irwin et al. 2012; Wilson et al. 2012; Ruddock et al. 2016). Hen Harriers cannot use closed-canopy forests for breeding or foraging, therefore the maturation of the existing forest estate threatens to deprive Hen Harriers of already scarce breeding habitat, while further increases in forest cover could also lead to increased habitat fragmentation and subsequently reduce the capacity of the landscape to support breeding pairs. Recreational activities were also strongly associated with survey squares containing Hen Harrier territories. Systematic reviews have demonstrated that recreational activities can negatively impact breeding birds (e.g. Steven et al. 2011; Larson et al. 2016) including above-ground foragers (Bötsch et al. 2017) and upland species such as Golden Plovers (*Pluvialis apricarius*; Finney et al. 2005). Thus, there exists the potential for disturbance of prospecting Hen Harriers early in the breeding season and/or foraging Hen Harriers once territories have been established.

Mammalian and avian predators were among the factors strongly associated with Hen Harrier territories. O'Donoghue (2010) attributed 55% of all nest failures in south and west

Ireland in 2007 and 2008 to predation events and foxes (*Vulpes vulpes*) have been observed depredating Hen Harrier chicks via remote-sensing camera traps (Irwin et al. 2012; Fernández-Bellon et al. 2017). Other potential predators of Hen Harrier nests in Ireland include Pine Marten (*Martes martes*), American Mink (*Neovison vison*), Stoat (*Mustela erminea*), Raven (*Corvus corax*) and Hooded Crow (*Corvus corone corvix*) (Picozzi, 1984; Fernández-Bellon et al., 2018a). These predators can have substantial negative impacts on ground-nesting birds (Paton 1994) as eggs and young chicks are particularly vulnerable to predation when parents are absent. Populations of generalist predators may be bolstered by changes in land-use and management, including afforestation and other forms of habitat fragmentation (e.g. Prestt, 1965; Haydon & Harrington, 2000; Chalfoun et al. 2002; Twining et al. 2019). However, data on the abundance and activity of upland predators in Ireland are scarce; efforts to investigate such may be of considerable benefit to the conservation of Hen Harriers.

It is notable that wind energy production was recorded more frequently within SPA boundaries than outside but was rarely recorded in survey squares that contained breeding Hen Harriers (3.3%) in this study, perhaps indicating avoidance of wind farms for breeding purposes (also see Wilson et al. 2017). Indeed, windfarm construction activity has been implicated in the desertion of traditional breeding sites in Ireland (O'Donoghue et al. 2011). The construction and operation of wind turbines can have both lethal and sub-lethal impacts on birds (Drewitt & Langston 2006; Marques et al. 2014; Wilson et al. 2015; Balotari-Chiebao et al. 2016; Smith & Dwyer 2016; Fernández-Bellon et al. 2018; Thaker et al. 2018). The Republic of Ireland is committed to EU targets on renewable energy including a national target of 40% electricity from renewables by 2020, which is likely to involve the construction of additional wind farms (DCCA 2010). Wind energy developments tend to be upland-focussed and future, large-scale expansion may pose a threat to breeding Hen Harriers. A bird

sensitivity mapping tool has been developed to guide the siting of future wind energy developments in Ireland in relation to the distribution of species of conservation concern, including Hen Harrier (McGuinness et al. 2015). However, there is as yet no mandatory obligation on developers to use this tool.

The timing of disturbance events may be a key consideration and many sources of disturbance may already be present at the onset of breeding, when pairs are establishing territories. Furthermore, other pressures such as peat extraction or illegal burning may not occur until after laying and, hence, can impact on parental care and, ultimately, breeding success. Current mitigation measures for Hen Harriers in Ireland adopt a reactive approach where circular ‘High Likelihood Nesting Areas’ (HNLA, formerly Red Areas) of high sensitivity to Hen Harriers, that contain nesting pairs and with a radius of 1.2 km, are added to the HNLA network when new breeding pairs are identified (NPWS 2015). Forestry operations that may cause disturbance are regulated within these HNLAs during the breeding season (Forest Service 2012). However, the protection afforded by HNLAs only applies to known pairs within the SPA network. Therefore, all other pairs that are outside of the SPA network (>50% of the breeding population; Ruddock et al. 2016), remain vulnerable to direct disturbance from forest management activities during the nesting season. Moreover, breeding Hen Harriers have been recorded travelling as far as 11 km from active nests (Irwin et al. 2012; Arroyo et al. 2014) and human activities and impacts in the wider landscape can have impacts on the physiology (Abbasi et al. 2017) and mortality (Ferrer & Hiraldo 1993) of birds. Human activities within the foraging range of breeding Hen Harriers could result in patch avoidance and/or stress-related responses in foraging birds, potentially keeping them away from the nest for longer periods of time and subsequently increasing chick vulnerability. It is possible that Hen Harriers in Ireland have the capacity to develop a tolerance for human activities during the breeding season. However, given their small

population size and conservation status, the precautionary principle suggests that human activities should be strictly regulated in areas of suitable Hen Harrier breeding and foraging habitat, particularly during key breeding months. Furthermore, pressures and their potential impacts on breeding Hen Harriers must be placed in a broader context that includes the timing of pressure occurrence, the composition of the wider landscape and the conservation of suitable habitat.

The pressures described herein represent potential disturbances to Hen Harriers throughout their breeding cycle and therefore may have important consequences for long-term population persistence or recovery. Recent research suggests that the same pressures impact another upland bird of prey, the Short-eared Owl (*Asio flammeus*), across their European range (Fernández-Bellon, unpubl. data). Thus, we recommend the following actions to enhance conservation benefits for Hen Harriers and other sensitive upland species and habitats: i) restrict forestry activities within the known hen Harrier range during the Hen Harrier breeding season (April – August) by using targeted surveys to detect Hen Harrier presence, thereby ensuring that forest management activities can be undertaken in areas that do not hold Hen Harriers during the summer months; ii) quantify the abundance and activity of upland predators and explore options for predator control, where appropriate; iii) discourage recreation and non-licensed forestry-related activities in areas known to hold Hen Harriers, throughout the breeding season, supported by a programme of community engagement, awareness-raising and upland signage; and iv) improve lines of communication between the relevant stakeholders so that potentially damaging activities can be identified at the earliest stages.

Failure to mitigate anthropogenic disturbances in upland areas of potentially suitable Hen Harrier breeding habitat, whether inside or outside of SPAs, could have negative consequences for this already vulnerable population. To date, none of the SPAs in the Hen

Harrier Natura 2000 network possess management plans, one of the key requirements of such sites, over a decade on from designation in 2007, and a Hen Harrier Threat Response Plan, initiated by the National Parks & Wildlife Service in the Republic of Ireland in 2016 with wide stakeholder consultation, has yet to be published. Furthermore, connecting multiple pressures is a key issue for conservation management, and Hen Harrier conservation policies must comprehensively account for cumulative anthropogenic impacts at regional level. Successful mitigation and management would represent a significant step towards the conservation of Hen Harriers in Ireland and serve as an example for upland conservation initiatives in Europe.

Acknowledgments

The authors thank the many people who collected data for the 2015 National Survey of Breeding Hen Harrier in Ireland and members of the stakeholder and scientific steering groups for the Supporting Hen Harriers in Novel Environments (SHINE) research project at UCC. The SHINE research project was funded by the Department of Agriculture, Food & the Marine. We also thank the Editor and two anonymous reviewers whose feedback helped improve this manuscript.

References

- Abbasi, N.A., Arukwe, A., Veerle, L.B.J., Eulaers, I., Mennilo, E., Ibor, O.R., Frantz, A., Covaci, A. & Malik, R.N. 2017. Oxidative stress responses in relationship to persistent organic pollutant levels in feathers and blood of two predatory bird species from Pakistan. *Sci. Total. Environ.* **580**: 26–33.
- Amar, A., Arroyo, B., Meek, E., Redpath, S. & Riley, H. 2008. Influence of habitat on breeding performance of Hen Harriers *Circus cyaneus* in Orkney. *Ibis* **150**: 400–404.

345 **Arroyo, B., Leckie, F., Amar, A., McCluskie, A. & Redpath, S.** 2014. Ranging behaviour
346 of Hen Harriers breeding in Special Protection Areas in Scotland. *Bird Study* **61**: 48–55.

347 **Balotari-Chiebao, F., Brommer, J.E., Niinimäki, T. & Laaksonen, T.** 2016. Proximity to
348 wind-power plants reduces the breeding success of the white-tailed eagle. *Anim. Conserv.* **19**:
349 265–272.

350 **Bötsch, Y., Tablado, Z. & Jenni, L.** 2017. Experimental evidence of human recreational
351 disturbance effects on bird-territory establishment. *Philos. Trans. R. Soc. Lond. B. Biol.*
352 *Sci.* **284**: 20170846.

353 **Burnham, K. & Anderson, D.** 2002. *Model selection and multi-model inference: a*
354 *practical information-theoretic approach*. New York: Springer.

355 **Carroll, C., Rohlf, D.J., Li, Y.-W., Hartl, B., Phillips, M.K. & Noss, R.F.** 2015.
356 Connectivity conservation and endangered species recovery: a study in the challenges of
357 defining conservation-reliant species: *Conserv. Lett.* **8**: 132–138.

358 **Chalfoun, A.D., Thompson, F.R. & Ratnaswamy, M.J.** 2002). Nest predators and
359 fragmentation: a review and meta-analysis. *Conserv. Biol.* **16**: 306–318.

360 **Ciuti, S., Northrup, J.M., Muhly, T.B., Simi, S., Musiani, M., Pitt, J.A. & Boyce, M.S.**
361 2012. Effects of humans on behaviour of wildlife exceed those of natural predators in a
362 landscape of fear. *PLOS ONE* **7**: p.e50611.

363 **Coetzee, B.W.T. & Chown, S.L.** 2016. A meta-analysis of human disturbance impacts on
364 Antarctic wildlife. *Biol. Rev.* **91**: 578–596.

365 **Colhoun, K. & Cummins, S.** 2013. Birds of conservation concern in Ireland. *Irish Birds* **9**:
366 523–544.

367 **Department of Communications, Climate Action & Environment [DCCAE].** 2010. *The*
368 *national renewable energy action plan: Submitted under Article 4 of Directive 2009/28/EC*.
369 Dublin, Ireland.

370 **Di Minin, E., Slotow, R., Hunter, L.T.B., Montesino Pouzols, F., Toivonen, T., Verburg,**
371 **P.H., Leader-Williams, N., Petracca, L. & Moilanen, A.** 2016. Global priorities for
372 national carnivore conservation under land use change. *Sci. Rep.* **6**: 23814.

373 **Drewitt, A.L. & Langston, R.H.W.** 2006. Assessing the impacts of wind farms on birds.
374 *Ibis* **148**: 29–42.

375 **Everitt, B., Landau, S. & Leese, M.** 2009. *Cluster analysis*. 4th ed. London : New York:
376 Arnold ; Oxford University Press.

377 **Fernández-Bellón, D., Irwin, S., Wilson, M. & O'Halloran, J.** 2015. Reproductive output
378 of Hen Harriers *Circus cyaneus* in relation to wind turbine proximity. *Irish Birds* **10**: 143–
379 150.

380 **Fernández-Bellón, D., Wilson, M., Irwin, S., Kelly, T.C., O'Mahony, B. & O'Halloran,**
381 **J.** 2018a. Video evidence of siblicide and cannibalism, movement of nestlings by adults, and
382 interactions with predators in nesting hen harriers. *J Raptor Res.* **52**: 393-399

383 **Fernández-Bellon, D., Wilson, M., Irwin, S. & O'Halloran, J.** 2018b. Effects of
384 development of wind energy and associated changes in land use on bird densities in upland
385 areas. *Conserv. Biol.* **33**: 413-422.

386 **Ferrer, M. & Hiraldo, F.** 1993. Evaluation of management techniques for the Spanish
387 Imperial Eagle. *Biol. Conserv.* **63**: 436-442.

388 **Finney, S.K., Pearce-Higgins, J.W. & Yalden, D.W.** 2005. The effect of recreational
389 disturbance on an upland breeding bird, the golden plover *Pluvialis apricaria*. *Biol. Conserv.*
390 **121**: 53–63.

391 **Galili, M.** 2015. dendextend: an R package for visualizing, adjusting, and comparing trees of
392 hierarchical clustering. *Bioinformatics* **31**: 3718–3720.

393 **González, L.M., Arroyo, B.E., Margalida, A., Sánchez, R. & Oria, J.** 2006. Effect of
394 human activities on the behaviour of breeding Spanish imperial eagles (*Aquila adalberti*):
395 management implications for the conservation of a threatened species. *Anim. Cons.* **9**: 85-93.

396 **Hayden, T. & Harrington, R.** 2000. *Exploring Irish mammals*. Town House and Country
397 House Ltd, Dublin.

398 **Hynes, S. & Buckley, C.** 2007. Recreational pursuits on marginal farm land: a discrete-
399 choice model of Irish farm commonage recreation. *Econ. Soc. Rev.* **38**: 63–84.

400 **Irwin, S., Wilson, M., Kelly, T.C., O'Donoghue, B., O'Mahony, B., Oliver, G., Cullen,**
401 **C., O'Donoghue, T. & O'Halloran, J.** 2008. Aspects of the breeding biology of Hen
402 Harriers *Circus cyaneus* in Ireland. *Irish Birds.* **8**: 331-334.

403 **Irwin, S., Wilson, M., O'Donoghue, B., O'Mahony, B., Kelly, T. & O'Halloran, J.** 2012.
404 *Optimum scenarios for Hen Harrier conservation in Ireland*. Cork: Department of
405 Agriculture, Food and the Marine by the School of Biological, Earth and Environmental
406 Sciences, University College Cork.

407 **Johnson, C.J., Boyce, M.S., Case, R.L., Cluff, H.D., Gau, R.J., Gunn, A. & Mulders, R.**
408 2005. Cumulative effects of human developments on Arctic wildlife. *Wildlife Monogr.* **160**:
409 1–36.

410 **Jolliffe, I.T. & Cadima, J.** 2016. Principal component analysis: a review and recent
411 developments. *Philos. Tran. R. S. A.* **374**: 20150202.

412 **Kareiva, P. & Marvier, M.** 2012. What is conservation science? *BioScience* **62**: 962–969.

413 **Kaufman, L. & Rousseeuw, P.J.** 1990. *Finding Groups in Data*. Hoboken, NJ, USA: John
414 Wiley & Sons, Inc.

415 **Kennedy, C.M., Miteva, D.A., Baumgarten, L., Hawthorne, P.L., Sochi, K., Polasky, K.,**
416 **Oakleaf, J.R., Uhlhorn, E.M. & Kiesecker, J.** 2016. Bigger is better: Improved nature
417 conservation and economic returns from landscape-level mitigation. *Sci. Adv.* **2**: e1501021–
418 e1501021.

419 **Kuhn, M.** 2017. *caret: Classification and regression training*. Available at: [https://CRAN.R-](https://CRAN.R-project.org/package=caret)
420 [project.org/package=caret](https://CRAN.R-project.org/package=caret)

421 **Larson, C.L., Reed, S.E., Merenlander, A.M. & Crooks, K.R.** 2016. Effects of recreation
422 on animals revealed as widespread through a global systematic review. *PLOS ONE* 11(12), p.
423 e0167259.

424 **Maechler, M., Rousseuw, P. Struyf, A., Hubert, M. & Hornik, K.** 2018. *cluster: Cluster*
425 *Analysis Basics and Extensions*. Available at: <https://CRAN.R-project.org/package=rgdal>.

426 **Marques, A.T., Batalha, H., Rodrigues, S., Costa, H., Pereira, M.J.R., Fonseca, C.,**
427 **Mascarenhas, M. & Bernardino, J.** 2014. Understanding bird collisions at wind farms: An
428 updated review on the causes and possible mitigation strategies. *Biol. Conserv.* **179**: 40–52.

429 **Mc Guinness, S., Muldoon, C., Tierney, N., Cummins, S., Murray, A., Egan, S. &**
430 **Crowe, O.** 2015. *Bird Sensitivity Mapping for Wind Energy Developments and Associated*
431 *Infrastructure in the Republic of Ireland*. BirdWatch Ireland, Kilcoole, Wicklow. #

432 **Mead, C.J.** 1973. Movements of British raptors. *Bird Study*. **20**: 259-286.

433 **Murgatroyd, M., Redpath, S.M., Murphy, S.G., Douglas, D.J., Saunders, R. & Amar, A.**
434 2019. Patterns of satellite tagged hen harrier disappearances suggest widespread illegal
435 killing on British grouse moors. *Nat. Commun.* **10**: 1094.

436 **National Parks & Wildlife Service [NPWS] 2015.** *Hen Harrier conservation and the*
437 *forestry sector in Ireland*. Dublin, Ireland: National Parks & Wildlife Service.

438 **O'Donoghue, B.** 2010. The Ecology and Conservation of Hen Harriers (*Circus cyaneus*) in
439 Ireland. PhD thesis. University College Cork.

440 **O'Donoghue, B., O'Donoghue, T.A. & King, F., 2011.**The Hen Harrier in Ireland:
441 conservation issues for the 21st century. *Biol. Env. Proc. R. Irish Acad.* **111B**: 83-93

442 **O'Riordan, M., Mahon, M. & McDonagh, J.** 2015. Power, discourse and participation in
443 nature conflicts: the case of turf cutters in the governance of Ireland's raised bog
444 designations. *J. Env. Policy Plann.* **17**:127-145.

445 **O'Rourke, E. & Kramm, N.** 2009. Changes in the management of the Irish Uplands: A
446 case-study from the Iveragh Peninsula. *Eur. Countryside* **1**: 53-66.

447 **Philip Whitfield, D., Ruddock, M. & Bullman, R.** 2008. Expert opinion as a tool for
448 quantifying bird tolerance to human disturbance. *Biol. Conserv.* **141**: 2708–2717.

449 **Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. & R Core Team.** 2017. *nlme: Linear and*
450 *Nonlinear Mixed Effects Models*. Available at: <https://CRAN.R-project.org/package=rgdal>.

451 **Prestt, I.** 1965. An enquiry into the recent breeding status of some of the smaller birds of
452 prey and crows in Britain. *Bird Study* **12**: 196-221.

453 **R Core Team** 2017. *R: A language and environment for statistical computing*. Vienna,
454 Austria.: R Foundation for Statistical Computing. Available at: <https://www.R-project.org/>.

455 **Redpath, S.M., Madders, M., Donnelly, E., Anderson, B., Thirgood, S., Martin, A.,**
456 **McLeod, D.** 1998. Nest site selection by Hen Harriers in Scotland. *Bird Study* **45**: 51–61.

457 **Redpath, S.M., Amar, A., Smith, A., Thompson, D.B. & Thirgood, S.** 2010. People and
458 nature in conflict: can we reconcile hen harrier conservation and game management. In
459 Baxter J. & Galbraith, C.A. (eds) *Species Management: Challenges and Solutions for the 21st*
460 *Century*, 335-350. Stationery Office Books, Edinburgh.

461 **Renou-Wilson, F., Bolger T., Bullock C., Convery F., Curry J., Ward S., Wilson D. &**
462 **Müller C.** 2011. *BOGLAND: Sustainable Management of Peatlands in Ireland – Final*
463 *Report*. Johnstown Castle, Co Wexford, Ireland: Environmental Protection Agency.

464 **Ruddock, M., Mee, A., Lusby, J., Nagle, T., O'Neill, S. & O'Toole, L.** 2016. *The 2015*
465 *National Survey of Breeding Hen Harrier in Ireland*. Irish Wildlife Manuals No. 93. National
466 Parks and Wildlife Service, Department of the Arts, Heritage and the Gaeltacht, Dublin,
467 Ireland.

468 **Ruddock, M. & Whitfield, D.P.** 2007. *A review of disturbance distances in selected bird*
469 *species*. Inverness, UK: Scottish Natural Heritage.

470 **Ruhlen, T.D., Abbott, S., Stenzel, L.E. & Page, G.W.** 2003. Evidence that human
471 disturbance reduces Snowy Plover chick survival. *J. Field Ornithol.* **74**: 300-305.

472 **Smart, J., Amar, A., Sim, I.M., Etheridge, B., Cameron, D., Christie, G. & Wilson, J.D.**
473 2010. Illegal killing slows population recovery of a re-introduced raptor of high conservation
474 concern—the red kite *Milvus milvus*. *Biol. Conserv.* **143**: 1278-1286.

475 **Smith, J.A. & Dwyer, J.F.** 2016. Avian interactions with renewable energy infrastructure:
476 An update. *The Condor* **118**: 411–423.

477 **Soule, M.E.** 1985. What is conservation biology? *BioScience* **35**: 727–734.

478 **Steven, R., Pickering, C. & Castley, J.G.** 2011. A review of the impacts of nature based
479 recreation on birds. *J. Env. Manage.* **92**: 2287–2294.

480 **Suzuki, R. & Shimodaira, H.** 2006. Pvcust: an R package for assessing the uncertainty in
481 hierarchical clustering. *Bioinformatics.* **22**: 1540–1542.

482 **Taylor, A.R. & Knight, R.L.** 2003. Wildlife responses to recreation and associated visitor
483 perceptions. *Ecol. Appl.* **13**: 951–963.

484 **Thaker, M., Zambre, A., Bhosale, H.** 2018. Wind farms have cascading impacts on
485 ecosystems across trophic levels. *Nat. Ecol. Evol.* **2**: 1854–1858.

486 **Twining, J.P., Montgomery, I., Fitzpatrick, V., Marks, N., Scantlebury, D.M. & Tosh,**
487 **D.G.** 2019. Seasonal, geographical, and habitat effects on the diet of a recovering predator
488 population: the European pine marten (*Martes martes*) in Ireland. *Euro. J Wildlife Res.* **65**:
489 51.

490 **Vangansbeke, P., Blondeel, H., Landuyt, D., De Frenne, P., Gorissen, L. & Herheyen, K.**
491 2017. Spatially combining wood production and recreation with biodiversity conservation.
492 *Biodivers. Conserv.* **26**: 3213–3239.

493 **Ward, J.H.** 1963. Hierarchical grouping to optimize an objective function. *J. Am. Stat.*
494 *Assoc.* **58**: 236–244.

- 495 **Watson, D.** 2017. *The Hen Harrier*. London, UK: Bloomsbury Natural History.
- 496 **Webber, A.F., Heath, J.A. & Fischer, R.A.** 2013. Human disturbance and stage- specific
497 habitat requirements influence snowy plover site occupancy during the breeding season. *Ecol.*
498 *Evol.* **3**: 853-863.
- 499 **White, C.M. & Thurow, T.L.** 1985. Reproduction of Ferruginous Hawks exposed to
500 controlled disturbance. *The Condor* **87**: 14–22.
- 501 **Wilcove, D.S., Rothstein, D., Phillips, A. & Losos, E.** 1998. Quantifying threats to
502 imperiled species in the United States. *BioScience* **48**: 607–615.
- 503 **Wilson, M.W., Fernández-Bellon, D., Irwin, S. & O’Halloran, J.** 2017. Hen Harrier
504 *Circus cyaneus* population trends in relation to wind farms. *Bird Study.* **64**: 20-29.
- 505 **Wilson, M.W., Irwin, S., Norriss, D.W., Newton, S.F., Collins, K., Kelly, T.C. &**
506 **O’Halloran, J.** 2009. The importance of pre- thicket conifer plantations for nesting Hen
507 Harriers *Circus cyaneus* in Ireland. *Ibis* **151**: 332-343.
- 508 **Wilson, M.W., O’Donoghue, B., O’Mahony, B., Cullen, C., O’Donoghue, T., Oliver, G.,**
509 **Ryan, B., Troake, P., Irwin, S., Kelly, T.C., Rotella, J.J. & O’Halloran, J.** 2012.
510 Mismatches between breeding success and habitat preferences in Hen Harriers *Circus*
511 *cyaneus* breeding in forested landscapes. *Ibis* **154**: 578-589.
- 512 **Young, J., Watt, A., Nowicki, P., Alard, D., Clitherow, J., Henle, K., Johnson, R.,**
513 **Laczko, E., McCaracken, D., Matouch, S., Niemela, J. & Richards, C.** 2005. Towards
514 sustainable land use: identifying and managing the conflicts between human activities and
515 biodiversity conservation in Europe. *Biodivers. Conserv.* **14**: 1641–1661.

Appendix I

Pressure codes and descriptions. Reproduced from Ruddock et al. (2016). Adapted from the EU Birds Directive reporting matrix (http://cdr.eionet.europa.eu/help/birds_art12).

Code	Description of pressure
A1	Modification of cultivation practices
A2	Agricultural intensification
A3	Mowing / cutting of grassland
A4	Abandonment / lack of mowing
A5	Intensive grazing
A6	Non-intensive grazing
A7	Abandonment of pastoral systems, lack of grazing
A8	Fertilisation (agricultural)
A9	Removal of hedges and copses or scrub
B1	Forest planting on open ground (increase in forest area, planting e.g. on grassland, heathland)
B2	Forest and plantation management & use
B3	Forest replanting (i.e. replanting on forest ground after clear-cutting)
B4	Forest clearance (clear-cutting, removal of all trees)
B5	Thinning of tree layer
B6	Fertilisation (forestry)
B7	Other forest activities (e.g. erosion due to forest clearing, fragmentation)
C1	Hand cutting of peat
C2	Mechanical removal of peat
C3	Wind energy production
D1	Paths, tracks, cycling tracks (includes non-paved forest roads)
D2	Roads, motorways (all paved/ tarred roads)
D3	Utility and service lines (e.g. power-lines, pipelines)
D4	Aircrafts or flightpaths
D5	Improved access to site
E1	Urbanisation, residential and commercial development
E2	Dispersed habitation (i.e. little or no human disturbance)
F1	Nest destruction
F2	Illegal killing (e.g. shooting, trapping, poisoning)
G1	Human intrusions and disturbances
G2	Outdoor sports and leisure activities, recreational activities
G3	Walking, horse-riding and non-motorised vehicles
G4	Motorised vehicles
G5	Off-road motorised driving
G6	Other outdoor sports and leisure activities
G7	Military manoeuvres
H1	Pollution (e.g. water pollution, fly-tipping)
J1	Natural fires
J2	Controlled burning (e.g. strip burning for grouse management)

- J3 Uncontrolled burning (e.g. widespread unmanaged or malicious burning)
 - J4 Modification of water levels or waterbodies
 - J5 Reduction or loss of specific habitat features (e.g. removal of hedgerows, deep heather, scrub, walls, drains)
 - J6 Reduction of prey availability
 - J7 Anthropogenic reduction of habitat connectivity (i.e. fragmentation such as by removal of large areas of habitat or creation of barriers between habitats)
 - K1 Interspecific faunal relations - predation (by other birds e.g. crows)
 - K2 Interspecific faunal relations - predation (by mammals e.g. foxes)
 - X No pressures recorded
 - O Other pressures not listed above; noted pressures included bracken encroachment, cattle, drainage, helicopter training, quarrying and shooting.
-

Appendix II

Summary data for all survey squares where the total number of pressures was >0. SPA = location of survey site relative to Special Protection Area (SPA) boundaries: 1 = inside; 0 = outside. PI = standardised Pressure Index (see main text).

Site number	Total number of pressures	Total number of observer visits	Total number of Hen Harrier territories	SPA	PI
1	1	5	0	0	0.01
2	10	45	0	0	0.02
3	22	16	0	0	0.10
4	8	10	0	0	0.06
5	10	9	0	0	0.08
6	23	8	0	0	0.21
7	5	3	0	0	0.12
8	4	8	0	0	0.04
9	12	6	0	0	0.15
10	34	17	1	0	0.15
11	14	14	3	0	0.07
12	16	14	1	0	0.08
13	19	22	1	0	0.06
14	9	2	0	0	0.33
15	2	3	0	0	0.05
16	4	3	0	0	0.10
17	9	12	1	0	0.06
18	17	35	2	1	0.04
19	9	11	2	0	0.06
20	2	3	0	0	0.05
21	6	30	2	0	0.01
22	7	12	0	0	0.04
23	8	10	1	0	0.06
24	3	5	1	0	0.04
25	6	43	1	0	0.01
26	15	13	1	1	0.09
27	23	78	1	1	0.02
28	10	8	0	0	0.09
29	4	8	0	0	0.04
30	3	13	0	0	0.02
31	3	6	0	0	0.04
32	18	31	0	0	0.04
33	9	9	0	0	0.07
34	20	78	3	1	0.02
35	9	29	0	0	0.02
36	2	6	0	0	0.02
37	2	2	0	0	0.07

38	6	3	0	0	0.15
39	6	5	0	0	0.09
40	5	82	2	1	0.00
41	7	10	1	1	0.05
42	1	44	1	0	0.00
43	13	142	7	1	0.01
44	5	3	0	0	0.12
45	16	21	0	0	0.06
46	3	3	0	0	0.07
47	3	2	0	0	0.11
48	24	8	0	0	0.22
49	3	1	0	0	0.22
50	5	2	0	0	0.19
51	6	11	0	0	0.04
52	19	10	0	0	0.14
53	9	9	0	0	0.07
54	6	2	0	0	0.22
55	26	5	0	0	0.39
56	1	8	0	0	0.01
57	65	23	1	1	0.21
58	95	32	2	1	0.22
59	6	1	0	0	0.44
60	17	3	0	0	0.42
61	122	60	2	1	0.15
62	59	11	0	0	0.40
63	17	11	1	0	0.11
64	6	2	0	0	0.22
65	33	17	1	0	0.14
66	56	40	2	1	0.10
67	68	62	2	1	0.08
68	2	5	0	0	0.03
69	13	79	5	1	0.01
70	17	60	4	1	0.02
71	17	4	0	0	0.31
72	81	50	2	0	0.12
73	1	5	0	0	0.01
74	54	25	0	0	0.16
75	40	21	2	1	0.14
76	8	34	2	1	0.02
77	2	31	0	0	0.00
78	2	43	2	1	0.00
79	33	17	0	0	0.14
80	9	8	0	0	0.08
81	4	1	0	0	0.30

82	2	2	0	0	0.07
83	40	50	2	0	0.06
84	5	12	1	0	0.03
85	54	12	0	0	0.33
86	5	7	0	0	0.05
87	48	36	2	1	0.10
88	2	1	0	0	0.15
89	113	143	5	0	0.06
90	3	1	0	0	0.22
91	74	21	3	0	0.26
92	14	9	0	0	0.12
93	150	60	2	1	0.19
94	19	11	1	0	0.13
95	53	25	2	0	0.16
96	35	15	2	1	0.17
97	73	14	1	0	0.39
98	11	11	0	0	0.07
99	21	26	1	1	0.06
100	5	11	1	0	0.03
101	11	17	1	0	0.05
102	20	5	0	0	0.30
103	15	4	1	0	0.28
104	15	25	0	0	0.04
105	23	10	0	0	0.17
106	6	5	0	0	0.09
107	4	3	0	0	0.10
108	13	11	0	0	0.09
109	5	20	2	0	0.02
110	8	17	1	1	0.03
111	31	15	1	0	0.15
112	85	15	3	0	0.42
113	7	3	0	0	0.17
114	2	9	0	0	0.02
115	26	9	0	0	0.21
116	5	54	2	1	0.01
117	14	4	0	0	0.26
118	2	3	0	0	0.05
119	16	24	0	0	0.05
120	15	9	0	0	0.12
121	6	8	0	0	0.06
122	9	11	0	0	0.06
123	6	8	0	0	0.06
124	4	6	0	0	0.05
125	11	4	0	0	0.20

126	14	7	0	0	0.15
127	8	9	0	0	0.07
128	8	4	0	0	0.15
129	27	19	0	0	0.11
130	10	16	0	0	0.05
131	10	16	0	0	0.05
132	4	2	0	0	0.15
133	13	10	0	0	0.10
134	54	4	0	0	1.00
135	2	1	0	0	0.15
136	14	12	1	1	0.09
137	9	13	0	0	0.05
138	18	10	0	0	0.13
139	6	3	0	0	0.15
140	86	79	1	0	0.08
141	9	8	1	0	0.08
142	23	21	3	0	0.08
143	1	3	0	0	0.02
144	100	23	1	0	0.32
145	1	3	0	0	0.02
146	1	16	0	0	0.00

Appendix III

Linear mixed-effects model results for pressures – expressed as Principal Components (PC) - associated with confirmed Hen Harrier territories (present/absent). Factors retained in the top subset of n models ($< \Delta 2$ AIC) are highlighted. Constituent pressures along with pressure codes and associated loadings (coefficients; in parentheses) are given. Pressure codes are taken and descriptions are abbreviated from those given in Ruddock et al. (2016; see Appendix I). Regression coefficients ($\beta \pm SE$) and significance of contributory PCs are given, where * = $p < 0.05$, ** = $p < 0.01$, and *** = $p < 0.001$. For constituent pressures in PC1-3, see Table 2 in the main text.

Principal Component (% variance explained)	Pressure	β	$\pm SE$	t	
PC1		0.025	0.007	3.52	**
PC2		-0.046	0.009	-5.05	***
PC3		0.044	0.011	-3.91	**
PC4 (5%)	Non-intensive grazing (A6; 0.51)	0.010	0.013	0.78	
	Agricultural fertilisation (A8; 0.32)				
	Urbanisation, residential and commercial development (E1; 0.50)				
PC5 (5%)	Hand cutting of peat (C1; -0.40)	-0.004	0.014	-0.27	
	Aircrafts or flightpaths (D4; -0.30)				
	Reduction of prey availability (J6; -0.52)				
	Other pressures not listed (O; -0.38)				
PC6 (4%)	Agricultural intensification (A2; 0.27)	-0.005	0.014	-0.38	
	Hand cutting of peat (C1; -0.25)				
	Roads, motorways (D2; 0.34)				
	Dispersed habitation, i.e. little or no human disturbance (E2; 0.37)				
	Off-road motorised driving (G5; -0.38)				
	Other outdoor sports and leisure activities (G6; 0.29)				
PC7 (4%)	Hand cutting of peat (C1; 0.29)	0.006	0.015	0.41	
	Dispersed habitation, i.e. little or no human disturbance (E2; 0.28)				
	Pollution (H1; 0.26)				
	Modification of water levels or waterbodies (J4; 0.27)				

Reduction of prey availability (J6; -0.27)

Other pressures not listed (O; -0.32)

Table 1. General Linear Model (GLM) results for regional differences in pressures on Hen Harrier breeding habitat - expressed as a Pressure Index (PI; log transformed). s = Special Protection Areas (SPA; inside/outside); r = confirmed territories (present/absent). Regression coefficients ($\beta \pm \text{SE}$) and significance of contributory variables are given, where $*$ = $p < 0.05$.

Model	Variable	t	$\beta (\pm \text{SE})$
PI $\sim r$	r	0.038	-0.39 \pm 0.19*
PI $\sim s$	s	0.028	-0.44 \pm 0.20*
PI $\sim r + s + r*s$	s	-0.515	-0.14 \pm 0.28
	r	-0.683	-0.16 \pm 0.23
	$s*r$	-1.128	-0.46 \pm 0.40

Table 2. Linear mixed-effects model results for pressures – expressed as Principal Components (PC) - associated with confirmed Hen Harrier territories (present/absent). Models were evaluated according to their Akaike’s Information Criterion (AIC) value. Factors retained in the top subset of n models ($< \Delta 2$ AIC) are highlighted. Constituent pressures along with pressure codes and associated loadings (coefficients; in parentheses) are given. Pressure codes are taken and descriptions are abbreviated from those given in Ruddock et al. (2016; see **Appendix I**). Regression coefficients ($\beta \pm SE$) and significance of contributory PCs are given, where * = $p < 0.05$, ** = $p < 0.01$, and *** = $p < 0.001$. For constituent pressures in PC4-7, see Appendix III.

Principal Component (% variance explained)	Pressure	β	$\pm SE$	t	
PC1 (21%)	Abandoned pastoral systems (A7; 0.29)	0.025	0.007	3.52	**
	Removal of hedges, copse and scrub (A9; 0.29)				
	Forest replanting (B3; 0.27)				
	Nest destruction (F1; 0.29)				
	Controlled burning (J2; 0.27)				
	Predation by birds (K1; 0.31)				
	Predation by mammals (K2; 0.26)				
PC2 (10%)	Forest management and use (B2; -0.28)	-0.046	0.009	-5.05	***
	Forest clearance (B4; -0.30)				
	Thinning of tree layer (B5; -0.26)				
	Paths, tracks, forest roads (D1; -0.35)				
	Roads, motorways (D2; -0.33)				
	Natural Fires (J1; -0.30)				
PC3 (7%)	Dispersed habitation (E2; -0.32)	0.044	0.011	-3.91	**
	Outdoor sport, leisure and recreation (G2; -0.35)				
	Walking, horse-riding, cycling (G3; -0.42)				
	Motorised vehicles (G4; -0.29)				
	Off-road driving (G5; -0.32)				
	Other outdoor sports and leisure (G6; -0.30)				

PC4	0.010	0.013	0.78
PC5	-0.004	0.014	-0.27
PC6	-0.005	0.014	-0.38
PC7	0.006	0.015	0.41

Figure 1. Maps of Ireland showing (a) survey squares and Special Protection Areas (SPAs, as indicated by grey polygons), (b) the total number of confirmed Hen Harrier territories per square, and (c) pressure indices derived from cumulative observations of pressures within each survey square. See Appendix II for square-specific data.

Figure 2. Relationships between pressures associated with potentially suitable breeding habitat for Hen Harriers. Pressure codes are taken and descriptions abbreviated from those given in Ruddock et al. (2016). Dashed grey rectangles indicate outermost clusters identified via the silhouette method and multiscale bootstrapping (10,000 iterations; Approximately Unbiased [AU] $p \leq 0.05$). \bigcirc = clusters supported at AU $p \leq 0.05$. For detailed pressure definitions, see **Appendix I**.



