

Title	Confidence of Irish dermatologists in caring for patients with skin of colour		
Authors	O'Connor, Cathal;Gallagher, C.;Bourke, John;Murphy, Michael		
Publication date	2021-08-16		
Original Citation	O'Connor, C., Gallagher, C., Bourke, J. and Murphy, M. (2021) 'Confidence of Irish dermatologists in caring for patients with skin of colour' [Letter to the Editor] Clinical and Experimental Dermatology. doi: 10.1111/ced.14897		
Type of publication	Other		
Link to publisher's version	10.1111/ced.14897		
Rights	© 2021, British Association of Dermatologists. Published by John Wiley & Sons, Inc. This is the peer reviewed version of the following item: O'Connor, C., Gallagher, C., Bourke, J. and Murphy, M. (2021) 'Confidence of Irish dermatologists in caring for patients with skin of colour' [Letter to the Editor] Clinical and Experimental Dermatology, doi: 10.1111/ced.14897, which has been published in final form at https://doi.org/10.1111/ced.14897. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.		
Download date	2025-06-01 19:37:48		
Item downloaded from	https://hdl.handle.net/10468/11823		





Confidence of Irish dermatologists in caring for patients with skin of colour

C. O'Connor, ^{1,2} C. Gallagher, ¹ J. Bourke, ^{1,2} M. Murphy^{1,2}

1 Department of Dermatology, South Infirmary Victoria University Hospital, Cork, Ireland

2 Department of Medicine, University College Cork, College Road, Cork, Ireland

Corresponding author: Dr Cathal O'Connor

Department of Dermatology, South Infirmary Victoria University Hospital, Cork, Ireland

Email: drcathaloconnor@gmail.com

Key words: skin of colour; ethnic dermatology; pigmented skin; dermatology training; medical education

Funding sources: none

Conflicts of interest: none

The first dermatology textbooks and atlases appeared in print in Western Europe at the turn of the 19th century, creating a publication bias towards dermatoses in lighter skin tones. Skin of colour (SOC), a term originating in North America, describes a genetically and culturally heterogenous group of people whose skin is more darkly pigmented than that of white Europeans. According to the 2016 Irish census, 7% of Irish people have SOC. ¹ The aim of this study was to assess Irish dermatologists' confidence with dermatoses in patients with SOC, and to identify opportunities to enhance training in dermatology in patients with SOC.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the <u>Version of Record</u>. Please cite this article as <u>doi:</u> 10.1111/CED.14897

This article is protected by copyright. All rights reserved

An online survey was distributed to all members of the Irish Association of Dermatology (IAD) by email. Questions related to demographic details, training in dermatology of SOC, and confidence with dermatology of SOC.

Of 140 members of the IAD, 56 members responded to the survey, equating to a response rate of 40%. Most respondents were female (87.5%) and white (98.2%). Registrars made up 51.8% of respondents, consultants 42.9%, and senior house officers 5.4%. Most (87.9%) respondents had predominantly trained on the island of Ireland (Table 1).

Only 23.2% of members reported receiving formal training in dermatology in SOC (Figure 1A). Half (50%) of members were 'not confident' or 'not at all confident' in diagnosing skin conditions (Figure 1B) and one-third (33.9%) were 'not confident' or 'not at all confident' in managing skin conditions (Figure 1C) in patients with SOC. Lack of exposure was cited as the biggest challenge to looking after patients with SOC by 60.7% of participants (Figure 1D), which was unsurprising as 67.9% of respondents estimated that <5% of their patients had SOC. Specific conditions that were noted to be difficult to diagnose or manage included (i) inflammatory dermatoses (n=29) such as atopic dermatitis and lupus, (ii) pigmentary disorders (n=16) such as melanocytic naevi and vitiligo, and (iii) keloid scarring (n=4). Suggested solutions to improving knowledge included dedicated study days, online resources, cultural training, and dedicated teaching clinics.

This study shows that Irish dermatologists have low confidence in diagnosing and managing conditions in people with SOC. This is consistent with a previous study that showed that only 56% of Australian dermatologists were confident in diagnosing and 75% confident in managing common dermatoses in patients with SOC. ² This discrepancy in confidence between skin types is important as patients with more darkly pigmented skin are more likely to have negative outcomes from dermatologic disease, including melanoma. 3 It has recently been shown that white skin is over-represented in images of Covid-19-associated-eruptions. 4 With growing awareness of racial equity in medicine, 5 efforts are being made to increase the availability of clinical images highlighting the appearance of dermatoses in SOC. Resources to improve education in SOC include the Skin Diversity Subcommittee of the British Association of Dermatologists, Skin Deep (https://dftbskindeep.com/), and the Skin of Colour Society (https://skinofcolorsociety.org/).

Ref

Irish dermatologists have low confidence with skin pathology in SOC, and specific training could reduce this disparity.

References

- 1. https://www.cso.ie/en/releasesandpublications/ep/p-cp8iter/p8iter/p8e/ (Accessed 23rd July 2021)
- 2. Rodrigues MA, Ross AL, Gilmore S, Daniel BS. Australian dermatologists' perspective on skin of colour: Results of a national survey. Australas J Dermatol. 2018;59(1):e23-e30. doi:10.1111/ajd.12556
- 3. Brady J, Kashlan R, Ruterbusch J, Farshchian M, Moossavi M. Racial Disparities in Patients with Melanoma: A Multivariate Survival Analysis. Clin Cosmet Investig Dermatol. 2021;14:547-550. Published 2021 May 24. doi:10.2147/CCID.S311694
- 4. Lester JC, Jia JL, Zhang L, Okoye GA, Linos E. Absence of images of skin of colour in publications of COVID-19 skin manifestations. Br J Dermatol. 2020;183(3):593-595. doi:10.1111/bjd.19258
- 5. Smith RJ, Oliver BU. Advocating for Black Lives—A Call to Dermatologists to Dismantle Institutionalized Racism and Address Racial Health Inequities. JAMA Dermatol. 2021;157(2):155–156. doi:10.1001/jamadermatol.2020.4392

Table

		n(%)
Gender	Male	7 (12.5%)
	Female	49 (87.5%)
Age	<30 years	11 (19.6%)
	30-45 years	28 (50%)
	>45 years	17 (30.4%)
Ethnicity	White Irish/British/European	55 (98.2%)
	Other*	1 (1.8%)
Grade	Consultant	24 (42.9%)
	Registrar	29 (51.8%)
	Senior House Officer	3 (5.4%)
Dermatology training scheme locations †	Republic of Ireland	43 (73.6%)
	Northern Ireland	8 (14.3)
	England (London)	3 (5.4%)
	England (outside London)	7 (12.5%)
	USA	2 (3.6%)
	Australia	1 (1.8%)
Estimated patients with SOC in clinics	<5%	37 (66.1%)
	5-10%	15 (26.8%)
SOC III CIIIIICS	>10%	4 (7.1%)

Table 1. Respondent characteristics.

Figure Legends


Figure 1A. Assessment of previous formal training in dermatology in skin of colour (n=56).

^{*}Not specified for confidentiality reasons †More than one may apply


Figure 1B. Assessment of confidence in diagnosing skin conditions in patients with skin of colour (n=56).

Figure 1C. Assessment of confidence in managing skin conditions in patients with skin of colour (n=56).


Figure 1D. Respondent rankings of challenges to care of patients with skin of colour (n=56).




ced_14897_f1a.jpg



ced_14897_f1b.jpg



ced_14897_f1c.jpg



ced_14897_f1d.jpg