

Title	Higher prevalence of unrecognized kidney disease at high altitude
Authors	Hurtado-Arestegui, Abdias;Plata-Cornejo, Raul;Cornejo, Arturo;Mas, Guiliana;Carbajal, Luz;Sharma, Shailendra;Swenson, Erik R.;Johnson, Richard J.;Pando, Jackelina
Publication date	2017-11-08
Original Citation	Hurtado-Arestegui, A., Plata-Cornejo, R., Cornejo, A., Mas, G., Carbajal, L., Sharma, S., Swenson, E. R., Johnson, R. J. and Pando, J. (2017) 'Higher prevalence of unrecognized kidney disease at high altitude', Journal of Nephrology, 31, pp. 263-269. doi:10.1007/s40620-017-0456-0
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1007/s40620-017-0456-0
Rights	© 2017, Italian Society of Nephrology. Published by Springer International Publishing AG. This is a post-peer-review, pre- copyedit version of an article published in Journal of Nephrology. The final authenticated version is available online at: http:// dx.doi.org/10.1007/s40620-017-0456-0
Download date	2024-09-26 10:44:23
Item downloaded from	https://hdl.handle.net/10468/5257



University College Cork, Ireland Coláiste na hOllscoile Corcaigh

# Higher Prevalence of Unrecognized Kidney Disease at High Altitude

Abdias Hurtado-Arestegui MD<sup>1, 2</sup>, Raul Plata-Cornejo MD<sup>3</sup>, Arturo Cornejo MD<sup>3</sup>, Guiliana Mas MD<sup>2</sup>, Luz Carbajal PhD<sup>2</sup>, Shailendra Sharma MD<sup>4</sup>, Erik R. Swenson MD<sup>5</sup>, Richard J. Johnson MD<sup>6</sup>, and Jackelina Pando MD<sup>7</sup>

<sup>1</sup>Division of Nephrology, Hospital Arzobispo Loayza,

<sup>2</sup> Cayetano Heredia University, Lima, Peru

<sup>3</sup>Instituto de Nefrología, La Paz, Bolivia

<sup>4</sup>Division of Renal Disease and Hypertension. The George Washington University, Washington DC, USA.

<sup>5</sup> Medical Service, Division of Pulmonary, Critical Care and Sleep Medicine, VA Puget Sound Health Care System, University of Washington, Seattle WA, USA

<sup>6</sup> Division of Renal Diseases and Hypertension, University of Colorado, Denver, CO, USA

<sup>7</sup> Department of Paediatrics and Child Health, University College Cork, Ireland

Informative title: risk for kidney disease at two different altitudes

Corresponding author: Abdias Hurtado-Arestegui

Address: Av. Tejada 215, Dpto: 202, Miraflores, Lima - Perú, Postal Code: Lima 18

Phone: 51 999858493 /511-3309361, Fax: 511-330936, email: ahurtadoa@hotmail.com

ORCID: 0000-0001-8850-7903

**Funding:** This study was partially supported by a grant N° 03-018 from the International Society of Nephrology (ISN) Global Outreach Research & Prevention Committee".

Conflict of Interest: The authors declare that they have no conflict of interest.

# Acknowledgements

We thank to Ronald Guillen MD, Marco Antonio Pariente MD, Claudia Ferrer MD Griselda Calisaya from Instituto de Nefrología, La Paz - Bolivia. Raquel Cancino MD, Luis Gonzales MD, Enrique Hernandez MD, Ivan Seminario MD, Jose Luis León MD from Hospital Nacional Arzobispo Loayza, Lima – Peru for their contribution for this work.

This study was partially supported by a grant N° 03-018 from the International Society of Nephrology (ISN) Global Outreach Research & Prevention Committee".

# Higher Prevalence of Unrecognized Kidney Disease at High Altitude

#### Abstract

**Background** High altitude renal syndrome has been described in populations with excessive erythrocytosis. We evaluated whether high altitude (HA) dwellers might be at increased risk for kidney disease.

**Methods:** We performed a cross-sectional study to investigate differences in prevalence of kidney function and metabolic syndrome in healthy subjects living at HA and sea level (SL) without any known history of hypertension, diabetes or chronic kidney disease.

**Results.** We examined 293 subjects, age 40 to 60 years, 125 from SL (154 m) and 168 HA (3640 m). HA subjects had higher serum creatinine, lower estimated glomerular function rate (eGFR) (69.5  $\pm$  15.2 vs 102.1  $\pm$  17.8 ml/min/1.73 m<sup>2</sup>, p <0.0001), with more proteinuria and higher hemoglobin concentrations compared to SL subjects. HA subjects had a lower prevalence of metabolic. Hemoglobin concentrations correlated inversely with eGFR in women and men (p = 0.001 and p= 0.03, respectively) among subjects living at HA. Using logistic regression analysis to compare subjects with eGFR < 90 versus > 90 ml/min/1.73m<sup>2</sup>, a lower eGFR was associated with gender (women) (OR adjusted: 5.65; 95% CI : 2.43-13.13, p = 0.001), high altitude (OR adjusted: 14.78, 95% CI : 6.46-33.79; p = 0.001), hemoglobin (OR adjusted: 1.68; 95% CI : 1.16-2.43; p = 0.001) and uric acid (OR adjusted: 1.93; 95% CI : 1.36-2.72; p = 0.001).

**Conclusions**: Subjects living at high altitude who are considered healthy have worse kidney function, a higher prevalence of proteinuria and a lower prevalence of metabolic syndrome compared to people living at SL.

Keywords: cardiovascular risk, high altitude, kidney function, proteinuria

#### Introduction

Chronic kidney disease (CKD) has many known causes, including diabetes, hypertension, glomerulonephritis and polycystic kidney disease. However, CKD may be caused by other mechanisms. We have hypothesized that living chronically at high altitude (HA) may be a risk factor for CKD related to the chronic exposure to hypoxia due to low ambient oxygen tension [1,2].

Recently we have reported that subjects living at HA are at risk for high altitude renal syndrome "HARS", a condition characterized by erythrocytosis, microalbuminuria, hypertension and hyperuricemia [3,4]. This is especially observed in subjects suffering from chronic mountain sickness (CMS), a form of altitude maladaptation characterized by excessive erythrocytosis (hematocrit > 65 %) and greater arterial hypoxemia (SaO<sub>2</sub>% < 85) [5]. While the pathogenesis of HARS is not fully understood, it may relate to glomerular hyperfiltration associated with polycythemia [6], or the effects of hyperuricemia on glomerular hemodynamics and tubular function [7,8].

Given these issues, we compared two similar ethnic populations: one living at sea level (Lima, Peru, altitude 154 m) versus one at high altitude (La Paz, Bolivia, altitude 3,640-4,500 m). We specifically wanted to compare 'healthy' populations who had no known hypertension, diabetes or CKD. Our hypothesis was that, even within an apparently healthy population and those without CMS, subjects chronically living at high altitude would show more evidence for kidney dysfunction.

# **Material and Methods**

# Study Design.

This was a cross-sectional study that evaluated subjects from La Paz, a city in Bolivia located at an altitude between 3,640 - 4,500 meters and Lima, a city in Peru, which is located at 154 meters above sea level. Both cities are capitals of developing countries with similar economic conditions and mixed European - Andean heritage.

We performed screening campaigns in the community of both cities, and the subjects were selected by convenience sampling if they fulfilled criteria of selection in the first interview. After the results of laboratory testing were obtained, additional subjects could be excluded based on fasting blood glucose levels of >125 mg/dl (n=19) and/or significantly elevated serum creatinine (> 1.4 mg/dl, n=8). The remaining 293 participants were aged 40 to 60 years, with no prior known history of hypertension, diabetes, chronic kidney disease, CMS or other chronic medical conditions. Informed consent was obtained from all individual participants included in the study. The exclusion criteria included a history of any systemic disease, long-term use of medications and incomplete information of the subject's condition or laboratory data. The protocol followed the principles of the Declaration of Helsinki. The study was approved by the ethics committee of the Hospital Nacional Arzobispo Loayza in Lima, Peru.

#### Evaluation

A detailed history was obtained from all including past medical history, family history of diabetes and kidney disease, as well as life style habits (alcohol and tobacco use and physical activity). We used the International Society of Nephrology form: "Program for Detection and Management of CKD, Hypertension, Diabetes and Cardiovascular Disease in Developing Countries", which was adapted to the needs of the study [9]. Physical examination included weight, height (after removal of shoes and heavy clothing), waist circumference (measured at the midpoint between the bottom edge of the last rib and the iliac crest taking as a reference the mid-axillary line and taken at the end of expiration) and hip circumference (measured using the maximum circumference at the level of the buttocks). Body mass index (BMI) and waist-hip ratio (WHR) were calculated. Blood pressure was measured following recommended procedures after sitting for 15 minutes [10], after which three readings were recorded at intervals of three minutes using a calibrated automatic blood pressure monitor OMROM, model HEM-705CP. Venous blood was obtained after 12 hours of fasting for total cholesterol, low density lipoprotein (LDL), high density lipoprotein (HDL), triglycerides, glucose, uric acid, hemoglobin, hematocrit, and creatinine with enzymatic methods. The data were processed in a computer model: STAT FAX 1935 PLUS, Roche Cobas C501. All coefficients of variation for original measurements were < 10%. Morning spot urine samples were evaluated for proteinuria and hematuria with a Dipstik test (Roche Diagnostics, Mannheim, Germany).

**Definitions:** 

Overweight was defined as BMI between 25-30 kg/m<sup>2</sup> and obesity as BMI > 30 kg/m<sup>2</sup>. Abdominal obesity was defined as WHR > 0.9 in men and > 0.85 in women. Estimated glomerular filtration rate (eGFR) was calculated using the Modification of Diet in Renal Disease (MDRD) four variables equation. Kidney Disease (KD) stages were classified according to Kidney Disease Outcomes Quality Initiative (KDOQI) [11]. Hematuria and proteinuria were defined as 1+ or greater on a dipstick test. The diagnosis of metabolic syndrome was based on the presence of 3 of the 5 following criteria: waist circumference > 94 cm in men and > 88 cm in women, triglyceride  $\geq$  150 mg/dl, high density lipoprotein HDL < 40 mg/dl in men and < 50 mg/dl in women, systolic blood pressure  $\geq$  130 mmHg and/or DBP  $\geq$  85 mmHg, fasting blood glucose  $\geq$  100 mg/dl [12]. The cardiovascular risk was calculated using the Framingham risk score [13]. This score addresses the following variables: age, sex, systolic blood pressure treated (medication and lifestyle changes) and untreated, BMI, smoking and diabetes. The consumption of alcohol and tobacco, and physical activity was recorded as present or absent.

**Sample Size.** We calculated a necessary sample size of 318 subjects to have 80 % power and a confidence interval of 95 % based on chronic KD prevalence of 10 % SL [14] and 19% for HA [15]. The sample calculation was obtained from the statistical package Stata v12 using the formula for comparison of two separate proportions.

**Statistical analysis**. An exploratory data analysis was done by obtaining measurements of central tendency such as: mean, standard deviation, median, minimum and maximum values for quantitative variables, and frequencies and percentages for qualitative variables. Before comparing means or medians, the distribution of the quantitative variables was evaluated using the Shapiro-Wilk test and the homogeneity of variance with Bartlett's test. To assess association between two qualitative variables in the bivariate analysis, Chi Square test or Fisher exact test were used respectively depending on the expected frequency number of more than 5 or not. Student's t-test was used for comparing two arithmetic means (when the distribution of the variable had a normal distribution), two-sample Wilcoxon rank-sum (Mann-Whitney) test was used for independent samples when the distribution of the variable was not normal. Multiple logistic regression was used to assess factors associated with renal disease (defined as CKD stage II, or an eGFR < 90 ml/min/1.73 m<sup>2</sup>). The level of statistical significance considered was 0.05, for a twotailed test. Statistical analysis was performed using the statistical package Stata v12.

#### Results

**Baseline characteristics of study subjects.** Of the 320 subjects selected, we excluded subjects with fasting glucose concentrations greater than 126 mg/dl who did not know whether they were diabetic (n = 19), and those with creatinine greater than 1.4 mg/dl (n = 8). Thus, 293 subjects were included in the final analysis (representing 92 % of the sample). Of these 125 were living at SL and 168 at HA. There were 145 men (65 SL, 80 HA) and 148 women (60 SL, 88 HA).

**Baseline characteristics of subjects living at SL and HA**. There were no significant differences in age or sex between the two groups. Compared to SL subjects, kidney function in the HA subjects was characterized by higher serum creatinine  $(1.05 \pm 0.14 \text{ vs } 0.77 \pm 0.16 \text{ mg/dl}, \text{ p} < 0.0001)$ , lower eGFR (69.5 ± 15.2 vs 102.2 ± 17.8 ml/min/1.73m<sup>2</sup>, p < 0.0001), higher percentage of patients with eGFR < 60 ml/min/1.73 m<sup>2</sup> (28.0 % vs 0 %, p < 0.001) and proteinuria (10.7 % vs 3.2%, p = 0.03).

HA subjects had higher hemoglobin concentrations ( $15.4 \pm 1.4$  vs  $13.9 \pm 1.2$  g/dl, p < 0.0001), HDL level ( $48.9 \pm 12.0$  vs  $44.9 \pm 10.9$  mg/dl, p < 0.002) and fasting triglycerides level ( $159.9 \pm 99.6$  vs  $123.4 \pm 55.7$  mg/dl, p < 0.001). In contrast, high altitude dwellers had less alcohol intake (40.5 % vs. 56.8 %, p < 0.05), lower systolic blood pressure ( $114.3 \pm 15.9$  vs  $129.7 \pm 17.2$  mmHg, p < 0.0001) and diastolic blood pressure ( $71.5 \pm 8.5$  vs  $76.3 \pm 10.0$  mm Hg, p < 0.0001), as well as lower weight ( $69.2 \pm 10.9$  vs  $73.2 \pm 14.2$ , p = 0.03), BMI ( $26.2 \pm 3.6$  vs  $28.9 \pm 4.3$  kg/m<sup>2</sup>, p < 0.0001), waist-hip ratio ( $0.96 \pm 0.04$  vs  $0.92 \pm 0.07$ , p < 0.0001), fasting serum glucose level ( $74.2 \pm 8.4$  vs  $93.1 \pm 10.9$  mg/dl, p < 0.0001), LDL level ( $126.2 \pm 41.3$  vs  $144.9 \pm 33.4$ , p < 0.0001), prevalence of metabolic syndrome (27.4 % vs 49.6 %, p < 0.0001) and Framingham risk score ( $0.91 \pm 1.15$  vs  $2.13 \pm 2.65$ , p < 0.001 (**Table 1**).

Values of hemoglobin, kidney function, metabolic syndrome and Framingham risk score in men and women living at SL and HA are shown in **Table 2**.

Multivariable logistic regression was used to compare subjects with evidence for stage 2 CKD or worse (eGFR 90 ml/min/ $1.73m^2$ ) versus those with normal renal function (defined as eGFR > 90 ml/min/ $1.73m^2$ ). After adjusting for age, sex, place of residence, hemoglobin, triglycerides, metabolic syndrome, uric acid, a lower eGFR was associated with being a woman (OR adjusted: 5.65; 95% CI: 2.43-13.13, p = 0.001), living at high altitude (OR adjusted: 14.78;

95% CI : 6.46-33.79; p = 0.001), having a higher hemoglobin (OR adjusted: 1.68; 95% CI: 1.16-2.43; p = 0.001) and for each unit of increase in uric acid (OR adjusted: 1.93; 95% CI: 1.36-2.72; p = 0.001), Table 3

#### Relationship between eGFR and hemoglobin in women and men.

In HA women and men, hemoglobin correlated inversely with eGFR (p = 0.001 and p = 0.03, respectively; r = -0.2, p = 0.01), but no significant correlation was observed in women or men living at SL (**Figures 1 and 2**).

# Discussion

In this cross-sectional study of 293 healthy ethnically similar participants from two cities of different altitude in South America, we found that individuals living at high altitude had significantly worse renal function than those living near sea level, amounting to a mean difference of eGFR of 30 percent. The worse kidney function and greater proteinuria observed at higher altitude occurred despite similar age and lower prevalence of metabolic syndrome compared to people living near sea level.

The prevalence of eGFR  $< 60 \text{ ml/min/1.73 m}^2$  and proteinuria was 28.0 % and 10.7 % respectively, among subjects living at HA and this was observed after excluding subjects with hypertension, diabetes, or known CKD. This prevalence is higher than values reported in Tibet, which are 2.1% for CKD and 2.4% for proteinuria [15]. One possible explanation is that Tibetans have been living at high altitude for many more generations than any groups in the Andes [16] and are therefore better adapted to the high altitude hypoxic environment, as evidenced by their higher arterial oxygen saturation and lesser hemoglobin concentration.

Potential mechanisms for the worse renal function at HA could be chronic hypoxia, leading to erythrocytosis. Increased red cell volume leads to reduced renal plasma flow (RPF) which eventually results in higher filtration fraction (FF) as glomerular filtration rate is preserved [6,17]. These characteristics resemble a model of chronic hyperfiltration, with its known consequences of glomerular damage and proteinuria [18]. The population at HA in our study did not have excessive erythrocytosis typical of CMS (hematocrit > 65%), nonetheless increased hemoglobin concentration was associated with worse kidney function.

The renal medulla is hypoxic with a range of PO<sub>2</sub> between 10-20 mmHg [2,19] and is highly susceptible to injury by hypoxia [20]. Medullary hypoxia is likely more severe among high altitude dwellers given their lower arterial oxygen saturation and no off-setting higher renal blood flow (2), thus further increasing the risk of renal injury. Indeed, hypoxia is a well-known risk factor for progressive CKD in various experimental animal models [19]. Similarly, subjects with kidney disease also have higher serum uric acid concentrations, which are known to be elevated in hypoxic conditions. While uric acid can function as an antioxidant [21], it can also stimulate oxidative stress in a variety of cell types, and can also cause endothelial dysfunction, and systemic and glomerular hypertension [22] thereby predisposing to the risk of CKD. We also found that HA subjects tended to have higher serum uric acid concentrations, although this was not statistically significantly different compared to the SL group. This may be due to the higher frequency of the metabolic syndrome in the SL group, which is also associated with hyperuricemia.

We also found a lower frequency of metabolic syndrome at HA. While it may be argued that the higher frequency of metabolic syndrome among SL dwellers might reflect different dietary or economic factors, the HA subjects were also urban residents in a large city (La Paz) with a comparable socioeconomic environment. Other factors may also be important for example, physical activity was more common among men but not women at HA, and might explain differences in weight. The metabolic responses to chronic hypoxia may also have protective effects on the development of the metabolic syndrome. HA residents are known to have lower serum glucose concentrations than inhabitants at SL [23]. Consistent with this observation, a lower incidence of diabetes has been observed at HA [24]. A lower fasting serum glucose concentration was observed in our study and contributes to a lower incidence of the metabolic syndrome in the population at HA. Furthermore, living at HA is also associated with higher HDL concentrations [25]. Not all studies, however, have found that the metabolic syndrome is lower at HA. A prior study from Peru found no difference in the metabolic syndrome between high altitude and sea level dwellers (22.2 % vs. 16.9 % respectively; p = 0.28) [26]. In another study, the metabolic syndrome was extremely high (60 %) among individuals living at high altitude with excessive erythrocytosis (mean age 61 yr) compared to a prevalence of 35 % among high altitude residents without erythrocytosis (mean age 55 yr) [27]. Another study of the Peruvian population also reported that subjects living at higher altitude had a metabolically healthier profile than those living at sea level [28].

The Framingham risk score is a quantitative method that estimates the likelihood of coronary heart disease over a period of time, and has been validated in various ethnic groups [29]. The population at HA (both men and women) had a lower Framingham score compared to the population living at sea level. The lower score in HA residents was primarily due to the higher HDL concentrations, lower prevalence of smoking, and lower blood pressure. Among women, the variables of age, cholesterol, HDL and smoking were similar, so that the lower Framingham score at HA was primarily due to a lower blood pressure. A study in Tibet reported a lower Framingham risk score than what we found in our Andean population: 16. 3% for men and 0.6% for women [30].

Our study has several limitations. First, it was a study of healthy individuals, and we did exclude subjects with known hypertension, diabetes or CKD. Therefore, it is not a study of overall prevalence of CKD. However, the striking finding that eGFR was 30 percent lower in the presumably healthy population at HA remains remarkable. Second, we did not assess for microalbuminuria. We also excluded subjects with hypertension, which may be part of the high altitude renal phenotype. However, by excluding subjects with hypertension we could also remove the possibility that the kidney disease was secondary to hypertension alone. A final limitation is that we failed to gather information on dietary history and measured the laboratory values only once. Nevertheless, the strength of the study is the robust differences identified in prevalence of kidney disease and metabolic syndrome.

In conclusion, apparently healthy subjects living at high altitude had a remarkably higher frequency of kidney disease and proteinuria independent of a history of hypertension or diabetes. This provides some of the strongest evidence to date that long-term living at high altitude carries an increased risk for kidney disease. The characteristics of kidney disease observed at high altitude may be a new form of CKD that might be mediated by chronic hypoxia and its effects. Given that there are more than 140 million people worldwide living at high altitude, further studies to better investigate the etiology and treatment of high altitude CKD are critical.

# Ethical approval:

"All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards."

#### References

- Hurtado A, Escudero E, Pando J, Sharma S, Johnson RJ. Cardiovascular and renal effects of chronic exposure to high altitude. Nephrol Dial Transplant. 2012 Dec;27 Suppl 4:iv11-16.
- Luks AM, Johnson RJ, Swenson ER. Chronic kidney disease at high altitude. J Am Soc Nephrol JASN. 2008 Dec;19(12):2262–71.
- Jefferson JA, Escudero E, Hurtado M-E, Kelly JP, Swenson ER, Wener MH, et al. Hyperuricemia, hypertension, and proteinuria associated with high-altitude polycythemia. Am J Kidney Dis. 2002 Jun;39(6):1135–42.
- Arestegui AH, Fuquay R, Sirota J, Swenson ER, Schoene RB, Jefferson JA, et al. High altitude renal syndrome (HARS). J Am Soc Nephrol. 2011 Nov;22(11):1963–8.
- León-Velarde F, Maggiorini M, Reeves JT, Aldashev A, Asmus I, Bernardi L, et al. Consensus statement on chronic and subacute high altitude diseases. High Alt Med Biol. 2005;6(2):147–57.
- Lozano R, Monge C. Renal function in high-altitude natives and in natives with chronic mountain sickness. J Appl Physiol. 1965 Sep;20(5):1026–7.
- Kang D-H, Nakagawa T, Feng L, Watanabe S, Han L, Mazzali M, et al. A role for uric acid in the progression of renal disease. J Am Soc Nephrol. 2002 Dec;13(12):2888–97.
- Ryu E-S, Kim MJ, Shin H-S, Jang Y-H, Choi HS, Jo I, et al. Uric acid-induced phenotypic transition of renal tubular cells as a novel mechanism of chronic kidney disease. Am J Physiol Renal Physiol. 2013 Mar 1;304(5):F471-480.
- International Society of Nephrology. Program for detection and management of chronic kidney disease, hypertension, diabetes, and cardiovascular disease in developing countries (KHDC). [Internet]. [cited 2012 Oct 10]. Available from: http://www.isn-online.org/isn/education/guidelines/isn/full/ed\_051027\_1.html.
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. JAMA. 2003 May 21;289(19):2560–72.
- 11. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. Am J Kidney Dis. 2002 Feb;39(2 Suppl 1):S1-266.

- б
- 12. Alberti KGMM, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the metabolic syndrome: a joint interim statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. Circulation. 2009 Oct 20;120(16):1640–5.
- Rupert Payne. Risk calculator graphs based on Joint British Societies risk prediction charts [Internet]. 2010
   [cited 2014 Nov 6]. Available from: http://cvrisk.mvm.ed.ac.uk/calculator/calc.asp
- 14. Hallan SI, Coresh J, Astor BC, Asberg A, Powe NR, Romundstad S, et al. International comparison of the relationship of chronic kidney disease prevalence and ESRD risk. J Am Soc Nephrol. 2006 Aug;17(8):2275–84.
- 15. Chen W, Liu Q, Wang H, Chen W, Johnson RJ, Dong X, et al. Prevalence and risk factors of chronic kidney disease: a population study in the Tibetan population. Nephrol Dial Transplant. 2011 May;26(5):1592–9.
- Beall CM. Andean, Tibetan, and Ethiopian patterns of adaptation to high-altitude hypoxia. Integr Comp Biol. 2006 Feb;46(1):18–24.
- Swenson ER. The kidneys and fluid balance. In: High Altitude: Human Adaptation to Hypoxia. Eds. Swenson ER and Bartsch P. New York: Springer Publications; 2014. 217-236 p.
- Brenner BM. Hemodynamically mediated glomerular injury and the progressive nature of kidney disease. Kidney Int. 1983 Apr;23(4):647–55.
- Shoji K, Tanaka T, Nangaku M. Role of hypoxia in progressive chronic kidney disease and implications for therapy. Curr Opin Nephrol Hypertens. 2014 Mar;23(2):161–8.
- 20. Evans RG, O'Connor PM. Initiation and progression of chronic kidney disease: can we definitively test the chronic hypoxia hypothesis? Hypertension. 2013 Nov;62(5):827–8.
- 21. Ames BN, Cathcart R, Schwiers E, et al. Uric acid provides an antioxidant defense in humans against oxidantand radical-caused aging and cancer: a hypothesis. Proc Natl Acad Sci U S A 1981; 78: 6858-6862.
- 22. Johnson RJ, Nakagawa T, Jalal D, Sánchez-Lozada LG, Kang D-H, Ritz E. Uric acid and chronic kidney disease: which is chasing which? Nephrol Dial Transplant. 2013 Sep;28(9):2221–8.
- 23. Castillo O, Woolcott OO, Gonzales E, Tello V, Tello L, Villarreal C, et al. Residents at high altitude show a lower glucose profile than sea-level residents throughout 12-hour blood continuous monitoring. High Alt Med Biol. 2007;8(4):307–11.

- б
- 24. Santos JL, Pérez-Bravo F, Carrasco E, Calvillán M, Albala C. Low prevalence of type 2 diabetes despite a high average body mass index in the Aymara natives from Chile. Nutr. 2001 Apr;17(4):305–9.
- 25. Domínguez Coello S, Cabrera De León A, Bosa Ojeda F, Pérez Méndez LI, Díaz González L, Aguirre-Jaime AJ. High density lipoprotein cholesterol increases with living altitude. Int J Epidemiol. 2000 Feb;29(1):65–70.
- 26. Baracco R, Mohanna S, Seclén S. A comparison of the prevalence of metabolic syndrome and its components in high and low altitude populations in peru. Metab Syndr Relat Disord. 2007;5(1):55–62.
- Miele CH, Schwartz AR, Gilman RH, Pham L, Wise RA, Davila-Roman VG, et al. Increased Cardiometabolic Risk and Worsening Hypoxemia at High Altitude. High Alt Med Biol. 2016 Jun;17(2):93–100.
- 28. Benziger CP, Bernabé-Ortiz A, Gilman RH, Checkley W, Smeeth L, Málaga G, et al. Metabolic Abnormalities Are Common among South American Hispanics Subjects with Normal Weight or Excess Body Weight: The CRONICAS Cohort Study. PloS One. 2015;10(11):e0138968.
- D'Agostino RB, Grundy S, Sullivan LM, Wilson P, CHD Risk Prediction Group. Validation of the Framingham coronary heart disease prediction scores: results of a multiple ethnic groups investigation. JAMA. 2001 Jul 11;286(2):180–7.
- 30. Sherpa LY, Deji, Stigum H, Chongsuvivatwong V, Luobu O, Thelle DS, et al. Lipid profile and its association with risk factors for coronary heart disease in the highlanders of Lhasa, Tibet. High Alt Med Biol. 2011;12(1):57–63.

Variables	Sea level		High	<b>p</b> *	
	Mean ± SD	n=125) 95 %CI	(n Mean ± SD	95%CI	
Age (years) <sup>1</sup>	$50.79 \pm 6.86$	49.57 - 52.00	$50.58 \pm 5.79$	49.70 - 51.46	0.679
Sex (men / women) <sup>4,a</sup>	65/60	0.39 - 0.56	80/88	0.44 - 0.60	0.458
Alcohol <sup>5,a</sup>	71 (56.8%)	0.47 - 0.65	68 (40.5%)	0.32 - 0.47	0.0001*
Smoking <sup>5,a</sup>	21 (16.8%)	0.10 - 0.23	17 (10.1%)	0.55 - 0.14	0.114
Physical Activity <sup>5,a</sup>	51 (40.8%)	0.32 - 0.49	73 (43.4%)	0.35 - 0.51	0.649
Systolic blood pressure (mmHg) <sup>3</sup>	129.71 ±17.22	126.66 - 132.76	$114.34 \pm 15.90$	111.92 – 116.76	0.0001*
Diastolic blood pressure (mmHg) <sup>3</sup>	$76.37 \pm 10.04$	74.59 - 78.15	$71.51\pm8.50$	70.22 - 72.81	0.0001*
Pulse (beats per min) <sup>1</sup>	$69.60 \pm 10.20$	67.79 - 71.40	$73.51 \pm 3.85$	72.93 - 74.10	0.0001*
Weight (kg) <sup>3</sup>	73.16 ± 14.16	70.65 - 75.67	$69.24 \pm 10.95$	67.57 – 70.91	0.030
Height (cm) <sup>3</sup>	$158.7\pm9.57$	157.00 - 160.39	$162.48\pm6.70$	167.46 - 163.50	0.0001*
Body mass index(kg/m <sup>2</sup> ) <sup>3</sup>	$28.90 \pm 4.29$	28.14 - 28.66	$26.18\pm3.61$	25.63 - 26.74	0.0001*
Waist circumference (cm) <sup>2</sup>	$102.84 \pm 9.28$	101-20 - 104.49	$101.04 \pm 10.47$	99.44 - 102.63	0.283
Hip circumference (cm) <sup>2</sup>	$99.58 \pm 8.94$	98.00 - 101.16	93.90 ± 13.82	91.79 - 96.01	0.0001*
Waist-hip ratio <sup>2</sup>	$0.96\pm0.04$	0.96 - 0.97	$0.92\pm0.07$	0.91 - 0.93	0.0001*
Hemoglobin (g/dl) <sup>1</sup>	13.91 ± 1.22	13.69 - 14.12	15.43 ± 1.36	15.23 - 15.64	0.0001*
Glucose (mg/dl) <sup>1</sup>	93.07 ± 10.94	91.13 - 95.01	$74.23 \pm 8.36$	75.95 - 75.50	0.0001*
Creatinine (mg/dl) <sup>3</sup>	$0.77 \pm 0.16$	0.74 - 0.80	$1.05\pm0.14$	1.03 – 1.08	0.0001*
eGRF (ml/min/1.73m <sup>2</sup> ) <sup>1</sup>	$102.15 \pm 17.82$	98.99 - 105.31	$69.52 \pm 15.18$	67.21 - 71.84	0.0001*
eGRF level (ml/min/1.73m <sup>2</sup> ) <sup>4,a</sup> > 90 60 - 89 < 60 Urine	96 (69.23%) 29 (30.77%) 0 (0.0%)		25 (14.88%) 96 (57.14%) 47 (27.98%)		0.001*
Hemoglobin (positive) Proteinuria (positive)	12 (9.6%) 4 (3.2%)		16 (9.5%) 18 (10.7%)		0.88 0.03
Total cholesterol (mg/dl) <sup>1</sup>	$216.90\pm41.63$	209.53 - 224.27	$209.32\pm54.13$	201.08 - 217.57	0.08
Low density lipoprotein (mg/dl) <sup>1</sup>	$144.86\pm33.40$	138.94 – 150.77	$126.22 \pm 41.28$	119.93 - 132.50	0.0001*

# Table 1 Clinical and laboratory characteristics of people living at Sea level and High altitude

High density lipoprotein (mg/dl) <sup>1</sup>	$44.92 \pm 10.90$	42.99 - 46.85	48.92 ± 11.97	47.10 - 50.75	0.002*
Triglycerides (mg/dl) <sup>1</sup>	$123.42\pm55.68$	113.56 - 133.28	$159.89\pm99.58$	144.72 - 175.06	0.001*
Uric acid (mg/dl) <sup>1</sup>	$4.92 \pm 1.18$	4.71 - 5.13	$5.11 \pm 1.43$	4.89 - 5.33	0.379
Metabolic syndrome <sup>4,a</sup>	62 (49.6%)	0.40 - 0.58	46 (27.38%)	0.20 - 0.34	0.0001*
Framingham risk score <sup>1</sup>	$2.13\pm2.65$	1.66 – 2.60	$0.91 \pm 1.15$	0.73 – 1.08	0.001*

(Mean ± Standard Deviation)

(N° and %  $\pm$  Standard Error)  $^{\rm a}$ 

1 Mann–Whitney test

2 Student's t-test with unequal variances

3 Student's t-test with equal variances

4 Chi-square test

5 Fisher's exact test

P\* Contrast Bonferroni

	men			women			
Variables	Sea level n = 65	High altitude n = 80	р	Sea level n = 60	High altitude n = 88	р	
Hemoglobin (g/dl) <sup>1</sup>	$14.6\pm0.8$	$15.5\pm0.9$	0.001	$13.1\pm1.1$	$15.3 \pm 1.6$	0.001	
Creatinine (mg/dl) <sup>3</sup>	$0.88 \pm 0.14$	$1.05\pm0.14$	0.001	$0.64 \pm 0.08$	$1.05\pm0.14$	0.001	
eGRF (ml/min/1.73m <sup>2</sup> ) <sup>1</sup>	99.2 ± 18.8	80.4 ± 12.6	0.001	105.3 ±16.2	$59.7\pm9.6$	0.001	
eGRF level (ml/min/1.73m <sup>2</sup> ) <sup>4</sup> > 90 60 - 89 < 60	45 (69.2%) 20 (30.8%) 0 (0.0%)	24 (30.0%) 54 (67.5%) 2 (2.5%)	0.001	51 (85%) 9 (15%) 0 (0.0%)	1 (1.1%) 42 (47.7%) 45 (51.1%)	0.001	
Metabolic syndrome <sup>4</sup>	32 (49.2%)	25 (31.3%)	0.04	30 (50%)	21 (23.9%)	0.001	
Framingham risk score <sup>1</sup>	$3.22\pm3.06$	$1.49 \pm 1.37$	0.001	$0.95 \pm 1.36$	$0.38\pm0.50$	0.001	

# Table 2 Characteristics of hemoglobin, kidney function, metabolic syndrome andFramingham score in men and women living at SL and HA

(Mean  $\pm$  standard deviation; n° and %)

1 Mann–Whitney test

2 Student's t-test with unequal variances

3 Student's t-test with equal variances

4 Chi-square test

Variables	OR	95%CI OR	р	OR adjusted	95%CI adjusted	р
Age (years)	1.00	0.97-1.04	0.64	1.02	0.97-1.08	0.34
Sex: Female	1.58	0.99-2.53	0.05	5.65	2.43-13.13	0.001
High altitude	20.86	11.39-38.21	0.001	14.78	6.46-33.79	0.001
Hemoglobin (g/dl)	2.22	1.76-2.80	0.001	1.68	1.16-2.43	0.001
Triglycerides (mg/dl)	1.00	1.00-1.00	0.001	1.00	0.99-1.00	0.51
Metabolic syndrome	0.54	0.33-0.87	0.01	0.48	0.21-1.11	0.08
Uric acid (mg/dl)	1.46	1.20-1.79	0.01	1.93	1.36-2.72	0.001

Table 3: Unadjusted and Multivariable-adjusted Odds of eGFR < 90 ml/min/1.73m2

Adjusted for age, sex, high altitude, hemoglobin, triglycerides, metabolic syndrome, uric acid. P-values calculated using logistic regression. OR, odds ratio; CI, confidence interval



eGFR Stage (ml/min/1.73m<sup>2</sup>)

Figure 1: Estimated glomerular filtration rate versus hemoglobin level at different altitude in women.

Box-plot diagram of the relationship between estimated glomerular filtration rate (eGFR) and hemoglobin level in women, at high altitude hemoglobin increased significantly with decreasing eGFR levels (p = 0.001; r = -0.2, p = 0.01), but no in women living at sea level. Statistical: Kruskal-Wallis, Spearman.



Figure 2: Estimated glomerular filtration rate versus hemoglobin level at different altitude in men

Box-plot diagram of the relationship between estimated glomerular filtration rate (eGFR) and hemoglobin level in men. at high altitude hemoglobin increased significantly with decreasing eGFR levels (p = 0.03: r = -0.2, p = 0.01), but no in men living at sea level. Statistical: Kruskal-Wallis, Spearman.