

Title	Incorporating host-parasite biotic factors in species distribution models: Modelling the distribution of the Castor Bean tick, <i>Ixodes ricinus</i>
Authors	McDonough, Sinead;Holloway, Paul
Publication date	2020-11-01
Original Citation	McDonough, S. and Holloway, P. (2020) 'Incorporating host-parasite biotic factors in species distribution models: Modelling the distribution of the Castor Bean tick, <i>Ixodes ricinus</i> ', Irish Geography, 53(2), pp. 105-125. doi: 10.2014/igj.v53i2.1416
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://irishgeography.ie/index.php/irishgeography/article/view/1416/1158 - 10.2014/igj.v53i2.1416
Rights	© 2020, Geographical Society of Ireland. Published by Routledge – Taylor & Francis Group. All rights reserved. This is an Accepted Manuscript of an item published by Taylor & Francis in Irish Geography in November 2020, available online: https://dx.doi.org/10.2014/igj.v53i2.1416
Download date	2026-10-07 11:16:08
Item downloaded from	https://hdl.handle.net/10468/11147

Incorporating host-parasite biotic factors in species distribution models: Modelling the distribution of the castor bean tick, *Ixodes ricinus*.

Abstract

Understanding where ticks are found, and the drivers of their geographic distributions is imperative for successful epidemiological precautions. Predictive models of tick distributions are often projected using solely abiotic (e.g., climate) variables, despite the strong biotic interaction that host species undoubtedly have with parasitic species. We used species distribution modelling to project the distribution of *Ixodes ricinus* in Ireland and the United Kingdom using different combinations of abiotic, biotic, and abiotic-biotic variables. We found that models parameterised solely on abiotic variables generally reported lower accuracy and ecological realism than models that incorporated biotic factors alongside climate. We also investigated representation of host distribution in models, testing four different methods (habitat suitability of individual hosts, presence-absence of individual hosts, ensembled habitat suitability, and ensembled presence-absence). Biotic representations of ensembled host distributions alongside abiotic variables reported the highest accuracy, with the variable representing host diversity (e.g., number of host species) the most important variable when measured using a jackknife test. Moreover, our results suggested how host distributions are represented (i.e., presence-absence, habitat suitability) greatly impacted results, with differences reported among habitat specialists and generalists. Results suggest that it is now imperative for projections of parasitic species to include a representation of biotic factors with host species. This research has improved our understanding of the drivers of tick distributions in a national context, and the investigation of biotic representation should foster discussion among researchers working in species distribution modelling and the wider biogeography discipline.

Keywords: biotic interactions; host species; parasitism; species distribution modelling; ticks

Introduction

Ticks are one of the most important global vectors in the transmission of disease (Kelly *et al.* 2001), known to carry a wider variety of infectious agents compared to other arthropods (Soenshine 1991). Within Europe *Ixodes ricinus* is the most common arthropod vector (Parola & Raoult, 2001), with the ability to transmit various pathogens, including *Borrelia burgdorferi* (Lyme disease - Pietzsch *et al.* 2005). Global incidence rates of Lyme disease vary, but most reported cases are found in Europe and North America, with research indicating increasing incidence rates (Mead 2018). In Ireland, varied incidence and infection rates across both space and time have been reported (Gray *et al.* 1996; McKeown and Garvey 2009; Cullen 2010; Lambert *et al.* 2019), with the crude incidence rate of Lyme neuroborreliosis progressively increasing towards the south and west of the country (HSE 2019). Reports in the UK have identified an increase from 693 cases in 2005 to 1,310 cases in 2016 (Lorenc *et al.* 2017). Due to increasing incidence rates and the spatial variability in reports, understanding where ticks are found, and the drivers of their geographic distributions is imperative for successful epidemiological precautions.

In Ireland and the UK studies have predominantly focused on reporting the presence and density of tick species at a national scale using fieldwork and location maps (Martyn 1988; Kirstein *et al.* 1997; Kelly *et al.* 2001; Dobson *et al.* 2011a; Medlock and Leach, 2015; Zintl *et al.* 2017). For example, both Kirstein *et al.* (1997) and Kelly *et al.* (2001) produced location maps indicating presence of tick species from field studies in Ireland, while in the UK, Martyn (1988) published the first atlas of tick distributions, with the tick surveillance scheme continued by Public Health England. Most recently, Zintl *et al.* (2017) reported 151 tick observations from 26 sites across Ireland. Such studies improve our understanding of the environmental drivers of tick abundance and densities, but are often restricted spatially to specific locations.

Species distribution models (SDMs) are one of the most important GIScience research areas in biogeography, and a powerful spatial ecological tool for studying the geographic distribution of plants, animals, and other taxa (Franklin 2009; Peterson *et al.* 2011). The modelling framework provides a methodology for researchers and practitioners to quantitatively assess the relationship between species distributions and environmental factors, having been widely used for various applications including predicting disease spread (Peterson *et al.* 2004; Crowl *et al.* 2008; Bhatt *et al.* 2013). In Europe, tick SDMs have primarily been undertaken at a continental scale. For example, Alkishe *et al.* (2017) modelled the current and future (2050 and

2070) distribution of *I. ricinus* based on several climatic factors and identified that distributions are expected to increase in area under all future climate scenarios. Similarly, Williams *et al.* (2015) identified a shift to higher latitudes for eight European tick species for 2050 and 2098. Several other global studies have undertaken distribution projections for ticks at regional (De Clercq *et al.* 2015; Raghavan *et al.* 2016; St John *et al.* 2016), continental (Springer *et al.* 2015; Rubel *et al.* 2016; Hahn *et al.* 2016; Sage *et al.* 2017), and even global (Alkishe *et al.* 2020) scales; all identifying the importance of climatic factors, with many outlining the potential risk associated with increased human exposure to such a prevalent disease vector.

Climatic factors are an important determinant of species distributions; however, they are not the sole driver. The ‘BAM’ framework (Figure 1), developed by Soberón and Peterson (2005), illustrates the individual and joint effects of the three most important factors in determining a species distribution. Biotic (**B**) factors represent interactions with other species (e.g., competition, parasitism), abiotic (**A**) factors represent the physiological tolerances of the species (e.g., temperature, precipitation), and movement (**M**) factors represent the ability of a species to access the habitats (e.g., dispersal, foraging). The importance of all three factors is well recognised, yet in the majority of the aforementioned studies projecting tick distributions, models were parameterised on only abiotic factors. The fact that ticks have a wide range of hosts, which often lack good-quality occurrence data, as well as uncertainty in how best to incorporate such data into model parameterisation has been cited as rationale for focusing on climatic factors when projecting tick distributions (Alkishe *et al.* 2020). Moreover, abiotic factors have often superseded biotic factors in SDMs due in part to the fact that abiotic factors often influence species at a broader spatial extent (Wisz *et al.* 2013; Miller and Holloway 2015). Despite this, distribution studies that have included biotic factors have observed increases in model accuracy (Araújo and Luoto 2007; Heikkinen *et al.* 2007; Kissling *et al.* 2008; Wisz *et al.* 2013; Raath *et al.* 2018), meaning distribution models could be improved by explicitly incorporating biotic factors.

[Figure 1 near here]

Studies have emerged that incorporate both abiotic and biotic factors when projecting tick distributions, often resulting in improved accuracy and ecological realism (Medlock *et al.* 2013; Donaldson *et al.* 2016; Estrada-Peña and de la Fuente 2017). For example, in a study predicting *Ornithodoros turicata* distribution in the continental USA, Donaldson *et al.* (2016) overlaid the predicted tick distribution with the distributions of 58 host species, and identified

91 areas of significant overlap that could identify principal hosts. Similarly, Estrada-Peña and de
92 la Fuente (2017) incorporated probabilities of occurrence of host species in their tick
93 distribution models, finding significant tick-host relationships among the complex
94 communities of host species. The importance of incorporating biotic factors as covariates in
95 any statistical models projecting species distributions is well established; however, it is perhaps
96 more pertinent for parasite species that have such a strong dependence on hosts (Wisz *et al.*
97 2013). Recent reviews (Wisz *et al.* 2013; Dormann *et al.* 2018; Blanchet *et al.* 2020) have
98 outlined several challenges associated with incorporating biotic factors within SDMs, one of
99 which is the method through which the biotic factors are represented. Raath *et al.* (2018)
100 recently compared four methods of incorporating biotic factors (host plant distribution) in
101 SDMs projecting the distribution of the African silk moth in sub-Saharan Africa. They found
102 that the method with which biotic factors were represented in the statistical model (e.g.,
103 presence-absence of individual hosts, habitat suitability of individual hosts, combined
104 presence-absence of all hosts, and combined habitat suitability of all hosts) greatly affected the
105 predictive ability of the models, yet there was inconsistency among representations, suggesting
106 a need for further research.

107 With the incorporation of biotic factors (i.e. host distributions) as environmental covariates in
108 statistical models shown to improve projections of tick distributions, coupled with the fact that
109 several tick SDM studies still neglect host distribution altogether, there persists a need to
110 investigate the methods of representing biotic factors within SDM, and how their incorporation
111 impacts the current geographic predictions of ticks. In this study, we investigated the role of
112 biotic (**B**) and abiotic (**A**) factors in determining the distribution of *I. ricinus* in Ireland and the
113 UK focusing on two specific research questions: 1) How do different ‘BAM’ scenarios (e.g.,
114 **A**, **B**, **BA**) affect the accuracy and ecological realism of *I. ricinus* projections in Ireland and the
115 UK? And 2) How do different representations of host species (e.g., **B**) affect the accuracy and
116 ecological realism of the projections?

117 **Methodology**

118 ***Data Collection***

119 We acquired 172 primary occurrence records for *I. ricinus* from the Global Biodiversity
120 Information Facility (GBIF 2020) across Ireland and the UK since 2000. Following GBIF data
121 cleaning recommendations (Chapman 2005), incomplete points and duplicates were removed,
122 resulting in 132 occurrence records for use in the statistical models. Data for 16 primary host

species were also obtained from GBIF (2019). Climate data was obtained from WorldClim (Fink and Hijmans *et al.*, 2017) and resampled using cubic convolution to 1000m. Corine land cover (EEA 2018) data was used to calculate the percentage of arable, forest, grass, pasture, urban, and water cover within a corresponding 1000m resolution.

Species Distribution Modelling

SDMs generate a prediction of habitat suitability in both environmental and geographic space. The most common response variables used are data representing species presence-absence or presence-only, which are coupled with a set of corresponding and georeferenced environmental variables. Species-environment relationships are then estimated using a range of statistical approaches. These species-environment relationships can be extrapolated in space and time generating a habitat suitability map that identifies the probability that the environmental conditions at that location are favourable for the study species. To create a categorical representation of habitat suitability, a threshold that delineates between presence-absence can be selected, with habitat suitability values above and below this predicted as present and absent, respectively. The choice of response data, environmental data, and statistical models are all related to hypotheses derived from the underlying niche concept. As outlined above, the choice of environmental variables can have substantial implications on model output. Here, we investigated three different ‘BAM’ scenarios. Each scenario incorporates different methodologies to derive environmental variables based on the ‘BAM’ diagram (Figure 1). Figure 2 provides a methodological workflow of the SDM steps used in this study.

Maximum entropy (MaxEnt) was chosen as the statistical method to model the species-environment relationships for all scenarios (Phillips *et al.* 2006). MaxEnt selects suitable environmental variables by measuring how well they delineate between the recorded presence and pseudo-absence observations. The method chooses the split in environmental variables (e.g., temperature) that records the purest split between values in the binary response variable (e.g., presence and pseudo-absence). This method has been found to outperform other presence-only methods in an extensive comparison study (Elith and Graham 2009). Moreover, MaxEnt is robust to variable collinearity in model training and the removal of highly correlated variables prior to model parameterisation has little impact due to the fact the algorithm removes redundant variables (Feng *et al.* 2019), meaning MaxEnt is a robust method to evaluate different combinations of multiple environmental variables. However, the potential for spurious results and overfitting still exists and results need to be underpinned with a

mechanistic understanding related to their contribution in determining tick distributions (Holloway *et al.* 2018). To do this, we implemented the jackknife test to measure variable importance and explore the impact of variable representation on results. In the final MaxEnt model, the maximum number of iterations was set to 5,000 to allow model convergence; the number of pseudo-absences was set at 10,000 (following Barbet-Massin *et al.* 2012), and the model incorporated only linear and quadratic features to avoid over-fitting.

To account for unequal survey coverage in the species data, each SDM was fit with a bias grid to control for any violation in the assumption of independence in the response data that would increase the likelihood of type I errors (Fourcarde *et al.* 2014). A bias grid is the equivalent of a sampling probability surface, where the cell values reflect sampling effort and provides a weight to the pseudo-absence data used in modelling. Following Elith *et al.* (2010), a Gaussian kernel density map of the occurrence locations within 10km was derived and divided by the weighted number of terrestrial cells in the neighbourhood in order to avoid the edge effects at the coastline. This was then rescaled so that values varied between 1 and 20 in order to control for the impact of extreme variation. Bias grids are presented in Supplementary Information 1 to provide an overview of the spatial bias in occurrence records.

[Figure 2 near here]

Scenario A

Abiotic factors represent the physiological tolerances of species (i.e. temperature, precipitation) meaning climate variables best represent these factors. WorldClim v2.1 has interpolations of observed data from ~1970-2000, with 19 bioclimatic variables derived from the monthly temperature and rainfall values (Fick and Hijmans 2017). The 15 abiotic variables used by Alkishe *et al.* (2017) were used within this study to allow for a direct comparison with recent European continental predictions, with variables 8-9 and 18-19 removed due to spatial artefacts. See Table 1 for variable code and description.

[Insert Table 1 near here]

Scenario B

Biotic factors represent interactions with other species (i.e. parasitism, competition). Ticks are dependent on host species, meaning in this study, the biotic factors consisted of the distributions of sixteen mammalian hosts. Of the 27 host species listed by Kelly *et al.* (2001), nine were excluded due to their domestication which would prevent reliable habitat suitability

186 predictions. Humans were excluded as a host species, due to the obvious clustering in urban
187 areas, and two other species (goat, mink) were excluded due to inadequate observations in
188 GBIF (2019). The final list of mammalian species consisted of; hedgehog, pygmy shrew,
189 rabbit, grey squirrel, red squirrel, bank vole, wood mouse, brown rat, fox, pine marten, stoat,
190 badger, fallow deer, red deer, sika deer, and hare. The distribution of these sixteen species was
191 predicted using the observed locations from GBIF (2019) since 2000, the fifteen abiotic
192 variables (Fink and Hijmans 2017), the land cover data (EEA 2018), and projected using
193 MaxEnt with the same settings and bias grids outlined above.

194 Recent research has indicated different predictive accuracies in using presence-absence of host
195 species compared to habitat suitability (Raath *et al.* 2018). Therefore, four methods of
196 incorporating biotic factors were investigated (Table 2). Both habitat suitability (hs.ind) and
197 presence-absence (pa.ind) of all sixteen species were incorporated as individual explanatory
198 variables, as well as an ensemble model (i.e., average or summation of multiple projections) of
199 both habitat suitability (hs.ens) and presence-absence (pa.ens). Thresholds for generating
200 binary presence-absence models from habitat suitability were calculated using maximum
201 sensitivity plus specificity.

202 [Insert Table 2 near here]

203 ***Scenario BA***

204 This scenario projects the potential distribution in the BAM framework. Both abiotic (i.e., the
205 fifteen bioclimatic variables) and biotic (i.e., distributions of the sixteen host species) variables
206 were used as the input variables in the statistical model. Again, all four methods of
207 incorporating biotic factors were included in different iterations of the statistical model.

208 ***Model Evaluation***

209 Model evaluation in SDM focuses on quantifying prediction accuracy to ascertain model
210 validity. To test our models, we withheld 20% of the 132 tick occurrence records from training
211 the statistical model for testing as is common practice in SDM research when independent test
212 data are lacking (Franklin 2009). This testing dataset was then used to evaluate the performance
213 of our models. We used three accuracy metrics that are best suited for presence-only data;
214 lowest possible threshold (LPT), minimum predicted area (MPA) and the Boyce Index (BI).
215 LPT is the value that results in zero omission errors, with higher values indicating a better
216 model. MPA represents the area encompassing a predefined proportion of observed species

occurrences (in this case 90 percent), with a lower value (measured in geographic area) representing a more parsimonious model (Engler et al. 2004). The BI implements a moving window analysis across the predicted values, using Spearman rank correlation coefficient to measure the monotonic increase in the predicted-to-expected frequency ratio with increasing habitat suitability (Boyce *et al.* 2002). This method was undertaken using the *ecospat* package (Broenniman *et al.* 2014) in R 3.6.1 (R Development Core Team 2008) using default settings, with values ranging from -1 to 1, with higher values representing more parsimonious models, and 0 indicating predictions indifferent from a random model.

Results

The projected tick distributions and subsequent model accuracies varied substantially across the different implementations of **A**, **B**, and **BA** (Figure 3, Table 3). The model parameterised only on **A** (Figure 3a) reported high probability (>0.5) for the south and west of Ireland, as well as Wales, the south, southwest and northwest of England, and eastern Scotland, with low probability (<0.5) of occurrence for the remainder of Ireland and eastern England. Models parameterised including representations of host species (**B**, **BA**) consistently projected the southwest of the UK as suitable, with much wider spatial variation elsewhere. When the percentage contribution and permutation importance of the host species was explored in models that incorporated biotic factors (Table 4), the distribution of sika deer and bank voles were continually high across all models (10-40%), as well as contributing to large gains in the final models (Supplementary Information 2), suggesting the distribution of these host species may be an important driver in determining tick distributions and supporting the high probability within the southwest regions of the study area. For **B** (pa.ens) and **B** (hs.ens) where only the ensembled representation of biotic factors were used in model parameterisation, clear overfitting of the data occurred, with large swaths of the study area projected as suitable (Figure 3d-e). The **BA** models appear to have much more localised areas of high suitability (>0.5) across all counties in Ireland and the UK, which most likely reflects a more ecologically realistic distribution.

[Table 3 near here]

[Figure 3 near here]

Certain species were reported as contributing heavily to the projections when measured using one representation, but not the other (Table 4, Supplementary Information 2). For example, rabbit and hare distributions had higher gains for models when measured using habitat

suitability compared to presence-absence (Supplementary Information 2), and contributed more to final models when represented as habitat suitability, with rabbits contributing 0% to the final models when represented as presence-absence (Table 4). Alternatively, red deer distributions contributed more when measured using presence-absence compared to habitat suitability (Table 4).

There was discrepancy in model validity when measured using the different accuracy metrics, with LPT, MPA, and BI indicating the most accurate models as B (hs.ens), BA (hs.ens), and BA (pa.ens) respectively (Table 3). LPT rewards correct prediction of presences, often resulting in a high commission error (i.e. high false positive rate), which aligns with the observed overfitting for models B (pa.ens) and B (hs.ens) (Figure 3d-e). MPA and BI both attempt to control omission errors (i.e. false negatives), with the BI considered the more robust metric due to its ability to measure the monotonic increase in the predicted-to-expected frequency ratio with increasing habitat suitability (Hirzel *et al.* 2006). Models BA (hs.ens) and BA (pa.ens) both reported BI values above 0.9 and low MPA values suggesting a combination of abiotic variables with only one variable representing host diversity reports the most valid projection of tick distributions. Jackknife testing was used to evaluate variable importance to the final models, with the ensemble representations of host species reporting the highest gain in model performance when used in isolation for the BA (pa.ens) parameterisations and the largest decrease in gain in model performance when removed from both the BA (pa.ens) and BA (hs.ens) parameterisations (Figure 4).

[Figure 4 near here]

[Table 4 near here]

Discussion

The aim of this study was to investigate the role of abiotic and biotic factors in determining tick distributions in Ireland and the UK, comparing methods of incorporating biotic variables into SDMs. From examining the models (Figure 3), it is evident that there are certain regions that are more suitable for *I. ricinus* when the roles of **B**, **A**, and **BA** are compared. The most suitable areas in Ireland appear to be in the south, west, southeast, and north, while in the UK the most suitable areas are the south-west region of England, Wales, and western Scotland, although we did identify high probabilities for suitable habitat in every county of Ireland and the UK (Figure 3). However, these areas do vary depending on the choice of environmental variables incorporated in the model. Previous continental scale projections of *I. ricinus* have

identified the importance of abiotic variables in determining both current and future distributions (e.g., Alkishe *et al.* 2017), with our result identifying the importance of mean annual temperature and precipitation of the driest month as important variables (Figure 4, Table 4, Supplementary Information 2); however, the results of this study also highlight the need to incorporate factors related to the distribution of host species.

The relatively lower accuracy of *I. ricinus* projections for models parameterised solely on abiotic (**A**) factors compared to both biotic and abiotic (**BA**) factors (Table 3) suggests that abiotic factors alone do not adequately capture the drivers of tick distributions. Due to the strong dependence of ticks on host species, we expected that tick distributions would exhibit positive relationships with host distributions. The presence of large mammals (e.g., sika deer, red deer) were often important in the final entropy models, while smaller mammals (e.g., bank voles) were also shown to be important to the relative percentage contribution and regularised training gain in the final models (Table 4, Supplementary Information 2). Improvements in model accuracy (Table 3) and ecological realism (Figure 3) corroborate other studies that have included explicit variables measuring biotic factors (e.g., Kissling *et al.* 2008; Donaldson *et al.* 2016), and our results suggest that to accurately and realistically ascertain the distribution of parasitic species, variables that account for such relationships must be included.

The models parameterised on the different representations of **B** varied greatly, highlighting the need for continued research into how such conceptualisations can affect results. We found that BA (pa.ens) was the most accurate when the BI was used as the accuracy metric (Table 3), corroborating the findings of Raath *et al.* (2018) who also found that a combined presence-absence map was the most accurate representation of **B** in their study of African silk moths. This variable represents the total number of projected host species in that grid (i.e., equivalent of host species richness). Subsequently, this model may outperform others due to its ability to incorporate information on all host species, rather than assuming dependence on only one or a handful of species. The areas predicted as highly suitable by this model (Figure 3e) correspond with areas that were projected to support 12-16 host species. Subsequently, a variable representing the host diversity in the model potentially captures the parasite-host relationships more realistically than individual habitat suitability maps of all host species or an average habitat suitability of all hosts. Moreover, when each variable was tested for its importance to the final model through the jackknife test, the ensemble representations of host species reported the highest gain in model performance when used in isolation for BA (pa.ens) parameterisations, and the largest decrease in gain in model performance when removed from

both the BA (pa.ens) and BA (hs.ens) parameterisations (Figure 4). Tick interactions with mammals such as sika deer (Kimura *et al.* 1995; Gray *et al.* 1999; Braticikov *et al.* 2019), red deer (Gray *et al.* 1999; Zintl *et al.* 2011; Razanske *et al.* 2019), foxes (Lappin 2016; D’Amico *et al.* 2017; Sándor *et al.* 2017) and small rodents (Cayol *et al.* 2017; Cull *et al.* 2018) are well established, meaning the dependence of *I. ricinus* on multiple host species is best represented through a measure of diversity and distribution rather than as a series of individual host distributions.

Unsurprisingly many host distributions were continuously important in final models, both through the combined representation of host diversity, but also when included as individual species (Table 4, Supplementary Information 2). Sika deer and bank voles were consistently reported as important variables in final models, with these species exhibiting a strong positive relationship with forest cover (results not shown). The importance of vegetation cover for tick distributions and abundance is well established (Dobson *et al.* 2011a, 2011b), meaning there is the potential that our models confounded the importance of host distribution with land cover. To investigate this, we parameterised a model using the abiotic variables and land cover (Supplementary Information 3). This model performed poorly (LPT 0.299, MPA 150787, BI 0.727) compared to models parameterised with host distributions, highlighting the importance of considering the full ecological niche requirements of host species when projecting tick distributions.

Moreover, the relative contribution and subsequent regularised training gain of biotic variables altered for individual species depending on whether the variable was represented as continuous habitat suitability or categorical presence-absence (Table 4, Supplementary Information 2). The preference for binary representations appeared to occur for habitat specialist species (e.g., red deer, wood mouse, bank vole, stoat), suggesting that delineating the most suitable habitat from lesser habitat into presence-absence improves projections of parasite-host relationships. A continuous representation favoured more habitat generalists (e.g., grey squirrel, rat, rabbit), in part due to their ubiquitous distributions, meaning binary representations predicted them as present throughout the study area. Therefore, for generalist species a continuous variable that captures nuances in the habitat suitability improves models when compared to a thresholded presence-absence map, although generalist species continually reported low contributions and gains to final models (Table 4, Supplementary Information 2). This should have important considerations for future SDM research when study species may only have one or two primary biotic interactions determining their potential distributions.

Our results highlight the substantial impact of user decisions in how biotic factors are represented (Tables 3-4, Figures 3-4), and subsequently we have refrained from making grandiose statements considering our projections to be the definitive distribution of *I. ricinus* in Ireland and the UK. Research using such approaches for epidemiologically important vector species must be aware of such uncertainties in model parameterisations and strive to incorporate or quantify these in projections (see Alkishe *et al.* 2020 for a comprehensive assessment of model uncertainty projecting global tick distributions). Similarly, while spatial bias in response data was controlled for by using a bias grid (Supplementary Information 1), the tick data (GBIF 2020) does vary spatially with a wider coverage of the UK (particularly England and Scotland) and a lower number of observations in Ireland and Wales, with a higher reporting of occurrence in coastal regions (i.e. four observations on the Aran Islands alone) . Future research collecting verified tick locations that are made available through open data repositories is needed to improve model calibration and tick predictions, and supports recent calls for more comprehensive surveys, particularly in Ireland (Zintl *et al.* 2017). Despite these caveats, we do report high accuracy values (Table 3), allowing confidence in the discussion of the important environmental drivers of *I. ricinus* distributions.

The importance of considering environmental variables, particularly in the context of the ‘BAM’ framework (Figure 1) is therefore imperative for distribution studies; however, we did not incorporate movement factors (**M**) in our models. **M** refers to the area that has been or will be accessible to a species within a certain timeframe. The majority of SDM research only incorporates movement when investigating response to climate change (Franklin 2010; Holloway *et al.* 2016; Holloway and Miller, 2017). Recent research on tick distributions incorporating movement has identified the role of temporal variation in mammal movements resulting from land cover, and that such movements can have implications for tick distributions (Martin *et al.* 2018). Similarly, Halsey and Miller (2018) developed a spatial agent-based model to explore host-tick interactions, highlighting the possibility to combine such approaches and better inform on host-tick movements and interactions. Such approaches combining SDM and agent-based models are beginning to emerge that could create finer scale distribution models that can account for all three ‘BAM’ factors (e.g., Holloway 2018), meaning future research should continue to explore the inter-linked relationships between biotic, abiotic, and movement variables.

Finally, there persists a need to explore habitat suitability of *I. ricinus* in conjunction with other measures of biodiversity and epidemiology. It is important to note that ground feeding species

of birds (e.g., blackbirds, redwings) have been found to be important hosts for *Ixodes* ticks, with some rural populations reportedly having infestation rates as high as 74% (Gregoire *et al.* 2002; Singh and Girshick 2003). We opted only to investigate the role of mammalian host species; however, future research should look to incorporate information on avian distributions when projecting the geographic ranges of ticks. Similarly, we should also note that high habitat suitability does not necessarily indicate presence of breeding populations of ticks or Lyme disease. For example, Talleklint and Jaenson (1996) found a significant relationship between tick density and infection rates (of Lyme disease) at lower tick densities, while Gray *et al.* (1999) found that tick infection rates were inversely related to tick abundance. Dobson *et al.* (2011a) also found spatial variability in the abundance of ticks across recreational areas in the UK, recording uniform presence across vegetation types, but higher densities in plots which contained trees. Healy and Bourke (2004) note that clusters of ticks and larvae should be used to identify breeding populations, which contradicts the predominant method in SDM of using binary representation of presence-absence. New methods, such as Poisson point process models, are emerging as suitable tools to predict the distribution and abundance of species from presence-only data (Warton and Shepherd, 2010; Renner and Warton 2013; Schank *et al.* 2017), and given the uncertain relationships identified, future studies should focus on developing predictions of both the distribution and density of ticks to inform epidemiological research.

Conclusion

Understanding the drivers of tick distributions is imperative for successful epidemiological precautions. Previous research has tended to focus only on the role of abiotic variables in determining tick distributions, and while research is beginning to emerge that highlights the importance of biotic variables (e.g., host distributions), there persists a need for the combined influence of **A** and **B** to be investigated together, as well as exploration into how best to represent biotic factors within SDM. Large differences in accuracy (Table 3) and area (Figures 3) were identified depending on the combination of ‘BAM’ variables used during model parameterisation. Representations of **B** varied substantially with results suggesting that a variable of host diversity captures the parasite-host relationships most accurately when coupled with abiotic factors (**BA**). The representation of host distribution also appeared to differ for habitat specialists and generalists (Table 4, Supplementary Information 2), suggesting use of a binary presence-absence conceptualisation and a continuous habitat suitability conceptualisation for specialist and generalist host species, respectively. There is now a

minimum requirement for any research into tick distributions to explicitly incorporate biotic factors, as excluding the host-parasite relationship masks the true distribution of species and does not capture an ecological realistic distribution of a species' habitat suitability.

Abstract

We would like to thank the anonymous reviewers and Ronan Foley for their comments and suggestions.

References

- Alkishe, A., Townsend Peterson, A., & Samy, A. (2017). Climate change influences on the potential geographic distribution of the disease vector tick *Ixodes ricinus*. *PLoS ONE*, 12 (12)
- Alkishe, A., Cobos, M.E., Townsend Peterson, A., & Samy, A.M. (2020). Recognising sources of uncertainty in disease vector ecological niche models: An example with the tick *Rhipicephalus sanguineus sensu lato*. *Perspectives in Ecology and Conservation*.
- Araújo, M.B. and Luoto, M., (2007). The importance of biotic interactions for modelling species distributions under climate change. *Global Ecology and Biogeography*, 16(6), pp.743-753.
- Barbet-Massin, M., F. Jiguet., C.H. Albert., and W. Thuiller. (2012). Selecting pseudo-absences for species distribution models: how, where and how many?. *Methods in Ecology and Evolution*, 3(2): 327-338.
- Bhatt, S., Gething, P.W., Brady, O.J., Messina, J.P., Farlow, A.W., Moyes, C.L., Drake, J.M., Brownstein, J.S., Hoen, A.G., Sankoh, O. and Myers, M.F., (2013). The global distribution and burden of dengue. *Nature*, 496(7446), p.504.
- Blanchet, F.G., Cazelles, K., Gravel, D. (2020). Co-occurrence is not evidence of ecological interactions. *Ecology Letters*, 23: 1050-1063.
- Boyce, M.S., P.R. Vernier., S.E. Nielsen., and F.K. Schmiegelow. (2002). Evaluating resource selection functions. *Ecological modelling*, 157(2): 281-300.
- Bratickov, M., Paulauskas, A., Rosef, O., Ražanskė, I., Gričiuvienė, L. and Radzijeuskaja, J., (2019). Prevalence and co-infection with tick-borne *Anaplasma phagocytophilum* and *Babesia*

spp. in red deer (*Cervus elaphus*) and roe deer (*Capreolus capreolus*) in Southern Norway. *International journal for parasitology: parasites and wildlife*. Amsterdam: Elsevier Ltd, 2019, Vol. 8.

Broenniman, O., B. Petitpierre., C. Randin., R. Engler., F. Breiner., M. D'Amen., L. Pellissier., J. Pottier., D. Pio., R. Garcia Mateo, *et al.* (2014). ecospat: Spatial ecology miscellaneous methods. R package version 1.0.

Cayol, C., Koskela, E., Mappes, T., Siukkola, A. and Kallio, E.R., (2017). Temporal dynamics of the tick *Ixodes ricinus* in northern Europe: epidemiological implications. *Parasites & vectors*, 10(1), p.166.

Chapman A. D. (2005) Principles and Methods of Data Cleaning - Primary Species and Species-Occurrence Data. version 1.0. Report for the Global Biodiversity Information Facility, Copenhagen. Available online at http://www.gbif.org/orc/?doc_id=1262

Crowl, T.A., Crist, T.O., Parmenter, R.R., Belovsky, G. and Lugo, A.E., (2008). The spread of invasive species and infectious disease as drivers of ecosystem change. *Frontiers in Ecology and the Environment*, 6(5), pp.238-246.

Cull, B., Pietzsch, M.E., Hansford, K.M., Gillingham, E.L. and Medlock, J.M., (2018). Surveillance of British ticks: An overview of species records, host associations, and new records of *Ixodes ricinus* distribution. *Ticks and tick-borne diseases*, 9(3), pp.605-614.

Cullen, E. (2010). Lyme disease and climate change. *Irish Medical Journal*. 103:1, 1-2.

D'Amico, G., Juránková, J., Tăbăran, F.A., Frgelecová, L., Forejtek, P., Matei, I.A., Ionică, A.M., Hodžić, A., Modrý, D. and Mihalca, A.D., (2017). Occurrence of ticks in the subcutaneous tissue of red foxes, *Vulpes vulpes* in Czech Republic and Romania. *Ticks and tick-borne diseases*, 8(2), pp.309-312.

De Clercq, E.M., Leta, S., Estrada-Peña, A., Madder, M., Adehan, S. and Vanwambeke, S.O., (2015). Species distribution modelling for *Rhipicephalus microplus* (Acari: Ixodidae) in Benin, West Africa: Comparing datasets and modelling algorithms. *Preventive veterinary medicine*, 118(1), pp.8-21.

Dobson., A.D.M., Taylor, J.L., & Randolph, S.E. (2011a). Ticks (*Ixodes ricinus*) abundance and seasonality at recreational sites in the UK: Hazards in relation to fine-scale habitat types revealed by complimentary sampling methods. *Ticks and Tick-borne Diseases*, 2: 67-74.

470 Dobson, A.D.M., Finnie, T.J.R., & Randolph, S.E. (2011b). A modified matrix model to
 471 describe the seasonal population ecology of the European tick *Ixodes ricinus*. *Journal of*
 472 *Applied Ecology*, 48: 1017-1028.

473 Donaldson, T.G., de Leon, A.A.P., Li, A.I., Castro-Arellano, I., Wozniak, E., Boyle, W.K.,
 474 Hargrove, R., Wilder, H.K., Kim, H.J., Teel, P.D. and Lopez, J.E., (2016). Assessment of the
 475 geographic distribution of *Ornithodoros turicata* (Argasidae): climate variation and host
 476 diversity. *PLoS neglected tropical diseases*, 10(2), p.e0004383.

477 Dormann, C.F., Bobrowski, M., Dehling, D.M., Harris, D.J., Hartig, F., Lischke, H., Moretti,
 478 M.D., Pagel, J., Pinkert, S., Schleuning, M. and Schmidt, S.I., (2018). Biotic interactions in
 479 species distribution modelling: 10 questions to guide interpretation and avoid false
 480 conclusions. *Global ecology and biogeography*, 27(9), pp.1004-1016.

481 EEA. (2018). "Corine Land Cover (CLC) 2018, Version 20b2." *Release Date: 21-12-*
 482 *2018*. European Environment Agency. [https://land.copernicus.eu/pan-european/corine-land-](https://land.copernicus.eu/pan-european/corine-land-cover/clc2018)
 483 [cover/clc2018](https://land.copernicus.eu/pan-european/corine-land-cover/clc2018)

484 Elith, J., and C.H. Graham. (2009). Do they? How do they? WHY do they differ? On finding
 485 reasons for differing performances of species distribution models. *Ecography*, 32(1): 66-77.

486 Elith, J., M. Kearney., and S. Phillips. (2010). The art of modelling range-shifting species.
 487 *Methods in Ecology and Evolution*, 1: 330–342

488 Engler, R., Guisan, A. and Rechsteiner, L., 2004. An improved approach for predicting the
 489 distribution of rare and endangered species from occurrence and pseudo-absence data. *Journal*
 490 *of applied ecology*, 41: 263-274.

491 Estrada-Peña, A. and de la Fuente, J., (2017). Host distribution does not limit the range of the
 492 tick *Ixodes ricinus* but impacts the circulation of transmitted pathogens. *Frontiers in Cellular*
 493 *and Infection Microbiology*, 7, p.405.

494 Feng, X., Park, D.S., Liang, Y., Pandey, R. and Papeş, M., 2019. Collinearity in ecological
 495 niche modeling: Confusions and challenges. *Ecology and evolution*, 9: 10365-10376.

496 Fourcade, Y., J.O. Engler., D. Rödder., and J. Secondi. (2014) Mapping Species Distributions
 497 with MAXENT Using a Geographically Biased Sample of Presence Data: A Performance
 498 Assessment of Methods for Correcting Sampling Bias, *PLoS One*, 9(5): e97122.

499 Franklin, J. (2009). *Mapping Species Distributions: Spatial Inference Prediction*. Cambridge:
500 Cambridge University Press.

501 Franklin, J. (2010). Moving beyond static species distribution models in support of
502 conservation biogeography. *Diversity and Distributions*, 16 (3): 321-330.

503 GBIF (20120) GBIF Occurrence Download <https://doi.org/10.15468/dl.zkqf26>

504 GBIF (2019) GBIF Occurrence Download <https://doi.org/10.15468/dl.wxxxwh9>

505 Gray, J.S., Cutler, S., Robertson, J. & O'Connell, S. (1996). Lyme disease in the Republic of
506 Ireland, p.6. In 'Who Workshop on Lyme Borreliosis diagnosis and surveillance'. Warsaw,
507 Poland 20th – 22nd June 1995.

508 Gray, J.S., Kirstein, F., Robertson, J.N., Stein, J. and Kahl, O., (1999). *Borrelia burgdorferi*
509 sensu lato in *Ixodes ricinus* ticks and rodents in a recreational park in south-western
510 Ireland. *Experimental & applied acarology*, 23(9), pp.717-729.

511 Gregoire, A., Faivre, B., Heeb, P. and Cezilly, F., (2002). A comparison of infestation patterns
512 by *Ixodes* ticks in urban and rural populations of the Common Blackbird *Turdus*
513 *merula*. *Ibis*, 144(4), pp.640-645.

514 Hahn, M.B., Jarnevich, C.S., Monaghan, A.J. and Eisen, R.J., (2016). Modeling the geographic
515 distribution of *Ixodes scapularis* and *Ixodes pacificus* (Acari: Ixodidae) in the contiguous
516 United States. *Journal of medical entomology*, 53(5), pp.1176-1191.

517 Halsey, S.J. and Miller, J.R., (2018). A spatial agent-based model of the disease vector *Ixodes*
518 *scapularis* to explore host-tick associations. *Ecological modelling*, 387, pp.96-106.

519 Healy, J.A.E. and Bourke, P., (2004). Field evidence for aggregating behaviour in the tick
520 *Ixodes ricinus* L. *Acarologia*, 44(1-2), pp.3-13.

521 Heikkinen, R.K., Luoto, M., Virkkala, R., Pearson, R.G. and Körber, J.H., (2007). Biotic
522 interactions improve prediction of boreal bird distributions at macro-scales. *Global Ecology*
523 *and Biogeography*, 16(6), pp.754-763.

524 Hirzel, A.H., G. Le Lay., V. Helfer., C. Randin., and A. Guisan. 2006. Evaluating the ability
525 of habitat suitability models to predict species presences. *Ecological Modelling*, 199(2): 142-
526 152.

527 Holloway, P., (2018). Simulating movement-related resource dynamics to improve species
528 distribution models: a case study with oilbirds in Northern South America. *The Professional*
529 *Geographer*, 70(4), pp.528-540.

530 Holloway, P., and Miller, J.A., (2017). A quantitative synthesis of the movement concepts
531 used in species distribution modelling. *Ecological Modelling*, 356: 91-103.

532 Holloway, P., Kudenko, D., and Bell, J. (2018). Dynamic selection of environmental
533 variables to improve the prediction of aphid phenology: A machine learning approach.
534 *Ecological Indicators*, 88; 512-521.

535 Holloway, P., Miller, J.A. and Gillings, S., (2016). Incorporating movement in species
536 distribution models: how do simulations of dispersal affect the accuracy and uncertainty of
537 projections?. *International Journal of Geographical Information Science*, 30(10), pp.2050-
538 2074.

539 HSE (2019). Primary Prevention and Surveillance of Lyme Borreliosis in Ireland. Report of
540 the Lyme Borreliosis Subcommittee of the Scientific Advisory Committee of the Health
541 Protection Surveillance Centre. Available at: [https://www.hpsc.ie/a-](https://www.hpsc.ie/a-z/vectorborne/lymedisease/informationforhealthcareprofessionals/)
542 [z/vectorborne/lymedisease/informationforhealthcareprofessionals/](https://www.hpsc.ie/a-z/vectorborne/lymedisease/informationforhealthcareprofessionals/). Accessed on 29th June
543 2020.

544 Kelly, T. C., Sleeman, D.P., Fennessy, G.J., Dillon, A., Walton, G.A. & Gray, J. (2001). The
545 mammal ticks (Acari: Ixodoidea) of Ireland: Their distribution and hosts. *Irish National*
546 *Journal*. 26:3, 63 – 70.

547 Kimura, K., Isogai, E., Isogai, H., Kamewaka, Y., Nishikawa, T., Ishii, N. and Fujii, N., (1995).
548 Detection of Lyme disease spirochetes in the skin of naturally infected wild sika deer (*Cervus*
549 *nippon yesoensis*) by PCR. *Appl. Environ. Microbiol.*, 61(4), pp.1641-1642.

550 Kirsten, F., Rijpkema, S., Molkenboer, M. & Gray, J.S., (1997). The distribution and
551 prevalence of *B. burgdorferi* genomospecies in *Ixodes ricinus* ticks in Ireland. *European*
552 *Journal of Epidemiology*. 13, 67.

553 Kissling, W.D., R. Field., and K. Böhning-Gaese. (2008). Spatial patterns of woody plant and
554 bird diversity: functional relationships or environmental effects?. *Global Ecology and*
555 *Biogeography*, 17(3): 327-339.

556 Lambert, J.S., Cook, M.J., Healy, J.E., Murtagh, R., Avramovic, G. and Lee, S.H., (2019).
557 Metagenomic 16S rRNA gene sequencing survey of *Borrelia* species in Irish samples of *Ixodes*
558 *ricinus* ticks. *PloS one*, 14(4).

559 Lappin, J.W.E., (2016). *Ixodid ticks: vectors for emerging diseases in Northern*
560 *Ireland* (Doctoral dissertation, Queen's University Belfast).

561 Lorenc, T., Jones-Diette, J., Blanchard, L., Sutcliffe, K., Stansfield, C., Harden, M., Sowden,
562 A. and Thomas, J., (2017). *Incidence and surveillance of Lyme disease*. London: EPPI-Centre,
563 Social Science Research Unit, UCL Institute of Education, University College London.

564 Martin, J., Vourc'h, G., Bonnot, N., Cargnelutti, B., Chaval, Y., Lourtet, B., Goulard, M.,
565 Hoch, T., Plantard, O., Hewison, A.M. and Morellet, N., (2018). Temporal shifts in landscape
566 connectivity for an ecosystem engineer, the roe deer, across a multiple-use
567 landscape. *Landscape ecology*, 33(6), pp.937-954.

568 Martyn, K.P. (1988) *Provisional Atlas of The Ticks (Ixodoidea) of the British Isles*. Biological
569 Records Centre Institute of Terrestrial Ecology, Abbots Ripton, Huntingdon.

570 McKeown, P. & Garvey, P. (2009). Lyme disease often under diagnosed says HPSC. *Epi-*
571 *Insight*. 10, 1 – 5.

572 Mead, P.S., (2018). Global Epidemiology of *Borrelia burgdorferi* Infections. *Lyme Disease:*
573 *an Evidence-based Approach*, p.76.

574 Medlock, J.M. and Leach, S.A., (2015). Effect of climate change on vector-borne disease risk
575 in the UK. *The Lancet Infectious Diseases*, 15(6), pp.721-730.

576 Medlock, J.M., Hansford, K.M., Bormane, A., Derdakova, M., Estrada-Peña, A., George, J.C.,
577 Golovljova, I., Jaenson, T.G., Jensen, J.K., Jensen, P.M. and Kazimirova, M., 2013. Driving
578 forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. *Parasites &*
579 *vectors*, 6:1.

580 Miller, J. A., and P. Holloway. (2015). Incorporating movement in species distribution models.
581 *Progress in Physical Geography*, 39(6): 837-849.

582 Parola, P. & Raoult, D. (2001). Tick-borne bacterial diseases emerging in Europe. *Clinical*
583 *Microbiology and Infection*. 7, 80-83.

584 Peterson, A.T., Bauer, J.T. and Mills, J.N., (2004). Ecologic and geographic distribution of
585 filovirus disease. *Emerging infectious diseases*, 10(1), p.40.

586 Peterson, A. T., J. Soberón., R.G. Pearson., R.P. Anderson., E. Martínez-Meyer., M.
587 Nakamura., and M.B. Araújo. (2011). Ecological niches and geographic distributions (MPB-
588 49). Princeton University Press.

589 Phillips, S. J., Anderson, R.P., & Schapire, R. E. (2006) Maximum entropy modelling of
590 species geographic distributions. *Ecological Modelling*, 190, 231-259

591 Pietzsch, M.E., Medlock, J.M, Jones, L., Avenell, D., Abbott, J., Harding, P. & Leach, S.
592 (2005). Distribution of *Ixodes ricinus* in the British Isles: Investigation of historical
593 records. *Medical and Veterinary Entomology*. 19, 306-314.

594 R Development Core Team (2008). R: A language and environment for statistical computing.
595 Vienna, Austria: R Foundation for Statistical Computing.

596 Raath, M.J., Le Roux, P.C., Veldtman, R. and Greve, M., (2018). Incorporating biotic
597 interactions in the distribution models of African wild silk moths (Gonometa species,
598 Lasiocampidae) using different representations of modelled host tree distributions. *Austral*
599 *ecology*, 43(3), pp.316-327.

600 Razanske, I., Rosef, O., Radzijeuskaja, J., Bratchikov, M., Gričiuvienė, L. and Paulauskas, A.,
601 (2019). Prevalence and co-infection with tick-borne *Anaplasma phagocytophilum* and *Babesia*
602 spp. in red deer (*Cervus elaphus*) and roe deer (*Capreolus capreolus*) in Southern
603 Norway. *International Journal for Parasitology: Parasites and Wildlife*, 8, pp.127-134.

604 Renner, I.W. and Warton, D.I., (2013). Equivalence of MAXENT and Poisson point process
605 models for species distribution modeling in ecology. *Biometrics*, 69(1), pp.274-281.

606 Raghavan, R.K., Goodin, D.G., Hanzlicek, G.A., Zolnerowich, G., Dryden, M.W., Anderson,
607 G.A. and Ganta, R.R., (2016). maximum entropy-based ecological niche model and bio-
608 climatic determinants of lone star tick (*Amblyomma americanum*) niche. *Vector-Borne and*
609 *Zoonotic Diseases*, 16(3), pp.205-211.

610 Rubel, F., Brugger, K., Pfeffer, M., Chitimia-Dobler, L., Didyk, Y.M., Leverenz, S., Dautel,
 611 H. and Kahl, O., (2016). Geographical distribution of *Dermacentor marginatus* and
 612 *Dermacentor reticulatus* in Europe. *Ticks and tick-borne diseases*, 7(1), pp.224-233.

613 Sage, K.M., Johnson, T.L., Teglas, M.B., Nieto, N.C. and Schwan, T.G., (2017). Ecological
 614 niche modeling and distribution of *Ornithodoros hermsi* associated with tick-borne relapsing
 615 fever in western North America. *PLoS neglected tropical diseases*, 11(10), p.e0006047.

616 Sándor, A.D., D'Amico, G., Gherman, C.M., Dumitrache, M.O., Domşa, C. and Mihalca, A.D.,
 617 (2017). Mesocarnivores and macroparasites: altitude and land use predict the ticks occurring
 618 on red foxes (*Vulpes vulpes*). *Parasites & vectors*, 10(1), p.173.

619 Schank, C.J., Cove, M.V., Kelly, M.J., Mendoza, E., O'Farrill, G., Reyna-Hurtado, R., Meyer,
 620 N., Jordan, C.A., González-Maya, J.F., Lizcano, D.J. and Moreno, R., (2017). Using a novel
 621 model approach to assess the distribution and conservation status of the endangered Baird's
 622 tapir. *Diversity and Distributions*, 23(12), pp.1459-1471.

623 Singh, S.K. and Girschick, H.J., (2003). Tick-host interactions and their immunological
 624 implications in tick-borne diseases. *Current science*, 85(9), pp.1284-1298.

625 Soberón, J. & Peterson, A. (2005). Interpretation of models of fundamental ecological niches
 626 and species' distributional areas. *Biodiversity Informatics*. 2:1-10

627 Soberón, J. (2007). Grinnellian and Eltonian niches and geographic distributions of species.
 628 *Ecology letters*, 10 (12): 1115-1123.

629 Soenshine, D.E. (1991) *Biology of ticks*. Oxford University Press, New York

630 Springer, Y.P., Jarnevich, C.S., Barnett, D.T., Monaghan, A.J. and Eisen, R.J., (2015).
 631 Modeling the present and future geographic distribution of the Lone Star Tick, *Amblyomma*
 632 *americanum* (Ixodida: Ixodidae), in the continental United States. *The American journal of*
 633 *tropical medicine and hygiene*, 93(4), pp.875-890.

634 St. John, H.K., Adams, M.L., Masuoka, P.M., Flyer-Adams, J.G., Jiang, J., Rozmajzl, P.J.,
 635 Stromdahl, E.Y. and Richards, A.L., (2016). Prevalence, distribution, and development of an
 636 ecological niche model of *Dermacentor variabilis* ticks positive for *Rickettsia*
 637 *montanensis*. *Vector-Borne and Zoonotic Diseases*, 16(4), pp.253-263.

638 Tälleklint, L. and Jaenson, T.G.T. (1996). Relationship between Ixodes ricinus density and
639 prevalence of infection with Borrelia-like spirochetes and density of infected ticks. *Journal of*
640 *Medical Entomology*. 33, 805–81

641 Warton, D.I. and Shepherd, L.C., (2010). Poisson point process models solve the “pseudo-
642 absence problem” for presence-only data in ecology. *The Annals of Applied Statistics*, 4(3),
643 pp.1383-1402.

644 Williams, H. W., Cross, D. E., Crump, H. L., Drost, C. J., & Thomas, C. J. (2015). Climate
645 suitability for European ticks: assessing species distribution models against null models and
646 projection under AR5 climate. *Parasites & vectors*, 8(1), 440.

647 Wisz, M. S., J. Pottier., W.D. Kissling., L. Pellissier., J. Lenoir., C.F. Damgaard., C.F.
648 Dormann., M.C. Forchhammer., J.A. Grythnes., A. Guisan., *et al.* (2013). The role of biotic
649 interactions in shaping distributions and realised assemblages of species: implications for
650 species distribution modelling. *Biological Reviews*, 88(1): 15-30.

651 Zintl, A., Finnerty, E.J., Murphy, T.M., de Waal, T. and Gray, J.S., (2011). Babesias of red
652 deer (*Cervus elaphus*) in Ireland. *Veterinary research*, 42(1), p.7.

653 Zintl, A., Moutailler, S., Stuart, P., Paredis, L., Dutraive, J., Gonzalez, E., O’Connor, J.,
654 Devillers, E., Good, B., OMuireagain, C. and De Waal, T., (2017). Ticks and Tick-borne
655 diseases in Ireland. *Irish veterinary journal*, 70(1), p.4.

656

657

658 **Tables**

659 Table 1. Variable information for the 19 bioclimatic variables acquired from Fick and Hijmans
660 (2017).

Code	Variable
Bio1	Annual mean temperature
Bio2	Mean diurnal range
Bio3	Isothermality
Bio4	Temperature seasonality
Bio5	Maximum temperature of warmest month
Bio6	Minimum temperature of coldest month
Bio7	Temperature annual range
Bio8	Mean temperature of wettest quarter
Bio9	Mean temperature of driest quarter
Bio10	Mean temperature of warmest quarter
Bio11	Mean temperature of coldest quarter
Bio12	Annual precipitation
Bio13	Precipitation of wettest month
Bio14	Precipitation of driest month
Bio15	Precipitation seasonality
Bio16	Precipitation of wettest quarter
Bio17	Precipitation of driest quarter
Bio18	Precipitation of warmest quarter
Bio19	Precipitation of coldest quarter

661

662 Table 2. Description of the biotic representations used within parameterised models

Biotic Representation	Explanation
hs.ind	Individual maps of habitat suitability for all 16 host species
pa.ind	Individual maps of presence-absence for all 16 host species
hs.ens	Ensemble map of habitat suitability projections. The mean habitat suitability of all 16 host species
pa.ens	Ensemble map of presence-absence projections. Sum of all presence-absence maps for the 16 host species. Equivalent of projected host diversity

663

Table 3. The three accuracy metrics used to evaluate model performance; lowest possible threshold (LPT), minimum predicted area (MPA) and the Boyce Index (BI). Three ‘BAM’ scenarios were explored; biotic (B), abiotic (A) and biotic-abiotic (BA) scenarios, and four representations of biotic interactions. The four representations are individual habitat suitability maps of all hosts (hs.ind), individual presence-absence maps of all hosts (pa.ind), ensemble map of mean habitat suitability for all hosts (hs.ens), and ensemble map representing summation of all binary presence-absence maps (pa.ens).

	LPT	MPA	BI
A	0.250	214066	0.896
B (HS.IND)	0.238	147349	0.769
B (PA.IND)	0.252	145331	0.727
B (HS.ENS)	0.540	241599	0.842
B (PA.ENS)	0.327	220647	0.879
BA (HS.IND)	0.278	112787	0.664
BA (PA.IND)	0.247	171757	0.873
BA (HS.ENS)	0.151	92740	0.925
BA (PA.ENS)	0.262	110689	0.934

673 Table 4: Percent Contribution (PC) and Permutation Importance (PI) of the different environmental variables used in the final models. The four
674 representations are individual habitat suitability maps of all hosts (hs.ind), individual presence-absence maps of all hosts (pa.ind), ensemble map
675 of average habitat suitability for all hosts (hs.ens), and ensemble map representing summation of all binary presence-absence maps (pa.ens). Biotic
676 representations using ensembles B (hs.ens) and B (pa.ens) are not included in table as they only contain one variable, which contributes 100% to
677 the final models. See Table 1 for a description of the abiotic (A) codes.

678

	Abiotic (A)		Biotic (hs.ind)		Biotic (pa.ind)		Biotic Abiotic (hs.ind)		Biotic Abiotic (pa.ind)		Biotic Abiotic (hs.ens)		Biotic Abiotic (pa.ens)	
	PC	PI	PC	PI	PC	PI	PC	PI	PC	PI	PC	PI	PC	PI
A (bio1)	27.3	19.4					0.2	0.0	0.5	11.3	20.6	18.4	11.1	17.5
A (bio2)	1.5	9.5					2.8	1.1	0.7	9.1	1.2	13.6	0.7	10.0
A (bio3)	9.9	0.4					5.3	4.6	15.8	0.0	15.2	0.0	21.8	0.3
A (bio4)	3.7	7.9					0.0	0.0	1.3	10.3	6.6	5.1	6.5	7.8
A (bio5)	1.0	10.5					0.0	0.0	0.3	5.2	3.3	1.4	2.1	7.5
A (bio6)	4.2	8.1					0.2	2.2	0.4	5.8	7.3	12.8	1.2	9.9
A (bio7)	0.0	0.0					0.0	0.0	0.0	0.0	0.3	0.0	0.0	0.0
A (bio10)	0.4	0.0					0.9	0.0	2.0	0.2	0.0	0.0	0.0	0.0
A (bio11)	0.0	0.0					0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0
A (bio12)	0.2	2.6					3.1	0.0	0.4	3.9	0.7	7.6	0.5	3.6
A (bio13)	0.3	0.0					0.6	6.1	0.4	2.4	0.4	0.0	0.0	0.0

A (bio14)	19.2	22.8					3.5	17.2	6.7	21.9	14.5	18.7	11.6	18.9
A (bio15)	14.8	1.3					0.4	1.8	1.3	0.4	7.6	0.0	5.1	0.0
A (bio16)	0.0	0.0					0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0
A (bio17)	17.7	17.6					1.0	16.1	4.8	18.5	8.5	14.8	9.2	17.2
B (ensemble)											13.8	7.5	30.2	7.4
B (fox)			0.2	0.0	0.0	0.0	1.9	0.8	0.0	0.0				
B (sika deer)			37.0	17.9	21.4	16.5	32.6	0.5	13.5	1.0				
B (red deer)			2.7	8.8	11.8	11.4	0.9	6.8	7.6	1.0				
B (fallow deer)			1.1	0.7	2.2	0.0	1.6	0.0	1.3	0.0				
B (hare)			7.1	15.4	4.4	5.5	1.9	2.3	1.9	0.0				
B (wood mouse)			0.0	0.0	5.9	5.9	0.3	0.0	4.4	0.8				
B (pygmy shrew)			2.0	4.2	1.2	2.3	0.1	0.7	0.2	0.0				
B (grey squirrel)			1.6	8.9	1.6	4.1	3.5	8.8	3.6	1.6				
B (red squirrel)			3.9	18.1	7.3	8.8	1.8	13.8	5.8	3.8				
B (bank vole)			15.0	16.2	33.3	27.6	12.8	14.5	21.2	1.2				
B (pine marten)			0.1	1.8	7.1	9.8	0.9	0.0	2.9	0.0				
B (rat)			3.0	6.0	1.3	1.8	2.5	0.0	0.1	0.0				
B (rabbit)			10.0	0.2	0.0	0.0	6.2	0.0	0.0	0.0				
B (stoat)			0.9	1.0	2.5	6.4	0.5	2.5	2.8	1.5				
B (badger)			0.1	0.7	0.0	0.0	0.4	0.0	0.0	0.0				
B (hedgehog)			15.2	0.0	0.0	0.0	13.4	0.0	0.0	0.0				


Figure Captions


Figure 1. The BAM diagram, which depicts the interaction between biotic (**B**), abiotic (**A**), and movement (**M**) factors. Three areas are depicted: **G** the geographical space within which analyses are developed, **G_o** = the occupied distributional area, and **G_i** = the invadable or potential distribution or **BA**. Modified from Soberon (2007).

Figure 2. Conceptual workflow of the methodological steps


Figure 3. Distribution maps for *Ixodes Ricinus* in Ireland and the UK. Models parameterized on a) abiotic variables, b) biotic (hs.ind), c) biotic (pa.ind), d) biotic (hs.ens), e) biotic (pa.ens), f) biotic-abiotic (hs.ind), g) biotic-abiotic (pa.ind), h) biotic-abiotic (hs.ens), and i) biotic-abiotic (pa.ens).

Figure 4. Regularised training gain from the jackknife test of variable importance. The environmental variable with the highest gain has the most useful information when models are parameterised solely on that variable. The environmental variable with the largest decrease in gain when omitted from model parameterisation has the most i






695



696



697

698 nformation that is not present in the other variables.