

Title	Some characterisations of concurrency points in a triangle
Authors	De, Prithwjit;MacHale, Des
Publication date	2025-02-17
Original Citation	De, P. and MacHale, D. (2025) 'Some characterisations of concurrency points in a triangle', The Mathematical Gazette, 109(574), pp.52-58. https://doi.org/10.1017/mag.2025.9
Type of publication	Article (peer-reviewed);journal-article
Link to publisher's version	10.1017/mag.2025.9
Rights	© 2025, The Authors. Published by Cambridge University Press on behalf of The Mathematical Association.
Download date	2026-09-16 08:37:07
Item downloaded from	https://hdl.handle.net/10468/17698



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Some Characterisations of Concurrency Points in a Triangle

In geometry, we sometimes prove that a certain configuration Q has property P . If it turns out that Q is the *only* configuration that has property P , then we say that P is a characterisation of Q . For example, the centre of a circle is a point in the plane which is equidistant from every point on the circumference of the circle. In fact, the centre is the *only* point in the plane which is equidistant from every point on the circumference of a given circle. Thus we say that the centre of a circle is characterised by this property.


In this paper we find characterisations of some concurrency points in a triangle.

Theorem 1

In a triangle ABC , let AX , BY and CZ be cevians concurrent at an internal point M . If

$$\frac{AM}{MX} = \frac{BM}{MY} = \frac{CM}{MZ} = k,$$

then M is the centroid of the triangle and $k = 2$.



Proof

It is easy to see that the ratio of the areas of two triangles with the same perpendicular height is the same as the ratio of the lengths of their bases. Thus, in the diagram, denoting the area of each of the six sub-triangles by the letters p , q , r , s , t and u , we have

$$k = \frac{AM}{MX} = \frac{p+q}{r} = \frac{t+u}{s};$$

$$k = \frac{BM}{MY} = \frac{r+s}{q} = \frac{t+u}{p};$$

and

$$k = \frac{CM}{MZ} = \frac{p+q}{u} = \frac{r+s}{t}.$$


Thus,

$$\frac{p+q}{r} = k = \frac{p+q}{u},$$

so $r = u$. Similarly, $s = p$ and $t = q$. So $u + t + s = r + p + q$ and $\text{Area}(\triangle ABX) = \text{Area}(\triangle AXC)$. Thus AX is a median of $\triangle ABC$. Similarly, BY and CZ are medians, so M is the centroid and $k = 2$.

Lemma 1

In a triangle ABC , let X be a point on BC such that $\frac{BX}{AB} = \frac{CX}{AC}$. Then AX is the bisector of $\angle BAC$.



Proof


If the bisector of $\angle BAC$ meets BC at X' , then

$$\frac{BX'}{AB} = \frac{CX'}{AC}.$$

Thus $BX' : CX' = AB : AC = BX : CX$ whence $X' = X$.

Theorem 2

Let AX , BY and CZ be cevians of a triangle ABC meeting at an internal point M . If $\frac{BX}{CX} = k \frac{AB}{AC}$; $\frac{CY}{AY} = k \frac{BC}{AB}$ and $\frac{AZ}{BZ} = k \frac{AC}{BC}$, then M is the incentre of the triangle and $k = 1$.



Proof

We have

$$\frac{BX}{CX} \cdot \frac{CY}{AY} \cdot \frac{AZ}{BZ} = 1$$

by Ceva's theorem since AX , BY and CZ are concurrent. Thus


$$1 = \left(k \frac{AB}{AC}\right) \left(k \frac{BC}{AB}\right) \left(k \frac{AC}{BC}\right) = k^3.$$

So $k = 1$ and by the lemma, AX , BY and CZ are the angle bisectors and M is the incentre of the triangle.

For the next case, we need another lemma.

Lemma 2

In a triangle ABC let X , Y and Z be internal points of the sides BC , CA and AB respectively such that the perpendiculars to the sides erected at X , Y and Z are concurrent at a point P . Then $AZ^2 + BX^2 + CY^2 = BZ^2 + CX^2 + AY^2$.



Proof

By Pythagoras,

$$\begin{aligned} AZ^2 &= AP^2 - PZ^2; & CX^2 &= PC^2 - PX^2; \\ BX^2 &= BP^2 - PX^2; & AY^2 &= AP^2 - PY^2; \\ CY^2 &= PC^2 - PY^2 & BZ^2 &= BP^2 - PZ^2; \end{aligned}$$

Adding gives the result.


It is clear that this result also holds if P is outside the triangle or on the perimeter, as the reader can see by drawing appropriate diagrams.

Theorem 3

In a triangle ABC let X , Y and Z be internal points of the sides BC , CA and AB respectively such that the perpendiculars to the sides erected at X , Y and Z are concurrent at a point P . If

$$\frac{AZ}{BZ} = \frac{BX}{CX} = \frac{CY}{AY} = k,$$

then $k = 1$ and P is the circumcentre of the triangle.

**Proof**

Denoting the segments AZ , BZ , BX , CX , CY , AY by z , r , x , p , y and q , respectively, for convenience, we have $x = kp$, $y = kq$ and $z = kr$. Then, by Lemma 2,

$$z^2 + x^2 + y^2 = r^2 + p^2 + q^2,$$

so

$$k^2(r^2 + p^2 + q^2) = r^2 + p^2 + q^2.$$

Hence $k = 1$, $p = x$, $q = y$ and $r = z$, forcing P to be the circumcentre of the triangle.

Recall that the Gergonne point of a triangle is the (internal) point of concurrency obtained by joining the vertices to the opposite points of contact of the inscribed circle.


Theorem 4

In a triangle ABC suppose that cevians AX , BY and CZ are concurrent at an internal point P . If

$$\frac{AZ}{AY} = \frac{BX}{BZ} = \frac{CY}{CX} = k,$$

then $k = 1$ and P is the Gergonne point of the triangle.

Proof



We have $AZ = k.AY$; $BX = k.BZ$; $CY = k.CX$. By Ceva's theorem,

$$AY.BZ.CX = AZ.BX.CY = (k.AY)(k.BZ)(k.CX) = k^3(AY.BZ.CX)$$

so $k^3 = 1$ forcing $k = 1$. Thus $AZ = AY$, $BX = BZ$ and $CX = CY$. Let $AZ = x$, $BX = y$ and $CX = z$. Then $x + y + z = s$, where $s = \frac{1}{2}(a + b + c)$ whence $x = s - a$, $y = s - b$ and $z = s - c$, and these are the tangent lengths to the incircle from the vertices. Hence P is the Gergonne point of the triangle.

We conclude by considering the orthocentre.

Theorem 5

In a triangle ABC suppose the cevians AX , BY and CZ are concurrent at a point P . If

$$AP^2 + BC^2 = BP^2 + CA^2 = CP^2 + AB^2$$

then P is the orthocentre of the triangle.

Proof

We use vectors. Let $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$ and $\mathbf{p}$ denote the position vectors of A , B , C and P with respect to an arbitrary origin O . Then the given condition reduces to

$$|\mathbf{p} - \mathbf{a}|^2 + |\mathbf{c} - \mathbf{b}|^2 = |\mathbf{p} - \mathbf{b}|^2 + |\mathbf{a} - \mathbf{c}|^2 = |\mathbf{p} - \mathbf{c}|^2 + |\mathbf{b} - \mathbf{a}|^2$$

whence

$$|\mathbf{p} - \mathbf{a}|^2 - |\mathbf{p} - \mathbf{b}|^2 = |\mathbf{a} - \mathbf{c}|^2 - |\mathbf{c} - \mathbf{b}|^2,$$

that is,

$$(2\mathbf{p} - \mathbf{a} - \mathbf{b}).(\mathbf{b} - \mathbf{a}) = (\mathbf{a} + \mathbf{b} - 2\mathbf{c}).(\mathbf{a} - \mathbf{b}).$$

So,

$$(\mathbf{p} - \mathbf{c}).(\mathbf{b} - \mathbf{a}) = 0$$

implying that CP is perpendicular to AB . In a similar manner we can show that BP is perpendicular to CA and AP is perpendicular to BC . Thus P is the orthocentre of the triangle.

The vector proof removes the obligation of considering three cases where the orthocentre is inside, outside or on the perimeter of the triangle.

Doubtless such characterisations exist for other triangle centres, which the reader might like to investigate.

Prithwijit De
Homi Bhabha Centre for Science Education
Tata Institute of Fundamental Research
V.N.Purav Marg, Mankhurd
Mumbai - 400088, India
email: prithwijit@hbcse.tifr.res.in, de.prithwijit@gmail.com

Des MacHale
School of Mathematical Sciences
University College Cork
Cork, Republic of Ireland
email: d.machale@ucc.ie