

|                      |  |
|----------------------|--|
| Title                | The role of intrinsic and extrinsic factors in shaping alternative migratory tactics and metabolic phenotypes in brown trout   |
| Authors              | Archer, Louise C.  |
| Publication date     | 2019-11  |
| Original Citation    | Archer, L. C. 2019. The role of intrinsic and extrinsic factors in shaping alternative migratory tactics and metabolic phenotypes in brown trout. PhD Thesis, University College Cork. |
| Type of publication  | Doctoral thesis  |
| Rights               | © 2019, Louise C. Archer. - <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>  |
| Download date        | 2025-01-13 20:39:48  |
| Item downloaded from | <a href="https://hdl.handle.net/10468/9996">https://hdl.handle.net/10468/9996</a>  |

Ollscoil na hÉireann, Corcaigh  
**National University of Ireland, Cork**



**The role of intrinsic and extrinsic factors in shaping  
alternative migratory tactics and metabolic  
phenotypes in brown trout**

Thesis presented by

**Louise Clair Archer**

B.Sc. University College Cork  
M.Sc. Imperial College London

for the degree of

**Doctor of Philosophy**

**University College Cork**

**School of Biological, Earth, and Environmental Sciences**

Head of School: Prof. Andrew Wheeler

Supervisors: Dr Thomas Reed and Prof. Phillip McGinnity

November 2019



# Contents

---

|  |            |
|--|------------|
| <b>Declaration.....</b>  | <b>ii</b>  |
| <b>Thesis summary .....</b>  | <b>iii</b> |
| <b>Acknowledgements.....</b>   | <b>v</b>   |
| <b>Thesis structure.....</b>   | <b>vi</b>  |
| <b>Chapter 1. General Introduction .....</b>   | <b>1</b>   |
| The phenomenon of facultative migration .....  | 1          |
| Proximate and ultimate drivers of migration.....   | 2          |
| Overview of life history variation in salmonines .....   | 4          |
| The importance of energy metabolism .....  | 9          |
| Objectives and overview of the thesis .....  | 11         |
| Additional reseach.....  | 12         |
| References .....   | 12         |
| <b>Chapter 2. The interplay between extrinsic and intrinsic factors in<br/>determining migration decisions in brown trout (<i>Salmo trutta</i>): An<br/>experimental study .....</b> | <b>21</b>  |
| Abstract .....   | 22         |
| Introduction.....  | 23         |
| Materials and methods.....   | 28         |
| Results .....  | 37         |
| Discussion .....   | 52         |
| Acknowledgements .....   | 63         |
| References .....   | 63         |
| Supporting Information.....  | 70         |
| <b>Chapter 3. Food and temperature stressors have opposing effects in<br/>determining flexible migration decisions in brown trout (<i>Salmo trutta</i>)<br/>.....</b>                | <b>78</b>  |
| Abstract .....   | 79         |
| Introduction.....  | 80         |
| Materials and Methods.....   | 84         |
| Results .....  | 92         |

|  |            |
|--|------------|
| Discussion .....   | 107        |
| Acknowledgements .....   | 115        |
| References .....   | 115        |
| Supporting Information.....  | 125        |
| <b>Chapter 4. Metabolic traits in brown trout (<i>Salmo trutta</i>) vary in response to food restriction and population background.....</b>  | <b>136</b> |
| Abstract .....   | 137        |
| Introduction.....  | 138        |
| Methods .....  | 143        |
| Results .....  | 152        |
| Discussion .....   | 160        |
| Acknowledgements .....   | 167        |
| References .....   | 168        |
| Supporting Information.....  | 175        |
| <b>Chapter 5. Developmental temperature and population factors affect variation and covariation of metabolic traits, with complex effects on growth rates in brown trout (<i>Salmo trutta</i>) .....</b> | <b>185</b> |
| Abstract .....   | 186        |
| Introduction.....  | 187        |
| Methods .....  | 192        |
| Results .....  | 202        |
| Discussion .....   | 213        |
| Acknowledgements .....   | 220        |
| References .....   | 220        |
| Supporting Information.....  | 231        |
| <b>Chapter 6. General Discussion.....</b>  | <b>240</b> |
| Overview of each chapter.....  | 240        |
| Proximate drivers of facultative migration.....  | 242        |
| Proximate and ultimate factors influencing metabolic traits .....  | 249        |
| Considerations and limitations of the research .....   | 252        |
| Conclusions and future directions.....   | 254        |
| References .....   | 256        |

## Declaration

---

This is to certify that the work I am submitting is my own and has not been submitted for another degree, at either University College Cork or elsewhere. All external references and sources are clearly acknowledged and identified within the contents. I have read and understood the regulations of University College Cork concerning plagiarism.

---

Louise Clair Archer

# Thesis summary

---

Variation among and within populations accounts for a considerable portion of phenotypic diversity produced in nature, and is instrumental to the structure and function of ecosystems. Understanding how, and why, intraspecific diversity persists is essential for predicting and managing the effects of global change, particularly because intraspecific variation may mediate diverse responses to changes in the environment. Alternative phenotypes – i.e. discrete phenotypic variation – can arise from a combination of proximate and ultimate mechanisms. Proximate mechanisms reflect how environmental factors shape phenotypic variation via intermediate physiological processes, which can themselves vary and be decomposed into genetic *versus* environmental components. Ultimate mechanisms concern the evolutionary function of a given phenotype. In this thesis, I explore how proximate and ultimate factors contribute to a particularly striking example of intraspecific diversity: alternative migratory tactics in brown trout *Salmo trutta*.

Brown trout are iconic for the variety of migratory life histories they exhibit; yet fundamental knowledge gaps remain regarding how environmental, physiological, and genetic factors integrate to underpin life history decisions among and within populations. In Chapter 2, I assessed how food restriction and population background influences the expression of migratory tactics in offspring from two populations that naturally differ in anadromy (i.e. sea-migration). Food restriction affected traits related to size and condition, and resulted in a higher frequency of anadromy in both populations, though populations varied in their responses according to the timing of food restriction treatments. While anadromy was overall more frequent in offspring from the naturally anadromous population, the expression of anadromous phenotypes in offspring from a non-anadromous population indicated that migratory tactics might emerge in response to unfavourable environmental conditions causing energetic limitation. In Chapter 3, I further considered proximate mechanisms by exploring how multiple environmental factors (food and temperature)

influence migration. Antagonistic effects of food restriction and increased temperature on condition and size-related traits were not translated at the level of migration tactics, where effects of food restriction and temperature were additive, but opposing (food restriction increased anadromy, whereas temperature decreased anadromy).

I explored how components of metabolic rate – a fundamental determinant of physiological status – varied according to food restriction and population background in Chapter 4. Standard metabolic rate (SMR) was lower in food-deprived fish, while SMR, maximum metabolic rate, and aerobic scope (AS) were higher in offspring from a naturally anadromous population compared to a non-anadromous population. Population-specific effects of food restriction on AS also emerged. I further addressed the causes and consequences of metabolic rate variation in Chapter 5, where I found metabolic traits varied according to both population background and temperature, with important consequences for growth rates (a key fitness-related trait that can shape life histories).

Collectively, these results contribute to our knowledge of how environmental and genetic factors underpin life-history diversity in terms of migratory tactics and physiology. Changes in environmental conditions will likely alter patterns of life-history diversity (mediated by changes in individual physiology) in ways that will also depend on population-specific factors. While predicting the impacts of multi-faceted environmental change will be complex, knowledge of the links between physiology, environment, and ultimately, life history, is crucial for conserving important biodiversity within brown trout, a species that is already in widespread decline due to pervasive global change.



# Acknowledgements

---

I would firstly like to thank my supervisors Tom Reed and Phil McGinnity for supporting, guiding, and encouraging me over the last four years. Tom, the genuine enthusiasm and enjoyment with which you approach research has been a real source of inspiration. You constantly challenged me to develop and improve, for which I am hugely grateful – I learnt so much from chats that inevitably lasted hours, but were instrumental in stimulating this research. Phil, your in-depth knowledge of all things trout has been invaluable to lean on.

This research would truly not have been possible without the incredible efforts of Steve Hutton, who was adept both at keeping our fish alive, and keeping me sane. I just can't thank you enough Steve for all of the long days, weekends, and late night emergencies in the tank room. Thanks also to Luke Harman for the massive work in designing, building, and maintaining our recirculation system. I always appreciated your willingness to lend a hand, or share advice on the intricacies of fish husbandry/rugby! I am also grateful to the UCC FishEye research group (especially Ronan, Rob, Peter, Jamie, Joe, and Tom Cross) for help in the lab and the tank room, or for general salmonid chat.

Our efforts at UCC build upon a wealth of salmonid research that has been underway in Ireland long before I arrived on the scene in Cork. Staff at the Marine Institute, Burrishoole and Inland Fisheries Ireland facilitated and assisted us in adding our bit. Co-authors Russell Poole (MI), Paddy Gargan (IFI), Michael O'Grady and Joe Kerry (UCC Food Science), and Steve McCormick (USGS) provided research support over the course of the PhD.

This process would have been so much harder without a fantastic group of friends and colleagues in the School of BEES. Adam, Amy, Enrico, Emma, Gavin, Iván, and Maria provided support, laughter, and ridicule in equal measure. My family and friends outside of UCC have been incredibly patient as I devoted much of the last few years to trout antics. Thanks for understanding and sticking with me. Darío, thank you for everything else – you kept me together.

## **Thesis structure**

---

The data chapters within this thesis are intended for publication in peer-reviewed journals and are thus written and formatted as standalone manuscripts. For ease of reading, figures and tables are embedded within the text and chapters are cross-referenced where appropriate, though each chapter can be read in isolation. I have indicated where chapters have already been submitted or published in peer-reviewed journals, and author contributions are listed at the beginning of each chapter.

# Chapter 1

---

## General Introduction

Much of the widespread diversity we observe in nature can be attributed to intra-specific phenotypic variation (Roff 1996). This intra-specific variation can be either continuous in kind, or discontinuous, i.e. discrete variation in morphological, behavioural or life history traits referred to as ‘alternative phenotypes’. Alternative phenotypes are often underpinned by developmental plasticity as polyphenisms (Suzuki and Nijhout 2008), and have attracted considerable attention in evolutionary biology, *e.g.* in the context of speciation, or evolution by genetic accommodation (West-Eberhard 2003; Crispo 2007; Oliveira *et al.* 2008). Alternative phenotypes also play key roles in community ecology (Bolnick *et al.* 2011) and have applied relevance for conservation and ecosystem management (Naish and Hard 2008).

A fundamental goal of evolutionary ecology is to determine the relationship between ecologically relevant phenotypic variation and underlying genetic variation, and in doing so understand how and why divergent phenotypes and life histories arise and persist (Debat and David 2001). This challenge is especially important in the context of pervasive global change, where predicting the outcomes of environmental change on natural populations is made more complex by the occurrence of alternative phenotypes. Phenotypic variation among individuals and populations may mediate diverse responses to environmental change, and understanding the effects of phenotypic diversity is crucial to successful management and conservation of natural populations (Schindler *et al.* 2010, 2015).

### 25 The phenomenon of facultative migration

Considerable phenotypic diversity can stem from migration, a spectacular, yet relatively common phenomenon evident in all major animal taxa (Swingland and Greenwood 1984). Optimal breeding and feeding habitats are often

separated in space and time, and thus migration is a pivotal factor shaping the temporal and spatial distribution of animals, with important consequences for ecosystem function and structure (Polis *et al.* 2004; Janetski *et al.* 2009). In addition to considerable inter-specific variation in migratory patterns, encompassing obligate migratory and non-migratory lifestyles, strikingly different migratory phenotypes are also evident within species (Dingle and Drake 2007), with migration usually occurring at specific life stages to exploit alternative foraging opportunities or to avoid unfavourable abiotic conditions (Chapman *et al.* 2011a). Intraspecific variation in migratory tendency can span populations that may be entirely migratory, others that may be solely resident, and others again that may be “partially migratory” and comprise a mix of migratory and resident phenotypes (Lack 1943; Lundberg 1987; Kaitala *et al.* 1993; Chapman *et al.* 2011a). Alternative phenotypes among and within populations might be determined purely by genetics, plasticity, or reflect a mix of genetic and environmental factors, thus appearing somewhat “flexible”. Facultative migration – where individuals show flexibility in tactic expression – is common among taxa (Chapman *et al.* 2011c, b), with well documented examples in birds (Lundberg 1988; Pulido *et al.* 1996; Newton 2008), ungulates (Ball *et al.* 2001; Cagnacci *et al.* 2011; Hebblewhite and Merrill 2011), zooplankton (Hansson and Hylander 2009), and fishes (Chapman *et al.* 2012; Dodson *et al.* 2013). However, despite the widespread occurrence of facultative migration, and its importance in ecosystem processes (Brodersen *et al.* 2008), the causes and consequences of the migration *versus* residency decision remain largely unresolved (Dingle and Drake 2007; Pulido 2011).

### **Proximate and ultimate drivers of migration**

The origin and persistence of intra-specific variation in migratory behaviour can be understood in terms of proximate and ultimate factors. Ultimate factors refer to the evolutionary reasons for, or phylogenetic patterns underpinning, migration, whereas proximate factors reflect how migration is triggered in response to environmental cues and the ontogeny of its expression. The proximate mechanisms underpinning migration are themselves evolvable, and

60 may be subject to selection in response to environmental factors. As such, alternative migratory phenotypes are likely underpinned by a complex genotype to phenotype mapping, where similar phenotypes can arise from different genotypes, or the same genotype can produce dramatically different phenotypes, via plasticity mediated by environmental cues (Via *et al.* 1995).

65 Facultative migration is often considered as a threshold trait (Roff 1996). As such, environmentally-triggered alternative migratory “tactics” are produced by a conditional strategy, where the optimal tactic for a given environment is conditional on intrinsic or extrinsic cues. Under the framework of the “environmentally cued threshold model” (Tomkins and Hazel 2007),

70 expression of alternative tactics is determined by the relationship between an environmentally-sensitive “status” trait (that may itself be influenced by genes *e.g.* physiological condition, or energy status) and a genetically variable threshold for said status trait. The inherited threshold can be thought of as a switch point. If an individual’s status trait exceeds the switch point during an

75 assessment period (or “decision window”), it remains resident and undergoes maturation. If the threshold for residency is not met, a migratory trajectory is adopted (though the timing of actual migration may be controlled by similar threshold mechanisms). Since environmental conditions are assumed to strongly influence potential status traits, environmentally-induced variation in

80 the status trait may cause alternative migratory tactic to arise from similar genetic thresholds via life-history plasticity. The context-dependent status trait can thus be considered as an environmental “cue”, related to the expression of alternative tactics by a “threshold reaction norm” (Tomkins and Hazel 2007; Piche *et al.* 2008; Pulido 2011; Buoro *et al.* 2012). Alternatively, since the

85 underlying threshold is evolvable (*i.e.* an ultimate mechanism), population or individual-level genetic variation in the threshold may result in similar environments producing different migratory phenotypes, as shown in blackcaps *Sylvia atricapilla* (Pulido *et al.* 1996) and Atlantic salmon *Salmo salar* (Piche *et al.* 2008).

90 The environmentally cued threshold model provides a useful framework for understanding how the frequency of migration *versus* residency for a given population depends on both the distribution of the status trait, and the distribution of the threshold values in the population (Tomkins and Hazel 2007). The prevalence of a given migratory phenotype can vary in the short  
 95 term in response to environmental conditions, and can shift over longer periods through evolution of the underlying threshold (Piche *et al.* 2008). In practice, however, it is not yet clear how environmental variation is translated into internal physiological signals, upon which individuals then base migratory decisions. Understanding the proximate determinants of phenotypic diversity  
 100 i.e. how genetic and environmental factors interactively shape life histories via intermediary physiological processes, is thus the overarching theme of this thesis.


### Overview of life history variation in salmonines

Fishes present many interesting examples of facultative migration, offering  
 105 valuable opportunities to gain insight into the mechanisms underlying alternative migratory tactics (AMTs) (Jonsson and Jonsson 1993; Chapman *et al.* 2012). Salmonines (salmons, trouts, and charrs) in particular display a multitude of diverse migratory life histories that vary among populations and individuals in terms of migration propensity, migration distance, and migration  
 110 destination (Dodson *et al.* 2013; Sloat *et al.* 2014; Kendall *et al.* 2014; Ferguson *et al.* 2019). Alternative reproductive tactics (ARTs) within male salmonines (*e.g.* the sneaker versus ‘bourgeois’ tactics) represent another interesting class of alternative phenotypes that are conceptually distinct from, but nonetheless related to, AMTs (Gross 1985, 1991; Hutchings and Myers 1994; Fleming 1996).  
 115 While freshwater spawning is obligate, a continuum of migration tactics is seen in salmonines, encompassing individuals that remain in natal streams/lakes for their entire life cycles (residency), to those that migrate to larger rivers (fluvial-adfluvial migration), lakes (potomodromy), or to marine environments (anadromy) before returning to spawn in natal freshwater systems (Klemetsen  
 120 *et al.* 2003; Ferguson *et al.* 2019; Nevoux *et al.* 2019). In several salmonines

(including brown trout *Salmo trutta*, the focal species of this thesis), populations may be dominated by either resident or migratory forms, but may also comprise a mixture of migratory phenotypes that can breed freely in sympatry (Chapman *et al.* 2012). Despite the considerable socioeconomic and cultural importance of various migratory phenotypes, (e.g. as a valued angling resource or a component of biodiversity), and their pivotal role in aquatic ecosystem dynamics (Naiman *et al.* 2002), the drivers of alternative life histories in salmonines remain largely unresolved (Harris and Milner 2008; Harris 2017). Such fundamental knowledge gaps limit our ability to manage and conserve these iconic species, which are in widespread decline due to anthropogenic pressures, most notably global change, in-stream barriers, and the development of aquaculture (Limburg and Waldman 2009).

The threshold reaction norm concept has been applied to understand migration decisions (or ARTs) in salmonines because their life histories appear largely compatible with the model framework (Hutchings and Myers 1994; Thorpe *et al.* 1998; Thériault *et al.* 2007). While offspring tend to show similar migratory tactics to their parents, either migratory phenotype can be produced from a given parental life history (Zimmerman and Reeves 2000; Berejikian *et al.* 2014). Since mounting evidence supports migration as a trait interactively determined by genotype and environment, i.e. under genotype-by-environmental control (Hutchings 2011), research has focused on identifying potential status traits, with inconclusive results. Previous studies have linked numerous aspects of physiological condition to migration tactics with equivocal support, including: body size (Thériault and Dodson 2003), body condition (Hecht *et al.* 2015), growth rates (Jonsson 1985), growth efficiency (Forseth *et al.* 1999; Morinville and Rasmussen 2003), metabolic rates (Sloat and Reeves 2014), and lipid stores (Jonsson and Jonsson 2005). Furthermore, the associations between these physiological traits and migration tendency have also been inconsistent. For example, fast growth and larger sizes have been both positively (Jonsson 1985; Acolas *et al.* 2012), and negatively (Morinville and Rasmussen 2003; McMillan *et al.* 2012) related to migration whilst others have

found no relationship (at a given age) (Thériault and Dodson 2003), or evidence for population-specific responses (Jonsson 1985).



**Figure 1:** Overview of alternative migratory tactics in brown trout, with an andromous life history highlighted in red, and a resident life history in blue. The inset shows the threshold mechanism underpinning migration decisions. A normal distribution of status trait values in a given environment is shown for a hypothetical population that displays facultative anadromy, with the mean threshold for residency represented by the dashed line.

Inconsistent results could occur if there are discrepancies in the timing of trait measurements (often occurring around the time of migration) and when the



migration *versus* residency decision is actually made. Migration often requires considerable morphological and physiological adjustments (Tanguy *et al.* 1994), meaning the migration decision window likely occurs many months in advance of the actual migration event. For example, future migrants must reach  
 160 sufficient sizes to successfully transition to the marine environment, where survival is size-dependent (Phillis *et al.* 2016; Armstrong *et al.* 2018). Thus, pre-migrants often display accelerated growth in advance of the migration period (Metcalf *et al.* 1995), while residents redirect energy reserves towards gonadal development and sexual maturation (Tocher 2003; Jonsson and Jonsson 2005).  
 165 Fewer studies have explored *when* the migratory decision might occur, but it is increasingly clear that size at migration may not reflect size (or other aspects of physiological condition) at the time when the decision was made. Indeed, there is evidence that migrants and residents can differ in potential status traits including growth rates (Beakes *et al.* 2010), metabolic costs (Morinville and  
 170 Rasmussen 2003), and condition (Hecht *et al.* 2015) up to a full year before migration takes place (Thorpe *et al.* 1998; Thorpe and Metcalfe 1998; Satterthwaite *et al.* 2009). The timing and nature of the decision window is largely unknown in brown trout (Ferguson *et al.* 2019), but is fundamental to our understanding of the pathways by which alternative tactics develop (Thorpe  
 175 *et al.* 1998)

While identification of underlying status traits has proved somewhat inconclusive, understanding the fitness costs and benefits of migration *versus* residency may help to clarify the factors that promote different migratory tactics. Since food abundance is often higher in the migratory destination  
 180 (larger rivers, lakes or the sea) compared to food-limited natal streams (Imre *et al.* 2005), migration might be favoured as an optimal tactic if increased growth rates are achieved in the new habitat, translating into higher fecundity and ultimately fitness (Brönmark *et al.* 2013). Such a scenario is supported by latitudinal clines in the prevalence of anadromy, where at high latitudes, higher  
 185 productivity of the marine environment relative to freshwaters has been linked to the evolution of the anadromous life history (Gross *et al.* 1988). However, any

feeding benefits must be traded off against the significant costs associated with migration, including considerable energetic expenditure (Stefansson *et al.* 2003), physiological stress (Peiman *et al.* 2017; Birnie-Gauvin *et al.* 2017), and  
 190 an increased risk of predation (Dieperink *et al.* 2002).

The balance between migratory costs and benefits is not necessarily equal among individuals, and is particularly likely to vary between the sexes (Hendry *et al.* 2004). While larger females have larger eggs and are more fecund (Fleming 1996; Quinn 2018), male reproductive success is limited by access to  
 195 mates (Fleming 1998), which is less size dependent because the success of ARTs (*e.g.* “sneaker” versus bourgeois, anadromous males) can be similar (Hutchings and Myers 1988; Foote *et al.* 1997; Young *et al.* 2013). Females thus often receive greater fitness benefits from migration (mediated by larger sizes at maturity) than males, reflected by general trends of female-biased migration (Nielsen *et al.* 2003; Rundio *et al.* 2012; Ohms *et al.* 2013; García-Vega *et al.* 2018; Kelson *et al.* 2019).  
 200

This balance of trade-offs may show further variation due to intraspecific differences in energetic uptake and output, with accumulating evidence to suggest that migration occurs in response to energetic limitation in the natal  
 205 environment (Forseth *et al.* 1999). The acquisition and allocation of energy resources in juveniles is strongly influenced by extrinsic environmental conditions, with the resulting individual energetic status (indicated by various potential status traits) determining life history trajectories in juveniles (Jonsson and Jonsson 1993). In particular, factors determining food availability have been  
 210 much linked to the frequency of migrants (O’Neal and Stanford 2011; Jones *et al.* 2015), *e.g.* increased competition at high population densities has been shown to influence adfluvial migration in brown trout (Olsson *et al.* 2006; Wysujack *et al.* 2009). Temperature appears to be an important abiotic factor affecting migration (Kendall *et al.* 2014), and cooler temperatures have been  
 215 found to increase maturation *in lieu* of migration in steelhead trout, perhaps due to changes in lipid deposition (Sloat and Reeves 2014). However, we currently lack a synthetic understanding of how variable environmental

conditions such as food and temperature might mediate migratory tactics among populations and individuals through effects on physiological condition or energetic status.

### **The importance of energy metabolism**

As the fundamental process governing an individual's energy budget, metabolism is likely to be profoundly linked to variation in migratory phenotypes. Standard metabolic rate (SMR) refers to the minimum energy required to sustain life, i.e. the costs of homeostasis and tissue maintenance in an inactive, unstressed, non-digestive organism (termed basal metabolic rate (BMR) in endotherms) (Chabot *et al.* 2016). Once size and age are accounted for, SMR can vary as much as three-fold among individuals of the same population (Burton *et al.* 2011; Konarzewski and Książek 2013). Such individual differences in SMR are integrally linked to variation in energy acquisition and allocation, making it a trait of particular interest in studies of life-history variation (Forseth *et al.* 1999; Burton *et al.* 2011; Metcalfe *et al.* 2016). Variation in SMR has been generally linked to differences in lifestyle among fishes (Killen *et al.* 2010), and intraspecific variation appears to be similarly linked to life-history differences. For example, in brown trout, juveniles with relatively higher baseline energetic demands subsequently adopted migratory tactics (Forseth *et al.* 1999). Migrating Atlantic salmon *S. salar* smolts were found to have higher SMR values than non-smolts deferring migration to a later age (Seppänen *et al.* 2010), and those with the highest recorded SMR also tended to migrate at younger ages (McCarthy 2000). Similarly, lower SMR was linked to a higher probability of freshwater maturation rather than migration in steelhead trout *Oncorhynchus mykiss* (Sloat and Reeves 2014).

While SMR defines the minimum energy demands, or metabolic “floor”, maximum metabolic rate (MMR) defines the upper limits of metabolism as the highest rate of aerobic metabolism (oxygen transport and ATP production) that can be achieved (Norin and Clark 2016). An individual's aerobic scope (AS) is bounded by SMR and MMR, and reflects the potential energy that can be directed towards key competing functions (*e.g.* activity/locomotion, growth,

digestion) once baseline energy requirements have been met (Guderley and  
 250 Pörtner 2010). MMR and AS appear to show similar levels of intraspecific  
 variation to SMR (Metcalf *et al.* 2016), but the reasons behind this variation are  
 somewhat unclear. Previous studies in salmonines have linked variation in  
 MMR and AS to swimming performance (Tudorache *et al.* 2007) and migration  
 distance and effort (Eliason *et al.* 2011), supporting it as a trait of ecological  
 255 relevance in facultatively migratory species. As such, MMR and AS may have  
 implications for life histories, and could contribute to the emergence and  
 persistence of alternative phenotypes. However, despite their relevance for life  
 history and phenotypic diversity (Metcalf *et al.* 2016; Biro *et al.* 2018), MMR  
 and AS are relatively under-researched traits in relation to facultative migration.  
 260 We know surprisingly little about how variation in, or indeed covariation  
 between, these metabolic traits relates to patterns of alternative migratory  
 tactics.

Since metabolic traits are strongly influenced by environmental conditions such  
 as temperature (Clarke and Johnston 1999; Brown *et al.* 2016), it is crucial to  
 265 also consider environmental factors when investigating the causes and  
 consequences of metabolic rate variation in facultatively migratory species. For  
 example, although empirical evidence suggests SMR is positively associated  
 with migration because of its influence on energy status (McCarthy 2000), this  
 relationship may be context-dependent (Burton *et al.* 2011), *i.e.* energetic  
 270 limitation may only occur at warm temperatures, or if there is insufficient food  
 to meet metabolic needs. Additionally, individuals or populations may also  
 differ in their metabolic responses to changing extrinsic conditions (Metcalf *et al.*  
*et al.* 2016; Norin and Metcalf 2019), further complicating the links between  
 metabolism, environment, and life history. Moreover, since metabolic rates are  
 275 partly determined by genetics (Pettersen *et al.* 2018) and are thus evolvable  
 (Wone *et al.* 2015; Sadowska *et al.* 2015), metabolic trait variation between  
 migratory phenotypes may arise due to the demands of a given life history. For  
 example, in populations that have a high incidence of migration (where the  
 thresholds for residency have presumably evolved to be relatively high),

280 selection may favour higher metabolic rates that improve migration performance (Dalziel *et al.* 2012a, b).

While metabolic components are assumed to have important implications for fitness, and thus shape life histories, empirical evidence linking metabolism to fitness-rated traits has provided mixed results. For example, higher metabolic  
 285 rates have been positively (McCarthy 2000) and negatively linked to growth rates (Norin and Malte 2011), whereas others again have found the optimal metabolic rate to vary depending on food availability (Reid *et al.* 2011; Auer *et al.* 2015a, b) and habitat (Reid *et al.* 2012). Thus, it seems increasingly likely that the benefits of a given metabolic phenotype may be context-specific, and  
 290 dependant on environmental conditions, or intrinsic factors related to population background and life history (Álvarez and Nicieza 2005; Robertsen *et al.* 2014). Further investigation into the links between metabolic trait variation, environmental conditions, and fitness will help to illuminate how these factors integrate to shape life histories.

## 295 **Objectives and overview of the thesis**

The overarching objective of this thesis was thus to explore the nexus between physiology, environment, and alternative life history tactics in a facultatively migratory species. In Chapter 2, I specifically aimed to investigate how expression of migratory tactics is mediated by interactions between food  
 300 availability and population background, and is underpinned by various aspects of physiological condition. I also explored whether the timing of food restriction influences migration, in order to better clarify when in early life the migratory decision might be undertaken, and to assess if populations varied in their responses to the timing of food restriction.

305 In Chapter 3, I extend the study of the proximate drivers of life history variation by exploring how co-occurring environmental factors collectively influence migratory tactics. Here, I aimed to test if food restriction and increased temperature interactively influence traits associated with physiological

condition, and to investigate whether these effects scale up to affect migratory  
 310 tactics in a similar fashion, in a facultatively migratory population.

Chapter 4 explores how intrinsic and extrinsic factors influence variation in  
 metabolic traits in facultatively migratory populations. Specifically, I aimed to  
 assess (i) how SMR, MMR, and AS vary in brown trout offspring from two  
 populations (that naturally differ in migration tendency); and (ii) to test how  
 315 metabolic traits respond to long-term conditions of food restriction. I also  
 explored if metabolic responses to food restriction are dependent on population  
 background and sex.

The objectives of Chapter 5 were to explore the causes and consequences of  
 variation in metabolic traits in response to increased temperature. Specifically,  
 320 I aimed to assess whether populations that differ in migratory tactics might also  
 show variable metabolic responses to temperature increases. I then explored the  
 implications of metabolic trait variation (in terms of differences between  
 populations, and variation within populations driven by plastic responses to  
 warming) for growth rates, a key-fitness related trait.

325 Chapter Six synthesises the results of the studies described above, and discusses  
 their contribution towards developing an integrated understanding of the links  
 between physiology, environment, and life history.

### **Additional research**

In addition to the chapters presented in this thesis, I have also been involved in  
 330 the following research during my studies:

Archer LC, Sohlström EH, Gallo B, Jochum M, Woodward G, Kordas RL, Rall  
 BC, and O’Gorman EJ. 2019. Consistent temperature dependence of functional  
 response parameters and their use in predicting population abundance. *Journal*  
*of Animal Ecology* **In Press**. <https://doi.org/10.1111/1365-2656.13060>.

### **References**

Acolas ML, Labonne J, Baglinière JL, and Roussel JM. 2012. The role of body size versus  
 growth on the decision to migrate: A case study with *Salmo trutta*.  
*Naturwissenschaften* **99**: 11–21.

- 340 Álvarez D and Nicieza AG. 2005. Is metabolic rate a reliable predictor of growth and survival of brown trout (*Salmo trutta*) in the wild? *Canadian Journal of Fisheries and Aquatic Sciences* **62**: 643–9.
- Armstrong JD, McKelvey S, Smith GW, Rycroft P, and Fryer RJ. 2018. Effects of individual variation in length, condition and run-time on return rates of wild-reared Atlantic salmon *Salmo salar* smolts. *Journal of Fish Biology* **92**: 569–78.
- 345 Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015a. The optimal combination of standard metabolic rate and aerobic scope for somatic growth depends on food availability. *Functional Ecology* **29**: 479–86.
- Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015b. Flexibility in metabolic rate confers a growth advantage under changing food availability.  
350 *Journal of Animal Ecology* **84**: 1405–11.
- Ball JP, Nordengren C, and Wallin K. 2001. Partial migration by large ungulates: Characteristics of seasonal moose *Alces alces* ranges in northern Sweden. *Wildlife Biology* **7**: 39–47.
- Beakes MP, Satterthwaite WH, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, and Mangel M. 2010. Smolt transformation in two California steelhead populations: Effects of temporal variability in growth. *Transactions of the American Fisheries Society* **139**: 1263–75.
- 355 Berejikian BA, Bush RA, and Campbell LA. 2014. Maternal control over offspring life history in a partially anadromous species, *Oncorhynchus mykiss*. *Transactions of the American Fisheries Society* **143**: 369–79.
- 360 Birnie-Gauvin K, Peiman KS, Larsen MH, Baktoft H, Aarestrup K, Willmore WG, and Cooke SJ. 2017. Oxidative stress and partial migration in brown trout (*Salmo trutta*). *Canadian Journal of Zoology* **95**: 829–35.
- Biro PA, Garland T, Beckmann C, Ujvari B, Thomas F, and Post JR. 2018. Metabolic scope as a proximate constraint on individual behavioral variation: Effects on personality, plasticity, and predictability. *The American Naturalist* **192**: 142–54.
- 365 Bolnick DI, Amarasekare P, Araújo MS, Bürger R, Levine JM, Novak M, Rudolf VHW, Schreiber SJ, Urban MC, and Vasseur DA. 2011. Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution* **26**: 183–92.
- 370 Brodersen J, Ådahl E, Brönmark C, and Hansson L-A. 2008. Ecosystem effects of partial fish migration in lakes. *Oikos* **117**: 40–6.
- Brönmark C, Hulthén K, Nilsson PA, Skov C, Hansson L-A, Brodersen J, and Chapman BB. 2013. There and back again: Migration in freshwater fishes. *Canadian Journal of Zoology* **92**: 467–79.
- 375 Brown JH, Gillooly JF, Allen AP, Savage VM, and West GB. 2016. Toward a metabolic theory of ecology. *Ecology*: 1771–89.
- Buoro M, Gimenez O, and Prévost E. 2012. Assessing adaptive phenotypic plasticity by means of conditional strategies from empirical data: The latent environmental threshold model. *Evolution* **66**: 996–1009.
- 380 Burton T, Killen SS, Armstrong JD, and Metcalfe NB. 2011. What causes intraspecific variation in resting metabolic rate and what are its ecological consequences? *Proceedings of the Royal Society B: Biological Sciences* **278**: 3465–73.

- Cagnacci F, Focardi S, Heurich M, Stache A, Hewison AJM, Morellet N, Kjellander P, Linnell JDC, Mysterud A, Neteler M, Delucchi L, Ossi F, and Urbano F. 2011. Partial migration in roe deer: Migratory and resident tactics are end points of a behavioural gradient determined by ecological factors. *Oikos* **120**: 1790–802.
- Chabot D, McKenzie DJ, and Craig JF. 2016. Metabolic rate in fishes: Definitions, methods and significance for conservation physiology. *Journal of Fish Biology* **88**: 1–9.
- Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011a. Partial migration: An introduction. *Oikos* **120**: 1761–3.
- Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011b. The ecology and evolution of partial migration. *Oikos* **120**: 1764–75.
- Chapman BB, Hulthén K, Blomqvist DR, Hansson L-A, Nilsson J-Å, Brodersen J, Nilsson PA, Skov C, and Brönmark C. 2011c. To boldly go: Individual differences in boldness influence migratory tendency. *Ecology Letters* **14**: 871–6.
- Chapman BB, Hulthén K, Brodersen J, Nilsson PA, Skov C, Hansson L-A, and Brönmark C. 2012. Partial migration in fishes: Causes and consequences. *Journal of Fish Biology* **81**: 456–78.
- Clarke A and Johnston NM. 1999. Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology* **68**: 893–905.
- Crispo E. 2007. The Baldwin effect and genetic assimilation: Revisiting two mechanisms of evolutionary change mediated by phenotypic plasticity. *Evolution* **61**: 2469–79.
- Dalziel AC, Ou M, and Schulte PM. 2012a. Mechanisms underlying parallel reductions in aerobic capacity in non-migratory threespine stickleback (*Gasterosteus aculeatus*) populations. *Journal of Experimental Biology* **215**: 746–59.
- Dalziel AC, Vines TH, and Schulte PM. 2012b. Reductions in prolonged swimming capacity following freshwater colonization in multiple threespine stickleback populations. *Evolution* **66**: 1226–39.
- Debat V and David P. 2001. Mapping phenotypes: Canalization, plasticity and developmental stability. *Trends in Ecology & Evolution* **16**: 555–61.
- Dieperink C, Bak BD, Pedersen L-F, Pedersen MI, and Pedersen S. 2002. Predation on Atlantic salmon and sea trout during their first days as postsmolts. *Journal of Fish Biology* **61**: 848–52.
- Dingle H and Drake VA. 2007. What is migration? *BioScience* **57**: 113–21.
- Dodson JJ, Aubin-Horth N, Thériault V, and Páez DJ. 2013. The evolutionary ecology of alternative migratory tactics in salmonid fishes: Alternative migratory tactics as threshold traits. *Biological Reviews* **88**: 602–25.
- Eliason EJ, Clark TD, Hague MJ, Hanson LM, Gallagher ZS, Jeffries KM, Gale MK, Patterson DA, Hinch SG, and Farrell AP. 2011. Differences in thermal tolerance among sockeye salmon populations. *Science* **332**: 109–12.
- Ferguson A, Reed TE, Cross TF, McGinnity P, and Prodöhl PA. 2019. Anadromy, potamodromy and residency in brown trout *Salmo trutta*: The role of genes and the environment. *Journal of Fish Biology* **95**: 692–718.



- Fleming IA. 1996. Reproductive strategies of Atlantic salmon: Ecology and evolution. *Reviews in Fish Biology and Fisheries* **6**: 379–416.
- 430 Fleming IA. 1998. Pattern and variability in the breeding system of Atlantic salmon (*Salmo salar*), with comparisons to other salmonids. *Canadian Journal of Fisheries and Aquatic Sciences* **55**: 59–76.
- Foote CJ, Brown GS, and Wood CC. 1997. Spawning success of males using alternative mating tactics in sockeye salmon, *Oncorhynchus nerka*. *Canadian Journal of Fisheries and Aquatic Sciences* **54**: 1785–95.
- 435 Forseth T, Nesje TF, Jonsson B, and Hårsaker K. 1999. Juvenile migration in brown trout: A consequence of energetic state. *Journal of Animal Ecology* **68**: 783–93.
- García-Vega A, Sanz-Ronda FJ, Fernandes Celestino L, Makrakis S, and Leunda PM. 2018. Potamodromous brown trout movements in the North of the Iberian Peninsula: Modelling past, present and future based on continuous fishway monitoring. *Science of The Total Environment* **640–641**: 1521–36.
- 440 Gross MR. 1985. Disruptive selection for alternative life histories in salmon. *Nature* **313**: 47–8.
- Gross MR. 1991. Salmon breeding behavior and life history evolution in changing environments. *Ecology* **72**: 1180–6.
- 445 Gross MR, Coleman RM, and McDowall RM. 1988. Aquatic productivity and the evolution of diadromous fish migration. *Science* **239**: 1291–3.
- Guderley H and Pörtner HO. 2010. Metabolic power budgeting and adaptive strategies in zoology: Examples from scallops and fish. *Canadian Journal of Zoology* **88**: 753–63.
- 450 Hansson L-A and Hylander S. 2009. Size-structured risk assessments govern *Daphnia* migration. *Proceedings of the Royal Society B: Biological Sciences* **276**: 331–6.
- Harris G. 2017. Sea Trout: Biology, Conservation and Management: Proceedings of the 2nd International Sea Trout Symposium. Leicestershire, UK: Matador.
- Harris G and Milner N. 2008. Sea Trout: Biology, Conservation and Management. Oxford, UK: Blackwells Scientific Publications.
- 455 Hebblewhite M and Merrill EH. 2011. Demographic balancing of migrant and resident elk in a partially migratory population through forage–predation tradeoffs. *Oikos* **120**: 1860–70.
- 460 Hecht BC, Hard JJ, Thrower FP, and Nichols KM. 2015. Quantitative genetics of migration-related traits in rainbow and steelhead trout. *G3: Genes, Genomes, Genetics* **5**: 873–89.
- Hendry AP, Bohlin T, Jonsson B, and Berg OK. 2004. To sea or not to sea? Anadromy v. non-anadromy in salmonids. In: Evolution illuminated - Salmon and their relatives. Edited by A.P. Hendry & S. C. Sterns. New York, NY: Oxford University Press.
- 465 Hutchings JA. 2011. Old wine in new bottles: Reaction norms in salmonid fishes. *Heredity* **106**: 421–37.
- Hutchings JA and Myers RA. 1988. Mating success of alternative maturation phenotypes in male Atlantic salmon, *Salmo salar*. *Oecologia* **75**: 169–74.

- Hutchings JA and Myers RA. 1994. The evolution of alternative mating strategies in variable environments. *Evolutionary Ecology* **8**: 256–268.
- Imre I, Grant JWA, and Cunjak RA. 2005. Density-dependent growth of young-of-the-year Atlantic salmon *Salmo salar* in Catamaran Brook, New Brunswick. *Journal of Animal Ecology* **74**: 508–16.
- Janetski DJ, Chaloner DT, Tiegs SD, and Lamberti GA. 2009. Pacific salmon effects on stream ecosystems: A quantitative synthesis. *Oecologia* **159**: 583–95.
- Jones DA, Bergman E, and Greenberg L. 2015. Food availability in spring affects smolting in brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 1694–9.
- Jonsson B. 1985. Life history patterns of freshwater resident and sea-run migrant brown trout in Norway. *Transactions of the American Fisheries Society* **114**: 182–94.
- Jonsson B and Jonsson N. 1993. Partial migration: Niche shift versus sexual maturation in fishes. *Reviews in Fish Biology and Fisheries* **3**: 348–65.
- Jonsson B and Jonsson N. 2005. Lipid energy reserves influence life-history decision of Atlantic salmon (*Salmo salar*) and brown trout (*S. trutta*) in fresh water. *Ecology of Freshwater Fish* **14**: 296–301.
- Kaitala A, Kaitala V, and Lundberg P. 1993. A theory of partial migration. *The American Naturalist* **142**: 59–81.
- Kelson SJ, Miller MR, Thompson TQ, O'Rourke SM, and Carlson SM. 2019. Do genomics and sex predict migration in a partially migratory salmonid fish, *Oncorhynchus mykiss*? *Canadian Journal of Fisheries and Aquatic Sciences* **76**: 2080–8.
- Kendall NW, McMillan JR, Sloat MR, Buehrens TW, Quinn TP, Pess GR, Kuzishchin KV, McClure MM, and Zabel RW. 2014. Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): A review of the processes and patterns. *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 319–42.
- Killen SS, Atkinson D, and Glazier DS. 2010. The intraspecific scaling of metabolic rate with body mass in fishes depends on lifestyle and temperature. *Ecology Letters* **13**: 184–93.
- Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, and Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): A review of aspects of their life histories. *Ecology of Freshwater Fish* **12**: 1–59.
- Konarzewski M and Książek A. 2013. Determinants of intra-specific variation in basal metabolic rate. *Journal of Comparative Physiology B* **183**: 27–41.
- Lack D. 1943. The problem of partial migration. *British Birds* **37**: 122–30.
- Limburg KE and Waldman JR. 2009. Dramatic declines in north Atlantic diadromous fishes. *BioScience* **59**: 955–65.
- Lundberg P. 1987. Partial bird migration and evolutionarily stable strategies. *Journal of Theoretical Biology* **125**: 351–60.
- Lundberg P. 1988. The evolution of partial migration in birds. *Trends in Ecology & Evolution* **3**: 172–5.

- McCarthy ID. 2000. Temporal repeatability of relative standard metabolic rate in juvenile Atlantic salmon and its relation to life history variation. *Journal of Fish Biology* **57**: 224–38.
- 515 McMillan JR, Dunham JB, Reeves GH, Mills JS, and Jordan CE. 2012. Individual condition and stream temperature influence early maturation of rainbow and steelhead trout, *Oncorhynchus mykiss*. *Environmental Biology of Fishes* **93**: 343–55.
- 520 Metcalfe NB, Taylor AC, and Thorpe JE. 1995. Metabolic rate, social status and life-history strategies in Atlantic salmon. *Animal Behaviour* **49**: 431–6.
- Metcalfe NB, Van Leeuwen TE, and Killen SS. 2016. Does individual variation in metabolic phenotype predict fish behaviour and performance? *Journal of Fish Biology* **88**: 298–321.
- 525 Morinville GR and Rasmussen JB. 2003. Early juvenile bioenergetic differences between anadromous and resident brook trout (*Salvelinus fontinalis*). *Canadian Journal of Fisheries and Aquatic Sciences* **60**: 401–10.
- Naiman RJ, Bilby RE, Schindler DE, and Helfield JM. 2002. Pacific salmon, nutrients, and the dynamics of freshwater and riparian ecosystems. *Ecosystems* **5**: 399–417.
- 530 Naish KA and Hard JJ. 2008. Bridging the gap between the genotype and the phenotype: Linking genetic variation, selection and adaptation in fishes. *Fish and Fisheries* **9**: 396–422.
- 535 Nevoux M, Finstad B, Davidsen JG, Finlay R, Josset Q, Poole R, Höjesjö J, Aarestrup K, Persson L, Tolvanen O, and Jonsson B. 2019. Environmental influences on life history strategies in partially anadromous brown trout (*Salmo trutta*, Salmonidae). *Fish and Fisheries* **In Press**.
- Newton I. 2008. The migration ecology of birds. Oxford, UK: Academic Press.
- Nielsen C, Aarestrup K, NØrum U, and Madsen SS. 2003. Pre-migratory differentiation of wild brown trout into migrant and resident individuals. *Journal of Fish Biology* **63**: 1184–96.
- 540 Norin T and Clark TD. 2016. Measurement and relevance of maximum metabolic rate in fishes. *Journal of Fish Biology* **88**: 122–51.
- 545 Norin T and Malte H. 2011. Repeatability of standard metabolic rate, active metabolic rate and aerobic scope in young brown trout during a period of moderate food availability. *Journal of Experimental Biology* **214**: 1668–75.
- Norin T and Metcalfe N. 2019. Ecological and evolutionary consequences of metabolic rate plasticity in response to environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**: 20180180.
- 550 Ohms HA, Sloat MR, Reeves GH, Jordan CE, and Dunham JB. 2013. Influence of sex, migration distance, and latitude on life history expression in steelhead and rainbow trout (*Oncorhynchus mykiss*). *Canadian Journal of Fisheries and Aquatic Sciences* **71**: 70–80.
- Oliveira RF, Taborsky M, and Brockmann HJ. 2008. Alternative reproductive tactics: An integrative approach. Cambridge, UK: Cambridge University Press.
- 555 Olsson IC, Greenberg LA, Bergman E, and Wysujack K. 2006. Environmentally induced migration: The importance of food. *Ecology Letters* **9**: 645–51.

- O'Neal SL and Stanford JA. 2011. Partial migration in a robust brown trout population of a Patagonian river. *Transactions of the American Fisheries Society* **140**: 623–35.
- 560 Peiman KS, Birnie-Gauvin K, Midwood JD, Larsen MH, Wilson ADM, Aarestrup K, and Cooke SJ. 2017. If and when: Intrinsic differences and environmental stressors influence migration in brown trout (*Salmo trutta*). *Oecologia* **184**: 375–84.
- Pettersen AK, Marshall DJ, and White CR. 2018. Understanding variation in metabolic rate. *Journal of Experimental Biology* **221**: jeb166876.
- 565 Phillis CC, Moore JW, Buoro M, Hayes SA, Garza JC, and Pearse DE. 2016. Shifting thresholds: Rapid evolution of migratory life histories in steelhead/rainbow trout, *Oncorhynchus mykiss*. *Journal of Heredity* **107**: 51–60.
- Piche J, Hutchings JA, and Blanchard W. 2008. Genetic variation in threshold reaction norms for alternative reproductive tactics in male Atlantic salmon, *Salmo salar*. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1571–5.
- 570 Polis GA, Power ME, and Huxel GR. 2004. Food webs at the landscape level. University of Chicago Press.
- Pulido F. 2011. Evolutionary genetics of partial migration – the threshold model of migration revis(it)ed. *Oikos* **120**: 1776–83.
- 575 Pulido F, Berthold P, and vanNoordwijk AJ. 1996. Frequency of migrants and migratory activity are genetically correlated in a bird population: Evolutionary implications. *Proceedings of the National Academy of Sciences of the United States of America* **93**: 14642–7.
- Quinn TP. 2018. The behaviour and ecology of pacific salmon and trout. Seattle, WA: University of Washington Press.
- 580 Reid D, Armstrong JD, and Metcalfe NB. 2011. Estimated standard metabolic rate interacts with territory quality and density to determine the growth rates of juvenile Atlantic salmon. *Functional Ecology* **25**: 1360–7.
- Reid D, Armstrong JD, and Metcalfe NB. 2012. The performance advantage of a high resting metabolic rate in juvenile salmon is habitat dependent. *Journal of Animal Ecology* **81**: 868–75.
- 585 Robertsen G, Armstrong JD, Nislow KH, Herfindal I, McKelvey S, and Einum S. 2014. Spatial variation in the relationship between performance and metabolic rate in wild juvenile Atlantic salmon. *Journal of Animal Ecology* **83**: 791–9.
- 590 Roff DA. 1996. The evolution of threshold traits in animals. *The Quarterly Review of Biology* **71**: 3–35.
- Rundio DE, Williams TH, Pearse DE, and Lindley ST. 2012. Male-biased sex ratio of nonanadromous *Oncorhynchus mykiss* in a partially migratory population in California. *Ecology of Freshwater Fish* **21**: 293–9.
- 595 Sadowska ET, Stawski Clare, Rudolf Agata, Dheyongera Geoffrey, Chrzęścik Katarzyna M., Baliga-Klimczyk Katarzyna, and Koteja Paweł. 2015. Evolution of basal metabolic rate in bank voles from a multidirectional selection experiment. *Proceedings of the Royal Society B: Biological Sciences* **282**: 20150025.
- Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, and Mangel M. 2009. Steelhead life history on California's Central Coast:
- 600

- Insights from a state-dependent model. *Transactions of the American Fisheries Society* **138**: 532–48.
- Schindler DE, Armstrong JB, and Reed TE. 2015. The portfolio concept in ecology and evolution. *Frontiers in Ecology and the Environment* **13**: 257–63.
- 605 Schindler DE, Hilborn R, Chasco B, Boatright CP, Quinn TP, Rogers LA, and Webster MS. 2010. Population diversity and the portfolio effect in an exploited species. *Nature* **465**: 609–12.
- Seppänen E, Piironen J, and Huuskonen H. 2010. Consistency of standard metabolic rate in relation to life history strategy of juvenile Atlantic salmon *Salmo salar*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **156**: 278–84.
- 610 Sloat MR, Fraser DJ, Dunham JB, Falke JA, Jordan CE, McMillan JR, and Ohms HA. 2014. Ecological and evolutionary patterns of freshwater maturation in Pacific and Atlantic salmonines. *Reviews in Fish Biology and Fisheries* **24**: 689–707.
- 615 Sloat MR and Reeves GH. 2014. Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences* **71**: 491–501.
- Stefansson SO, Björnsson BT, Sundell K, Nyhammer G, and McCormick SD. 2003. Physiological characteristics of wild Atlantic salmon post-smolts during estuarine and coastal migration. *Journal of Fish Biology* **63**: 942–55.
- 620 Suzuki Y and Nijhout HF. 2008. Genetic basis of adaptive evolution of a polyphenism by genetic accommodation. *Journal of Evolutionary Biology* **21**: 57–66.
- Swingland IR and Greenwood PJ. 1984. Ecology of animal movement. Oxford, UK: Clarendon Press.
- 625 Tanguy JM, Ombredane D, Baglinière JL, and Prunet P. 1994. Aspects of parr-smolt transformation in anadromous and resident forms of brown trout (*Salmo trutta*) in comparison with Atlantic salmon (*Salmo salar*). *Aquaculture* **121**: 51–63.
- 630 Thériault V and Dodson JJ. 2003. Body size and the adoption of a migratory tactic in brook charr. *Journal of Fish Biology* **63**: 1144–59.
- Thériault V, Garant D, Bernatchez L, and Dodson JJ. 2007. Heritability of life-history tactics and genetic correlation with body size in a natural population of brook charr (*Salvelinus fontinalis*). *Journal of Evolutionary Biology* **20**: 2266–77.
- 635 Thorpe JE, Mangel M, Metcalfe NB, and Huntingford FA. 1998. Modelling the proximate basis of salmonid life-history variation, with application to Atlantic salmon, *Salmo salar* L. *Evolutionary Ecology* **12**: 581–99.
- Thorpe JE and Metcalfe NB. 1998. Is smolting a positive or a negative developmental decision? *Aquaculture* **168**: 95–103.
- 640 Tocher DR. 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11**: 107–84.
- Tomkins JL and Hazel W. 2007. The status of the conditional evolutionarily stable strategy. *Trends in Ecology & Evolution* **22**: 522–8.

- 645 Tudorache C, Blust R, and Boeck GD. 2007. Swimming capacity and energetics of  
migrating and non-migrating morphs of three-spined stickleback *Gasterosteus*  
*aculeatus* L. and their ecological implications. *Journal of Fish Biology* **71**: 1448–  
56.
- Via S, Gomulkiewicz R, De Jong G, Scheiner SM, Schlichting CD, and Van Tienderen  
650 PH. 1995. Adaptive phenotypic plasticity: Consensus and controversy. *Trends*  
*in Ecology & Evolution* **10**: 212–7.
- West-Eberhard MJ. 2003. Developmental plasticity and evolution. Oxford, UK: Oxford  
University Press.
- Wone BWM, Madsen P, Donovan ER, Labocha MK, Sears MW, Downs CJ, Sorensen  
655 DA, and Hayes JP. 2015. A strong response to selection on mass-independent  
maximal metabolic rate without a correlated response in basal metabolic rate.  
*Heredity* **114**: 419–27.
- Wysujack K, Greenberg LA, Bergman E, and Olsson IC. 2009. The role of the  
environment in partial migration: Food availability affects the adoption of a  
migratory tactic in brown trout *Salmo trutta*. *Ecology of Freshwater Fish* **18**: 52–  
660 9.
- Young B, Conti DV, and Dean MD. 2013. Sneaker “jack” males outcompete dominant  
“hooknose” males under sperm competition in Chinook salmon (*Oncorhynchus*  
*tshawytscha*). *Ecology and Evolution* **3**: 4987–97.
- 665 Zimmerman CE and Reeves GH. 2000. Population structure of sympatric anadromous  
and nonanadromous *Oncorhynchus mykiss*: Evidence from spawning surveys  
and otolith microchemistry. *Canadian Journal of Fisheries and Aquatic Sciences*  
**57**: 2152–62.

## Chapter 2

---

# The interplay between extrinsic and intrinsic factors in determining migration decisions in brown trout (*Salmo trutta*): An experimental study

Louise C. Archer<sup>\*1,2</sup>, Stephen Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, Michael N. O'Grady<sup>3</sup>, Joseph P. Kerry<sup>3</sup>, W. Russell Poole<sup>4</sup>, Paddy Gargan<sup>5</sup>, Philip McGinnity<sup>1,4</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.

<sup>2</sup> Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland

<sup>3</sup> Food Packaging Group, School of Food and Nutritional Sciences, University College Cork, Cork, Ireland.

<sup>4</sup> Marine Institute, Furnace, Newport, Co. Mayo, Ireland.

<sup>5</sup> Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.

**\*Correspondence:** Louise Archer (l.archer@umail.ucc.ie)

**Key words:** climate change, partial migration, anadromy, aquatic, brown trout, genotype by environment, *Salmo trutta*, proximate drivers

**Published in:** Archer LC, Hutton SA, Harman L, O'Grady MN, Kerry JP, Poole WR, Gargan P, McGinnity P, and Reed TE. 2019. The interplay between extrinsic and intrinsic factors in determining migration decisions in brown trout (*Salmo trutta*): An experimental study. *Frontiers in Ecology and Evolution* 7: 222.

**Contributions:** LA, TR, PMG, and RP conceived the study. LA, SH, TR, and LH collected data and contributed to experimental design. LA conducted statistical analysis and led the manuscript writing. All authors contributed to interpretation of results and revisions of the manuscript.

## Abstract

Many species are capable of facultative migration, but the relative roles of extrinsic versus intrinsic factors in generating diverse migratory tactics remain unclear. Here we explore the proximate drivers of facultative migration in brown trout in an experimental laboratory setting. The effects of reduced food, as a putative environmental cue, were examined in two populations: one that exhibits high rates of anadromy (sea-migration) in nature, and one that does not exhibit anadromy in nature. Juveniles derived from wild-caught parents were reared for two years under four environmental treatments: low food in years 1 and 2 (Low-Low); high food in years 1 and 2 (High-High), low food in year 1 and high in year 2 (Low-High), and vice versa (High-Low). Food restriction had a significant effect on migratory tactics, with the frequency of smolts (juveniles choosing migration) highest in the Low-Low treatment in both populations. No individuals became smolts in the High-High treatment, and intermediate smolting rates were observed in the Low-High and High-Low treatments. Higher overall smolting rates in the naturally anadromous population suggested an inherited component to anadromy/migration decisions, but both populations showed variability in migratory tactics. Importantly, some fish from the naturally non-anadromous population became smolts in the experiment, implying the capacity for migration was lying ‘dormant’, but they exhibited lower hypo-osmoregulatory function than smolts from the naturally anadromous population. Tactic frequencies in the naturally anadromous population were more affected by food in the 2<sup>nd</sup> year, while food in the 1<sup>st</sup> year appeared more important for the naturally non-anadromous population. Migratory tactics were also related to sex, but underpinned in both sexes by growth in key periods, size and energetic state. Collectively these results reveal how migration decisions are shaped by a complex interplay between extrinsic and intrinsic factors, informing our ability to predict how facultatively migratory populations will respond to environmental change.



## 30 Introduction

Intraspecific phenotypic variation accounts for much of the diversity of form and function in nature (Roff 1996). Understanding the mechanisms generating and maintaining divergent phenotypes and life histories within and among populations is thus a fundamental goal of evolutionary ecology, with applied  
 35 relevance to conservation and wildlife management (Naish and Hard 2008). A particularly striking example of alternative phenotypes is the phenomenon of facultative migration, whereby individuals within a population vary in their migratory tendencies. Facultatively migratory populations can comprise a mixture of migrant and resident individuals (sometimes called ‘partial  
 40 migration’), with migration at specific life stages occurring typically to take advantage of alternative foraging opportunities or avoid adverse abiotic (*e.g.* climatic) conditions (Chapman *et al.* 2011a). Despite its widespread occurrence across taxa and regions, fundamental gaps still exist in our understanding of proximate and ultimate drivers of facultative migration. In particular, there is a  
 45 dearth of studies addressing how facultatively migratory species respond to environmental change (Doswald *et al.* 2009; Chapman *et al.* 2011b), limiting our ability to generalise about the impacts of anthropogenic factors on migratory species and to effectively manage their populations.

Polymorphisms such as facultative migration are potentially underpinned by a  
 50 complex mapping between genotype and phenotype, *i.e.* phenotypic similarity can arise from different genotypes, or the same genotypes can produce dramatically different phenotypes through plasticity mediated by environmental cues (Roff 1996). As such, migration and residency have often been considered as environmentally-triggered alternative phenotypes/tactics  
 55 produced by an evolvable conditional strategy, where optimal tactic choice in a given context is conditional on extrinsic or intrinsic cues (Chapman *et al.* 2011b). This interplay between proximate and ultimate drivers of conditional strategies has been formalised as the so-called ‘environmentally cued threshold model’ (Tomkins and Hazel 2007). Within this framework, alternative tactics  
 60 are controlled by an environmentally-sensitive status trait (*e.g.* physiological

condition, energy state) and an inherited threshold, or ‘switch point’, which is assumed to be genetically variable. An individual assesses their status trait and, for example, adopts a resident tactic if it exceeds their inherited switch point, otherwise it switches to a migratory tactic. Individual physiological  
65 condition/energy state is strongly influenced by the environment, and so the assessed status trait can vary relative to the intrinsic threshold depending on external conditions; for this reason, the status trait can be thought of as an ‘environmental cue’ and the step function relating tactic expression to cue as a ‘threshold reaction norm’ (Tomkins and Hazel 2007; Piche *et al.* 2008; Pulido  
70 2011; Buoro *et al.* 2012). There is some evidence for genetic variation in thresholds for alternative tactics, *e.g.*, in blackcaps *Sylvia atricapilla* (Pulido *et al.* 1996) and Atlantic salmon *Salmo salar* (Piche *et al.* 2008), but detailed understanding of how external environmental variation is translated into internal physiological signals, on which migratory decisions are then based, is  
75 lacking.

Salmonine fishes (salmons, trouts and charrs) are excellent models for disentangling causes of facultative migration as they display wide variation across a continuum of migratory strategies, coupled with obligate freshwater spawning (Klemetsen *et al.* 2003; Ferguson *et al.* 2019). Individuals can remain  
80 in freshwater post hatching for their entire life cycle, either staying in their natal stream or lake (residency tactic) or undertaking an adfluvial migration that takes them to a larger river or lake (potamodromous tactic) (Dodson *et al.* 2013; Ferguson *et al.* 2019). Facultative anadromy is an extreme form of this conditional migration strategy, where some individuals adopt the residency  
85 tactic whilst others from the same population undertake a marine migration (involving anywhere from tens to thousands of kilometres of directed movement between freshwater and saltwater). This is followed by a period of marine or estuarine feeding and growth (from months to years), before returning to spawn in natal streams (Jonsson and Jonsson 1993). Populations  
90 can contain both resident and migratory (anadromous or potamodromous) forms, or be dominated by one life history type (Chapman *et al.* 2012). Both

forms can breed freely in sympatry, and although offspring tend to track the tactics of their parents, either life history can be produced from a given migratory phenotype (Zimmerman and Reeves 2000; Berejikian *et al.* 2014).

95 Such flexibility indicates an interplay between genetic predisposition and environmental conditions experienced i.e., genotype by environment interactions, underpinning facultative migration (Hutchings 2011).

The threshold reaction norm framework has been useful in understanding migratory decisions in salmonines (Hutchings and Myers 1994; Thorpe *et al.* 100 1998; Thériault *et al.* 2007). If during a key decision window, an individual's status trait exceeds their predetermined threshold, the fish adopts a residency tactic leading to maturation in freshwater; if not, maturation is deferred in favour of migration (Dodson *et al.* 2013; Kendall *et al.* 2014; Ferguson *et al.* 2017). However, the proximate factors on which individuals base the migration 105 decision remain unclear. Previous studies have focused on a range of aspects of physiological state/energy status that may influence migratory tactics such as body size (Thériault and Dodson 2003), lipid reserves (Jonsson and Jonsson 2005), body condition (Hecht *et al.* 2015), growth (Jonsson 1985), growth efficiency (Forseth *et al.* 1999; Morinville and Rasmussen 2003), and 110 metabolism (Sloat and Reeves 2014). While body size is often used as a surrogate for, or argued to itself be, the status trait triggering alternative migratory tactics, the associations here have been varied and inconclusive. Larger sizes and faster growth rates have been associated with early age at migration (Jonsson 1985), whereas others have found no size-based differences 115 between migrants and non-migrants at a given age (Thériault and Dodson 2003), or conversely found larger sizes (and higher lipid reserves) to be associated with freshwater maturation *in lieu* of anadromy (McMillan *et al.* 2012). These inconsistencies could reflect species' specific responses, and thus require further exploration to establish potential status traits for a given species. 120 Studies might also be inconclusive because size is typically measured sometime after the migratory decision itself, perhaps at the parr-to-smolt transformation stage, and size at migration may not accurately reflect size when the decision

was made. For example, residents may have meanwhile diverted energy into maturation and gonadal development at the expense of somatic growth (Tocher  
 125 2003), while migrants may undergo accelerated growth as the migration itself approaches (Metcalf 1998).

Moreover, there may be at least two separate threshold decisions: an early one determining whether a fish will migrate *per se* or not, and a later one determining whether fish on a migratory trajectory actually migrate this year or  
 130 defer migration to an older age (Ferguson *et al.* 2019). Size may be the cue used for the second decision, given that survival on entry to the sea or a lake is typically positively related to size (Klemetsen *et al.* 2003; Phillis *et al.* 2016). Yet, size at the migration point may be unrelated to, or inconsistently related to, the status trait triggering the initial migration decision, which could occur  
 135 considerably earlier than the point at which migrants and resident become phenotypically distinguishable (Beakes *et al.* 2010). Identifying the key proximate drivers of migration is therefore complicated by the fact that the exact time windows for each of these putative decisions may not be known *a priori*, while correlations among physiological, energy status and growth traits  
 140 may be variable across ontogeny or contexts. In the particular case of facultative anadromy, sea-migration requires a suite of adjustments in preparation for life in saltwater and therefore the physiological remodelling process, which includes changes in osmoregulation, colouration, and body shape (Tanguy *et al.* 1994), is likely to begin sometime in advance of the migratory period. The  
 145 existence of early ‘decision windows’ that initiate divergent life-history trajectories in salmonine fishes (Thorpe *et al.* 1998; Thorpe and Metcalfe 1998) has some empirical support; for example, body condition of anadromous *Oncorhynchus mykiss* was found to be significantly lower than resident counterparts within a year of hatching and a full 12 months prior to emigration  
 150 (Hecht *et al.* 2015).

Although the proximate drivers of migration in salmonines are unresolved, there is some consensus that potamodromous or anadromous migratory tactics are promoted by energetic limitation in natal rivers, which prevents fish

reaching the inherited physiological threshold for maturation as residents  
 155 (Kendall *et al.* 2014). Energetic limitation can arise through an interplay  
 between environmental factors and intrinsic physiological state; for example, if  
 freshwater food resources are insufficient to support growth rates or metabolic  
 demands, then migration could be triggered that takes the fish to a better  
 feeding environment such as the sea or a large lake (O'Neal and Stanford 2011;  
 160 Sloat and Reeves 2014; Jones *et al.* 2015). Food limitation arising from  
 competition at high population densities has also been shown to increase the  
 proportion of adfluvial migratory brown trout, whereas low population  
 densities have been associated with residency and maturation (Olsson *et al.*  
 2006; Wysujack *et al.* 2009). It remains largely unknown, however, during  
 165 which ontogenetic stages food limitation is most important to migration  
 decisions.

Brown trout (*Salmo trutta*) are an interesting model for understanding  
 facultative migration as they exhibit highly variable strategies, with some  
 individuals/populations remaining resident in their natal stream their entire  
 170 lives, while others migrate to a larger river, a lake, an estuary, or the sea (Jonsson  
 and Jonsson 1993; Klemetsen *et al.* 2003; Cucherousset *et al.* 2005; Ferguson *et al.*  
 2019). Here we present the results of an experimental laboratory study of  
 brown trout that involved F1 progeny of wild-caught parents from two  
 populations that exhibit divergent migratory life-histories in nature. Our  
 175 primary aim was to explore the interaction between intrinsic proximal factors  
 (which may encompass both inherited and non-inherited variation) and the  
 extrinsic environment in generating alternative migratory tactics in brown  
 trout. Specifically, we aimed to: (i) assess the relative importance of food  
 availability and inherited differences between populations in determining  
 180 alternative migratory tactics; (ii) determine whether food restriction was more  
 important in the first year or second year of freshwater rearing; (iii) test for  
 differences between our two populations in their response to food restriction  
 and its timing, which may be indicative of genotype-by-environment  
 interactions influencing tactic frequencies, and (iv) explore associations

185 between status traits (length, weight, condition factor) and migratory tactics. We expected that food restriction would increase the frequency of the migratory tactic overall. While we expected migratory tactic frequencies to vary overall between fish from our two population backgrounds, we also anticipated that the naturally non-anadromous stock might produce migratory phenotypes  
 190 when subjected to reduced food, given that migration may only be expressed under certain environmental conditions (Roff 1996; Pulido 2011).

## Materials and methods

### *Study populations*

Wild-origin brown trout brood stock were obtained by seine netting from the  
 195 Burrishoole (53° 57' N: 09° 35' W) and Erriff (53° 37' 0.00" N: 09° 40' 17.10" W) catchments in the west of Ireland in November 2015. Burrishoole brood stock were caught in Lough Bunaveela (46 ha, Figure S1) in the headwaters of the catchment. A local population of non-anadromous trout remain resident in Lough Bunaveela for most of their lifecycle, bar very short-distance directed  
 200 movements (on the order of 10s to 100s of metres) between the lake and two spawning rivers (one inflowing to the lake, the other outflowing). No obvious genetic structure at neutral microsatellite markers is evident between these spawning rivers, implying trout from Lough Bunaveela comprise a single panmictic population (R. Finlay, *pers. comm.*). A large run of sea trout (typically  
 205 2000+ anadromous recruits annually) occurred in the Burrishoole catchment up to 30 years ago. The Burrishoole anadromous trout run collapsed in the late 1980s, coinciding with sea-lice outbreaks following the establishment of salmon aquaculture farms in the downstream estuary. The exact spawning locations of the historic anadromous individuals within the Burrishoole catchment remain  
 210 uncertain, and we cannot exclude the potential for some anadromous fish having contributed to the Bunaveela population before the anadromous population collapse. Nevertheless, despite Bunaveela spawning streams being accessible to anadromous migrants, there is little to no evidence that the Bunaveela population produced anadromous trout historically or recently

215 (Poole *et al.* 2007; Magee 2017) and we thus consider it a population that rarely, if ever, expresses anadromy.

Erriff brood stock were caught in Tawnyard Lough, a small upland lake (56 ha) on the western side of the Erriff catchment (the National Salmonid Index catchment) that is fed by a primary inflowing stream, the Glendavoch River and  
 220 a number of smaller tributaries (Figure S1). The vast majority of trout spawned in the Glendavoch River are believed to disperse as fry or parr to Tawnyard Lough (a distance of a few hundred metres to a few kilometres, depending on how far up the Glendavoch River spawning occurred), although a small fraction remain permanently resident in the natal stream (P. Gargan, *pers. comm.*). A  
 225 large run of out-migrating anadromous juveniles (in the range of five hundred to three thousand smolts per year over the last 30 years) is enumerated annually in a trap at the outflow of Tawnyard Lough (Gargan *et al.* 2016). The remaining fish never go to sea but instead spend several years growing in the lake, before returning to spawn in the Glendavoch River and smaller tributaries once  
 230 mature. Brood stock from the Tawnyard population used in this experiment putatively comprised a mix of anadromous and non-anadromous fish, assumed to represent naturally occurring frequencies of anadromous and non-anadromous tactics (see Table S1 for details of brood stock), with local expertise indicating that the Tawnyard population in general shows high rates of  
 235 anadromy (P. Gargan, *pers comm.*). In summary, we consider the Tawnward population to have a strong migratory/anadromous background, and the Bunaveela population to have essentially no (recent) anadromous background and to exhibit only limited local movements. For ease of reading, juveniles derived from Tawnyard parents are hereafter referred to simply as the  
 240 “anadromous-background” population and juveniles from Bunaveela parents as the “non-anadromous background” population.

### *Fish rearing*

Females were stripped of eggs, and the eggs of each female were divided into two batches, each fertilised by the milt of a single male from the same source  
 245 population (i.e. Tawnyard or Bunaveela; see Table S1 for full details on crossing).

Fertilised eggs were then incubated in standard Heath trays in a hatchery facility located within the Burrishoole catchment. Surviving unfed fry (two to three weeks prior to exogenous feeding) were transferred to a rearing facility at University College Cork (Aquaculture and Fisheries Development Centre).

250 While transitioning to exogenous feeding, fry were held in 100L growth tanks on a recirculating aquaculture system (RAS) with bio filtration, and fed *ad libitum* to satiation using commercially available trout pellets (Skretting Ltd, Norway). The populations were kept separately in two 100L tanks during this initial rearing phase and maintained under a natural temperature regime

255 regulated by a single conditioning unit. Once the fry had transitioned to exogenous feeding (June 2016), they were fed *ad libitum* with commercial trout pellets for a period of two months. All fish experienced the same constant photoperiod regime (12 hours of light and 12 of dark) during this initial rearing phase.

260 In September 2016, fish were randomly allocated into four 100L tanks in the same RAS as described above (two tanks for Tawnyard and two tanks for Bunaveela), at which point the experimental phase began and food manipulations were initiated (see next section for experimental treatments). A random subset of fish ( $n = 200$  per population) were given individual identifier

265 tags using unique colour combinations of visible implant elastomer tags (Northwest Marine Technology Ltd., USA). To facilitate growth, in December 2016 the fry were transferred (within their experimental groups) to 520L growth tanks in a larger RAS in the same aquaculture facility. Continuous through flow of water prevented any waste accumulation in tanks, with returning water

270 passed to a central holding sump and treated via mechanical filtration, protein skimming, bio filtration, and ozone and UV sterilisation. Water quality in the system was monitored weekly, and levels of pH, nitrate, nitrite, and ammonia were within acceptable ranges for optimal fish health. During the experimental phase, the fish experienced a seasonally-changing photoperiod and temperature

275 regime typical of the west of Ireland, simulated via an automated lighting system of LED lights (BioLumen, UK) above each tank and a single conditioning



unit. Negligible natural mortality occurred during the experimental phase but to maintain total biomass in the RAS at acceptable levels from a water quality perspective, fish were randomly culled ( $n = 120$  in total across all tanks) over  
 280 the course of the two years of tank rearing, with equal fish densities maintained between food treatments. Fish that were prematurely culled were excluded from all analyses. Full details on the stripping, crossing and rearing procedures are given in Supplementary Information.

### *Experimental design*

285 The experimental phase ran for a 22 month period, from September 2016 to June 2018, with all fish humanely euthanized at the end of the experiment under licence (the study and all associated procedures were carried out with ethical approval from Health Products Regulatory Authority (HPRA) Ireland, under HPRA project license AEI9I30/P034, and HPRA individual licenses  
 290 AEI9I30/I087, AEI9I30/I200, AEI9I30/I201 and AEI9I30/I202).

To investigate the relative importance of the extrinsic environment (food supply) and intrinsic inherited factors (population-of-origin) in determining migratory tactics, juveniles from the anadromous and non-anadromous background populations were divided evenly and allocated randomly across  
 295 four tanks receiving water from the same recirculating source, each experiencing a different feeding regime over the experimental phase. Populations were kept separately for the duration of the study ( $n = 90$  per feeding treatment per population, at the beginning of the experimental phase). Great care was taken to ensure that all measured variables other than feeding  
 300 regime (fish densities, temperature, photoperiod, lux, flow rates) were constant across the tanks. The four feeding regime treatments were designed to test the effects of food restriction in the early versus late periods of this experimental phase, with each period corresponding to approximately 11 months (chosen because similar periods of *c.* 9 months have been reported to alter adfluvial  
 305 migration rates in trout (Olsson *et al.* 2006)). These four food regimes were as follows: (i) *High-High* treatment: fish fed recommended daily pellet rations for optimal growth in both periods, calculated as a percentage of their body weight

and adjusted for seasonally-changing temperatures (Skretting Ltd, Norway); (ii) *Low-Low* treatment: fish fed 25% of recommended optimal rations in both  
 310 periods; (iii) *High-Low* treatment: fish fed 100% of optimal daily rations in the first period and 25% of optimal daily ration in the second period; and (iv) *Low-High* treatment: fish fed 25% of optimal daily rations in the first period and 100% of optimal daily ration in the second period. A value of 25% of optimum levels was chosen for the *Low* feeding regime because similar reductions have  
 315 previously been shown to reduce the frequency of the resident tactic in adfluvial brown trout (Wysujack *et al.* 2009). Rations were reduced down to 25% of optimal gradually over a four-week period, to minimise stress. Within each food treatment, absolute rations were adjusted according to manufacturer's instructions (see Table S2) on a monthly basis to account for changes in body  
 320 mass and temperature (i.e. there was no variation in daily rations within months, within groups).

#### *Life history determination and data collection*

In the spring of 2017 and 2018 (March – June in year one and year two of the experimental phase of the study), fish were routinely assessed for morphological  
 325 indicators of 'smoltification': the series of morphological, physiological and behavioural changes that is generally considered a precursor to downstream migration of juvenile salmonids (Tanguy *et al.* 1994). Here we use 'smolt' to simply mean a fish showing external morphological features consistent with preparing for a migration, and we used saltwater tolerance tests (see below) to  
 330 further assess physiological aspects of smoltification. We visually assessed morphological smoltification (silvered flanks/loss of parr marks, pronounced lateral line, colourless fins and fusiform shape) according to Tanguy *et al.* (1994). No fish matched the morphological criteria of smolts in the spring of 2017, the very earliest point at which we expected any smoltification (Poole *et al.* 2007; Gargan *et al.* 2016). Individuals that matched the morphological  
 335 criteria for smolts in spring 2018 were transferred to saltwater at 30 ppt for 24 hours to assess their hypo-osmoregulation as a further indicator of anadromy capacity. We used 30 ppt salinity [following Tanguy *et al.* (1994)] because trout

often spend large amounts of time in brackish water/estuaries when migrating,  
 340 hence trout smolts are typically less saltwater tolerant than other salmonids *e.g.*  
 Atlantic salmon (Urke *et al.* 2010). After the 24-hour immersion in saltwater, a  
 period proposed to induce hypo-osmoregulation in euryhaline species (Schultz  
 and McCormick 2012), fish were euthanised with an overdose of MS-222 and a  
 blood sample was taken from the caudal vasculature using a 21G needle and a  
 345 2.6ml heparinised syringe. Blood samples were transferred to 2 ml epindorpha  
 and centrifuged at 8000 rpm for 3 minutes. The plasma aliquot was then  
 siphoned off and stored at -80 °C before being measured for plasma chloride  
 concentration as an indicator of hypo-osmoregulatory ability.

All fish, whether identified morphologically as smolts or non-smolts, were  
 350 dissected to visually determine sex and maturation status according to gonad  
 development. Males were classed as sexually mature if they had enlarged white  
 testes or had running milt. Males that had visible testes that were moderately  
 enlarged but not running milt were classed as maturing. Females were classed  
 as mature or maturing if the body cavity contained identifiable eggs. Fish with  
 355 immature gonads, or that could not be identified as either male or female by  
 visual inspection were classed as immature at the time of sampling, and their  
 genotypic sex was later determined using a microsatellite sex marker (P.  
 Prodöhl, unpublished). In the wild, the natural spawning period for these brown  
 trout populations is in late autumn/early winter, and the migratory period is in  
 360 the spring (Poole *et al.* 2007; Gargan *et al.* 2016). Fish showing signs of maturity  
 in freshwater without having first gone to sea, were considered to be on a non-  
 anadromous trajectory, while smolts migrating to sea in a given spring were all  
 immature. Fish in our experiment were thus classed as smolts (migratory tactic)  
 if they were morphologically assessed as smolts and were immature, and were  
 365 classed as mature (freshwater maturation tactic) if they were mature or  
 maturing at the time of sampling. Fish that were classed as immature, but did  
 not have morphological indicators of smoltification, were considered to have an  
 unknown life history tactic at the time of sampling. A small number of fish (n =  
 12) had significant skin/fin damage at the time of sampling, and were excluded

370 from the analysis. Whole body lipid content (%) was measured for all smolts, and for a random sample of mature fish ( $n = 111$ ), using a SMART Trac 5 system (CEM GmnH, Kamp-Lintfort, Germany) of integrated microwave heating and nuclear resonance on homogenised samples.

### *Statistical analysis*

375 To assess whether food treatment and population influenced life history tactics (Aims 1 and 2), we constructed generalized linear models (GLMs) with a logit link function and binary life-history response variables. One GLM was created to predict smolt status (binary response: 1 = smolt, 0 = non-smolt) using the *brglm* package in R (Kosmidis 2019) to account for separation in the data (no  
380 smolts recorded in the *High-High* treatment) (Heinze and Schemper 2002). A second GLM was created to predict maturation (binary response: 1 = mature or maturing, 0 = immature). Categorical explanatory variables in both of these GLMs included food treatment (*High-High*, *Low-High*, *Low-Low*, *High-Low*), population (anadromous-background versus non-anadromous-background),  
385 and sex (male or female) as predictors. We constructed a third GLM to test for treatment/population effects on likelihood of being classed as “unassigned” (i.e. not having expressed a migratory/resident phenotype by the end of the study (binary response: 1 = unassigned, 0 = smolt or mature). We included an interaction term between food treatment and population to determine if life  
390 history responses in each population were similar under the different food regimes (Aim 3). To test whether food restriction was more important in the early or late rearing periods (Aim 2), we conducted Tukey post-hoc tests (with Bonferroni correction applied for multiple tests) of all possible pairwise comparisons among the levels of food treatment using the *emmeans* package in  
395 R (Lenth 2019). Overall, one expects the strongest difference in life-history tactics to be found between the *High-High* and *Low-Low* treatments. If the effects of food restriction are additive and the timing of food restriction does not matter, then one expects life-history tactics in the *Low-High* and *High-Low* treatments to be intermediate between the *High-High* and *Low-Low* treatments,  
400 and not significantly different from each other. Conversely, if food restriction is

more important in the first period, then one expects tactic frequencies in the *Low-High* treatment to be closer to those in the *Low-Low* treatment (and the *High-Low* treatment should be more similar to the *High-High* treatment), while if food restriction is more important in the second period, the *High-Low* treatment should be closer to the *Low-Low* treatment and the *Low-High* treatment to the *High-High*. To further explore factors influencing variation in saltwater tolerance (Aims 1-3) – a key component of life-history tactics – we constructed a linear model (normal errors) with plasma chloride concentration as the continuous response, and population, food treatment, sex, and an interaction between population and food treatment included as predictors.

To address Aim 4, we explored factors influencing variation in the length, weight and condition factor of fish at different measurement time points across the study period within a mixed-effects modelling framework [*nlme* package, (Pinheiro *et al.* 2019)]. Measurement time points were September and November in 2016, February, April, June, July, September, and December in 2017, and April 2018. Condition factor was calculated as Fulton's *K* where:

$$Condition\ (K) = \frac{mass\ (g)}{fork\ length\ (cm)^3} \times 100$$

For the subsequent analyses of status traits, we created a new categorical variable called 'life-history tactic' with two levels: migratory (i.e. immature smolts) or mature/maturing (hereafter simply called mature). Fish which were neither classified as migratory nor mature (unassigned fish) were not included in the status trait analyses, as it could not be determined which life history trajectory they might adopt (i.e. these fish could have displayed either migratory or mature tactics the following spring (a full three years after hatching), but the experiment was terminated the previous spring (two years after hatching)). In addition to life-history tactics, month (continuous variable), population (categorical variable with two levels), food treatment (categorical variable with four levels) and sex (categorical variable with two levels) were included as fixed effects, and individual identity was included as a random effect to account for multiple measurements on some individuals. We included an interaction

between life-history tactics and month (to test whether individuals on different life-history trajectories diverged through time in their length/weight/condition factor), an interaction between life-history tactics and population (to test whether average differences in length/weight/condition factor between the two tactics was similar across the two populations), and an interaction between population and food treatment (to test whether the effects of food regime were similar across populations). Temporal autocorrelation of the response variable was accounted for by modelling an autoregressive error structure as a first order lag function of month. Separate models were constructed each for length, weight and condition factor and normal errors were assumed in each case.

We also explored factors influencing variation in final length,  $K$  and whole body lipids (i.e. the final measurements for these status traits at the end of the study) in a mixed effects modelling framework, where life-history tactics, food treatment, population and sex were included as fixed effects, and date of terminal sample (categorical variable with 11 sampling dates) was modelled as a random effect. We included two interaction terms (life-history tactics  $\times$  population, and food treatment  $\times$  population), to explore whether the patterns for each population were similar across tactics and food treatments, respectively. Separate models were constructed each for length,  $K$  and whole body lipids and normal errors were assumed in each case. Marginal  $R^2$  values for mixed effect models were calculated using the *MuMIn* package in R (Barton 2018).

For all of the above models, statistical significance at a 5% alpha level of predictor variables was assessed using likelihood ratio tests (LRT), and non-significant interaction terms were omitted so the main effects could be interpreted.

Finally, to assess whether variation in growth was associated with life-history tactics (Aim 4), we compared growth trajectories of migratory and mature fish by fitting three typical models of fish growth: the von-Bertalanffy growth curve, the Gompertz growth curve and a logistic growth curve. The logistic growth curve best described the data according to AIC ( $\Delta AIC = 0$ ), and was used for all

further growth trajectory analysis. The logistic growth equation models asymptotic growth as:

$$L = \frac{L_{\infty}}{1 + e^{(-g_i(T-I))}}$$

Where  $L$  is fork length,  $L_{\infty}$  is asymptotic fork length (cm),  $g_i$  is the growth rate (cm/day),  $T$  is time (days) and  $I$  is the inflection point. The logistic model was fitted using non-linear least squares to length data collected on individually-identifiable fish during the experiment, with separate models fitted for smolts and mature fish. As non-linear least squares regression is sensitive to starting values of parameters, the model was fitted using the *nls\_multstart* function from the *nls.multstart* package in R (Padfield and Matheson 2018). This allowed for starting values for each parameter to be randomly selected from a bounded distribution over 1000 iterations of the model, with the best available model then selected by AIC. To determine the fit of the most parsimonious model to our data, we bootstrapped with replacement 10,000 times and constructed 95% confidence intervals from the bootstrapped fits.

All analysis was carried out in R version 3.5.3 (R Core Team 2019), and all statistical models were checked against assumptions of the given model (independence, non-normality of residuals, heteroscedasticity and multicollinearity).

## Results

### *Life-history tactics*

By the end of the experimental phase, a total of 567 fish had been categorised as either smolts, i.e. putatively migratory (n=36 females and n=18 males) or non-smolts (n=277 females and n=236 males). All of the smolts were by definition immature, and 15.52% of the non-smolt females and 28.39% of the non-smolt males were immature. See Table 1 for a full breakdown of life-history tactics by population background, food treatment and sex. The proportion of smolts varied according to food treatment and population (Figure 1). Highest proportions of smolts were seen in the *Low-Low* food treatment, in which

26.56% of the anadromous-background population, and 15.71% of the non-anadromous background population, were classified as smolts. The lowest rates of smolting were found in the *High-High* food treatment, in which no fish from either population were categorised as smolts. Intermediate smolting rates were  
 495 observed in the other two treatments, with 6.45% of fish from the anadromous-background population and 13.75% of fish from the non-anadromous background population classified as smolts in the *Low-High* treatment, and 15.87% and 1.22% of fish from each population, respectively, classified as smolts in the *High-Low* treatment.

500 **Table 1:** Percentage of brown trout (n = 567, F1 offspring of wild trout from two population backgrounds) classed as smolts (i.e. migratory tactic) or non-smolts (mature or immature) after two years of experimental tank-rearing. Values correspond to percentages for each category, broken down by sex, of the total number of fish per tank (where each tank corresponds to a given population  
 505 background by food treatment combination, i.e. a single row in the table). Sample size (n) given in brackets after the %.

| Treatment        | Population background | % Smolts (n)  |              | % Non-smolts (n) |               |               |               |
|------------------|-----------------------|---------------|--------------|------------------|---------------|---------------|---------------|
|                  |                       | Female        | Male         | Mature           |               | Immature      |               |
|                  |                       |               |              | Female           | Male          | Female        | Male          |
| <i>Low-Low</i>   | Anadromous            | 23.44<br>(15) | 3.13<br>(2)  | 25.00<br>(16)    | 35.94<br>(23) | 3.10<br>(2)   | 9.38<br>(6)   |
| <i>Low-High</i>  | Anadromous            | 4.84<br>(3)   | 1.61<br>(1)  | 50.00<br>(31)    | 32.26<br>(20) | 1.61<br>(1)   | 9.68<br>(6)   |
| <i>High-Low</i>  | Anadromous            | 11.11<br>(7)  | 4.76<br>(3)  | 39.68<br>(25)    | 34.92<br>(22) | 4.76<br>(3)   | 4.76<br>(3)   |
| <i>High-High</i> | Anadromous            | 0.00<br>(0)   | 0.00<br>(0)  | 48.44<br>(31)    | 45.31<br>(29) | 1.56<br>(1)   | 4.69<br>(3)   |
| <i>Low-Low</i>   | Non-anadromous        | 2.86<br>(2)   | 12.86<br>(9) | 44.29<br>(31)    | 14.29<br>(10) | 1.43<br>(1)   | 24.29<br>(17) |
| <i>Low-High</i>  | Non-anadromous        | 10.00<br>(8)  | 3.75<br>(3)  | 42.50<br>(34)    | 28.75<br>(23) | 7.50<br>(6)   | 7.50<br>(6)   |
| <i>High-Low</i>  | Non-anadromous        | 1.22<br>(1)   | 0.00<br>(0)  | 35.37<br>(29)    | 17.07<br>(14) | 20.73<br>(17) | 25.61<br>(21) |
| <i>High-High</i> | Non-anadromous        | 0.00<br>(0)   | 0.00<br>(0)  | 45.12<br>(37)    | 34.15<br>(28) | 14.63<br>(12) | 6.10<br>(5)   |




The probability of smolting was described by a GLM retaining food treatment ( $\chi^2 = 44.57$ ,  $df = 3$ ,  $p < 0.001$ ), population ( $\chi^2 = 3.46$ ,  $df = 1$ ,  $p = 0.063$ ), sex ( $\chi^2 = 4.40$ ,  $df = 1$ ,  $p = 0.036$ ), and an interaction between food treatment and population (LRT for the model with and without interaction term:  $\chi^2 = 11.66$ ,  $df = 3$ ,  $p = 0.009$ ). Overall across the two populations, there appeared to be an additive effect of food treatment on the probability of smolting – that is, the percentages of smolts in the *Low-High* and *High-Low* treatments were similar, and approximately intermediate to the percentages in the *Low-Low* and *High-High* treatments, when population was ignored (Figure 1). However, when population was taken into account, the life-history response to food treatment varied by population and appeared to be non-additive within each population (Table 2, Figure 1). Fish from the anadromous-background population exhibited a relatively high percentage of smolts (15.87%) under the *High-Low* treatment that was closer to the *Low-Low* treatment (26.56% smolts) than to the *High-High* treatment (0% smolts) and post-hoc comparisons of *High-Low* against *Low-Low* were not significant ( $p = 0.377$ ). The opposite was true for the anadromous-background population in the *Low-High* treatment (6.45% smolts) with significant post-hoc comparisons of *Low-Low* and *Low-High* ( $p = 0.016$ ). In contrast, fish from the non-anadromous-background population exhibited a relatively high percentage of smolts (13.75%) under the *Low-High* treatment that was closer to the *Low-Low* treatment (15.71% smolts) than to the *High-High* treatment (0% smolts) (post-hoc contrasts between *Low-High* and *Low-Low* were non-significant,  $p = 0.994$ ), while the opposite was true for this population in the *High-Low* treatment (1.22% smolts) (post-hoc contrasts between *High-Low* and *Low-Low* were significant,  $p = 0.042$ ). This implies that food restriction was more important in the second period for fish from the anadromous-background population, while food restriction in the first period was more important for the non-anadromous-background fish.

Maturation tactics in freshwater were also significantly affected by food treatment ( $\chi^2 = 33.03$ ,  $df = 3$ ,  $p < 0.001$ ), population ( $\chi^2 = 12.14$ ,  $df = 1$ ,  $p < 0.001$ ), sex ( $\chi^2 = 4.54$ ,  $df = 1$ ,  $p = 0.033$ ) but there was no significant interaction between

food treatment and population (LRT for the model with and without interaction  
 540 term:  $\chi^2 = 5.31$ ,  $df = 3$ ,  $p = 0.150$ ). Food restriction had a negative effect on  
 maturation probability, in direct contrast to food restriction effects on smolting  
 rates. Fish in the *Low-Low* food treatment had the lowest probability of  
 maturing, ( $p < 0.001$ , Table 2) and the highest rates of maturity were observed  
 in the *High-High* food treatment, ( $p < 0.001$ , Table 2). Fish from the  
 545 anadromous-background population were significantly more likely to mature  
 than fish from the non-anadromous-background population in all food  
 treatments ( $p = 0.001$ , Table 2). See Table 2 for all parameter estimates and  
 associated standard errors. The probability of having been unassigned a life  
 history showed similar patterns to maturation tactics, and was similarly  
 550 significantly affected by food treatment ( $\chi^2 = 16.95$ ,  $df = 3$ ,  $p = 0.001$ ), population  
 ( $\chi^2 = 30.74$ ,  $df = 1$ ,  $p < 0.001$ ) and sex ( $\chi^2 = 16.21$ ,  $df = 1$ ,  $p < 0.001$ ), see Table 2.  
 The interaction between food treatment and population was marginally not  
 significant (LRT for the model with and without interaction term:  $\chi^2 = 7.75$ ,  $df =$   
 1,  $p = 0.052$ ).


555 We found a significant effect of population on plasma chloride levels of fish  
 classified as smolts ( $F = 9.47$ ,  $df = 1,48$ ,  $p = 0.003$ ), but the interaction term  
 between population and food treatment was not significant (LRT for model with  
 and without interaction term:  $F = 1.39$ ,  $df = 2$ ,  $p = 0.259$ ). Fish from the  
 anadromous-background population had significantly lower plasma chloride  
 560 concentrations than non-anadromous-background fish ( $p = 0.003$ , Table 3,  
 Figure 2). There was no significant effect of food treatment ( $F = 2.95$ ,  $df = 2,48$ ,  
 $p = 0.062$ ) or sex ( $F = 0.01$ ,  $df = 1,48$ ,  $p = 0.991$ ) on plasma chloride levels (Table  
 3).



**Figure 1:** Proportion of brown trout (n = 567, F1 offspring of wild trout from two population backgrounds) classed as smolts after two years of tank rearing under varying food restriction treatments. Food treatment is denoted in the format “food in year one - food in year two”, where “high” refers to optimal food rations and “low” refers to 25% of optimal rations. P-values shown are Tukey post-hoc pairwise comparisons across all levels of food treatment for each population.

**Table 2:** Parameter estimates with associated standard errors (SE) for two binomial generalised linear models (GLM) predicting smolt (migratory) probability (dummy coded: smolt = 1, non-smolt = 0) and freshwater maturation (dummy coded: mature/maturing = 1, immature = 0) in brown trout (n = 567). The reference level of each factor is in brackets, i.e. effects in both models were contrasted against female fish from the anadromous-population background in the *Low-Low* food treatment. Statistical significance was assessed at  $p < 0.05$ .

| Effect   | Estimate | SE   | t-value | p-value |
|--|----------|------|---------|---------|
| <i>GLM of probability of smoltification:</i>         |          |      |         |         |
| Intercept (Low-Low, female, Anadromous background)   | -0.71    | 0.31 | -2.28   | 0.022   |
| Food: Low-High                                       | -1.61    | 0.57 | -2.83   | 0.005   |
| High-Low   | -0.66    | 0.44 | -1.49   | 0.136   |
| High-High  | -3.87    | 1.45 | -2.66   | 0.008   |
| Population: Non-anadromous background                | -0.63    | 0.43 | -1.47   | 0.142   |
| Sex: Male  | -0.63    | 0.31 | -2.06   | 0.039   |
| Low-High : Non-anadromous background                 | 1.38     | 0.73 | 1.90    | 0.058   |
| High-Low: Non-anadromous background                  | -1.75    | 0.99 | -1.77   | 0.077   |
| High-High: Non-anadromous background                 | 0.33     | 2.06 | 0.16    | 0.873   |
| <i>GLM of probability of maturation:</i>             |          |      |         |         |
| Intercept (Low-Low, female, Anadromous background)   | 0.97     | 0.24 | 4.12    | < 0.001 |
| Food: Low-High                                       | 0.78     | 0.27 | 2.90    | 0.004   |
| High-Low   | 0.10     | 0.25 | 0.42    | 0.676   |
| High-High  | 1.43     | 0.30 | 4.78    | < 0.001 |
| Population: Non-anadromous background                | -0.68    | 0.20 | -3.43   | 0.001   |
| Sex: Male  | -0.41    | 0.19 | -2.13   | 0.033   |
| <i>GLM of probability of "unassigned" phenotype:</i> |          |      |         |         |
| Intercept (Low-Low, female, Anadromous background)   | -2.77    | 0.33 | -8.29   | < 0.001 |
| Food: Low-High                                       | -0.44    | 0.34 | -1.28   | 0.201   |
| High-Low   | 0.67     | 0.30 | 2.24    | 0.025   |
| High-High  | -0.36    | 0.33 | -1.08   | 0.279   |
| Population: Non-anadromous background                | 1.32     | 0.25 | 5.18    | < 0.001 |
| Sex: Male  | 0.92     | 0.23 | 4.01    | < 0.001 |



**Figure 2:** Plasma chloride concentration (mmol/L) after 24 hr saltwater immersion of brown trout smolts (migratory tactic,  $n = 54$ ) derived from two population backgrounds. The median is represented by the white horizontal lines in each box.

**Table 3:** Parameter estimates with associated standard errors (SE) for the linear model testing effects of population, sex and food treatment on plasma chloride concentration (mmol/L) of brown trout classified as smolts ( $n = 54$ ). The reference level of each factor is in brackets, i.e. effects were contrasted against female fish from the anadromous-population background in the *Low-Low* food treatment. Note that no individuals were classed as having adopted the anadromous tactic in the *High-High* food treatment, and this category was dropped for this analysis. Statistical significance was assessed at  $p < 0.05$ .

| Effect   | Estimate | SE   | <i>t</i> -value | <i>p</i> -value |
|--|----------|------|-----------------|-----------------|
| Intercept (Low-Low, female, Anadromous background) | 148.81   | 3.85 | 38.68           | < 0.001         |
| Population   | 17.22    | 5.60 | 3.08            | 0.003           |
| Food: Low-High                                     | -9.99    | 5.85 | -1.71           | 0.094           |
| Food: High-Low                                     | 8.36     | 5.97 | 1.40            | 0.168           |
| Sex: Male  | 0.06     | 5.15 | 0.01            | 0.991           |

#### *Factors explaining variation in status traits at different time points*

At the time at which the food treatments were first applied, fish from both populations were in similar condition ( $F = 0.41$ ,  $df = 1,137$ ,  $p = 0.523$ ), however, anadromous-background fish were heavier ( $F = 17.14$ ,  $df = 1,137$ ,  $p < 0.001$ ) and longer ( $F = 16.31$ ,  $df = 1,137$ ,  $p < 0.001$ ) than non-anadromous-background fish. A mixed model analysis indicated further divergence in these status traits over the study period that was related to life-history tactics, food treatment, and population effects (Table 4, Figure 3). The models for length (marginal  $R^2 = 0.77$ ), weight (marginal  $R^2 = 0.62$ ), and  $K$  (marginal  $R^2 = 0.35$ ) retained a significant interaction between food treatment and population, and a significant interaction between life-history tactics and month (Table 4). Sex did not have a significant effect on length ( $\chi^2 = 0.024$ ,  $df = 1$ ,  $p = 0.877$ ), weight ( $\chi^2 = 0.050$ ,  $df = 1$ ,  $p = 0.823$ ), or condition factor ( $\chi^2 = 0.082$ ,  $df = 1$ ,  $p = 0.774$ ). After accounting for growth between measurement periods, (i.e. the fixed effect of measurement period), smolts tended to be shorter, lighter and have lower condition than mature fish (Table S3). The differences in length, weight and  $K$  were similar for both populations (an interaction between population and life-

history tactics was not retained in any of the final models, see Table 4). The significant interaction between food treatment and population indicated that fish from the anadromous-background were larger, and heavier (but in similar condition) than fish from the non-anadromous-background under both *High-Low* and *High-High* treatments (Table S3). However, in the *Low-Low* and *Low-High* treatments, there were negligible differences in length, weight and *K* between populations (Table S3). The significant interaction between month and life-history tactics indicated that changes in length, weight and *K* through time varied between smolts and mature fish. Mature fish tended to increase in length and weight quicker (Figure 3b, Table S3), while smolts tended to be in worse condition (lower *K*) earlier (Figure 3C Table S3). See Table S3 for all model outputs.

*Factors explaining variation in final values for status traits*

At the end of the study, fish differed in length, condition and lipid content according to food treatment, life-history tactics and population (Figure 4). The model describing length (marginal  $R^2 = 0.50$ ) retained a significant interaction between food treatment and population (Table 5) but did not indicate a significant effect of life-history tactics ( $\chi^2 = 2.83$ ,  $df = 1$ ,  $p = 0.093$ ), or sex ( $\chi^2 = 0.005$ ,  $df = 1$ ,  $p = 0.947$ ). The models describing condition (marginal  $R^2 = 0.56$ ) and whole body lipids (marginal  $R^2 = 0.73$ , Table 5) each retained an interaction between population and food treatment (Table 5), and included a significant effect of life-history tactics on condition ( $\chi^2 = 64.58$ ,  $df = 1$ ,  $p < 0.001$ ), and whole body lipids ( $\chi^2 = 7.71$ ,  $df = 1$ ,  $p = 0.005$ ). Sex did not have a significant effect on condition ( $\chi^2 = 3.43$ ,  $df = 1$ ,  $p = 0.064$ ) or whole body lipids ( $\chi^2 = 2.18$ ,  $df = 1$ ,  $p = 0.140$ ). Overall, smolts were of similar length to mature fish at the end of the study (Figure 4), but tended to be in poorer condition ( $p < 0.001$ , Table S4) and have slightly higher whole body lipids ( $p = 0.008$ , Table S4). We detected an interactive effect of food treatment and population, where fish from the anadromous-background population were larger than fish from the non-anadromous-background population, but similar under *Low-Low* food conditions (Table S4). However, non-anadromous-background fish were overall


in better condition ( $p = 0.011$ , Table S4) and had higher whole body lipids ( $p < 0.001$ , Table S4), and these differences between populations were strongest under conditions of *Low-Low* food (Table S4, Figure 4). The lack of significant interactions between life-history tactics and population in the models for length,  $K$ , and whole body lipids indicated that differences between populations were similar for both mature fish and smolts (Table 5). See Table S4 for all model outputs.

#### *Growth rate differences*

The somatic growth of fish during the experiment was well described by a logistic growth model. Initial model fitting indicated the most parsimonious model included separate growth parameters for smolts and mature fish. Mature fish had higher intrinsic growth rates ( $g_i = 0.0050$ ,  $SE = 0.0006$ ,  $p < 0.001$ ), a smaller asymptotic size ( $L_\infty = 25.44$ ,  $SE = 0.86$ ,  $p < 0.001$ ), and a lower point of inflection ( $I = 172.7$ ,  $SE = 13.8$ ,  $p < 0.001$ ) than smolts, where  $g_i = 0.0039 \pm SE$  0.0009 ( $p < 0.001$ ),  $L_\infty = 27.31 \pm SE$  4.13 ( $p < 0.001$ ) and  $I = 305.7 \pm SE$  89.9 ( $p = 0.001$ ). Mature individuals were relatively larger earlier in life than smolts, and had faster overall growth (Figure 5).

Growth differences between the two populations were also identified, where fish from the anadromous-background population were relatively larger earlier in the study than fish from the non-anadromous-background population, and grew faster (Figure 6). Anadromous-background fish had higher intrinsic growth rates ( $g_i = 0.0045$ ,  $SE = 0.0009$ ,  $p < 0.001$ ), similar asymptotic size ( $L_\infty = 26.83$ ,  $SE = 1.68$ ,  $p < 0.001$ ), and a lower point of inflection ( $I = 184.1$ ,  $SE = 26.9$ ,  $p < 0.001$ ) than non-anadromous-background fish, where  $g_i = 0.0043 \pm SE$  0.0007 ( $p < 0.001$ ),  $L_\infty = 26.45 \pm SE$  1.65 ( $p < 0.001$ ), and  $I = 236.3 \pm SE$  32.9 ( $p < 0.001$ ).






**Figure 3:** Trajectories of (a) length, (b) mass, and (c) condition factor ( $K$ ) of brown trout offspring (derived from wild-caught parents from two populations) that were classed as either smolt (migratory tactic) or freshwater maturing (non-migratory/resident) tactic. AB = anadromous-background population; non-AB = non-anadromous-background population. Mean values (with associated standard errors) are shown for measurements taken at key time points over the course of two years of tank rearing.

66o **Table 4:** Results of the mixed effect model analysis for length, weight and condition factor (*K*) trajectories of brown trout in the experiment with life-history classed as either smolts (i.e. migratory) or freshwater mature across the study period. The results of the model selection procedure on interaction terms are given, and the selected model for each response is highlighted in bold. The models included a random effect of individual identity and a first-order autoregressive correlation structure with respect to month was also modelled.


| Model   | df | AIC   | logLik | <i>L</i> -ratio | <i>p</i> -value |
|---|----|-------|--------|-----------------|-----------------|
| Length ~ month*life-history + population*life-history + population*food + sex   | 16 | 4277  | -2122  |                 |                 |
| <b>Length ~ month*life-history + population*food + sex</b>                      | 15 | 4276  | -2123  | 1.52            | 0.218           |
| Length ~ month + life-history + population*food + sex                           | 14 | 4279  | -2125  | 4.31            | 0.038           |
| Length ~ month + life-history + population + food + sex                         | 11 | 4306  | -2142  | 33.31           | < 0.001         |
| Weight ~ month*life-history + population*life-history + population*food + sex   | 16 | 10229 | -5099  |                 |                 |
| <b>Weight ~ month*life-history + population*food + sex</b>                      | 15 | 10228 | -5099  | 0.51            | 0.474           |
| Weight ~ month + life-history + population*food + sex                           | 14 | 10245 | -5109  | 19.37           | < 0.001         |
| Weight ~ month + life-history + population + food + sex                         | 11 | 10263 | -5120  | 23.45           | < 0.001         |
| <i>K</i> ~ month*life-history + population*life-history + population*food + sex | 16 | -1524 | 778    |                 |                 |
| <b><i>K</i> ~ month* life-history + population*food + sex</b>                   | 15 | -1525 | 778    | 0.86            | 0.354           |
| <i>K</i> ~ month + life-history + population*food + sex                         | 14 | -1514 | 771    | 12.77           | < 0.001         |
| <i>K</i> ~ month + life-history + population + food + sex                       | 11 | -1488 | 755    | 331.89          | < 0.001         |




**Figure 4:** Effects of food treatment on final (a) length, (b) condition factor ( $K$ ), and (c) whole body lipids at the end of the experimental study (Spring 2018) of brown trout offspring classed as either smolts (migratory) or freshwater maturing (non-migratory/resident). Offspring were derived from wild-caught parents from an anadromous-background population (AB) and a non-anadromous-background population (non-AB). The median is represented by the white horizontal lines in each box.

665 **Table 5:** Results of the mixed effect model analysis for length, condition factor (*K*), and whole body lipids of brown trout (life-history classed as either smolts or freshwater mature) at the end of the experimental study period. The results of the model selection procedure on interaction terms are given, and the selected model for each response is highlighted in bold. The models included a random effect of sample date.

| Model   | df | AIC  | logLik | <i>L</i> -ratio | <i>p</i> -value |
|---|----|------|--------|-----------------|-----------------|
| Length ~life-history*population + population*food + sex     | 13 | 2074 | -1024  |                 |                 |
| <b>Length ~ life-history + population*food + sex</b>        | 12 | 2076 | -1026  | 3.87            | 0.05            |
| Length ~ life-history + population + food + sex             | 9  | 2093 | -1037  | 22.98           | < 0.001         |
| <i>K</i> ~ life-history *population + population*food + sex | 13 | -798 | 411.91 |                 |                 |
| <b><i>K</i> ~ life-history + population*food + sex</b>      | 12 | -800 | 411.90 | 0.01            | 0.922           |
| <i>K</i> ~ life-history + population + food + sex           | 9  | -786 | 402.11 | 19.59           | < 0.001         |
| Lipids ~life-history *population + population*food + sex    | 13 | 489  | -231.4 |                 |                 |
| <b>Lipids ~ life-history + population*food + sex</b>        | 12 | 487  | -231.6 | 0.46            | 0.500           |
| Lipids ~ life-history + population + food + sex             | 9  | 503  | -242.6 | 21.94           | < 0.001         |



**Figure 5:** Growth curves, based on length measurements spanning two years of experimental tank-rearing, of brown trout classed as either smolt (migratory) or freshwater maturing (resident) in Spring 2018. Fitted lines are based on the best-fitting parameters from the logistic growth model, fitted using non-linear least squares regression. Shaded areas represent the 95% confidence intervals constructed by bootstrapping for 10,000 iterations.



**Figure 6:** Growth curves, based on length measurements spanning two years of experimental tank rearing, of brown trout derived from two population backgrounds (anadromous or non-anadromous). Fitted lines are based on the best-fitting parameters from the logistic growth model, fitted using non-linear least squares regression. Shaded areas represent the 95% confidence intervals constructed by bootstrapping for 10,000 iterations.

670

## Discussion

Salmonine fishes exhibit some of the most striking examples of animal migration, but uncertainty still surrounds the mechanisms by which alternative migratory tactics can be expressed, or inhibited, across salmonine populations.

675 A principle aim of our study was to assess the importance of food availability at  
different time points during early ontogeny in determining migratory/life-  
history tactics in two populations of brown trout. Food reduction across almost  
two years led to increased rates of smolting (migratory tactic) in fish from both  
population backgrounds, whilst no fish were classed as having adopted the  
680 migratory tactic in either population after two years of experiencing high food,  
i.e. optimal rations (Figure 1). Migratory/life-history tactics were also influenced  
by population background, consistent with an inherited component to  
migratory/life-history decisions – fish derived from a naturally anadromous  
population were more often classed as smolts in our experiment, while offspring  
685 derived from a naturally non-anadromous population were more often classed  
as non-smolts, or having undergone freshwater maturation consistent with a  
residency tactic. Intriguingly, the populations responded differently to the  
timing of food restriction, with fish from an anadromous population  
background seemingly having been more affected by food restriction in their  
690 second year, whilst fish from a non-anadromous population background were  
more affected by food restriction in their first year. Females were more likely  
than males to become smolts under all food treatments. Collectively, these  
results indicate both extrinsic (food-driven) and intrinsic effects (related to  
population background, sex, and other individual-level attributes) on  
695 migratory/life-history tactics in brown trout, that may interact in complex ways  
and influence how populations respond in the wild to changing environmental  
conditions.

Differences in growth and body condition were apparent from an early stage  
between fish adopting different life-history/migratory tactics, and were  
700 maintained across the full (almost two-year) duration of the study. These  
differences were in turn also driven by both extrinsic and intrinsic effects.  
Extrinsic effects were evidenced by the fact that large differences in fork length,  
mass, body condition, and whole body lipids were apparent between fish reared  
under different food treatments, which in turn contributed to fish adopting  
705 different life-history tactics via phenotypic plasticity. Intrinsic differences

among individuals in ‘status traits’ clearly also contributed to migratory/life-history outcomes, given that differences in body size, condition, and lipids were apparent between populations, and between fish from each population that adopted different tactics *within* each food treatment – where the external environment was the same. Such intrinsic variation within and between populations could reflect inherited genetic effects, inherited non-genetic effects (*e.g.* parental effects, epigenetic inheritance), or non-inherited differences driven by early-life environmental influences that have a relatively long-lasting effect on phenotype (Burton and Metcalfe 2014). Expanding our approach to incorporate even earlier life stages (*e.g.* post-hatching/fry) could further illuminate how factors in early life influence life history.

#### *Extrinsic factors*

The observed increases in smolting in the face of food restriction, together with decreases in maturation, suggested that the reduction in food supply prevented individuals from meeting an intrinsic (*e.g.* genetically determined) threshold for residency and maturity in freshwater, which is in agreement with previous studies (Olsson *et al.* 2006; Wysujack *et al.* 2009; O’Neal and Stanford 2011; Jones *et al.* 2015). Indeed, the absence of any smolts under conditions of high food supply was surprising, particularly within fish from the Tawnyard population (anadromous-background), which has a naturally high frequency of anadromy in the wild (Gargan *et al.* 2016). This suggests that, in nature, a large number of fish in the Tawnyard system must typically experience relatively low food availability as freshwater juveniles, as otherwise anadromy rates (broadly estimated as 50 – 60% in the wild population) would be lower in the wild. Moreover, the balance of fitness cost and benefits of migration in the system must be such that natural selection has caused a relatively high threshold for residency to evolve (an ultimate mechanism; Hazel *et al.* 1990; Tomkins and Hazel 2007; Pulido 2011), meaning a minority of Tawnyard fish in the wild typically surpass their intrinsic freshwater maturation threshold and the anadromous tactic is more frequent.



Manipulation of the timing of food reduction revealed that life-history responses of a given population to environmental change might depend on the point during ontogeny at which the change is experienced. This could come about via two non-mutually exclusive mechanisms: populations could exhibit  
 740 variation in sensitivity to cues experienced during given fixed “decision windows”, and/or the timing of the decision windows themselves may vary across populations. In our study, food restriction in the first year (*Low-High* treatment) was a more important driver of smolting rates than food in the second year (*High-Low*) for fish from the non-anadromous-background  
 745 population, whereas food in the second year was more important for the anadromous-background population. This was an intriguing outcome, and hints at a complex interplay between extrinsic environmental and intrinsic or population-specific factors. The apparently greater importance of food restriction in the first year for the non-anadromous-background population  
 750 could perhaps be related to lower intrinsic growth rates in this population in the wild. Given their low potential growth rates, individuals in the non-anadromous-background population might be constrained to make a life-history decision (i.e. choose future migration or residency) early in life in order to divert energy intake towards meeting the associated demands of the chosen  
 755 tactic. Because residents must accumulate sufficient lipid reserves to be converted into reproductive tissue before spawning (McMillan *et al.* 2012), in the wild, Bunaveela fish may have experienced selection for adopting a maturation trajectory relatively early in order to allow sufficient time for growth and energy accumulation, with early decision windows evolving as a  
 760 consequence. In contrast, fish from the anadromous-background population with higher intrinsic growth potential may be less constrained in this regard, and may defer choice of migratory tactics to the second year of life, or indeed have flexibly reversible life-history trajectories where, for example, fish choosing residency based on high food in year one may switch to migratory  
 765 tactics in response to low food in year two. There is some evidence for conditions in the second year of life being a key driver of migratory tactics in a

naturally facultatively anadromous brown trout population to support this (Cucherousset *et al.* 2005).

Coupled with a later “decision window”/higher sensitivity to conditions in year  
 770 two, a naturally high intrinsic growth propensity in the anadromous-  
 background population could have facilitated high levels of compensatory  
 growth when receiving optimal food resources in year two in the *Low-High*  
 treatment. If growth, or some aspect of energy usage related to growth such as  
 body condition, is used as a cue for migratory tactic choice, this may then have  
 775 translated into more individuals from this population meeting their threshold  
 for maturation in the *Low-High* treatment. Strong compensatory responses after  
 periods of food restriction have been observed in salmonids in general, and  
 interestingly, the compensatory response has often appeared to be directed  
 towards restoring body condition, rather than size. Nicieza and Metcalfe (1997)  
 780 found food restricted fish recovered similar condition to controls within a year  
 of food supply restoration, and Álvarez and Nicieza (2005) further found a  
 compensatory response that resulted in restoration of condition and energy  
 status rather than skeletal growth in brown trout post food restriction.

Alternatively, we cannot rule out the presence of multiple migration versus  
 785 residency decision windows, that re-occur annually or more frequently,  
 whereby an individual repeatedly re-assesses its status trait relative to its  
 inherited freshwater maturation threshold and can remain ‘undecided’ at the  
 first or even second windows, though there is little empirical evidence for this.  
 A simpler explanatory model is that there is a single, initial decision  
 790 determining migration versus residency, and then subsequent decision  
 windows occur for fish on each trajectory (migrants and resident) related to the  
 timing of expression of the adopted life-history tactic, where for example  
 migrants must decide at what age to actually migrate (determined by pressures  
 of size-dependent sea survival), or indeed *where* to migrate (Ferguson *et al.*  
 795 2019). Similarly, a resident individual must also decide *when* to mature (Thorpe  
*et al.* 1998; Thorpe and Metcalfe 1998), a decision shown to be affected by lipid  
 reserves in Atlantic salmon (Rowe *et al.* 1991; Jonsson and Jonsson 1993, 2005)

and possibly triggered by similar threshold type mechanisms in brown trout. These timing decisions could be further influenced by extrinsic environmental conditions, giving rise to a temporal continuum of migration and maturation tactics. This may explain why some fish in our study were classified as having an undetermined life history (neither smolt nor mature) by the spring of year two: these individuals may simply have been delaying expression of a migratory or freshwater maturing phenotype until the following year. These *caveats* must be born in mind when interpreting our experimental results, as the life-history tactic frequencies we measured in year 2 could be indicative of age-specific tactic frequencies, rather than overall rates of migration versus residency across all ages. However, the basic conclusions were the same in the GLMs where the data were analysed as either smolt versus non-smolt, or immature versus mature, giving us confidence that the patterns reflect the migration decision *per se*.

#### *Variation in status traits underpinning alternative tactics*

Size-based differences between migrating individuals (those classified at the end of the study as smolts) and resident fish (those classified at the end of the study as mature) were established relatively early, with differences in weight, length, and condition that were maintained during the course of the study. The early divergence in physiological condition between migrants and residents supports the energy limitation scenario, where fish adopt migration as a result of failing to meet the necessary condition in early life to mature as residents in freshwater (Jonsson and Jonsson 1993). Maturing fish reached an apparent size asymptote earlier than migrating fish (i.e. had smaller inflection point in Figure 5, and were larger earlier in the study). Size appears to be a potential status trait that regulates, or correlates with factors regulating early sexual maturation, as has been documented in Atlantic salmon, where anadromous males are smaller than their counterparts that mature early in freshwater as so-called ‘precocious parr’ (Whalen and Parrish 1999; Garant *et al.* 2002). However, although body size has been suggested as a major component of the status (cueing) trait for anadromy in brook trout (Thériault *et al.* 2007), the divergence in mass and

condition we found in our study suggests that other factors beyond size also  
 830 contribute to the maturation versus migration/anadromy decision. It seems  
 increasingly likely that a suite of interlinked physiological components is  
 assessed (*e.g.* overall energetic status or rate of change in energy), and no single  
 trait controls the migratory/anadromy decision. Genetic covariance between  
 life history traits such as growth, size, metabolism, and other morphological  
 835 traits further suggests that migration decisions are associated with a suite of  
 inter-linked phenotypic traits (Doctor *et al.* 2014; Hecht *et al.* 2015).

Fish on a migratory trajectory here appeared to maintain growth rates during  
 the experiment (and had a higher inflection point), such that they were similar  
 in length to mature fish by the end of our study. Constant, or even accelerated  
 840 growth in pre-migratory fish (Metcalf 1998) has been explained by size-  
 dependent survival at sea (Klemetsen *et al.* 2003) due to better osmoregulation  
 ability (Finstad and Ugedal 1998) and reduced predation of larger anadromous  
 individuals (Dill 1983; Jonsson *et al.* 2017). Interestingly here, although skeletal  
 growth (*i.e.* length) was maintained, migratory fish were considerably lighter  
 845 and in worse condition than mature fish at the end of study, which suggests that  
 once on a migratory trajectory, resources were primarily allocated to meeting a  
 size-based threshold for surviving actual migration. The maintenance of growth  
 rates in migrants as such does not contradict the energy limitation scenario, but  
 rather suggests that migratory fish redirect what resources they obtain into  
 850 becoming large enough to survive the migration, at a cost to their overall body  
 condition.

The diminished body condition of migratory individuals was not, however,  
 reflected in levels of whole-body lipids at the end of the study. Contrary to our  
 expectations, migratory fish had marginally higher levels of whole body lipids  
 855 than mature fish. Lipid storage has been identified previously as an important  
 precursor of maturation in fish (Tocher 2003) and an indicator of a residency  
 life history in salmonids (Sloat and Reeves 2014; Sloat *et al.* 2014 and references  
 therein). The unexpected trend we observed in lipids may have been a  
 consequence of measuring lipids during the smolt migration period, at which

860 stage fish that have initiated maturation might have already converted some of their energy stores into gonadal tissue, and hence show depleted lipids levels relative to migrants (Tocher 2003; Sloat and Reeves 2014). Alternatively, higher lipid levels in migrants could reflect accumulation of reserves, as either a bet-hedging strategy if resources in the migration destination are uncertain, or to  
 865 fuel the migration journey itself (Stefansson *et al.* 2003). Pre-migratory “fattening” strategies are relatively common in migratory birds (Piersma *et al.* 2005) but less so in salmonines (Jonsson and Jonsson 2005).

### *Intrinsic factors*

We had predicted that the two populations in our study would show variability  
 870 in adopting migratory tactics across all food restriction scenarios and indeed, overall, the probability of smolting was higher in the anadromous background population than in the non-anadromous population. Moreover, higher hypo-osmoregulatory function (lower plasma chloride concentration) was documented in smolts from the former population relative to the latter,  
 875 implying that smolts from the anadromous-background population were physiologically better prepared for transition to marine conditions. In contrast, although some fish from the non-anadromous-background population were classified as smolts in the experiment, these putative smolts exhibited relatively lower saltwater tolerance. A potential explanation for the reduced hypo-  
 880 osmoregulatory function of non-anadromous-background smolts might be that they are poorly adapted to saltwater given their lack of (recent) evolutionary exposure to marine conditions. Relaxed selection leading to degradation of hypo-osmoregulation has similarly been observed in non-anadromous populations of landlocked Atlantic salmon (Nilsen *et al.* 2008; McCormick *et al.* 2019) and alewife *Alosa pseudoharengus* (Velotta *et al.* 2014, 2015).  
 885 Alternatively, reduced saltwater tolerance could be evidence of an emerging migration continuum whereby putative smolts may have chosen a potamodromous (freshwater migratory) tactic and hence were unprepared physiologically for transitioning to saltwater. Nevertheless, the causal  
 890 mechanisms underpinning anadromy and potamodromy are proposed to be

similar, *e.g.* reduced food availability has previously been reported to increase adfluvial migration in freshwater brown trout transplanted to streams of high population density (Olsson *et al.* 2006). All brown trout in Ireland presumably have anadromous ancestral origins, since they would have had to recolonise the island after the Last Glacial Maximum via the sea (Ferguson *et al.* 2019). It thus seems more likely that the capacity for anadromy (or at least migration), albeit somewhat deteriorated in terms of saltwater tolerance, lay dormant in the Bunaveela fish, with anadromy re-expressed under experimental conditions of energy limitation.

The putative re-emergence of an anadromous life history in our Bunaveela fish is of particular interest from a fisheries management perspective, as it suggests the capacity for anadromy (or at least migration) may lie dormant within apparently resident populations. Such populations may thus have the potential to contribute to the restoration of anadromous stocks that have experienced widespread reductions, as evidenced by Gargan *et al.* (2006) in two formerly anadromous populations that suffered collapses. Anadromous phenotypes arising from resident genotypes have similarly been documented in *O. mykiss* (Kelson *et al.* 2019), and from common garden experiments with lake resident *O. mykiss* which were formally anadromous but were prevented from migrating by impassable dams or waterfalls (Thrower *et al.* 2004). These findings make sense within the framework of the conditional threshold model (Tomkins and Hazel 2007), where environmental factors can affect life history tactic frequency by changing the distribution of the realised physiological state relative to inherited switch points (a proximate mechanism). Environmental factors could also drive longer term changes in tactic frequency via natural selection acting to shift the genotypic distribution of underlying switch points (an ultimate mechanism) (Hazel *et al.* 1990; Tomkins and Hazel 2007; Pulido 2011); for example, if survival or growth at sea is poor then migration may become less prevalent in the population if residents attain higher overall relative fitness than migrants. Within the Burrishoole system, the establishment of an Atlantic salmon farm in the estuary was implicated in the collapse of the

anadromous life history from this catchment over a period of 30 years due to high rates of sea lice transmission (Poole *et al.* 1996, 2007). Reduced marine survival rates may have imposed strong selection against anadromy, and hence  
 925 caused the evolution of lower mean threshold values for freshwater maturation within the Burrishoole catchment as a whole. Our current results are consistent with this evolutionary explanation, in that we demonstrated heritable differences (or at least phenotypic differences among genetically divergent populations in a common garden experiment) – a pre-requisite for evolutionary  
 930 responses. However, they also show that phenotypic plasticity can drive changes in migratory tactics, which may contribute to observed life-history changes in natural populations (Gargan *et al.* 2006; Sandlund and Jonsson 2016).

Early-life differences in length and mass between the two populations may  
 935 proximately cause different anadromy propensities, as has been seen in brook trout, where size of juvenile fish was negatively related to probability of future residency (Thériault *et al.* 2007). Interestingly, though our populations differed in size early in the study (before food restriction), they were in similar condition at this time, suggesting that both populations had similar energy intake *versus*  
 940 output, at least initially. Higher intrinsic growth rates in the anadromous background population may have increased the likelihood of eventual energetic limitation in freshwater, thus reducing relative condition and increasing anadromy propensity (exemplified in our *Low-Low* food treatment). Conversely, when food resources are in ample supply, high intrinsic growth rates could  
 945 hasten freshwater maturity instead of anadromy in this population (*c.f.* the scenario of optimal food resources in our study). Such variability in migratory tactics is a feature of salmonines in general [*e.g.* “retirement” from anadromy in Dolly Varden *Salvelinus malma* (Bond *et al.* 2015)], which may buffer species from increasing anthropogenic pressures in the marine environment (Russell *et*  
 950 *al.* 2012).

### Conclusions

Collectively, the results of this study show that the adoption of migratory tactics in brown trout involves an interplay between inherited components and environmentally cued physiological condition, in line with previous salmonines studies (Chapman *et al.* 2012; Dodson *et al.* 2013; Kendall *et al.* 2014). The differences we observed in population responses to food restriction and its timing suggest a complex relationship between intrinsic and extrinsic factors that may allow for a continuum of migratory tactics to exist. These population differences, together with the fact that putative anadromy emerged within offspring of a naturally non-anadromous population, emphasise that a range of life-history outcomes are possible even within a single species, which can contribute to so-called portfolio effects that cushion the species as a whole from rapidly changing environmental conditions (Schindler *et al.* 2015). Although our study offers some important insight into how extrinsic and intrinsic factors interactively shape life-history tactics, we have only considered one element of the freshwater environment here. Future studies should expand to consider how other proximate drivers such as temperature, which influences a range of physiological and life-history traits in salmonines (Satterthwaite *et al.* 2010; McMillan *et al.* 2012; Sloat and Reeves 2014; Doctor *et al.* 2014; Kendall *et al.* 2014), govern migratory tactics in fish from different genetic backgrounds. Moreover, it is now important to expand this approach into natural systems using, for example, common garden or reciprocal transplant experiments, to assess whether these findings hold up under real world complexities.

Finally, our results have important implications for the conservation of facultatively migratory species, which are in global decline due to in-stream barriers, habitat degradation, climate change, overfishing and the expansion of aquaculture (Costello 2009; Limburg and Waldman 2009). Knowledge of how extrinsic and intrinsic factors affect fish migratory tactics may aid in successful management and restoration of facultatively migratory populations, and in doing so maintain important intraspecific biocomplexity, which offers increased resilience to effects of global change (Schindler *et al.* 2015).



## Acknowledgements

The authors would like to thank Brian Clarke, Deirdre Cotter, members of the FishEyE team at UCC, and the staff of Inland Fisheries Ireland and the Marine  
 985 Institute for obtaining brood stock and for assistance in fish rearing, along with Robert Wynne, Ronan O'Sullivan, Peter Moran and Adam Kane for assistance in fish husbandry, and Jamie Coughlan for genotyping work. This research was supported by an ERC Starting Grant (639192-ALH) and an SFI ERC Support Award awarded to TER. PMcG was supported in part by grants from Science  
 990 Foundation Ireland (15/IA/3028 & 16/BBSRC/3316) and by grant-in-aid (RESPI/FS/16/01) from the Marine Institute (Ireland) as part of the Marine Research Programme by the Irish Government. We thank the Associate Editor and two reviewers for comments that improved the manuscript.

## References

- 995 Álvarez D and Nicieza AG. 2005. Compensatory response 'defends' energy levels but not growth trajectories in brown trout, *Salmo trutta* L. *Proceedings of the Royal Society of London B: Biological Sciences* **272**: 601–7.
- Barton K. 2018. MuMIn: Multi-Model Inference. R package version 1.42.1. <https://CRAN.R-project.org/package=MuMIn>.
- 1000 Beakes MP, Satterthwaite WH, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, and Mangel M. 2010. Smolt transformation in two California steelhead populations: Effects of temporal variability in growth. *Transactions of the American Fisheries Society* **139**: 1263–75.
- 1005 Berejikian BA, Bush RA, and Campbell LA. 2014. Maternal control over offspring life history in a partially anadromous species, *Oncorhynchus mykiss*. *Transactions of the American Fisheries Society* **143**: 369–79.
- Bond MH, Miller JA, and Quinn TP. 2015. Beyond dichotomous life histories in partially migrating populations: Cessation of anadromy in a long-lived fish. *Ecology* **96**: 1899–910.
- 1010 Buoro M, Gimenez O, and Prévost E. 2012. Assessing adaptive phenotypic plasticity by means of conditional strategies from empirical data: The latent environmental threshold model. *Evolution* **66**: 996–1009.
- Burton T and Metcalfe NB. 2014. Can environmental conditions experienced in early life influence future generations? *Proceedings of the Royal Society of London B: Biological Sciences* **281**: 20140311.
- 1015 Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011a. Partial migration: An introduction. *Oikos* **120**: 1761–3.
- Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011b. The ecology and evolution of partial migration. *Oikos* **120**: 1764–75.

- 1020 Chapman BB, Hulthén K, Brodersen J, Nilsson PA, Skov C, Hansson L-A, and Brönmark C. 2012. Partial migration in fishes: Causes and consequences. *Journal of Fish Biology* **81**: 456–78.
- Costello MJ. 2009. How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *Proceedings of the Royal Society B: Biological Sciences* **276**: 3385–94.
- 1025 Cucherousset J, Ombredane D, Charles K, Marchand F, and Baglinière J-L. 2005. A continuum of life history tactics in a brown trout (*Salmo trutta*) population. *Canadian Journal of Fisheries and Aquatic Sciences* **62**: 1600–10.
- Dill LM. 1983. Adaptive flexibility in the foraging behavior of fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **40**: 398–408.
- 1030 Doctor K, Berejikian B, Hard JJ, and VanDoornik D. 2014. Growth-mediated life history traits of steelhead reveal phenotypic divergence and plastic response to temperature. *Transactions of the American Fisheries Society* **143**: 317–33.
- Dodson JJ, Aubin-Horth N, Thériault V, and Páez DJ. 2013. The evolutionary ecology of alternative migratory tactics in salmonid fishes: Alternative migratory tactics as threshold traits. *Biological Reviews* **88**: 602–25.
- 1035 Doswald N, Willis SG, Collingham YC, Pain DJ, Green RE, and Huntley B. 2009. Potential impacts of climatic change on the breeding and non-breeding ranges and migration distance of European Sylvia warblers. *Journal of Biogeography* **36**: 1194–208.
- 1040 Ferguson A, Reed TE, Cross TF, McGinnity P, and Prodöhl PA. 2019. Anadromy, potamodromy and residency in brown trout *Salmo trutta*: The role of genes and the environment. *Journal of Fish Biology* **95**: 692–718.
- Ferguson A, Reed TE, McGinnity P, and Prodöhl P. 2017. Anadromy in brown trout (*Salmo trutta*): A review of the relative roles of genes and environmental factors and the implications for management and conservation. In: *Sea Trout: Science and Management - Proceedings of the 2nd International Sea Trout Symposium*. Matador, Leicestershire, UK.
- 1045 Finstad B and Ugedal O. 1998. Smolting of sea trout *Salmo trutta* L. in northern Norway. *Aquaculture* **168**: 341–9.
- Forseth T, Nesje TF, Jonsson B, and Hårsaker K. 1999. Juvenile migration in brown trout: A consequence of energetic state. *Journal of Animal Ecology* **68**: 783–93.
- Garant D, Fontaine P-M, Good SP, Dodson\* JJ, and Bernatchez L. 2002. The influence of male parental identity on growth and survival of offspring in Atlantic salmon (*Salmo salar*). *Evolutionary Ecology Research* **4**: 537–49.
- 1055 Gargan P, Kelly F, Shephard S, and Whelan K. 2016. Temporal variation in sea trout *Salmo trutta* life history traits in the Erriff River, western Ireland. *Aquaculture Environment Interactions* **8**: 675–89.
- Gargan P, Roche W, Frode G, and Ferguson A. 2006. Characteristics of sea trout (*Salmo trutta* L.) stocks from the Owengowla and Invermore Fisheries, western Ireland, and recent trends in marine survival. In: *Sea Trout: Biology, Conservation and Management*. Oxford, UK: Blackwells Scientific Publications.
- 1060

- Hazel WN, Smock R., and Johnson M. D. 1990. A polygenic model for the evolution and maintenance of conditional strategies. *Proceedings of the Royal Society of London Series B: Biological Sciences* **242**: 181–7.
- Hecht BC, Hard JJ, Thrower FP, and Nichols KM. 2015. Quantitative genetics of migration-related traits in rainbow and steelhead trout. *G3: Genes, Genomes, Genetics* **5**: 873–89.
- Heinze G and Schemper M. 2002. A solution to the problem of separation in logistic regression. *Statistics in Medicine* **21**: 2409–19.
- Hutchings JA. 2011. Old wine in new bottles: Reaction norms in salmonid fishes. *Heredity* **106**: 421–37.
- Hutchings JA and Myers RA. 1994. The evolution of alternative mating strategies in variable environments. *Evolutionary Ecology* **8**: 256–268.
- Jones DA, Bergman E, and Greenberg L. 2015. Food availability in spring affects smolting in brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 1694–9.
- Jonsson B. 1985. Life history patterns of freshwater resident and sea-run migrant brown trout in Norway. *Transactions of the American Fisheries Society* **114**: 182–94.
- Jonsson B and Jonsson N. 1993. Partial migration: Niche shift versus sexual maturation in fishes. *Reviews in Fish Biology and Fisheries* **3**: 348–65.
- Jonsson B and Jonsson N. 2005. Lipid energy reserves influence life-history decision of Atlantic salmon (*Salmo salar*) and brown trout (*S. trutta*) in fresh water. *Ecology of Freshwater Fish* **14**: 296–301.
- Jonsson B, Jonsson M, and Jonsson N. 2017. Influences of migration phenology on survival are size-dependent in juvenile Atlantic salmon (*Salmo salar*). *Canadian Journal of Zoology* **95**: 581–7.
- Kelson SJ, Miller MR, Thompson TQ, O'Rourke SM, and Carlson SM. 2019. Do genomics and sex predict migration in a partially migratory salmonid fish, *Oncorhynchus mykiss*? *Canadian Journal of Fisheries and Aquatic Sciences* **76**: 2080–8.
- Kendall NW, McMillan JR, Sloat MR, Buehrens TW, Quinn TP, Pess GR, Kuzishchin KV, McClure MM, and Zabel RW. 2014. Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): A review of the processes and patterns. *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 319–42.
- Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, and Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): A review of aspects of their life histories. *Ecology of Freshwater Fish* **12**: 1–59.
- Kosmidis I. 2019. brglm: Bias Reduction in Binary-Response Generalized Linear Models. R package version 0.6.2, <URL: <https://cran.r-project.org/package=brglm>>.
- Lenth R. 2019. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.3.5.1. <https://CRAN.R-project.org/package=emmeans>.
- Limburg KE and Waldman JR. 2009. Dramatic declines in north Atlantic diadromous fishes. *BioScience* **59**: 955–65.

- Magee J. 2017. A comparison of population structuring and genetic stock identification of brown trout (*Salmo trutta*) displaying distinct migratory strategies. In: PhD thesis, pp277. Queens University Belfast, Belfast, NI.
- 1110 McCormick SD, Regish AM, Ardren WR, Björnsson BT, and Bernier NJ. 2019. The evolutionary consequences for seawater performance and its hormonal control when anadromous Atlantic salmon become landlocked. *Scientific Reports* **9**: 968.
- 1115 McMillan JR, Dunham JB, Reeves GH, Mills JS, and Jordan CE. 2012. Individual condition and stream temperature influence early maturation of rainbow and steelhead trout, *Oncorhynchus mykiss*. *Environmental Biology of Fishes* **93**: 343–55.
- 1120 Metcalfe NB. 1998. The interaction between behavior and physiology in determining life history patterns in Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* **55**: 93–103.
- Morinville GR and Rasmussen JB. 2003. Early juvenile bioenergetic differences between anadromous and resident brook trout (*Salvelinus fontinalis*). *Canadian Journal of Fisheries and Aquatic Sciences* **60**: 401–10.
- 1125 Naish KA and Hard JJ. 2008. Bridging the gap between the genotype and the phenotype: Linking genetic variation, selection and adaptation in fishes. *Fish and Fisheries* **9**: 396–422.
- Nicieza AG and Metcalfe NB. 1997. Growth compensation in juvenile Atlantic salmon: Responses to depressed temperature and food availability. *Ecology* **78**: 2385–400.
- 1130 Nilsen TO, Ebbesson LOE, Kiilerich P, Björnsson BTh, Madsen SS, McCormick SD, and Stefansson SO. 2008. Endocrine systems in juvenile anadromous and landlocked Atlantic salmon (*Salmo salar*): Seasonal development and seawater acclimation. *General and Comparative Endocrinology* **155**: 762–72.
- 1135 Olsson IC, Greenberg LA, Bergman E, and Wysujack K. 2006. Environmentally induced migration: The importance of food. *Ecology Letters* **9**: 645–51.
- O’Neal SL and Stanford JA. 2011. Partial migration in a robust brown trout population of a Patagonian river. *Transactions of the American Fisheries Society* **140**: 623–35.
- 1140 Padfield D and Matheson G. 2018. nls.multstart: Robust Non-Linear Regression using AIC Scores. R package version 1.0.0. <https://CRAN.R-project.org/package=nls.multstart>.
- Phillis CC, Moore JW, Buoro M, Hayes SA, Garza JC, and Pearse DE. 2016. Shifting thresholds: Rapid evolution of migratory life histories in steelhead/rainbow trout, *Oncorhynchus mykiss*. *Journal of Heredity* **107**: 51–60.
- 1145 Piche J, Hutchings JA, and Blanchard W. 2008. Genetic variation in threshold reaction norms for alternative reproductive tactics in male Atlantic salmon, *Salmo salar*. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1571–5.
- 1150 Piersma T, Pérez-Tris J, Mouritsen H, Bauchinger U, and Bairlein F. 2005. Is there a “migratory syndrome” common to all migrant birds? *Annals of the New York Academy of Sciences* **1046**: 282–93.

- Pinheiro J, Bates D, DebRoy S, Sarkar D, and R Core Team. 2019. nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-139. URL <https://CRAN.R-project.org/package=nlme>.
- 1155 Poole WR, Dillane M, DeEyto E, Rogan G, McGinnity P, and Whelan K. 2007. Characteristics of the Burrishoole sea trout population: Census, marine survival, enhancement and stock-recruitment relationship, 1971-2003. In: Sea Trout: Biology, Conservation and Management (Eds: G Harris and N Milner). Oxford, UK: Blackwells Scientific Publications.
- 1160 Poole WR, Whelan KF, Dillane MG, Cooke DJ, and Matthews M. 1996. The performance of sea trout, *Salmo trutta* L., stocks from the Burrishoole system western Ireland, 1970-1994. *Fisheries Management and Ecology* **3**: 73-92.
- Pulido F. 2011. Evolutionary genetics of partial migration – the threshold model of migration revisited. *Oikos* **120**: 1776-83.
- 1165 Pulido F, Berthold P, and vanNoordwijk AJ. 1996. Frequency of migrants and migratory activity are genetically correlated in a bird population: Evolutionary implications. *Proceedings of the National Academy of Sciences of the United States of America* **93**: 14642-7.
- 1170 R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Roff DA. 1996. The evolution of threshold traits in animals. *The Quarterly Review of Biology* **71**: 3-35.
- 1175 Rowe DK, Thorpe JE, and Shanks AM. 1991. Role of fat stores in the maturation of male Atlantic salmon (*Salmo salar*) parr. *Canadian Journal of Fisheries and Aquatic Sciences* **48**: 405-13.
- Russell IC, Aprahamian MW, Barry J, Davidson IC, Fiske P, Ibbotson AT, Kennedy RJ, Maclean JC, Moore A, Otero J, Potter T (E CE), and Todd CD. 2012. The influence of the freshwater environment and the biological characteristics of Atlantic salmon smolts on their subsequent marine survival. *ICES Journal of Marine Science: Journal du Conseil*: fsr208.
- 1180 Sandlund OT and Jonsson B. 2016. Life history plasticity: Migration ceased in response to environmental change? *Ecology of Freshwater Fish* **25**: 225-33.
- 1185 Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, and Mangel M. 2010. State-dependent life history models in a changing (and regulated) environment: Steelhead in the California Central Valley. *Evolutionary Applications* **3**: 221-43.
- Schindler DE, Armstrong JB, and Reed TE. 2015. The portfolio concept in ecology and evolution. *Frontiers in Ecology and the Environment* **13**: 257-63.
- 1190 Schultz ET and McCormick SD. 2012. Euryhalinity in an evolutionary context. In: Fish physiology. Cambridge, Massachusetts: Academic Press.
- Sloat MR, Fraser DJ, Dunham JB, Falke JA, Jordan CE, McMillan JR, and Ohms HA. 2014. Ecological and evolutionary patterns of freshwater maturation in Pacific and Atlantic salmonines. *Reviews in Fish Biology and Fisheries* **24**: 689-707.
- 1195 Sloat MR and Reeves GH. 2014. Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus*

- mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences* **71**: 491–501.
- 1200 Stefansson SO, Björnsson BT, Sundell K, Nyhammer G, and McCormick SD. 2003. Physiological characteristics of wild Atlantic salmon post-smolts during estuarine and coastal migration. *Journal of Fish Biology* **63**: 942–55.
- Tanguy JM, Ombredane D, Baglinière JL, and Prunet P. 1994. Aspects of parr-smolt transformation in anadromous and resident forms of brown trout (*Salmo trutta*) in comparison with Atlantic salmon (*Salmo salar*). *Aquaculture* **121**: 51–63.
- 1205 Thériault V and Dodson JJ. 2003. Body size and the adoption of a migratory tactic in brook charr. *Journal of Fish Biology* **63**: 1144–59.
- Thériault V, Garant D, Bernatchez L, and Dodson JJ. 2007. Heritability of life-history tactics and genetic correlation with body size in a natural population of brook charr (*Salvelinus fontinalis*). *Journal of Evolutionary Biology* **20**: 2266–77.
- 1210 Thorpe JE, Mangel M, Metcalfe NB, and Huntingford FA. 1998. Modelling the proximate basis of salmonid life-history variation, with application to Atlantic salmon, *Salmo salar* L. *Evolutionary Ecology* **12**: 581–99.
- Thorpe JE and Metcalfe NB. 1998. Is smolting a positive or a negative developmental decision? *Aquaculture* **168**: 95–103.
- 1215 Thrower FP, Hard JJ, and Joyce JE. 2004. Genetic architecture of growth and early life-history transitions in anadromous and derived freshwater populations of steelhead. *Journal of Fish Biology* **65**: 286–307.
- Tocher DR. 2003. Metabolism and functions of lipids and fatty acids in teleost fish. *Reviews in Fisheries Science* **11**: 107–84.
- 1220 Tomkins JL and Hazel W. 2007. The status of the conditional evolutionarily stable strategy. *Trends in Ecology & Evolution* **22**: 522–8.
- Urke HA, Koksvik J, Arnekleiv JV, Hindar K, Kroglund F, and Kristensen T. 2010. Seawater tolerance in Atlantic salmon, *Salmo salar* L., brown trout, *Salmo trutta* L., and *S. salar* × *S. trutta* hybrids smolt. *Fish Physiology and Biochemistry* **36**: 845–53.
- 1225 Velotta JP, McCormick SD, O'Neill RJ, and Schultz ET. 2014. Relaxed selection causes microevolution of seawater osmoregulation and gene expression in landlocked Alewives. *Oecologia* **175**: 1081–92.
- 1230 Velotta JP, McCormick SD, and Schultz ET. 2015. Trade-offs in osmoregulation and parallel shifts in molecular function follow ecological transitions to freshwater in the Alewife. *Evolution* **69**: 2676–88.
- Whalen KG and Parrish DL. 1999. Effect of maturation on parr growth and smolt recruitment of Atlantic salmon. *Canadian Journal of Fisheries and Aquatic Sciences* **56**: 79–86.
- 1235 Wysujack K, Greenberg LA, Bergman E, and Olsson IC. 2009. The role of the environment in partial migration: Food availability affects the adoption of a migratory tactic in brown trout *Salmo trutta*. *Ecology of Freshwater Fish* **18**: 52–9.
- 1240 Zimmermann CE and Reeves GH. 2000. Population structure of sympatric anadromous and nonanadromous *Oncorhynchus mykiss*: Evidence from spawning surveys

and otolith microchemistry. *Canadian Journal of Fisheries and Aquatic Sciences* **57**: 2152–62.

## Supporting Information

---

### **The interplay between extrinsic and intrinsic factors in determining flexible decisions in brown trout (*Salmo trutta*): An experimental study**

Louise C. Archer<sup>\*1,2</sup>, Stephen Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, Michael N. O'Grady<sup>3</sup>, Joseph P. Kerry<sup>3</sup>, W. Russell Poole<sup>4</sup>, Paddy Gargan<sup>5</sup>, Philip McGinnity<sup>1,4</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> *School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.*


<sup>2</sup> *Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland*

<sup>3</sup> *Food Packaging Group, School of Food and Nutritional Sciences, University College Cork, Cork, Ireland.*

<sup>4</sup> *Marine Institute, Furnace, Newport, Co. Mayo, Ireland.*

<sup>5</sup> *Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.*





**Figure S1:** Locations in the west of Ireland of brown trout brood stock collected by seine netting in winter 2015, and used to produce F1 offspring for an experimental tank-rearing study. Fish with a non-anadromous-population background were offspring of brown trout collected from (a) Bunaveela Lough in the Burrishoole catchment (no natural occurrence of anadromy). Fish with an anadromous-population background were offspring of brood stock collected in (b) Tawnyard Lough in the Erriff catchment (a high natural frequency of anadromy).

**Table SI:** Brood stock crossing design for producing F1 offspring of wild-origin brown trout collected by seine netting from two populations (Bunaveela Lough and Tawnyard Lough) in the west of Ireland. Each female was stripped of eggs, which were then fertilised by the milt of two males from the same population and incubated in a hatchery facility within the Burrishoole catchment. Offspring produced from the Bunaveela population were considered to have a non-anadromous-population background, and offspring produced from the Tawnyard population were considered to have an anadromous-population background.

| Stripping date | Population | Catchment   | Sex    | ID   | Male cross #1 | Male cross #2 |
|----------------|------------|-------------|--------|------|---------------|---------------|
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF1  | BM1           | BM2           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF2  | BM1           | BM2           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF3  | BM3           | BM4           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF4  | BM3           | BM4           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF5  | BM5           | BM6           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF6  | BM5           | BM6           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF7  | BM7           | BM8           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF8  | BM7           | BM8           |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF9  | BM9           | BM10          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF10 | BM9           | BM10          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF11 | BM11          | BM12          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF12 | BM11          | BM12          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF13 | BM13          | BM15          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF14 | BM13          | BM15          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF15 | BM17          | BM18          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF16 | BM17          | BM18          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF17 | BM19          | BM20          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF18 | BM19          | BM20          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF19 | BM22          | BM23          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF20 | BM22          | BM23          |
| 22/12/2015     | Bunaveela  | Burrishoole | Female | BF21 | BM25          | BM26          |
| 27/11/2015     | Tawnyard   | Erriff      | Female | TF4  | TM6           | TM7           |
| 27/11/2015     | Tawnyard   | Erriff      | Female | TF5  | TM6           | TM7           |
| 27/11/2015     | Tawnyard   | Erriff      | Female | TF6  | TM9           | TM10          |
| 10/12/2015     | Tawnyard   | Erriff      | Female | TF7  | TM12          | TM13          |

**Table S2:** Feeding guidelines (as % body mass fed per day), used to adjust absolute daily feeding rations on a monthly basis according to body mass and rearing temperature of brown trout reared in a recirculating aquaculture system for 22 months. Type of feed (based on fish size) is listed in italics (Skretting, Norway).

| Fish mass (g)     | Rearing temperature |      |      |       |       |       |       |       |
|-------------------|---------------------|------|------|-------|-------|-------|-------|-------|
|                   | 4 °C                | 6 °C | 8 °C | 10 °C | 12 °C | 14 °C | 16 °C | 18 °C |
| <i>Nutra Parr</i> |                     |      |      |       |       |       |       |       |
| 2 - 6.5g          | 1.6                 | 1.9  | 2.2  | 2.6   | 3     | 3.4   | 3.8   | NA    |
| 6.5 - 8g          | 1.5                 | 1.8  | 2.1  | 2.5   | 2.9   | 3.3   | 3.7   | NA    |
| <i>Horizon</i>    |                     |      |      |       |       |       |       |       |
| 8 - 12g           | 1.21                | 1.53 | 1.73 | 2.1   | 2.42  | 2.78  | 3.06  | 3.57  |
| 12 - 21g          | 1.21                | 1.36 | 1.57 | 1.87  | 2.16  | 2.55  | 2.84  | 3.34  |
| <i>Elite FR</i>   |                     |      |      |       |       |       |       |       |
| 21 - 35g          | 1.22                | 1.47 | 1.71 | 1.95  | 2.2   | 2.57  | 2.93  | 3.42  |
| 35 - 80g          | 0.96                | 1.2  | 1.32 | 1.56  | 1.8   | 2.16  | 2.4   | 2.76  |
| 80 - 150g         | 0.68                | 0.88 | 1.08 | 1.28  | 1.48  | 1.68  | 1.88  | 2.08  |
| 150 - 300g        | 0.52                | 0.72 | 0.92 | 1.12  | 1.32  | 1.48  | 1.64  | 1.8   |
| 300 - 500g        | 0.46                | 0.61 | 0.76 | 0.94  | 1.1   | 1.24  | 1.38  | 1.52  |
| 500g +            | 0.34                | 0.48 | 0.62 | 0.76  | 0.89  | 1.02  | 1.16  | 1.3   |

**Table S3:** Parameter estimates with associated standard errors (SE) for mixed model analyses testing the effects of life-history tactics, population-background, food treatment, and sex on the fork length (cm), weight (g), and condition factor ( $K$ ) trajectories of brown trout over two years of experimental tank rearing. Measurements were taken at key periods denoted by “month”. Effects were contrasted against mature female fish from the anadromous-background in the *Low-Low* treatment. Statistical significance was assessed at  $p < 0.05$ .

| Effect  | Estimate | SE    | t-value | p-value |
|---|----------|-------|---------|---------|
| <i>Mixed model for length</i>                                     |          |       |         |         |
| Intercept   | 7.103    | 0.387 | 18.336  | < 0.001 |
| Month   | 0.699    | 0.013 | 54.865  | < 0.001 |
| Life-history: Smolt   | -2.310   | 0.596 | -3.876  | < 0.001 |
| Population: Non-anadromous-background                             | -0.252   | 0.438 | -0.576  | 0.565   |
| Food: Low-High  | 3.322    | 0.429 | 7.736   | < 0.001 |
| High-Low  | 2.646    | 0.415 | 6.382   | < 0.001 |
| High-High   | 5.434    | 0.425 | 12.798  | < 0.001 |
| Sex: Male   | -0.033   | 0.215 | -0.154  | 0.878   |
| Month: Smolt  | 0.067    | 0.032 | 2.068   | 0.039   |
| Non-anadromous-background: Low-High                               | -0.774   | 0.600 | -1.289  | 0.198   |
| Non-anadromous-background: High-Low                               | -2.382   | 0.630 | -3.781  | < 0.001 |
| Non-anadromous-background: High-High                              | -3.066   | 0.594 | -5.160  | < 0.001 |
| <i>Mixed model for weight</i>                                     |          |       |         |         |
| Intercept   | -73.06   | 8.11  | -9.01   | < 0.001 |
| Month   | 9.47     | 0.28  | 34.24   | < 0.001 |
| Life-history: Smolt   | 22.61    | 12.69 | 1.78    | 0.075   |
| Population: Non-anadromous-background                             | -1.36    | 9.07  | -0.15   | 0.881   |
| Food: Low-High  | 76.57    | 8.87  | 8.63    | < 0.001 |
| High-Low  | 37.75    | 8.56  | 4.41    | < 0.001 |
| High-High   | 114.86   | 8.78  | 13.09   | < 0.001 |
| Sex: Male   | -0.99    | 4.45  | -0.22   | 0.824   |
| Month: Smolt  | -3.08    | 0.70  | -4.39   | < 0.001 |
| Non-anadromous-background: Low-High                               | -19.85   | 12.43 | -1.60   | 0.111   |
| Non-anadromous-background: High-Low                               | -38.59   | 13.05 | -2.96   | 0.003   |
| Non-anadromous-background: High-High                              | -57.67   | 12.32 | -4.68   | < 0.001 |
| <i>Mixed model for condition factor (Fulton's <math>K</math>)</i> |          |       |         |         |
| Intercept   | 1.193    | 0.016 | 76.858  | < 0.001 |
| Month   | -0.004   | 0.001 | -5.341  | < 0.001 |
| Life-history: Smolt   | -0.001   | 0.027 | -0.025  | 0.980   |
| Population: Non-anadromous-background                             | 0.023    | 0.016 | 1.430   | < 0.001 |
| Food: Low-High  | 0.123    | 0.016 | 7.923   | < 0.001 |
| High-Low  | 0.070    | 0.015 | 4.746   | < 0.001 |
| High-High   | 0.139    | 0.015 | 9.034   | < 0.001 |

|                                      |        |       |        |         |
|--------------------------------------|--------|-------|--------|---------|
| Sex: Male                            | 0.002  | 0.008 | 0.285  | 0.776   |
| Month: Smolt                         | -0.006 | 0.002 | -3.564 | < 0.001 |
| Non-anadromous-background: Low-High  | -0.036 | 0.022 | -1.644 | 0.101   |
| Non-anadromous-background: High-Low  | -0.081 | 0.023 | -3.461 | 0.001   |
| Non-anadromous-background: High-High | 0.041  | 0.022 | 1.838  | 0.067   |

---

**Table S4:** Parameter estimates with associated standard errors (SE) for mixed model analysis testing the effects of life-history tactics, population-background, food treatment, and sex on the final fork length (cm), condition factor ( $K$ ) and whole body lipids (%) of brown trout after two years of experimental tank-rearing. Effects were contrasted against mature female fish from the anadromous-background in the *Low-Low* treatment. Statistical significance was assessed at  $p < 0.05$ .

| Effect   | df  | Estimate | SE    | t-value | p-value |
|--|-----|----------|-------|---------|---------|
| <i>Response: Length</i>                        |     |          |       |         |         |
| Intercept                                      | 437 | 20.00    | 0.41  | 48.58   | < 0.001 |
| Life-history: Smolt                            | 437 | -0.59    | 0.35  | -1.66   | 0.097   |
| Population: Non-anadromous-background          | 437 | 0.38     | 0.54  | 0.70    | 0.487   |
| Food: Low-High                                 | 437 | 5.06     | 0.44  | 11.39   | < 0.001 |
| High-Low                                       | 437 | 2.41     | 0.55  | 4.42    | < 0.001 |
| High-High                                      | 437 | 6.83     | 0.54  | 12.56   | < 0.001 |
| Sex: Male                                      | 437 | 0.01     | 0.22  | 0.07    | 0.947   |
| Non-anadromous-background: Low-High            | 437 | -1.30    | 0.63  | -2.05   | 0.040   |
| Non-anadromous-background: High-Low            | 437 | -2.14    | 0.75  | -2.86   | 0.004   |
| Non-anadromous-background: High-High           | 437 | -3.90    | 0.76  | -5.17   | < 0.001 |
| <i>Response: Condition factor (Fulton's K)</i> |     |          |       |         |         |
| Intercept                                      | 437 | 1.079    | 0.018 | 60.095  | < 0.001 |
| Life-history: Smolt                            | 437 | -0.121   | 0.015 | -7.948  | < 0.001 |
| Population: Non-anadromous-background          | 437 | 0.061    | 0.024 | 2.559   | 0.011   |
| Food: Low-High                                 | 437 | 0.195    | 0.019 | 10.212  | < 0.001 |
| High-Low                                       | 437 | 0.048    | 0.024 | 2.007   | 0.045   |
| High-High                                      | 437 | 0.192    | 0.024 | 8.063   | < 0.001 |
| Sex: Male                                      | 437 | -0.017   | 0.009 | -1.831  | 0.068   |
| Non-anadromous-background: Low-High            | 437 | -0.066   | 0.027 | -2.439  | 0.015   |
| Non-anadromous-background: High-Low            | 437 | -0.078   | 0.032 | -2.404  | 0.017   |
| Non-anadromous-background: High-High           | 437 | 0.025    | 0.033 | 0.742   | 0.458   |
| <i>Response: Whole body lipids</i>             |     |          |       |         |         |
| Intercept                                      | 144 | 5.38     | 0.21  | 26.21   | < 0.001 |
| Life-history: Smolt                            | 144 | 0.51     | 0.19  | 2.69    | 0.008   |
| Population: Non-anadromous-background          | 9   | 2.59     | 0.29  | 8.83    | < 0.001 |
| Food: Low-High                                 | 144 | 3.18     | 0.29  | 11.07   | < 0.001 |

|                                      |     |       |      |       |         |
|--------------------------------------|-----|-------|------|-------|---------|
| High-Low                             | 144 | 0.84  | 0.27 | 3.17  | 0.002   |
| High-High                            | 144 | 3.31  | 0.32 | 10.29 | < 0.001 |
| Sex: Male                            | 144 | -0.25 | 0.17 | -1.43 | 0.155   |
| Non-anadromous-background: Low-High  | 144 | -1.41 | 0.42 | -3.39 | 0.001   |
| Non-anadromous-background: High-Low  | 144 | -2.09 | 0.48 | -4.32 | < 0.001 |
| Non-anadromous-background: High-High | 144 | -1.36 | 0.65 | -2.10 | 0.037   |

---

## Chapter 3

---

### Food and temperature stressors have opposing effects in determining flexible migration decisions in brown trout (*Salmo trutta*)

Louise C. Archer<sup>\*1,2</sup>, Stephen A. Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, Stephen D. McCormick<sup>3</sup>, Michael N. O'Grady<sup>4</sup>, Joseph P. Kerry<sup>4</sup>, W. Russell Poole<sup>5</sup>, Patrick Gargan<sup>6</sup>, Philip McGinnity<sup>1,5</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.

<sup>2</sup> Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland.

<sup>3</sup> U.S. Geological Survey, Leetown Science Centre, S.O. Conte Anadromous Fish Research Laboratory, Turners Falls, MA, 01376, USA.

<sup>4</sup> Food Packaging Group, School of Food and Nutritional Sciences, University College Cork, Cork, Ireland.

<sup>5</sup> Marine Institute, Furnace, Newport, Co. Mayo, Ireland.

<sup>6</sup> Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.

**\*Correspondence:** Louise Archer (l.archer@umail.ucc.ie)

**Key words:** climate change, partial migration, anadromy, aquatic, brown trout, multiple stressors, *Salmo trutta*, antagonistic interaction

**Submitted:** Archer LC, Hutton SA, Harman L, McCormick, SD, O'Grady MN, Kerry JP, Poole WR, Gargan P, McGinnity P, and Reed TE. 2019. Food and temperature stressors have opposing effects in determining flexible migration decisions in brown trout (*Salmo trutta*). *Global Change Biology* **In review**.

**Contributions:** LA, TR, and PMG conceived the study. LA, SH, and LH collected data and contributed to experimental design. LA conducted statistical analysis and led the manuscript writing. All authors contributed to interpretation of results and revisions of the manuscript.



## Abstract

In an era of rapid global change, organisms in natural systems are exposed to a multitude of stressors that are likely to co-occur, with uncertain impacts. Here, we explore the individual and cumulative effects of co-occurring environmental stressors on the striking, yet poorly understood, phenomenon of facultative migration. We reared offspring of a brown trout population that naturally demonstrates facultative anadromy (sea-migration), under different environmental stressor treatments and measured life-history responses in terms of migratory tactics and freshwater maturation rates. Juvenile fish were exposed to reduced food availability, temperatures elevated to 1.8 °C above natural conditions, or both treatments in combination over 18 months of experimental tank rearing. When considered in isolation, reduced food had negative effects on the size, mass and condition of fish across the experiment, and we detected variable effects of warm temperatures (negative effects on size and mass, but positive effect on lipids). However, when combined with food restriction, temperature effects on these traits were less pronounced, implying antagonistic dual stressor effects on morphological traits. The two stressors combined additively, but had opposing effects on life-history tactics: migration increased and maturation rates were lower under low food conditions, whereas the opposite was true in the warm temperature treatment. Not all fish had expressed maturation or migration tactics by the end of the study, and the frequency of these “unassigned” fish was higher in food restriction treatments, but lower in warm treatments. Fish choosing migration over freshwater maturation tended to be smaller and in poorer condition (but were similarly sized to unassigned fish). We further detected effects of food restriction on hypo-osmoregulatory function of migrants that may influence the fitness benefits of the migratory tactic at sea. We also highlight that responses to multiple stressors may vary depending on the response considered. Collectively, our results indicate contrasting effects of environmental stressors on life-history trajectories in a facultatively migratory species.

## Introduction

Against the backdrop of rapid global change, organisms are increasingly exposed to a variety of pressures stemming from anthropogenic activities (Sanderson *et al.* 2002). Temperature increases, habitat degradation, pollution, exploitation, and land use changes are examples of pressures, or “stressors”, that contribute to recent patterns of population and biodiversity decline, and altered ecosystem functioning (Walther *et al.* 2002; Parmesan 2006). While much research has addressed the effects of individual stressors, in practice, stressors rarely occur in isolation, and it is imperative we also understand the combined effects of multiple stressors (Breitburg *et al.* 1998) in order to better forecast and manage species’ responses to global change (Côté *et al.* 2016). This is not necessarily straightforward however, with co-occurring stressors – defined here as biotic or abiotic changes beyond the range typically experienced under natural conditions (Breitburg *et al.* 1999; Crain *et al.* 2008) – potentially acting additively, synergistically, or antagonistically (Folt *et al.* 1999; Crain *et al.* 2008).

Although a growing body of research is now expanding beyond single-stressor approaches, uncertainty still surrounds the net effects of co-occurring stressors, and empirical studies have provided mixed results. Meta-analyses suggest that synergistic effects dominate in the marine sphere (Crain *et al.* 2008; Harvey *et al.* 2013; Przeslawski *et al.* 2015), but antagonistic effects (Byrne and Przeslawski 2013) and additive effects (O’Gorman *et al.* 2012) have also been reported. In freshwater systems, which are particularly sensitive to multi-faceted change (Ormerod *et al.* 2010), antagonistic effects of multiple stressors are most prevalent (Jackson *et al.* 2016).

The effects of stressors may be highly context-specific. For example, increased temperature is a stressor likely to be experienced almost universally across natural systems, yet the impacts of warming at the individual level can range from positive to negative depending on whether optimum performance temperatures are exceeded (Huey and Stevenson 1979; Sinclair *et al.* 2016). Moreover, stressor effects can differ depending on the trait/response, or the level of organisation considered *e.g.* warming can increase individual metabolic

and feeding rates, but may reduce survival, cause population/species extinctions (Petchey *et al.* 1999; Fussmann *et al.* 2014) or alter community stability due to long-term changes in species' interaction strengths (Rall *et al.* 65 2010). Predicting stressor effects at multiple levels is likely to be additionally challenging when more than one stressor is involved (Galic *et al.* 2018).

An understudied aspect of multiple stressors is how effects at the individual level shape life-history trajectories, which in turn may mediate how stressors scale up to influence higher-level (*e.g.* population, community, ecosystem) 70 processes. One fundamental decision that many animals face, which is associated with a broad range of life-history and eco-evolutionary consequences, is whether to migrate or not. Facultative migration – the phenomenon whereby individuals retain the capacity to adopt either a migratory or a non-migratory life-style – is common across many animal taxa 75 (Lack 1943; Swingland and Greenwood 1984; Lundberg 1987; Kaitala *et al.* 1993; Chapman *et al.* 2011), with well documented examples in birds (Berthold and Querner 1982; Lundberg 1988; Pulido *et al.* 1996; Newton 2008), ungulates (Ball *et al.* 2001; Cagnacci *et al.* 2011; Hebblewhite and Merrill 2011), zooplankton (Hansson and Hylander 2009), and fishes (Northcote and Ward 1985; Jonsson 80 1985; Chapman *et al.* 2012; Dodson *et al.* 2013). Environmental stresses such as limited food or inclement temperatures often appear to play a role in driving individuals to migrate (Chapman *et al.* 2012). Alternative migratory phenotypes have often been considered within the framework of the “environmentally cued threshold model” (Tomkins and Hazel 2007; Piche *et al.* 2008; Pulido 2011; 85 Buoro *et al.* 2012), in which tactic frequencies are controlled by the relationship between an environmentally-sensitive status trait (*e.g.* physiological condition or energy status) and an inherited threshold, assumed to be genetically variable. If the status trait exceeds the threshold, or “switch point”, residency occurs; if not, migratory tactics are adopted. Environmental factors are likely to be 90 important drivers of migratory tactics at both proximate and ultimate levels, yet few studies have addressed how facultatively migratory species respond to

pressures arising from environmental change (Doswald *et al.* 2009), either in isolation, or when stressors act in combination.

Salmonine fishes (trouts, salmons, and charrs) represent an excellent group to study multiple stressor effects (McGinnity *et al.* 2009; de Eyto *et al.* 2016). In facilitating obligate freshwater spawning, salmonines display a suite of migratory phenotypes, encompassing residents that remain in natal streams their entire lives, individuals that migrate to larger rivers or lakes (potomodromy), and others still that undertake a marine migration before returning to fresh water to spawn (anadromy) (Klemetsen *et al.* 2003; Dodson *et al.* 2013; Ferguson *et al.* 2019; Nevoux *et al.* 2019). The migration *versus* residency decision represents a trade-off, where the benefits of migration (avoiding abiotic or biotic stresses in the natal stream, exploiting better food resources in the new environment, which translate into higher growth and thus higher fecundity or mating success) must be balanced against the costs (energetic demands of migration, physiological stress of changing environments/habitats, a potentially increased risk of predation) (Kendall *et al.* 2014). Environmental conditions in natal fresh waters can interact with intrinsic physiological traits to determine alternative migratory tactics, *e.g.* if food resources cannot support growth or metabolism in early life, the resulting energetic deficit may promote migration. Food resources have been shown to directly (Davidsen *et al.* 2014; Jones *et al.* 2015; Archer *et al.* 2019) and indirectly influence migration (Olsson *et al.* 2006; Wysujack *et al.* 2009; O'Neal and Stanford 2011). Similarly, because temperature profoundly influences biological processes (Gillooly *et al.* 2001; Dell *et al.* 2011), temperature effects on physiological status traits/energetic balance likely make it an important environmental determinant of migratory decisions. Higher temperatures have been associated with increased anadromy *in lieu* of maturation in steelhead trout *Oncorhynchus mykiss* (Sloat and Reeves 2014), but under experimental conditions of constant food supply. Warming is likely to be accompanied by reductions in freshwater macroinvertebrate abundance (Durance and Ormerod

2007), with potentially synergistic effects if elevated metabolic demands induced by warming are compounded by low food availability.

Environmental stressors may also act to affect the performance of individuals once a migratory decision has been made. Although migration potentially confers many benefits on individuals, the ensuing costs mean that environmental conditions experienced prior to out-migration might further affect the future success of migrants, both during migration and in the subsequent environment (river, lake, or sea) (McCormick *et al.* 2009a). Sea-migration in particular requires substantial physiological remodelling for transitioning to salt water, and the changes in physiological regulation of ions, colouration, and body shape (collectively termed “smolting”, (Tanguy *et al.* 1994)) necessitate an expensive investment by individuals that is likely to begin long before the migration is initiated, and hence may be affected by environmental stressors. Empirical evidence suggests that once the migratory decision is made, migrants divert resources towards accelerated growth (Metcalf 1998). Smolt survival at sea is positively related to size (Ward and Slaney 1988) and, as such, favourable freshwater conditions may produce larger and more successful migrants, with associated fitness benefits altering the migration-residency trade off.

Here, we present the results of an experimental laboratory study of physiology, migration and maturation of brown trout, using the F1 progeny of wild-caught parents from a population that exhibits facultative migration in nature. We aimed to explore if, and how, life-history decisions are shaped by individual and interactive effects of two putative extrinsic environmental stressors. Specifically, we aimed to: (i) determine the effects of food restriction and elevated temperature on a range of physiological traits (size, mass, condition, lipids); (ii) assess the life-history consequences, in terms of migratory tactics and maturation decisions, of these stressors (both in isolation and combined); and (iii) explore how environmental stressors affect future migratory performance. We expected that food reduction and elevated temperature would each increase the prevalence of migratory tactics, with potentially synergistic

effects when the stressors were combined. We also expected stressors to negatively affect future migratory capacity of anadromous individuals by  
 155 reducing potential for fast growth (and thus reducing smolt viability), or by inhibiting osmoregulation in salt water.

## Materials and Methods

### *Study population and fish rearing*

Brown trout brood stock from a wild population were caught by seine-netting  
 160 in November 2015 in Tawnyard Lough, an upland lake of 56 ha in the Erriff catchment (National Salmonid Index Catchment) in the west of Ireland (53° 37' 0.00" N: 09° 40' 17.10" W). Tawnyard Lough is fed primarily by the Glendavoch river, and a number of smaller tributaries (Figure S1). Brown trout primarily spawn within the Glendavoch River and move downstream as fry or parr to  
 165 Tawnyard Lough, a distance of a few hundred metres to kilometres, depending on where spawning occurred. Tawnyard Lough produces a large run of out-migrating anadromous juveniles (smolts), with annual estimates of 500 to 3000 smolts enumerated at the outflow of the Lough over the last 30 years (Gargan *et al.* 2016). An unknown proportion of the population remain within the lake,  
 170 and undergo several years of freshwater growth before returning to the natal stream to spawn. While the migration phenotypes of the brood stock could not be determined unambiguously in this study (because external signs of prior migration are not completely unambiguous in this system), we assumed that the frequencies of migratory/resident phenotypes among our brood stock was  
 175 broadly representative of naturally occurring frequencies, given that brood stock were obtained haphazardly. Moreover, our goals in this study were not to test explicitly for inherited variation in migration tactics, but rather to explore proximate (environmental) drivers. The *caveat* must be kept in mind, however, that any environmental effects we document are contingent on the genotypic  
 180 composition of our sample.

Each ripe female ( $n = 7$ ) was stripped of eggs, which were then split into two batches, and each batch was fertilised by the milt of a single male ( $n = 10$ ) (*i.e.* each female was mated to two males, creating fourteen full-sib families). Due to

constraints in obtaining brood stock and variation in the timing of when females were ripe to be stripped, eggs were fertilised on three occasions in November and December 2015 (see Table S1 fertilisation/ brood stock crossing details). Fertilised eggs from each crossing were incubated separately in standard Heath trays in the nearby Burrishoole hatchery. Prior to exogenous feeding, fry from the late November/early December fertilisations (hereafter “Nov-Dec fertilisation group”) were transferred to a rearing facility at University College Cork (Aquaculture and Fisheries Development Centre). Here, families were mixed, and fry were held in a 100 L growth tank on a recirculating aquaculture system (RAS), maintained by a conditioning unit at a seasonally varying temperature regime approximating that of the catchment from which the brood stock were obtained (i.e. the annual cycle in mean weekly temperatures in the nearby Burrishoole catchment for the preceding three years, see Figure S2). Note that in winter 2015, simulated water temperatures were higher than the source catchment due to initial logistical constraints in achieving low temperatures. Fry were fed *ad libitum* with commercially available trout pellets (Skretting Ltd., Norway) to facilitate their transition to exogenous feeding (by June 2016), and were then fed *ad libitum* until the experimental phase began. Fish experienced a constant photoperiod (12:12 hours of light: dark) during this initial rearing phase. Due to logistical constraints, an additional group of fry from the first fertilisation event (termed “Nov fertilisation group”) was maintained at a natural temperature regime in a flow-through tank at the Burrishoole hatchery facility, where they transitioned to exogenous feeding via *ad libitum* feeding with the same trout pellets. Fry from the Nov group were transported to the UCC rearing facility in May 2016, and reared in a 100 L tank, in the same RAS and under the same conditions as the Nov-Dec fertilisation group. Due to size differences (Nov group fish were larger than Nov-Dec group fish by the beginning of the experiment) the two fertilisation groups were reared in separate tanks for the duration of the study to prevent emergence of feeding hierarchies.

*Experimental treatments*

215 The study and all associated procedures were carried out with ethical approval from Health Products Regulatory Authority (HPRA) Ireland, under HPRA project license AEI9I30/P034, and individual licenses AEI9I30/1087, AEI9I30/I200, AEI9I30/I201 and AEI9I30/I202). The experimental phase ran for an 18-month period from December 2016 to June 2018 with all fish humanely  
 220 euthanized at the end of the experiment.

Juvenile brown trout were randomly allocated across eight 520L tanks at the end of November 2016 (initial n = 140 per tank for Nov-Dec group, and n = 35 per tank (filled to 203L) for Nov group), each assigned one of two temperature treatments and one of two food treatments. Water flowed continually through  
 225 tanks to prevent the build-up of waste, returned to a central sump via mechanical filtration, and was treated with protein skimming, biofiltration and UV skimming. Weekly monitoring of water quality indicated that levels of pH, nitrate, nitrite, and ammonia were well within acceptable ranges for fish health. Mortality during the experimental phase was minimal (4%). To avoid  
 230 compromising water quality with excessive biomass, fish were culled haphazardly (n = 229 in total across all treatments) over the two years of tank rearing, with equal densities in terms of fish numbers (fish per L) maintained across treatment groups and equal biomass densities (g per L) maintained between fertilisation groups. Fish culled in this manner were not included in  
 235 the analyses.

To explore the individual and interactive effects of food restriction and temperature in determining migratory tactics, food and temperature treatments were applied in isolation and in combination for both fertilisation groups for the duration of the experimental phase. The two food treatments  
 240 were: (i) high food treatment: fish fed recommended daily rations for optimal growth calculated as a percentage of their body mass and adjusted for seasonally-changing temperatures (Skretting Ltd., Norway); and (ii) low food treatment: fish fed 25% of optimal daily rations. A value of 25% of optimal daily rations was chosen for the low food treatment because similar restrictions have



245 previously been shown to reduce the frequency of residency in adfluvial brown  
 trout (Wysujack *et al.* 2009). Food was dispensed during daylight hours via  
 automatic feeders (Arvo-Tec Oy, Huutokoski, Finland) located above each tank.  
 Feeders delivered regular pulses of food (lasting two seconds), with the  
 frequency of pulses adjusted according to food treatment (i.e. fewer pulses for  
 250 low food treatments). The two temperature regimes were achieved by passing  
 water through one of two conditioning units that maintained two temperature  
 treatments as follows : (i) cool temperature treatment: temperatures matching  
 the natural, seasonally-varying, temperature regime for the Erriff catchment;  
 and (ii) warm temperature treatment: temperatures elevated by  $1.8^{\circ}\text{C} \pm 0.55$   
 255 (SD) above the cool temperature treatment. The cool treatment ranged from  
 $5.9 - 16.4^{\circ}\text{C}$  (mean temperature =  $10.9^{\circ}\text{C} \pm 3.2$  SD) and the warm treatment  
 ranged from  $7.5 - 18.2^{\circ}\text{C}$  (mean temperature =  $12.7^{\circ}\text{C} \pm 3.2$  SD). An increase of  
 $1.8^{\circ}\text{C}$  for the warm temperature treatment was chosen because this is in line  
 with increases of  $1 - 3^{\circ}\text{C}$  projected to occur under climate change scenarios  
 260 (IPCC 2014). While both treatments remained within sub-lethal ranges for  
 brown trout (Forseth *et al.* 2009; Jonsson and Jonsson 2009), the warm  
 temperature treatment was considered “stressful” because the maximum  
 temperatures in the warm treatment approached upper thermal growth limits  
 for brown trout ( $18.7^{\circ}\text{C}$ ). Optimal temperatures for growth have been estimated  
 265 to be between  $13.1 - 13.9^{\circ}\text{C}$  (Elliott and Hurley 2000; Hari *et al.* 2006; Elliott  
 and Elliott 2010; Kovach *et al.* 2016). Temperatures in the warm treatment  
 remained above this for twice as long as those in the cool treatment, which  
 remained closer to optimal growth conditions. Food rations were reduced over  
 four weeks and temperature was increased by  $0.5^{\circ}\text{C}$  per week to minimise  
 270 stress. Within each treatment, absolute rations were adjusted on a monthly  
 basis to account for changes in body mass and temperature.

#### *Data collection and life-history determination*

Within each food and temperature treatment combination, 25 – 30 fish per  
 fertilisation group were lightly anaesthised with MS-222 and marked with  
 275 unique colour combinations of visible implant elastomer (VIE) tags (Northwest

Marine Technology Ltd., Washington, USA), allowing for re-identification of individuals. Fork length and mass were measured at key periods throughout the study [Late-November in 2016 (prior to initiation of treatments), February, April, June, July, September, December in 2017, and April in 2018].

280 All fish were checked weekly for morphological indicators of smolting from March to June in each of 2017 and 2018, the period corresponding to the natural migratory period in the wild for this population (Gargan *et al.* 2016). Wild smolts typically migrate out of the Erriff system aged between 1+ and 4+, with the large majority doing so aged 2+ or 3+ (approximately equal numbers of  
285 each) (Gargan *et al.* 2016). Smolting is a precursor to downstream migration in several salmonines, and comprises a number of morphological, behavioural and physiological changes. We used the following morphological indicators to assess smolting [following Tanguy *et al.* 1994]: silvering/loss of parr marks; pronounced lateral line (*i.e.* clearly visible and raised to the touch); transparent  
290 fins; and fusiform shape (pointed snout, slimmer body, and elongated caudal peduncle) (Riddell and Leggett 1981; Hard *et al.* 1999; Villar-Guerra *et al.* 2019). Fish that clearly matched three of these criteria were classed as smolts. In spring 2017, no fish matched the morphological criteria for smolts. In spring 2018, fish that matched the morphological criteria for smolts were transferred to salt  
295 water at 30 ppt for 24 hours to assess hypo-osmoregulatory function. Seawater “challenges” are used as an indicator of anadromy capacity, where the ability to regulate ion concentrations (*e.g.* to maintain plasma chloride concentration) in sea water is a measure of saltwater tolerance, or physiological “readiness” of smolts for seawater entry (Clarke 1982; McCormick *et al.* 1998; Schultz and  
300 McCormick 2012; McCormick 2012). Fish were monitored during this period to ensure that any fish showing signs of failing the challenge (loss of equilibrium) could be removed and euthanised (though no fish showed signs of failure in our study). After the 24-hour seawater challenge, fish were euthanised with an overdose of MS-222. The mass and fork length of each individual was recorded,  
305 and a blood sample was obtained from the caudal vasculature using a 21 gauge needle and a 2.6ml heparinised syringe. Blood samples were centrifuged at

8000 rpm for 3 minutes, and the plasma aliquot was siphoned off, stored at -80 °C and later measured for plasma chloride concentration as an indicator of osmoregulatory performance. Four to six gill filaments were placed in 100µl of  
 310 ice-cold SEI buffer (150 mmol l<sup>-1</sup> sucrose, 10 mmol l<sup>-1</sup> EDTA, 50 mmol l<sup>-1</sup> imidazole, pH 7.3) and frozen at -80 °C for later measurement of gill Na<sup>+</sup>/K<sup>+</sup>-ATPase (NKA) activity.

All fish (classed morphologically as smolts or non-smolts) were dissected in spring 2018 to visually determine sex and maturation status based on gonad  
 315 development. Mature males had enlarged white testes or running milt. Maturing males had visible or moderately enlarged testes but no running milt. Mature females had visible eggs in the body cavity. Immature fish (unconfirmed sex at the time of sampling) were later genotyped to determine genotypic sex using a microsatellite sex marker. The natural spawning period for the wild  
 320 population-of-origin is in late autumn/early winter, and the migratory period is in the spring (Gargan *et al.* 2016). Since freshwater maturation generally precludes migration in brown trout (Jonsson 1985; Dellefors and Faremo 1988; Dębowski and Dobosz 2016) any fish showing signs of maturing without having migrated to sea are considered to be on a non-anadromous trajectory, while  
 325 smolts which undertake marine migrations are immature. We thus classed fish as smolts (migratory tactic) if they matched the morphological criteria for smolts and were immature. Fish were classed as mature (residency tactic) if they showed signs of maturation at the time of sampling. Fish that were immature and did not match the morphological criteria for smolts had an unknown life  
 330 history at the time of sampling and were classed as “unassigned”. Whole body lipid content (%) was measured for all smolts and a random sample of mature (n = 107) and unassigned (n =19) fish using a CEM Smart trac5 system of integrated heating and nuclear resonance (CEM Corporation, Matthews, NC, USA) on individual homogenised fish samples (Toussaint *et al.* 2002; Keeton *et al.* 2003; Nielsen *et al.* 2005). Plasma chloride concentration was measured by  
 335 coulometric titration using a Jenway PCLM3 chloride meter (FishVet Group, Oranmore, Ireland) for all smolts and a random sample of non-smolts (n = 107

mature fish and  $n = 18$  unassigned fish). Gill NKA activity was measured following McCormick *et al.* (2009) for a random sample of smolts and non-smolts ( $n = 25$  smolts,  $n = 135$  mature fish and  $n = 22$  unassigned fish).

### *Statistical analysis*

To test if food and temperature acted as stressors at the level of individual traits underpinning migration (Aim 1), we explored factors affecting fork length, mass and condition of fish across the study period within a mixed effects modelling framework (*nlme* package (Pinheiro *et al.* 2019)). We calculated condition factor as:

$$\text{Condition factor} = \frac{\text{mass (g)}}{\text{fork length (cm)}^b} \times 100$$

Where  $b$  is the slope estimated from the linear relationship between  $\log(\text{mass})$  and  $\log(\text{fork length})$  (Bolger and Connolly 1989). The mixed effects models included time (continuous variable representing weeks since start of experiment), a quadratic term for time (to account for non-linearity of traits through time), food treatment, temperature treatment, fertilisation group, and sex as fixed effects, and individual identity as a random effect to account for multiple measurements on some individuals. We included an interaction between food treatment and temperature treatment (to test for synergistic or antagonistic effects of food and temperature), and a three-way interaction (food treatment  $\times$  temperature treatment  $\times$  time) to test whether trajectories diverged through time according to treatment combination. To compare single stressor effects with combined stressor effects, we carried out pairwise comparisons across all levels of the food  $\times$  temperature interaction using Tukey post-hoc tests (*emmeans* package (Lenth 2019)).

To test whether trait trajectories were similar in mature fish and smolts, we created additional mixed effects models with time, a quadratic term for time, migratory tactics (categorical variable, two levels: smolt/mature), sex, and fertilisation group as fixed effects, an interaction between time and migratory tactics, and a random effect of individual identity. We excluded unassigned fish

in this comparison of “status” traits, as we could not determine their life-history trajectory, *i.e.* some of the unassigned fish may have been on a migratory trajectory, but were deferring actual migration until a later age. For all of the  
 370 above models, temporal autocorrelation of the response variable was accounted for by modelling an autoregressive error structure as a first order lag function of time. Separate models were constructed for z-standardised measures of length, mass and condition, and normal errors were assumed in each case.

We similarly used mixed effects models (normal errors) to explore factors  
 375 influencing variation in final measurements of traits (z-standardised length, condition, and whole body lipids) at the end of the study. We included food treatment, temperature treatment, fertilisation group, sex, and an interaction between food treatment and temperature treatment as fixed effects, and date of terminal sample (categorical variable with 8 sampling dates) as a random effect.  
 380 Additional mixed effects models tested for differences in final measurements of status traits according to migratory tactics (migratory tactic, sex and fertilisation group included as fixed effects, and terminal sample date as a random effect).

To test if food and temperature treatments affected life-history tactics (Aim 2),  
 385 we built three generalized linear models (GLMs) with a logit-link function and binary life-history response variables. One GLM predicted smolt status (binary response: 1 = smolt, 0 = non-smolt), a second GLM predicted maturation (binary response: 1 = mature, 0 = immature), and a third GLM tested for differences in expression of any life-history tactics by the end of the study (*i.e.* by age 2+,  
 390 second year of life) (binary response: 1 = unassigned, 0 = smolt or mature). All GLMs included the categorical variables: food treatment, temperature treatment, sex, fertilisation group, and an interaction between food and temperature treatments.

To explore variation in osmoregulatory performance, we first tested for  
 395 differences in gill NKA activity (log transformed) and plasma chloride concentration between smolts and non-smolts using mixed effects models (normal errors). Each model included life-history tactic, fertilisation group, and

sex as fixed effects, and terminal sampling date as a random effect. We retained the “unassigned” life-history class in these analyses to determine if unassigned  
 400 fish showed signs of hypo-osmoregulatory capacity in salt water relative to mature fish, suggesting that these unassigned fish were in fact on a smolting trajectory but were yet to express morphological indicators of smolting. Post-hoc pairwise comparisons between life-history tactics were carried out using the *emmeans* package (Lenth 2019).

405 We explored the mechanisms underpinning osmoregulatory capacity by fitting a GLM (normal errors) to gill NKA activity as a function of size-corrected plasma chloride concentration in smolts and non-smolts. Because hypo-osmoregulatory capacity generally increases with size in brown trout (Finstad and Ugedal 1998), we corrected for size in this analysis by using the residuals of  
 410 the linear relationship between plasma chloride and fork length. Finally, we assessed how food and temperature treatments influenced osmoregulatory capacity of smolts (Aim 3) using mixed effects models. Separate models (normal errors) were constructed for z-standardised gill NKA activity and plasma chloride concentration, with food treatment, temperature treatment, sex,  
 415 fertilisation group, fork length (a covariate to correct for body size effects), and a food  $\times$  temperature interaction included as fixed effects, and terminal sample date as a random effect.

Marginal  $R^2$  values for mixed effect models were calculated using the *MuMIn* package in R (Barton 2019). We used likelihood ratio tests (LRT) to assess  
 420 statistical significance of predictor variables for all models at a 5% alpha level, and non-significant interaction terms were excluded to interpret main effects. Analysis was carried out in R version 3.5.3 (R Core Team 2019), and all models were checked against assumptions of the given model (independence, non-normality of residuals, heteroscedasticity and multicollinearity).

## 425 **Results**

At the termination of the experiment, 349 fish were assigned a life-history tactic (30 smolts and 319 mature fish) and 76 fish were classed as “unassigned” (Table

1). The frequency of smolting varied by food and temperature treatments. Overall, the proportion of smolts (averaged across fertilisation groups) was  
 430 highest in the low food-cool temperature treatment (18.9 %), with the lowest proportion in the high food-warm temperature treatment (0.9 %), and intermediate proportions in low food-warm temperature (4.6 %) and high food-cool temperature treatments (3.8 %).

The proportion of mature fish (averaged across fertilisation groups) was highest  
 435 in the high food-warm temperature treatment (92.5 %), followed by similar proportions in high food-cool temperature (75.2 %) and low food-warm temperature treatments (75.0 %). Maturation was lowest in the low food-cool temperature treatment (57.6 %).

**Table 1:** Percentage of brown trout ( $n = 425$ ), F1 offspring of wild trout, classed  
 440 as smolts (i.e. migratory tactic) or non-smolts (mature or unassigned) after two years of experimental tank-rearing. Offspring were derived from brood stock gametes fertilised in November and December 2015 (coded here as early November = “Nov” and late November/early December = “Nov-Dec”). Values correspond to percentages for each category (broken down by sex) of the total  
 445 number of fish per tank (where each tank corresponds to a given food treatment by temperature regime combination, i.e. a single row in the table). Sample size ( $n$ ) given in brackets after the %.

| Food,<br>Temperature | Fertilisation<br>Group | % Smolts<br>(n) |            | % Mature<br>(n) |              | % Unassigned<br>(n) |              |
|----------------------|------------------------|-----------------|------------|-----------------|--------------|---------------------|--------------|
|                      |                        | Female          | Male       | Female          | Male         | Female              | Male         |
| Low, Cool            | Nov                    | 6.1<br>(2)      | 3.0<br>(1) | 48.5<br>(16)    | 6.1<br>(2)   | 6.1<br>(2)          | 30.3<br>(10) |
| Low, Cool            | Nov-Dec                | 15.1<br>(11)    | 8.2<br>(6) | 19.2<br>(14)    | 39.7<br>(29) | 8.2<br>(6)          | 9.6<br>(7)   |
| Low, Warm            | Nov                    | 0<br>(0)        | 0<br>(0)   | 51.5<br>(17)    | 36.4<br>(12) | 3.0<br>(1)          | 9.9<br>(3)   |
| Low, Warm            | Nov-Dec                | 2.7<br>(2)      | 4.0<br>(3) | 41.3<br>(31)    | 28.0<br>(21) | 2.7<br>(2)          | 21.3<br>(16) |
| High, Cool           | Nov                    | 2.9<br>(1)      | 0<br>(0)   | 32.4<br>(11)    | 32.4<br>(11) | 5.9<br>(2)          | 26.5<br>(9)  |
| High, Cool           | Nov-Dec                | 2.8<br>(2)      | 1.4<br>(1) | 43.7<br>(31)    | 36.6<br>(26) | 2.8<br>(2)          | 12.7<br>(9)  |
| High, Warm           | Nov                    | 0<br>(0)        | 0<br>(0)   | 61.8<br>(21)    | 23.5<br>(8)  | 2.9<br>(1)          | 11.8<br>(4)  |
| High, Warm           | Nov-Dec                | 1.4<br>(1)      | 0<br>(0)   | 48.6<br>(35)    | 47.2<br>(34) | 0<br>(0)            | 2.8<br>(2)   |

*Morphological trait trajectories*

450 Physiological trait trajectories diverged through time in response to food treatment, temperature treatment, and fertilisation group (Figure 1A, Table 2). The models for length (marginal  $R^2 = 0.68$ ) and condition factor (marginal  $R^2 = 0.33$ ) retained significant interactions between food treatment and temperature treatment, food treatment and time, and temperature treatment and time

455 (Table 2). The model for mass (marginal  $R^2 = 0.61$ ) retained a significant time  $\times$  food  $\times$  temperature interaction (Table 2). Fertilisation group had a significant effect on length ( $\chi^2 = 57.17$ ,  $df = 1$ ,  $p < 0.001$ ), mass ( $\chi^2 = 24.49$ ,  $df = 1$ ,  $p < 0.001$ ), and condition factor ( $\chi^2 = 8.73$ ,  $df = 1$ ,  $p = 0.003$ ), with fish in the Nov fertilisation group tending to be larger and heavier than those in the Nov-Dec

460 group, and in marginally lower condition. There was no significant effect of sex on length ( $\chi^2 = 0.73$ ,  $df = 1$ ,  $p = 0.394$ ), mass ( $\chi^2 = 2.01$ ,  $df = 1$ ,  $p = 0.156$ ) or condition factor ( $\chi^2 = 0.29$ ,  $df = 1$ ,  $p = 0.591$ ) across the study. When food and temperature stressors were experienced in isolation (*i.e.* a single treatment applied) fish receiving the low food treatment were smaller (post hoc

465 comparison of low food-cool temperature *versus* high food-cool temperature treatment:  $t$ -value = 12.06,  $p < 0.001$ ), lighter ( $t$ -value = 13.26,  $p < 0.001$ ) and in poorer condition ( $t$ -value = 10.74,  $p < 0.001$ ). Fish in the warm temperature treatment were also smaller (warm temperature-high food *versus* cool temperature-high food treatment:  $t$ -value = 3.23,  $p = 0.007$ ), lighter ( $t$ -value =


470 3.66,  $p = 0.002$ ), but in similar condition ( $t$ -value = 1.41,  $p = 0.495$ ) (Figure 2A, B, C). The positive interaction term between food treatment and temperature treatment indicated that effects of combined stressor treatments on length, mass, and  $K$  were less than we might expect based off their effects in isolation.

Fish also varied in length (marginal  $R^2 = 0.55$ ), mass (marginal  $R^2 = 0.36$ ) and

475 condition factor (marginal  $R^2 = 0.16$ ) trajectories according to migratory tactics, with smolts tending to be smaller than mature fish across the duration of the study period ( $\chi^2 = 15.55$ ,  $df = 1$ ,  $p < 0.001$ ). The significant interaction between migratory tactics and time for mass and condition factor (Figure 1B, Table 3)



indicated smolts gained less mass, with lower condition trajectories (Figure 2D,E,F).




**Figure 1:** Coefficient estimates ( $\pm$  associated standard errors) of mixed effects models describing z-standardised length, mass and condition trajectories of brown trout offspring (derived from wild-caught parents from a facultatively anadromous population) that were (A) exposed to food and temperature stressors and (B) classed as adopting either smolt (migratory tactic) or freshwater maturing (non-migratory/resident) tactics after 18 months of tank rearing.

485 **Table 2:** Results of the mixed effect model analysis for length, mass and condition trajectories of brown trout exposed to food and temperature stressors. The results of the model selection procedure on interaction terms are given, and the selected model for each response is highlighted in bold. The models included a random effect of individual identity and a first-order autoregressive correlation structure with respect to time (weeks of experimental treatment) was also modelled.

| Model   | df | AIC    | logLik  | L-ratio | p-value |
|---|----|--------|---------|---------|---------|
| Length ~ time*food*temperature + time <sup>2</sup> + fertilisation + sex                                      | 14 | 1222.8 | -597.4  |         |         |
| <b>Length ~ time*food + time*temperature + time<sup>2</sup> + food*temperature + fertilisation + sex</b>      | 13 | 1222.5 | -598.3  | 1.8     | 0.186   |
| Length ~ time*food + time <sup>2</sup> + food*temperature + fertilisation + sex                               | 12 | 1229.7 | -602.8  | 9.1     | 0.003   |
| Length ~ time*temperature + time <sup>2</sup> + food*temperature + time <sup>2</sup> + fertilisation + sex    | 12 | 1422.2 | -699.1  | 201.6   | < 0.001 |
| Length ~ time*food + time*temperature + time <sup>2</sup> + fertilisation + sex                               | 12 | 1236.1 | -606.1  | 15.6    | < 0.001 |
| <b>Mass ~ time*food*temperature + time<sup>2</sup> + fertilisation + sex</b>                                  | 14 | 1667.1 | -819.6  |         |         |
| Mass ~ time*food + time*temperature + time <sup>2</sup> + food*temperature + fertilisation + sex              | 13 | 1672.6 | -823.3  | 7.5     | 0.006   |
| Condition ~ time*food*temperature + time <sup>2</sup> + fertilisation + sex                                   | 14 | 3023.4 | -1497.7 |         |         |
| <b>Condition ~ time*food + time*temperature + time<sup>2</sup> + food*temperature + fertilisation + sex</b>   | 13 | 3022.4 | -1498.2 | 0.9     | 0.337   |
| Condition ~ time*food + time <sup>2</sup> + food*temperature + fertilisation + sex                            | 12 | 3029.8 | -1502.9 | 9.4     | 0.002   |
| Condition ~ time*temperature + time <sup>2</sup> + food*temperature + time <sup>2</sup> + fertilisation + sex | 12 | 3059.6 | -1517.8 | 39.3    | < 0.001 |
| Condition ~ time*food + time*temperature + time <sup>2</sup> + fertilisation + sex                            | 12 | 3027.3 | -1501.7 | 7.0     | 0.008   |

490 **Table 3:** Results of the mixed effect model analysis for length, mass and condition trajectories of brown trout in the experiment with life-history classed as either smolts (i.e. migratory) or freshwater mature across the study period. The results of the model selection procedure on interaction terms are given, and the selected model for each response is highlighted in bold. The models included a random effect of individual identity and a first-order autoregressive correlation structure with respect to time (weeks of experimental treatment) was also modelled.

| Model   | df | AIC    | logLik  | L-ratio | p-value |
|---|----|--------|---------|---------|---------|
| Length ~ time*life-history + time <sup>2</sup> + sex + fertilisation          | 10 | 1300.6 | -640.3  |         |         |
| <b>Length ~ time + life- history + sex + fertilisation</b>                    | 9  | 1298.9 | -640.4  | 0.2     | 0.637   |
| Mass ~ time*life-history + time <sup>2</sup> + sex + fertilisation            | 10 | 1836.0 | -908.0  |         |         |
| Mass ~ time + life- history + time <sup>2</sup> + sex + fertilisation         | 9  | 1843.6 | -912.8  | 9.7     | 0.002   |
| <b>Condition ~ time*life-history + time<sup>2</sup> + sex + fertilisation</b> | 10 | 2674.4 | -1327.2 |         |         |
| Condition ~ time + life- history + time <sup>2</sup> sex + fertilisation      | 9  | 2678.8 | -1330.4 | 6.4     | 0.011   |



**Figure 2:** Trajectories of fork length, mass, and condition of brown trout offspring (derived from wild-caught parents from a facultatively anadromous population) under different food treatments and temperature treatments (A, B, C), and classed according to life-history tactics (D, E, and F). Food and temperature treatments are denoted in the format “Food-Temperature” (High or Low food, and Warm or Cool temperature) and life histories were classed as either smolt (migratory tactics) or mature (non-migratory). Week 0 = end of November 2015, when fish were 10 to 11 months old (Nov-Dec and Nov group, respectively).


495

#### *Morphological traits at the end of the study*

At the end of the experiment, fish varied in length, condition and whole body lipids depending on food treatment, temperature treatment, life-history tactics and fertilisation group (Figure 3A, B). The models describing final length (marginal  $R^2 = 0.48$ ) and condition (marginal  $R^2 = 0.38$ ) each retained a significant interaction between food treatment and temperature treatment but the model describing whole body lipids (marginal  $R^2 = 0.41$ ) did not (Table 4). We detected significant negative main effects of food treatment ( $\chi^2 = 63.44$ ,  $df = 1$ ,  $p < 0.001$ ) but positive effects of temperature treatment ( $\chi^2 = 3.91$ ,  $df = 1$ ,  $p$

505 = 0.048) on lipid levels (Figure 3A). The significant positive interaction term (Figure 3A) indicated an antagonistic effect of food and temperature treatments on length and condition (Figure 3C, D, and E). Fertilisation group significantly affected length ( $\chi^2 = 4.56$ ,  $df = 1$ ,  $p = 0.033$ ) and condition ( $\chi^2 = 5.15$ ,  $df = 1$ ,  $p = 0.023$ ). Fish in the Nov fertilisation group tended to be larger but in poorer  
 510 condition, with similar lipid levels ( $\chi^2 = 0.02$ ,  $df = 1$ ,  $p = 0.880$ ) to those in the Nov-Dec group. There was no significant effect of sex on length ( $\chi^2 = 0.14$ ,  $df = 1$ ,  $p = 0.712$ ), condition ( $\chi^2 = 2.60$ ,  $df = 1$ ,  $p = 0.107$ ) or lipids ( $\chi^2 = 1.91$ ,  $df = 1$ ,  $p = 0.167$ ).

Life-history tactics significantly affected final length ( $\chi^2 = 4.80$ ,  $df = 1$ ,  $p = 0.036$ ),  
 515 final condition ( $\chi^2 = 19.62$ ,  $df = 1$ ,  $p < 0.001$ ), and final lipids ( $\chi^2 = 13.87$ ,  $df = 1$ ,  $p = 0.002$ ). Overall, smolts were smaller than mature fish, with lower condition values, and higher lipid levels (Figure 3B, C, D, E). Smolts and unassigned fish were similarly sized ( $\chi^2 = 0.35$ ,  $df = 1$ ,  $p = 0.554$ ), with similar lipid levels ( $\chi^2 = 1.49$ ,  $df = 1$ ,  $p = 0.222$ ), though smolts had lower condition values ( $\chi^2 = 07.48$ ,  $df$   
 520 = 1,  $p = 0.006$ ) (Figure S3).



**Figure 3:** Coefficient estimates ( $\pm$  standard errors) from mixed effect models describing effects of (A) food treatment and temperature treatment and (B) migratory tactics on z-standardised final measures of length, condition, and whole body lipids of brown trout offspring classed as either smolts (migratory) or freshwater mature (non-migratory/resident) at the end of the experimental study (Spring 2018). Median values of (C) length, (D) condition, and (E) whole body lipids are represented by the white horizontal lines in each box in (C), (D), and (E). Food and temperature treatments are denoted in the format “Food-Temperature” (High or Low food, and Warm or Cool temperature). Note that only one smolt was recorded in the High-Warm treatment, and thus there is no corresponding white line for the median in the High-Warm treatment.

525 **Table 4:** Results of the mixed effect model analysis for length, condition, and whole body lipids of brown trout exposed to food and temperature stressors at the end of the experimental study period. The results of the model selection procedure on interaction terms are given, and the selected model for each response is highlighted in bold. The models included a random effect of terminal sample date.

| Model   | df | AIC    | logLik | <i>L</i> -ratio | <i>p</i> -value |
|---|----|--------|--------|-----------------|-----------------|
| <b>Length ~ food*temperature + fertilisation + sex</b>    | 8  | 901.5  | -422.8 |                 |                 |
| Length ~ food + temperature + fertilisation + sex         | 7  | 916.5  | -451.3 | 16.98           | < 0.001         |
| <b>Condition ~ food*temperature + fertilisation + sex</b> | 8  | 1034.0 | -509.0 |                 |                 |
| Condition ~ food + temperature + fertilisation + sex      | 7  | 1036.9 | -511.5 | 4.94            | 0.026           |
| Lipids ~ food*temperature + fertilisation + sex           | 8  | 375.5  | -179.8 |                 |                 |
| <b>Lipids ~ food + temperature + fertilisation + sex</b>  | 7  | 375.7  | -180.8 | 2.18            | 0.140           |


*Migratory tactics*

The model describing the probability of smolting had significant effects of food treatment ( $\chi^2 = 16.50$ ,  $df = 1$ ,  $p < 0.001$ ), temperature treatment ( $\chi^2 = 14.08$ ,  $df = 1$ ,  $p < 0.001$ ), fertilisation group ( $\chi^2 = 7.09$ ,  $df = 1$ ,  $p = 0.008$ ) and sex ( $\chi^2 = 4.34$ ,  $df = 1$ ,  $p = 0.037$ ). The interaction between food treatment and temperature treatment was not significant (LRT for model with and without interaction term:  $\chi^2 = 0.02$ ,  $df = 1$ ,  $p = 0.882$ ). Food restriction increased the probability of smolting whereas the warm temperature treatment decreased the probability of smolting (Figure 4A, B). Males were less likely to smolt than females, and fish in the Nov fertilisation group were less likely to smolt than those in the Nov-Dec group (Figure 4A, B).

The model describing the probability of maturing also had significant effects of food treatment ( $\chi^2 = 19.13$ ,  $df = 1$ ,  $p < 0.001$ ), temperature treatment ( $\chi^2 = 17.49$ ,  $df = 1$ ,  $p < 0.001$ ), sex ( $\chi^2 = 15.90$ ,  $df = 1$ ,  $p < 0.001$ ), but the effect of fertilisation group was not significant ( $\chi^2 = 1.04$ ,  $df = 1$ ,  $p = 0.308$ ). The interaction between food treatment and temperature treatment was not significant (LRT for model with and without interaction term:  $\chi^2 = 0.99$ ,  $df = 1$ ,  $p = 0.319$ ). In contrast to effects on smolting, the high food treatment increased the probability of maturing, as did the warm temperature treatment (Figure 4C, D). Males were less likely to mature than females in all treatments (Figure 4C, D).

The model describing the probability of being unassigned a life-history tactic included significant effects of food treatment ( $\chi^2 = 5.62$ ,  $df = 1$ ,  $p = 0.018$ ), temperature treatment ( $\chi^2 = 4.91$ ,  $df = 1$ ,  $p = 0.027$ ), sex ( $\chi^2 = 34.05$ ,  $df = 1$ ,  $p < 0.001$ ) and fertilisation group ( $\chi^2 = 7.69$ ,  $df = 1$ ,  $p = 0.006$ ). The interaction between food treatment and temperature treatment was not significant (LRT for model with and without interaction term:  $\chi^2 = 3.31$ ,  $df = 1$ ,  $p = 0.069$ ). Fish were significantly more likely to be unassigned a life history in either the low food or cool temperature treatments, as were males, and fish from the Nov fertilisation group (Table 5).





**Figure 4:** Co-efficient estimates ( $\pm$  95% confidence intervals) of GLMs describing probability of adopting (A, B) migratory and (C, D) maturation tactics in brown trout ( $n = 425$ , F1 offspring of wild trout from naturally facultatively anadromous population). Fish were classed as smolts or maturing after 18 months of tank rearing under varying food restriction and temperature treatments.

**Table 5:** Parameter estimates with associated standard errors (SE) for three binomial generalised linear models (GLM) predicting smolt (migratory) probability (dummy coded: smolt = 1, non-smolt = 0), freshwater maturation (dummy coded: mature/maturing = 1, immature = 0), and “unassigned” life-history tactics (dummy coded: unassigned = 1, smolt/mature = 0) in brown trout (n = 425). The reference level of each factor is in brackets, i.e. effects in all models were contrasted against female fish from the Nov-Dec fertilisation in the High food and Cool temperature treatment. Statistical significance was assessed at  $p < 0.05$ .

| Effect  | Estimate | SE    | t-value | p-value |
|---|----------|-------|---------|---------|
| <i>GLM of probability of smolting:</i>                        |          |       |         |         |
| Intercept (High-Cool, Female, Nov-Dec fertilisation)          | -2.559   | 0.494 | -5.176  | < 0.001 |
| Food: Low   | 1.811    | 0.513 | 3.533   | < 0.001 |
| Temperature: Warm   | -1.621   | 0.481 | -3.372  | 0.001   |
| Fertilisation group: Nov                                      | -1.341   | 0.569 | -2.358  | 0.018   |
| Sex: Male   | -0.849   | 0.417 | -2.037  | 0.042   |
| <i>GLM of probability of maturation:</i>                      |          |       |         |         |
| Intercept (High-Cool, Female, Nov-Dec fertilisation)          | 1.879    | 0.284 | 6.625   | < 0.001 |
| Food: Low   | -1.054   | 0.248 | -4.242  | < 0.001 |
| Temperature: Warm   | 1.004    | 0.247 | 4.073   | < 0.001 |
| Fertilisation group: Nov                                      | -0.261   | 0.255 | -1.023  | 0.306   |
| Sex: Male   | -0.963   | 0.248 | -3.888  | < 0.001 |
| <i>GLM of probability of being unassigned a life history:</i> |          |       |         |         |
| Intercept (High-Cool, Female, Nov-Dec fertilisation)          | -2.923   | 0.366 | -7.986  | < 0.001 |
| Food: Low   | 0.644    | 0.276 | 2.337   | 0.019   |
| Temperature: Warm   | -0.601   | 0.274 | -2.192  | 0.028   |
| Fertilisation group: Nov                                      | 0.784    | 0.281 | 2.789   | 0.005   |
| Sex: Male   | 1.655    | 0.311 | 5.320   | < 0.001 |

*Osmoregulatory performance*

570 Gill NKA activity varied according to life history ( $\chi^2 = 56.74$ ,  $df = 2$ ,  $p < 0.001$ ), but was not affected by sex ( $\chi^2 = 1.28$ ,  $df = 1$ ,  $p = 0.258$ ) or fertilisation group ( $\chi^2 = 0.72$ ,  $df = 1$ ,  $p = 0.397$ ). Post hoc testing showed smolts had significantly higher NKA activity than mature fish ( $t = -7.41$ ,  $df = 172$ ,  $p < 0.001$ ) and unassigned fish ( $t = 5.15$ ,  $df = 172$ ,  $p < 0.001$ ; Figure 5A). Similarly, plasma chloride concentration

575 was significantly affected by life history ( $\chi^2 = 52.14$ ,  $df = 2$ ,  $p < 0.001$ ), with no significant effect of sex ( $\chi^2 = 2.75$ ,  $df = 1$ ,  $p = 0.097$ ) or fertilisation group ( $\chi^2 = 2.03$ ,  $df = 1$ ,  $p = 0.154$ ). Smolts had significantly lower plasma chloride concentration after saltwater exposure than mature fish ( $t = 5.56$ ,  $df = 144$ ,  $p < 0.001$ ) and unassigned fish ( $t = -6.77$ ,  $df = 144$ ,  $p < 0.001$ ; see Figure 5B). Size-


580 corrected plasma chloride concentration decreased significantly with gill NKA activity in smolts ( $\chi^2 = 14.18$ ,  $df = 1$ ,  $p < 0.001$ , Figure 5C), however there was no significant relationship between size-corrected plasma chloride concentration and gill NKA activity in non-smolts ( $\chi^2 = 1.79$ ,  $df = 1$ ,  $p = 0.180$ , Figure 5D).

After accounting for the significant effect of body size ( $\chi^2 = 5.97$ ,  $df = 1$ ,  $p =$


585  $0.015$ ), the model describing plasma chloride concentration in smolts (marginal  $R^2 = 0.49$ ) did not retain a significant food  $\times$  temperature treatment interaction (LRT:  $\chi^2 = 0.26$ ,  $df = 1$ ,  $p = 0.610$ ). We detected a significant main effect of food treatment on plasma chloride concentration ( $\chi^2 = 5.29$ ,  $df = 1$ ,  $p = 0.021$ ), where the high food treatment was associated with lower chloride values (Figure 6A,

590 B). There was no significant effect of temperature treatment ( $\chi^2 = 2.26$ ,  $df = 1$ ,  $p = 0.133$ ), sex ( $\chi^2 = 1.60$ ,  $df = 1$ ,  $p = 0.205$ ) or fertilisation group ( $\chi^2 = 2.77$ ,  $df = 1$ ,  $p = 0.096$ ) on chloride concentrations. Mixed model analysis indicated non-significant effects of fork length ( $\chi^2 = 0.06$ ,  $df = 1$ ,  $p = 0.814$ ), food treatment ( $\chi^2 = 0.03$ ,  $df = 1$ ,  $p = 0.862$ ), temperature treatment ( $\chi^2 = 0.85$ ,  $df = 1$ ,  $p = 0.358$ ),

595 sex ( $\chi^2 = 2.47$ ,  $df = 1$ ,  $p = 0.116$ ) and fertilisation group ( $\chi^2 = 3.53$ ,  $df = 1$ ,  $p = 0.060$ ) on gill NKA activity in smolts (marginal  $R^2 = 0.20$ , Figure 6A). Overall, this indicates positive direct effects food treatment (independent of size) on saltwater tolerance of smolts, which were not reflected in gill NKA activity. See Tables S6 and S7 for parameter estimates from the mixed effect models.



**Figure 5:** Variation in (A) gill NKA activity, and (B) plasma chloride concentration of brown trout – classed as smolts (migratory tactics) and mature (non-migratory), or unassigned a life-history tactic – after 24 hours in salt water (30 ppt salinity). Size-corrected plasma chloride concentration was negatively related to Gill NKA activity in (C) smolts (dashed line = 95% confidence interval) but there was no relationship in (D) non-smolts (mature and unassigned fish).



**Figure 6:** Coefficient estimates ( $\pm$  95% confidence intervals) from the mixed effects models describing z-standardised (A) gill NKA activity and plasma chloride concentration of brown trout smolts after a 24 hours in salt water (30 ppt salinity). Variation in plasma chloride concentration of smolts according to food treatment is shown in (B), where the white lines in each box represent the median. Note that only one smolt was recorded in the High-Warm treatment, and thus there is no corresponding white line for the median in the High-Warm treatment.

## Discussion

Accelerating global change is exposing ecosystems to a multitude of co-occurring stressors, the implications of which are uncertain, particularly for migratory populations. Here we showed that food restriction increased the occurrence of a migratory phenotype (smolts), but conversely elevated temperature reduced smolting rates in favour of increased freshwater maturation (a phenotype consistent with a residency tactic). The observed effects on life-history were underpinned by complex, interactive effects of these putative stressors on underlying status traits associated with migratory decisions.

*Effects of multiple stressors on underlying morphological traits*

While stressors applied in isolation generally appeared to have negative effects on morphological traits, the effects of warming were less pronounced than food restriction, and varied depending on the response considered. Though fish at higher temperatures were smaller than their counterparts in the cool treatment, they maintained similar condition trajectories, and indeed had higher lipid stores at the end of the study, suggesting temperature may alter patterns of energy allocation in different ways to food restriction. Smaller sizes and higher lipid stores might arise if investment into gonadal development is prioritised over somatic growth earlier when environmental conditions appear favourable for early growth (*i.e.* the warm temperature treatments) (Jonsson *et al.* 2013), supported by the high prevalence of mature fish in warm treatments. Interestingly, the cumulative effects of food restriction and temperature were less than expected, based on their effects in isolation, suggesting complex antagonistic interactions between the stressors, whereby increased temperatures reduced body size and mass at high, but not low, food levels. The dampened response to temperature when combined with food restriction could perhaps be explained by metabolic rate depression under low food conditions, which has previously been documented in food-limited brown trout (Auer *et al.* 2015, 2016). This, together with overall reduced consumption rates, may simply have swamped any effects of temperature on growth in the low food treatments. Indeed, bioenergetics modelling of stream-dwelling rainbow trout growth has indicated growth may be more affected by factors influencing food intake rates (such as reduced overall food availability) than by direct effects of temperature, particularly during warmer summer months (Railsback and Rose 1999).

*Effects of multiple stressors on migration*

The antagonistic effects of food restriction and higher temperatures on physiological traits were not apparent at the level of migratory tactics. Indeed, opposing (additive) effects of these putative stressors on migratory phenotypes seemed initially to be counterintuitive. While an increase in the migratory tactic in response to food limitation is in line with previous work (Olsson *et al.* 2006;

Wysujack *et al.* 2009; O’Neal and Stanford 2011; Jones *et al.* 2015; Archer *et al.* 2019), surprisingly, a temperature increase of 1.8 °C above the natural  
 645 temperature regime of the source catchment reduced smolting rates. An energy limitation scenario (where an environmental stressor may act to prevent individuals from reaching genetically determined maturation thresholds) was supported in our results at the level of the status traits. Future migrants (*i.e.* smolts) were consistently smaller than fish that matured in fresh water (*i.e.* residents) and differences in mass and condition trajectories indicated migrants were energetically deficient (relative to mature fish). Energetic limitation appeared to be associated with low food availability, but less so with warmer temperatures.

Warmer temperatures have been proposed to generally increase the frequency  
 655 of migrants through energetic limitation, if associated elevated metabolic demands are not offset by increased energetic intake (Sloat and Reeves 2014; Kendall *et al.* 2014). However, warming can have a range of context-dependent impacts on patterns of energy acquisition and allocation in salmonines, which in turn may lead to a diversity of effects on life histories. For example, warmer  
 660 temperatures are associated with higher levels of growth up until some thermal optimum (Jensen 1990; Forseth and Jonsson 1994; Elliott *et al.* 1995; Ojanguren *et al.* 2001; Jonsson *et al.* 2013), but bioenergetics modelling shows that optimal temperatures for growth are negatively related to daily ration amount and body size (Beauchamp 2009). Thus, higher temperatures could either increase or  
 665 decrease average somatic growth, depending on food supply, the current distribution of fish sizes, and proximity to thermal growth optima. High somatic growth, along with high body condition and lipids, has been linked to increased freshwater maturation in facultatively migratory salmonines (Jonsson and Jonsson 1993; Dodson *et al.* 2013; Hecht *et al.* 2015), but other studies have  
 670 found that faster growing juveniles may be more likely to migrate at earlier ages and smaller sizes because they are more energetically constrained by limited food availability (owing to much higher metabolic costs) than slower growers (Forseth *et al.* 1999). Moreover, migration tendency is linked to the relative

productivity of marine and freshwater habitats, with anadromy more prevalent  
 675 in areas where the marine environment offers better opportunities for feeding  
 and growth (*e.g.* in higher latitudes) (Gross *et al.* 1988). Although the  
 temperature stressor we simulated in our study is in line with projected climate  
 warming scenarios of 1–3 °C (IPCC 2014), our warm temperature treatment  
 remained largely within the optimal temperature range for growth in brown  
 680 trout of 13–17 °C (Elliott *et al.* 1995; Elliott and Hurley 2000; Ojanguren *et al.*  
 2001) (maximum temperature in the warm treatments was 18.2 °C). It is  
 therefore likely that warmer temperatures did not tip most individuals into an  
 energetic deficit, thus fish were more likely to mature, rather than to smolt, in  
 the warm treatments.

685 Bioenergetic modelling of migratory variation in steelhead trout has suggested  
 that reductions in food resources can be mediated or exacerbated by water  
 temperatures to alter expression of life histories (Benjamin *et al.* 2013). Few  
 studies have empirically tested the cumulative effects of food supply and  
 temperature on migratory tactics, but from our study, it appears these two  
 690 environmental stressors may act additively, rather than synergistically, at least  
 for populations that are well within their thermal limits. Moreover, the positive  
 effect of temperature on maturation, coupled with negative temperature effects  
 on the frequencies of smolts and unassigned fish, indicates here that warming  
 acts to hasten the expression of life histories, driving earlier maturation instead  
 695 of migration. Similar changes in life-history dynamics have been predicted in  
 partially migratory masu salmon (*O. masou*), where favourable early growth  
 conditions associated with warming promoted maturation over migration and  
 caused an overall decline in life-history diversity (Morita *et al.* 2014).

Antagonistic effects of temperature and food on physiological traits (presumed  
 700 to underpin migratory decisions) were not translated at the level of migratory  
 tactics (where the putative stressors combined additively). Our results provide  
 additional evidence that multiple stressors can alter ecological responses in  
 unexpected ways, sometimes termed “ecological surprises” (Paine *et al.* 1998).  
 This suggests that effects of stressors can vary depending on the level of



705 organisation, or indeed the response, that is measured (Galic *et al.* 2018). It also underscores how environmental factors may affect migratory decisions directly, and not solely through environmentally-induced changes in putative cueing traits. For example, temperature can affect gene expression with long-lasting consequences for future behaviour and life history (Jonsson and Jonsson 2019).

710 There is some evidence to support that warming, in particular, can directly alter life-history tactics *e.g.* temperature hastened maturation at smaller sizes in nine-spined sticklebacks (*Pungitius pungitius*) independently of temperature-mediated growth (Kuparinen *et al.* 2011). Changes in somatic growth or energy allocation due to antagonistic effects of stressors therefore adds an additional

715 layer of complexity to our ability to infer future migratory tactics from patterns of juvenile growth.

#### *Early life conditions affect migration propensity*

The effects of fertilisation group on migratory propensity was an unexpected outcome of our study. While the relatively small numbers of brood stock used

720 makes it difficult to draw conclusions regarding differences between fertilisation groups, which most likely stem from inherited genetic effects or epigenetic/parental effects, differences might nonetheless reflect non-inherited variation arising from early-life environment that has knock-on effects for future phenotype (Burton and Metcalfe 2014). Moreover, differences in the

725 timing of readiness for reproduction/fertilisation also indicates differences among brood stock (*e.g.* spawning site in the wild) that may have translated into genetic or parental effects, rather than an effect of fertilisation date *per se*. Although both fertilisation groups experienced the same food restriction and temperature treatments, phenotypic differences that were established before

730 the application of treatments continued throughout the experimental phase of the study (*e.g.* fish from the Nov fertilisation group, whilst larger, tended to be in poorer condition), supporting genetic/ parental effects as factors which may be equally as important as downstream environmental conditions. Interestingly, individuals in the earlier fertilisation group were more likely to be

735 classed as “unassigned” by the end of the study. Delayed phenotypic expression

of migratory tactics can be reconciled with the existence of multiple decision windows, where an initial window determines the overall migration *versus* residency decision but then subsequent windows determine the actual age at which migrants become smolts, and residents mature (Ferguson *et al.* 2019).

740 Age at smolting, and age at maturation, may involve similar threshold mechanisms as the overarching migration decision but perhaps with different status traits playing the role of ‘trigger’; *e.g.* size or growth rates may determine age at smolting in Atlantic salmon but lipid levels may determine age at first maturation (Rowe *et al.* 1991; Jonsson and Jonsson 1993, 2005). Complex

745 environmental stressor effects on these various cues could then increase or decrease overall life-history diversity through temporal variation in migration or maturation patterns. In the Erriff system from which our brood stock was obtained, seaward-migration of wild fish is typically undertaken by smolts at two or three years after hatching (Gargan *et al.* 2016), with potentially up to half

750 of smolts migrating at age 3+. Similarly, although the age distribution of mature residents is unknown for the Erriff system, maturation at ages of 3+ and older is likely. As such, the patterns we observed might have reflected stressor effects on age at migration or age at maturity, in addition to effects on tactic choice *per se*. At least in relation to food restriction, we have no reason to expect that the

755 increased smolting rates we observed in our low food treatments at age 2+ would have been counter-balanced by increased maturation rates at age 3+ of the remaining unassigned fish; if anything, these were likely to have been simply delaying actual smolting until an older age, and therefore larger size (given that larger smolts are more likely to survive the critical transition to the marine

760 environment). In other words, smolting rate differences measured across ages 2+ and 3+ combined, if the experiment had been continued for an additional year, were likely to have been even more pronounced between food treatments. It is less obvious whether the same can be said for temperature effects, but we have no *a priori* reason to expect that age 3+ smolting rates would respond in

765 the opposite direction to sustained higher temperatures than age 2+ smolting rates.

*Stressor effects on future migratory capacity*

A final aim of our study was to assess if exposure to multiple stressors influenced the capacity of migratory individuals to successfully transition to salt water (an indicator of future migratory success). Smolts generally showed heightened hypo-osmoregulatory performance relative to non-smolts (for a given fish size as indexed by reduced plasma chloride levels following saltwater exposure), which was associated with increased Na<sup>+</sup>, K<sup>+</sup>-ATPase activity, a key enzyme involved in ion regulation. This is in agreement with many previous studies in salmonines that have described high NKA activity in smolts, which is related directly to ability to maintain homeostasis in seawater (McCormick *et al.* 1998, 2009b, 2013; Nilsen *et al.* 2007). We had expected the addition of stressors might further influence the hypo-osmoregulatory performance of smolts through negative effects on size, irrespective of enzyme activity, and indeed, we did detect a size dependency in plasma chloride levels, which was not reflected in gill NKA activity. Size-dependent increases in salinity tolerance that are independent of the size-dependent smolt decision have also been previously established for salmonines (McCormick *et al.* 1998). Thus, though the overall frequency of smolting was lower in optimal growth conditions (*i.e.* high food regimes in our study), the small number of smolts that were produced under these good growing conditions were larger and in better condition than their counterparts exposed to less favourable conditions. Intriguingly, we detected an additional negative effect of food restriction on hypo-osmoregulatory performance that was independent of size. Food limitation could potentially contribute to competitive interactions between individuals, emergence of dominance hierarchies, and generally heightened physiological stress, with implications for seawater tolerance and survival of brown trout (Pickering 1989; Sigholt and Finstad 1990; Liebert and Schreck 2006; Midwood *et al.* 2014). Collectively, these results suggest that food restriction may act as a stressor to migrants, which may have negative impacts for survival at sea (Ward and Slaney 1988), and possibly reduce the overall fitness of the migratory life history.

*Implications and considerations*

Considerable uncertainty still surrounds how environment and genetics integrate, via mediating physiological traits, to influence complex life-history patterns of facultatively migratory species (Crozier *et al.* 2008). Here, we focused on two stressors that are likely to occur in synchrony based on projections of global change, with important implications for an experimentally reared population of trout that can be extrapolated to wild populations, though some caveats should be noted. Whilst macroinvertebrate abundance and size are indeed projected to shrink with rising temperatures across the range of brown trout distribution (Durance and Ormerod 2007), in natural systems trout have been shown to track shifts in prey community assemblies (Woodward *et al.* 2010) or even become more selective in diet as temperatures increase (O’Gorman *et al.* 2016). Furthermore, given that our study population originated from a relatively cool catchment, it is possible that temperature increases in similar systems will primarily serve to increase fish growth/energetic status and promote residency, although this very much depends on how other key factors such as food supply, flow rates, biotic factors also respond to climate change. Any reductions in anadromy would likely alter the transfer of nutrients between freshwater and marine systems, with consequences for wider ecosystem processes (Doughty *et al.* 2016). If warming results in overall decreases in life-history diversity within populations (*e.g.* Benjamin *et al.* 2013; Morita *et al.* 2014) that are coherent over broad spatial scales, this could lead to a reduction in the “portfolio effect” in salmonines, whereby maintaining a range of phenotypic diversity buffers aggregations of populations and even entire species from changing environmental conditions (Schindler *et al.* 2015).

Expanding our approach to additional populations, including those that are closer to their thermal limits, *e.g.* in southern Europe, may alter the patterns we observed here (Almodóvar *et al.* 2012). There is some evidence for genotype by temperature interactions in key phenotypic traits in salmonines (Doctor *et al.* 2014) but the role of intrinsic factors relative to environmental context requires

further exploration (Ferguson *et al.* 2017). Common garden or reciprocal transplant style experiments in the wild would help to elucidate the mechanisms underpinning responses to multiple stressors in migratory species, whilst also incorporating the complexities of natural systems, such as changes in prey community structure, or abiotic correlates of warming (*e.g.* reduced oxygen/flow) (Clews *et al.* 2010). Nevertheless, our study has important implications for the development of management strategies to conserve facultatively migratory salmonines, a culturally iconic group in global decline due to aquaculture expansion, habitat degradation, and climate change (Costello 2009; Limburg and Waldman 2009).

### Acknowledgements

The authors would like to thank Brian Clarke, Deirdre Cotter, members of the FishEyE team at UCC, and the staff of Inland Fisheries Ireland and the Marine Institute for obtaining brood stock and for assistance in fish rearing, along with Robert Wynne, Ronan O'Sullivan, Peter Moran and Adam Kane for assistance in fish husbandry. This research was supported by an ERC Starting Grant (639192-ALH) and an SFI ERC Support Award awarded to TER. PMcG was supported in part by grants from Science Foundation Ireland (15/IA/3028 & 16/BBSRC/3316) and by grant-in-aid (RESPI/FS/16/01) from the Marine Institute (Ireland) as part of the Marine Research Programme by the Irish Government. We thank Dan Hall for his assistance in running the NKA activity assay and Jamie Coughlan for genotyping work. Adam Kane and Darío Fernández-Bellón provided useful comments on an earlier version of this manuscript. Any use of product or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

### References

- Almodóvar A, Nicola GG, Ayllón D, and Elvira B. 2012. Global warming threatens the persistence of Mediterranean brown trout. *Global Change Biology* **18**: 1549–60.
- Archer LC, Hutton SA, Harman L, O'Grady MN, Kerry JP, Poole WR, Gargan P, McGinnity P, and Reed TE. 2019. The interplay between extrinsic and intrinsic factors in determining migration decisions in brown trout (*Salmo trutta*): An experimental study. *Frontiers in Ecology and Evolution* **7**: 222.

- 860 Auer SK, Salin K, Anderson GJ, and Metcalfe NB. 2016. Flexibility in metabolic rate and activity level determines individual variation in overwinter performance. *Oecologia* **182**: 703–12.
- Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015. Flexibility in metabolic rate confers a growth advantage under changing food availability.
- 865 *Journal of Animal Ecology* **84**: 1405–11.
- Ball JP, Nordengren C, and Wallin K. 2001. Partial migration by large ungulates: Characteristics of seasonal moose *Alces alces* ranges in northern Sweden. *Wildlife Biology* **7**: 39–47.
- Barton K. 2019. MuMIn: Multi-Model Inference. R package version 1.43.6.
- 870 <https://CRAN.R-project.org/package=MuMIn>.
- Beauchamp DA. 2009. Bioenergetic ontogeny: Linking climate and mass-specific feeding to life-cycle growth and survival of salmon. *American Fisheries Society Symposium* **70**: 1–19.
- Benjamin JR, Connolly PJ, Romine JG, and Perry RW. 2013. Potential effects of changes in temperature and food resources on life history trajectories of juvenile *Oncorhynchus mykiss*. *Transactions of the American Fisheries Society* **142**: 208–20.
- 875
- Berthold P and Querner U. 1982. Partial migration in birds: Experimental proof of polymorphism as a controlling system. *Experientia* **38**: 805–6.
- Bolger T and Connolly PL. 1989. The selection of suitable indices for the measurement and analysis of fish condition. *Journal of Fish Biology* **34**: 171–82.
- 880
- Breitburg DL, Baxter JW, Hatfield CA, Howarth RW, Jones CG, Lovett GM, and Wigand C. 1998. Understanding effects of multiple stressors: Ideas and challenges. In: Pace ML, Groffman PM (Eds). *Successes, Limitations, and Frontiers in Ecosystem Science*. New York, NY: Springer New York.
- 885
- Breitburg DL, Sanders JG, Gilmour CC, Hatfield CA, Osman RW, Riedel GF, Seitzinger SP, and Seitzinger SP. 1999. Variability in responses to nutrients and trace elements, and transmission of stressor effects through an estuarine food web. *Limnology and Oceanography* **44**: 837–63.
- Buoro M, Gimenez O, and Prévost E. 2012. Assessing adaptive phenotypic plasticity by means of conditional strategies from empirical data: The latent environmental threshold model. *Evolution* **66**: 996–1009.
- 890
- Burton T and Metcalfe NB. 2014. Can environmental conditions experienced in early life influence future generations? *Proceedings of the Royal Society of London B: Biological Sciences* **281**: 20140311.
- 895
- Byrne M and Przeslawski R. 2013. Multistressor impacts of warming and acidification of the ocean on marine invertebrates' life histories. *Integrative and Comparative Biology* **53**: 582–96.
- Cagnacci F, Focardi S, Heurich M, Stache A, Hewison AJM, Morellet N, Kjellander P, Linnell JDC, Mysterud A, Neteler M, Delucchi L, Ossi F, and Urbano F. 2011. Partial migration in roe deer: Migratory and resident tactics are end points of a behavioural gradient determined by ecological factors. *Oikos* **120**: 1790–802.
- 900
- Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011. The ecology and evolution of partial migration. *Oikos* **120**: 1764–75.

- 905 Chapman BB, Hulthén K, Brodersen J, Nilsson PA, Skov C, Hansson L-A, and Brönmark C. 2012. Partial migration in fishes: Causes and consequences. *Journal of Fish Biology* **81**: 456–78.
- Clarke WC. 1982. Evaluation of the seawater challenge test as an index of marine survival. *Aquaculture* **28**: 177–83.
- 910 Clews E, Durance I, Vaughan IP, and Ormerod SJ. 2010. Juvenile salmonid populations in a temperate river system track synoptic trends in climate. *Global Change Biology* **16**: 3271–83.
- Costello MJ. 2009. How sea lice from salmon farms may cause wild salmonid declines in Europe and North America and be a threat to fishes elsewhere. *Proceedings of the Royal Society B: Biological Sciences* **276**: 3385–94.
- 915 Côté IM, Darling ES, and Brown CJ. 2016. Interactions among ecosystem stressors and their importance in conservation. *Proceedings of the Royal Society B: Biological Sciences* **283**: 20152592.
- Crain CM, Kroeker K, and Halpern BS. 2008. Interactive and cumulative effects of multiple human stressors in marine systems. *Ecology Letters* **11**: 1304–15.
- 920 Crozier LG, Hendry AP, Lawson PW, Quinn TP, Mantua NJ, Battin J, Shaw RG, and Huey RB. 2008. Potential responses to climate change in organisms with complex life histories: Evolution and plasticity in Pacific salmon. *Evolutionary Applications* **1**: 252–70.
- 925 Davidsen JG, Daverdin M, Sjørnsen AD, Rønning L, Arnekleiv JV, and Koksvik JI. 2014. Does reduced feeding prior to release improve the marine migration of hatchery brown trout *Salmo trutta* smolts? *Journal of Fish Biology* **85**: 1992–2002.
- 930 Dębowski P and Dobosz S. 2016. Influence of parental life history on maturation and smoltification in brown trout (*Salmo trutta* L.). *Archives of Polish Fisheries* **24**: 177–86.
- Dell AI, Pawar S, and Savage VM. 2011. Systematic variation in the temperature dependence of physiological and ecological traits. *Proceedings of the National Academy of Sciences* **108**: 10591–6.
- 935 Dellefors C and Faremo U. 1988. Early sexual maturation in males of wild sea trout, *Salmo trutta* L., inhibits smoltification. *Journal of Fish Biology* **33**: 741–9.
- Doctor K, Berejikian B, Hard JJ, and VanDoornik D. 2014. Growth-mediated life history traits of steelhead reveal phenotypic divergence and plastic response to temperature. *Transactions of the American Fisheries Society* **143**: 317–33.
- 940 Dodson JJ, Aubin-Horth N, Thériault V, and Páez DJ. 2013. The evolutionary ecology of alternative migratory tactics in salmonid fishes: Alternative migratory tactics as threshold traits. *Biological Reviews* **88**: 602–25.
- Doswald N, Willis SG, Collingham YC, Pain DJ, Green RE, and Huntley B. 2009. Potential impacts of climatic change on the breeding and non-breeding ranges and migration distance of European Sylvia warblers. *Journal of Biogeography* **36**: 1194–208.
- 945 Doughty CE, Roman J, Faurby S, Wolf A, Haque A, Bakker ES, Malhi Y, Dunning JB, and Svenning J-C. 2016. Global nutrient transport in a world of giants. *Proceedings of the National Academy of Sciences* **113**: 868–73.

- 950 Durance I and Ormerod SJ. 2007. Climate change effects on upland stream macroinvertebrates over a 25-year period. *Global Change Biology* **13**: 942–57.
- Elliott JM and Elliott JA. 2010. Temperature requirements of Atlantic salmon *Salmo salar*, brown trout *Salmo trutta* and Arctic charr *Salvelinus alpinus*: Predicting the effects of climate change. *Journal of Fish Biology* **77**: 1793–817.
- 955 Elliott JM and Hurley MA. 2000. Daily energy intake and growth of piscivorous brown trout, *Salmo trutta*. *Freshwater Biology* **44**: 237–45.
- Elliott JM, Hurley MA, and Fryer RJ. 1995. A new, improved growth model for brown trout, *Salmo trutta*. *Functional Ecology* **9**: 290–8.
- 960 Eyto E de, Dalton C, Dillane M, Jennings E, McGinnity P, O'Dwyer B, Poole R, Rogan G, and Taylor D. 2016. The response of North Atlantic diadromous fish to multiple stressors, including land use change: A multidecadal study. *Canadian Journal of Fisheries and Aquatic Sciences* **73**: 1759–69.
- Ferguson A, Reed TE, Cross TF, McGinnity P, and Prodöhl PA. 2019. Anadromy, potamodromy and residency in brown trout *Salmo trutta*: The role of genes and the environment. *Journal of Fish Biology* **95**: 692–718.
- 965 Ferguson A, Reed TE, McGinnity P, and Prodöhl P. 2017. Anadromy in brown trout (*Salmo trutta*): A review of the relative roles of genes and environmental factors and the implications for management and conservation. In: Sea Trout: Science and Management - Proceedings of the 2nd International Sea Trout Symposium. Matador, Leicestershire, UK.
- 970 Finstad B and Ugedal O. 1998. Smolting of sea trout *Salmo trutta* L. in northern Norway. *Aquaculture* **168**: 341–9.
- Folt CL, Chen CY, Moore MV, and Burnaford J. 1999. Synergism and antagonism among multiple stressors. *Limnology and Oceanography* **44**: 864–77.
- 975 Forseth T and Jonsson B. 1994. The growth and food ration of piscivorous brown trout (*Salmo trutta*). *Functional Ecology* **8**: 171–7.
- Forseth T, Larsson S, Jensen AJ, Jonsson B, Näslund I, and Berglund I. 2009. Thermal growth performance of juvenile brown trout *Salmo trutta*: No support for thermal adaptation hypotheses. *Journal of Fish Biology* **74**: 133–49.
- 980 Forseth T, Nesje TF, Jonsson B, and Hårsaker K. 1999. Juvenile migration in brown trout: A consequence of energetic state. *Journal of Animal Ecology* **68**: 783–93.
- Fussmann KE, Schwarzmüller F, Brose U, Jousset A, and Rall BC. 2014. Ecological stability in response to warming. *Nature Climate Change* **4**: 206–10.
- 985 Galic N, Sullivan LL, Grimm V, and Forbes VE. 2018. When things don't add up: Quantifying impacts of multiple stressors from individual metabolism to ecosystem processing. *Ecology Letters* **21**: 568–77.
- Gargan P, Kelly F, Shephard S, and Whelan K. 2016. Temporal variation in sea trout *Salmo trutta* life history traits in the Erriff River, western Ireland. *Aquaculture Environment Interactions* **8**: 675–89.
- 990 Gillooly JF, Brown JH, West GB, Savage VM, and Charnov EL. 2001. Effects of size and temperature on metabolic rate. *Science* **293**: 2248–51.
- Gross MR, Coleman RM, and McDowall RM. 1988. Aquatic productivity and the evolution of diadromous fish migration. *Science* **239**: 1291–3.



- Hansson L-A and Hylander S. 2009. Size-structured risk assessments govern *Daphnia* migration. *Proceedings of the Royal Society B: Biological Sciences* **276**: 331–6.
- Hard JJ, Winans GA, and Richardson JC. 1999. Phenotypic and genetic architecture of juvenile morphometry in Chinook salmon. *Journal of Heredity* **90**: 597–606.
- Hari RE, Livingstone DM, Siber R, Burkhardt-Holm P, and Güttinger H. 2006. Consequences of climatic change for water temperature and brown trout populations in Alpine rivers and streams. *Global Change Biology* **12**: 10–26.
- Harvey BP, Gwynn-Jones D, and Moore PJ. 2013. Meta-analysis reveals complex marine biological responses to the interactive effects of ocean acidification and warming. *Ecology and Evolution* **3**: 1016–30.
- Hebblewhite M and Merrill EH. 2011. Demographic balancing of migrant and resident elk in a partially migratory population through forage–predation tradeoffs. *Oikos* **120**: 1860–70.
- Hecht BC, Hard JJ, Thrower FP, and Nichols KM. 2015. Quantitative genetics of migration-related traits in rainbow and steelhead trout. *G3: Genes, Genomes, Genetics* **5**: 873–89.
- Huey RB and Stevenson RD. 1979. Integrating thermal physiology and ecology of ectotherms: A discussion of approaches. *Integrative and Comparative Biology* **19**: 357–66.
- IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. In: Geneva, Switzerland.
- Jackson MC, Loewen CJG, Vinebrooke RD, and Chimimba CT. 2016. Net effects of multiple stressors in freshwater ecosystems: A meta-analysis. *Global Change Biology* **22**: 180–9.
- Jensen AJ. 1990. Growth of young migratory brown trout *Salmo trutta* correlated with water temperature in Norwegian rivers. *Journal of Animal Ecology* **59**: 603–14.
- Jones DA, Bergman E, and Greenberg L. 2015. Food availability in spring affects smolting in brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 1694–9.
- Jonsson B. 1985. Life history patterns of freshwater resident and sea-run migrant brown trout in Norway. *Transactions of the American Fisheries Society* **114**: 182–94.
- Jonsson B and Jonsson N. 1993. Partial migration: Niche shift versus sexual maturation in fishes. *Reviews in Fish Biology and Fisheries* **3**: 348–65.
- Jonsson B and Jonsson N. 2005. Lipid energy reserves influence life-history decision of Atlantic salmon (*Salmo salar*) and brown trout (*S. trutta*) in fresh water. *Ecology of Freshwater Fish* **14**: 296–301.
- Jonsson B and Jonsson N. 2009. A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of Fish Biology* **75**: 2381–447.
- Jonsson B and Jonsson N. 2019. Phenotypic plasticity and epigenetics of fish: Embryo temperature affects later-developing life-history traits. *Aquatic Biology* **28**: 21–32.

- Jonsson B, Jonsson N, and Finstad AG. 2013. Effects of temperature and food quality on age and size at maturity in ectotherms: An experimental test with Atlantic salmon. *Journal of Animal Ecology* **82**: 201–10.
- Kaitala A, Kaitala V, and Lundberg P. 1993. A theory of partial migration. *The American Naturalist* **142**: 59–81.
- Keeton JT, Hafley BS, Eddy SM, Moser CR, McManus BJ, and Leffler TP. 2003. Rapid determination of moisture and fat in meats by microwave and nuclear magnetic resonance analysis. *Journal of AOAC International* **86**: 1193–202.
- Kendall NW, McMillan JR, Sloat MR, Buehrens TW, Quinn TP, Pess GR, Kuzishchin KV, McClure MM, and Zabel RW. 2014. Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): A review of the processes and patterns. *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 319–42.
- Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, and Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): A review of aspects of their life histories. *Ecology of Freshwater Fish* **12**: 1–59.
- Kovach RP, Muhlfeld CC, Al-Chokhachy R, Dunham JB, Letcher BH, and Kershner JL. 2016. Impacts of climatic variation on trout: A global synthesis and path forward. *Reviews in Fish Biology and Fisheries* **26**: 135–51.
- Kuparinen A, Cano JM, Loehr J, Herczeg G, Gonda A, and Merilä J. 2011. Fish age at maturation is influenced by temperature independently of growth. *Oecologia* **167**: 435–43.
- Lack D. 1943. The problem of partial migration. *British Birds* **37**: 122–30.
- Lenth R. 2019. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.3.5.1. <https://CRAN.R-project.org/package=emmeans>.
- Liebert AM and Schreck CB. 2006. Effects of acute stress on osmoregulation, feed intake, IGF-I, and cortisol in yearling steelhead trout (*Oncorhynchus mykiss*) during seawater adaptation. *General and Comparative Endocrinology* **148**: 195–202.
- Limburg KE and Waldman JR. 2009. Dramatic declines in north Atlantic diadromous fishes. *BioScience* **59**: 955–65.
- Lundberg P. 1987. Partial bird migration and evolutionarily stable strategies. *Journal of Theoretical Biology* **125**: 351–60.
- Lundberg P. 1988. The evolution of partial migration in birds. *Trends in Ecology & Evolution* **3**: 172–5.
- McCormick SD. 2012. 5 - Smolt physiology and endocrinology. In: McCormick SD, Farrell AP, Brauner CJ (Eds). *Fish physiology*. Cambridge, Massachusetts: Academic Press.
- McCormick SD, Hansen LP, Quinn TP, and Saunders RL. 1998. Movement, migration, and smolting of Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* **55**: 77–92.
- McCormick SD, Lerner DT, Monette MY, Nieves-Puigdollér K, Kelly JT, and Björnsson BT. 2009a. Taking it with you when you go: How perturbations to the freshwater environment, including temperature, dams, and contaminants, affect marine survival of salmon. *Challenges For Diadromous Fishes in a*

- 1085        *Dynamic Global Environment American Fisheries Society Symposium* **69**: 195–214.
- McCormick SD, Regish AM, and Christensen AK. 2009b. Distinct freshwater and seawater isoforms of Na<sup>+</sup>/K<sup>+</sup>-ATPase in gill chloride cells of Atlantic salmon. *Journal of Experimental Biology* **212**: 3994–4001.
- 1090        McCormick SD, Regish AM, Christensen AK, and Björnsson BT. 2013. Differential regulation of sodium–potassium pump isoforms during smolt development and seawater exposure of Atlantic salmon. *Journal of Experimental Biology* **216**: 1142–51.
- 1095        McGinnity P, Eleanor J, Elvira deEyto, Allott Norman, Samuelsson Patrick, Rogan Gerard, Whelan Ken, and Cross Tom. 2009. Impact of naturally spawning captive-bred Atlantic salmon on wild populations: Depressed recruitment and increased risk of climate-mediated extinction. *Proceedings of the Royal Society B: Biological Sciences* **276**: 3601–10.
- 1100        Metcalfe NB. 1998. The interaction between behavior and physiology in determining life history patterns in Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* **55**: 93–103.
- Midwood J, Larsen M, Boel M, Jepsen N, Aarestrup K, and Cooke S. 2014. Does cortisol manipulation influence outmigration behaviour, survival and growth of sea trout? A field test of carryover effects in wild fish. *Marine Ecology Progress Series* **496**: 135–44.
- 1105        Morita K, Tamate T, Kuroki M, and Nagasawa T. 2014. Temperature-dependent variation in alternative migratory tactics and its implications for fitness and population dynamics in a salmonid fish. *Journal of Animal Ecology* **83**: 1268–78.
- 1110        Nevoux M, Finstad B, Davidsen JG, Finlay R, Josset Q, Poole R, Höjesjö J, Aarestrup K, Persson L, Tolvanen O, and Jonsson B. 2019. Environmental influences on life history strategies in partially anadromous brown trout (*Salmo trutta*, Salmonidae). *Fish and Fisheries* **In Press**.
- Newton I. 2008. The migration ecology of birds. Oxford, UK: Academic Press.
- 1115        Nielsen D, Hyldig G, Nielsen J, and Nielsen HH. 2005. Lipid content in herring (*Clupea harengus* L.)—influence of biological factors and comparison of different methods of analyses: Solvent extraction, Fatmeter, NIR and NMR. *LWT - Food Science and Technology* **38**: 537–48.
- 1120        Nilsen TO, Ebbesson LOE, Madsen SS, McCormick SD, Andersson E, Björnsson BT, Prunet P, and Stefansson SO. 2007. Differential expression of gill Na<sup>+</sup>,K<sup>+</sup>-ATPase $\alpha$  - and  $\beta$ -subunits, Na<sup>+</sup>,K<sup>+</sup>,2Cl<sup>-</sup> cotransporter and CFTR anion channel in juvenile anadromous and landlocked Atlantic salmon *Salmo salar*. *Journal of Experimental Biology* **210**: 2885–96.
- Northcote TG and Ward FJ. 1985. Lake resident and migratory smelt, *Retropinna retropinna* (Richardson), of the lower Waikato River system, New Zealand. *Journal of Fish Biology* **27**: 113–129.
- 1125        O’Gorman EJ, Fitch JE, and Crowe TP. 2012. Multiple anthropogenic stressors and the structural properties of food webs. *Ecology* **93**: 441–8.
- O’Gorman EJ, Ólafsson ÓP, Demars BOL, Friberg N, Guðbergsson G, Hannesdóttir ER, Jackson MC, Johansson LS, McLaughlin ÓB, Ólafsson JS, Woodward G, and

- 1130 Gíslason GM. 2016. Temperature effects on fish production across a natural thermal gradient. *Global Change Biology* **22**: 3206–20.
- Ojanguren AF, Reyes-Gavilán FG, and Braña F. 2001. Thermal sensitivity of growth, food intake and activity of juvenile brown trout. *Journal of Thermal Biology* **26**: 165–70.
- 1135 Olsson IC, Greenberg LA, Bergman E, and Wysujack K. 2006. Environmentally induced migration: The importance of food. *Ecology Letters* **9**: 645–51.
- O’Neal SL and Stanford JA. 2011. Partial migration in a robust brown trout population of a Patagonian river. *Transactions of the American Fisheries Society* **140**: 623–35.
- 1140 Ormerod SJ, Dobson M, Hildrew AG, and Townsend CR. 2010. Multiple stressors in freshwater ecosystems. *Freshwater Biology* **55**: 1–4.
- Paine RT, Tegner MJ, and Johnson EA. 1998. Compounded perturbations yield ecological surprises. *Ecosystems* **1**: 535–45.
- Parmesan C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* **37**: 637–69.
- 1145 Petchey OL, McPhearson PT, Casey TM, and Morin PJ. 1999. Environmental warming alters food-web structure and ecosystem function. *Nature* **402**: 69.
- Piche J, Hutchings JA, and Blanchard W. 2008. Genetic variation in threshold reaction norms for alternative reproductive tactics in male Atlantic salmon, *Salmo salar*. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1571–5.
- 1150 Pickering AD. 1989. Environmental stress and the survival of brown trout, *Salmo trutta*. *Freshwater Biology* **21**: 47–55.
- Pinheiro J, Bates D, DebRoy S, Sarkar D, and R Core Team. 2019. nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-139. URL <https://CRAN.R-project.org/package=nlme>.
- 1155 Przeslawski R, Byrne M, and Mellin C. 2015. A review and meta-analysis of the effects of multiple abiotic stressors on marine embryos and larvae. *Global Change Biology* **21**: 2122–40.
- Pulido F. 2011. Evolutionary genetics of partial migration – the threshold model of migration revis(it)ed. *Oikos* **120**: 1776–83.
- 1160 Pulido F, Berthold P, and vanNoordwijk AJ. 1996. Frequency of migrants and migratory activity are genetically correlated in a bird population: Evolutionary implications. *Proceedings of the National Academy of Sciences of the United States of America* **93**: 14642–7.
- 1165 R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- Railsback SF and Rose KA. 1999. Bioenergetics modelling of stream trout growth: Temperature and food consumption effects. *Transactions of the American Fisheries Society* **128**: 241–56.
- 1170 Rall BC, Vucic-Pestic O, Ehnes RB, Emmerson M, and Brose U. 2010. Temperature, predator–prey interaction strength and population stability. *Global Change Biology* **16**: 2145–57.

- Riddell BE and Leggett WC. 1981. Evidence of an adaptive basis for geographic variation in body morphology and time of downstream migration of juvenile Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* **38**: 308–20.
- Rowe DK, Thorpe JE, and Shanks AM. 1991. Role of fat stores in the maturation of male Atlantic salmon (*Salmo salar*) parr. *Canadian Journal of Fisheries and Aquatic Sciences* **48**: 405–13.
- Sanderson EW, Jaiteh M, Levy MA, Redford KH, Wannebo AV, and Woolmer G. 2002. The human footprint and the last of the wild: The human footprint is a global map of human influence on the land surface, which suggests that human beings are stewards of nature, whether we like it or not. *BioScience* **52**: 891–904.
- Schindler DE, Armstrong JB, and Reed TE. 2015. The portfolio concept in ecology and evolution. *Frontiers in Ecology and the Environment* **13**: 257–63.
- Schultz ET and McCormick SD. 2012. Euryhalinity in an evolutionary context. In: Fish physiology. Cambridge, Massachusetts: Academic Press.
- Sigholt T and Finstad B. 1990. Effect of low temperature on seawater tolerance in Atlantic salmon (*Salmo salar*) smolts. *Aquaculture* **84**: 167–72.
- Sinclair BJ, Marshall KE, Sewell MA, Levesque DL, Willett CS, Slotsbo S, Dong Y, Harley CDG, Marshall DJ, Helmuth BS, and Huey RB. 2016. Can we predict ectotherm responses to climate change using thermal performance curves and body temperatures? *Ecology Letters* **19**: 1372–85.
- Sloat MR and Reeves GH. 2014. Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences* **71**: 491–501.
- Swingland IR and Greenwood PJ. 1984. Ecology of animal movement. Oxford, UK: Clarendon Press.
- Tanguy JM, Ombredane D, Baglinière JL, and Prunet P. 1994. Aspects of parr-smolt transformation in anadromous and resident forms of brown trout (*Salmo trutta*) in comparison with Atlantic salmon (*Salmo salar*). *Aquaculture* **121**: 51–63.
- Tomkins JL and Hazel W. 2007. The status of the conditional evolutionarily stable strategy. *Trends in Ecology & Evolution* **22**: 522–8.
- Toussaint CA, Médale F, Davenel A, Fauconneau B, Haffray P, and Akoka S. 2002. Determination of the lipid content in fish muscle by a self-calibrated NMR relaxometry method: Comparison with classical chemical extraction methods. *Journal of the Science of Food and Agriculture* **82**: 173–8.
- Villar-Guerra D del, Larsen MH, Baktoft H, Koed A, and Aarestrup K. 2019. The influence of initial developmental status on the life-history of sea trout (*Salmo trutta*). *Scientific Reports* **9**: 1–13.
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M, Hoegh-Guldberg O, and Bairlein F. 2002. Ecological responses to recent climate change. *Nature* **416**: 389.

- Ward BR and Slaney PA. 1988. Life history and smolt-to-adult survival of Keogh River steelhead trout (*Salmo gairdneri*) and the relationship to smolt size. *Canadian Journal of Fisheries and Aquatic Sciences* **45**: 1110–22.
- 1220 Woodward G, Dybkjær JB, Ólafsson JS, Gíslason GM, Hannesdóttir ER, and Friberg N. 2010. Sentinel systems on the razor's edge: Effects of warming on Arctic geothermal stream ecosystems. *Global Change Biology* **16**: 1979–91.
- 1225 Wysujack K, Greenberg LA, Bergman E, and Olsson IC. 2009. The role of the environment in partial migration: Food availability affects the adoption of a migratory tactic in brown trout *Salmo trutta*. *Ecology of Freshwater Fish* **18**: 52–9.

## Supporting Information

---

### **Food and temperature stressors have opposing effects in determining flexible migration decisions in brown trout (*Salmo trutta*)**

Louise C. Archer<sup>\*1,2</sup>, Stephen A. Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, Stephen D. McCormick<sup>3</sup>, Michael N. O'Grady<sup>4</sup>, Joseph P. Kerry<sup>4</sup>, W. Russell Poole<sup>5</sup>, Patrick Gargan<sup>6</sup>, Philip McGinnity<sup>1,5</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> *School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.*


<sup>2</sup> *Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland.*

<sup>3</sup> *U.S. Geological Survey, Leetown Science Centre, S.O. Conte Anadromous Fish Research Laboratory, Turners Falls, MA, 01376, USA.*

<sup>4</sup> *Food Packaging Group, School of Food and Nutritional Sciences, University College Cork, Cork, Ireland.*


<sup>5</sup> *Marine Institute, Furnace, Newport, Co. Mayo, Ireland.*

<sup>6</sup> *Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.*




**Figure S1:** Location in the west of Ireland of brown trout brood stock collected by seine netting in winter 2015, and used to produce F1 offspring for an experimental tank-rearing study. Fish used in the study were offspring of brood stock collected in Tawnyard Lough in the Erriff catchment (a population with a high natural frequency of anadromy).





**Figure S2:** Weekly temperatures (open circles) for the three years preceding the study, as recorded at the Marine Institute long-term monitoring station at Newport, Mayo (west of Ireland), close to the Erriff catchment where brood stock used in the experiment originated. Red and blue lines are the mean weekly temperatures (averaged across tanks) for the cool and warm treatments in the laboratory experiment. The grey line represents the thermal growth optimum for trout (13.9 °C) (Elliott and Hurley 2000).



**Figure S3:** Final measures of (A) length, (B) condition, and (C) whole body lipids of brown trout offspring classed as either smolts (migratory) or unassigned a life-history tactic at the end of the experimental study. Median values of length, condition, and whole body lipids are represented by the white horizontal lines in each box in (A), (B), and (C).

**Table SI:** Brood stock crossing design for producing F1 offspring of wild-origin brown trout collected by seine netting from a population in Tawnyard Lough in the west of Ireland. Each female was stripped of eggs, which were then fertilised by the milt of two males from the same population and incubated in a hatchery facility within the Burrishoole catchment. Fish were subsequently reared in two separate groups according to fertilisation date, where “November fertilisation group” refers to offspring produced from crosses made on 12<sup>th</sup> November 2015, and “Nov-Dec fertilisation group” refers to offspring produced from crosses on 27<sup>th</sup> November and 10<sup>th</sup> December 2015.

| Stripping date | Fertilisation Group | Catchment         | Sex    | ID  | Male cross #1 | Male cross #2 |
|----------------|---------------------|-------------------|--------|-----|---------------|---------------|
| 12/11/2015     | Nov                 | Erriff (Tawnyard) | Female | TF1 | TM1           | TM2           |
| 12/11/2015     | Nov                 | Erriff (Tawnyard) | Female | TF2 | TM1           | TM2           |
| 12/11/2015     | Nov                 | Erriff (Tawnyard) | Female | TF3 | TM3           | TM4           |
| 27/11/2015     | Nov-Dec             | Erriff (Tawnyard) | Female | TF4 | TM5           | TM6           |
| 27/11/2015     | Nov-Dec             | Erriff (Tawnyard) | Female | TF5 | TM5           | TM6           |
| 27/11/2015     | Nov-Dec             | Erriff (Tawnyard) | Female | TF6 | TM7           | TM8           |
| 10/12/2015     | Nov-Dec             | Erriff (Tawnyard) | Female | TF7 | TM9           | TM10          |

**Table S2:** Parameter estimates with associated standard errors (SE) for mixed model analyses testing the effects of food treatment, temperature treatment, fertilisation group, and sex on the fork length (cm), mass (g), and condition trajectories of brown trout over 18 months of experimental tank rearing. Measurements were taken at key periods denoted by “time”. Effects were contrasted against female fish from the Nov-Dec fertilisation group in the high food and cool temperature treatment. Responses are z-standardised and statistical significance was assessed at  $p < 0.05$ .

| Effect                                  | Estimate | SE   | t-value | p-value |
|---|----------|------|---------|---------|
| <i>Mixed model for length</i>           |          |      |         |         |
| Intercept (Female, Nov-Dec, Cool, High) | 0.48     | 0.07 | 7.28    | < 0.001 |
| Time                                    | 0.87     | 0.02 | 51.23   | < 0.001 |
| Time <sup>2</sup>                       | -0.22    | 0.01 | -17.45  | < 0.001 |
| Sex: Male                               | -0.04    | 0.05 | -0.85   | 0.397   |
| Fertilisation group: Nov                | 0.42     | 0.06 | 7.53    | < 0.001 |
| Temperature: Warm                       | -0.24    | 0.08 | -3.23   | 0.001   |
| Food: Low                               | -0.90    | 0.07 | -12.06  | < 0.001 |
| Time × Warm temperature                 | 0.06     | 0.02 | 3.02    | 0.003   |
| Time × Low food                         | -0.32    | 0.02 | -16.08  | < 0.001 |
| Low food × Warm temperature             | 0.42     | 0.11 | 3.96    | < 0.001 |
| <i>Mixed model for mass</i>             |          |      |         |         |
| Intercept (Female, Nov-Dec, Cool, High) | 0.58     | 0.07 | 8.18    | < 0.001 |
| Time                                    | 0.89     | 0.03 | 32.55   | < 0.001 |
| Time <sup>2</sup>                       | -0.13    | 0.02 | -8.09   | < 0.001 |
| Sex: Male                               | -0.08    | 0.05 | -1.41   | 0.159   |
| Fertilisation group: Nov                | 0.28     | 0.06 | 4.93    | < 0.001 |
| Temperature: Warm                       | -0.29    | 0.08 | -3.66   | < 0.001 |
| Food: Low                               | -1.03    | 0.08 | -13.26  | < 0.001 |
| Time × Warm temperature                 | -0.01    | 0.04 | -0.15   | 0.880   |
| Time × Low food                         | -0.62    | 0.04 | -16.05  | < 0.001 |
| Low food × Warm temperature             | 0.41     | 0.11 | 3.69    | < 0.001 |
| Time × Low food × Warm temperature      | 0.15     | 0.06 | 2.72    | 0.007   |
| <i>Mixed model for condition</i>        |          |      |         |         |
| Intercept (Female, Nov-Dec, Cool, High) | 0.76     | 0.09 | 8.36    | < 0.001 |
| Time                                    | -0.29    | 0.04 | -7.88   | < 0.001 |
| Time <sup>2</sup>                       | -0.17    | 0.03 | -5.83   | < 0.001 |
| Sex: Male                               | 0.04     | 0.07 | 0.54    | 0.593   |
| Fertilisation group: Nov                | -0.21    | 0.07 | -2.94   | 0.003   |
| Temperature: Warm                       | -0.14    | 0.10 | -1.41   | 0.160   |
| Food: Low                               | -1.02    | 0.09 | -10.74  | < 0.001 |
| Time × Warm temperature                 | 0.13     | 0.04 | 3.08    | 0.002   |
| Time × Low food                         | -0.27    | 0.04 | -6.31   | < 0.001 |
| Low food × Warm temperature             | 0.35     | 0.13 | 2.64    | 0.009   |

**Table S3:** Parameter estimates with associated standard errors (SE) for mixed model analyses testing the effects of life-history tactics, fertilisation group, and sex on the fork length (cm), weight (g), and condition trajectories of brown trout classed as smolts (migratory) or mature (non-migratory) over 18 months of experimental tank rearing. Measurements were taken at key periods denoted by “time”. Effects were contrasted against mature female fish from the Nov-Dec fertilisation group. Responses are z-standardised and statistical significance was assessed at  $p < 0.05$ .

| Effect                              | Estimate | SE   | t-value | p-value |
|-------------------------------------|----------|------|---------|---------|
| <i>Mixed model for length</i>       |          |      |         |         |
| Intercept (Female, Nov-Dec, Mature) | 0.18     | 0.06 | 3.06    | 0.002   |
| Time                                | 0.74     | 0.01 | 52.94   | < 0.001 |
| Time <sup>2</sup>                   | -0.23    | 0.02 | -13.91  | < 0.001 |
| Life-history: Smolt                 | -0.48    | 0.12 | -3.93   | < 0.001 |
| Sex: Male                           | -0.07    | 0.07 | -1.06   | 0.290   |
| Fertilisation group: Nov            | 0.33     | 0.07 | 4.48    | < 0.001 |
| <i>Mixed model for mass</i>         |          |      |         |         |
| Intercept (Female, Nov-Dec, Mature) | 0.16     | 0.08 | 2.18    | 0.029   |
| Time                                | 0.64     | 0.02 | 32.27   | < 0.001 |
| Time <sup>2</sup>                   | -0.13    | 0.02 | -5.95   | < 0.001 |
| Life-history: Smolt                 | -0.54    | 0.15 | -3.47   | 0.001   |
| Sex: Male                           | -0.10    | 0.08 | -1.19   | 0.234   |
| Fertilisation group: Nov            | 0.19     | 0.09 | 2.11    | 0.036   |
| Time × Smolt                        | -0.20    | 0.06 | -3.10   | 0.002   |
| <i>Mixed model for condition</i>    |          |      |         |         |
| Intercept (Female, Nov-Dec, Mature) | 0.42     | 0.09 | 4.81    | < 0.001 |
| Time                                | -0.32    | 0.03 | -11.99  | < 0.001 |
| Time <sup>2</sup>                   | -0.16    | 0.03 | -4.63   | < 0.001 |
| Life-history: Smolt                 | -0.80    | 0.16 | -4.87   | < 0.001 |
| Sex: Male                           | 0.02     | 0.09 | 0.18    | 0.859   |
| Fertilisation group: Nov            | -0.32    | 0.10 | -3.37   | 0.001   |
| Time × Smolt                        | -0.24    | 0.09 | -2.53   | 0.012   |

**Table S4:** Parameter estimates with associated standard errors (SE) for mixed model analysis testing the effects of food treatment, temperature treatment, fertilisation group, and sex on the final fork length (cm), condition, and whole body lipids (%) of brown trout after 18 months of experimental tank-rearing. Effects were contrasted against female fish from the Nov-Dec fertilisation group in the high food and cool temperature treatment. Responses are z-standardised and statistical significance was assessed at  $p < 0.05$ .

| Effect                                       | Estimate | SE   | t-value | p-value |
|--|----------|------|---------|---------|
| <i>Mixed model for length</i>                |          |      |         |         |
| Intercept (Female, Nov-Dec, High, Cool)      | 0.61     | 0.14 | 4.50    | < 0.001 |
| Sex: Male                                    | 0.02     | 0.07 | 0.37    | 0.714   |
| Fertilisation group: Nov                     | 0.40     | 0.19 | 2.12    | 0.078   |
| Temperature: Warm                            | -0.20    | 0.09 | -2.16   | 0.031   |
| Food: Low                                    | -1.53    | 0.16 | -9.68   | < 0.001 |
| Low food x Warm temperature                  | 0.55     | 0.13 | 4.13    | < 0.001 |
| <i>Mixed model for condition</i>             |          |      |         |         |
| Intercept (Female, Nov-Dec, High, Cool)      | 0.80     | 0.15 | 5.31    | < 0.001 |
| Sex: Male                                    | -0.12    | 0.08 | -1.60   | 0.110   |
| Fertilisation group: Nov                     | -0.45    | 0.20 | -2.25   | 0.065   |
| Temperature: Warm                            | -0.01    | 0.11 | -0.02   | 0.987   |
| Food: Low                                    | -1.36    | 0.18 | -7.65   | < 0.001 |
| Low food x Warm temperature                  | 0.35     | 0.16 | 2.21    | 0.027   |
| <i>Mixed model for whole body lipids (%)</i> |          |      |         |         |
| Intercept (Female, Nov-Dec, High, Cool)      | 0.65     | 0.16 | 4.08    | < 0.001 |
| Sex: Male                                    | -0.17    | 0.13 | -1.36   | 0.177   |
| Fertilisation group: Nov                     | 0.03     | 0.17 | 0.15    | 0.887   |
| Temperature: Warm                            | 0.25     | 0.13 | 1.95    | 0.054   |
| Food: Low                                    | -1.25    | 0.16 | -7.84   | < 0.001 |

**Table S5:** Parameter estimates with associated standard errors (SE) for mixed model analysis testing the effects of life-history tactics, fertilisation group, and sex on the final fork length (cm), condition and whole body lipids (%) of brown trout after 18 months of experimental tank-rearing. Effects were contrasted against mature female fish from the Nov-Dec fertilisation group. Responses are z-standardised and statistical significance was assessed at  $p < 0.05$ .

| Effect                                       | Estimate | SE   | t-value | p-value |
|--|----------|------|---------|---------|
| <i>Mixed model for length</i>                |          |      |         |         |
| Intercept (Mature, Female, Nov-Dec)          | -0.11    | 0.27 | -0.40   | 0.688   |
| Life-history: Smolt                          | -0.28    | 0.13 | -2.08   | 0.038   |
| Sex: Male                                    | 0.03     | 0.07 | -0.35   | 0.726   |
| Fertilisation group: Nov                     | 0.44     | 0.52 | 0.84    | 0.435   |
| <i>Mixed model for condition</i>             |          |      |         |         |
| Intercept (Mature, Female, Nov-Dec)          | 0.27     | 0.23 | 1.17    | 0.241   |
| Life-history: Smolt                          | -0.73    | 0.17 | -4.40   | < 0.001 |
| Sex: Male                                    | -0.14    | 0.09 | -1.51   | 0.133   |
| Fertilisation group: Nov                     | -0.48    | 0.43 | -1.11   | 0.309   |
| <i>Mixed model for whole body lipids (%)</i> |          |      |         |         |
| Intercept (Mature, Female, Nov-Dec)          | -0.23    | 0.35 | -0.67   | 0.504   |
| Life-history: Smolt                          | 0.61     | 0.17 | 3.67    | < 0.001 |
| Sex: Male                                    | -0.17    | 0.14 | -1.26   | 0.209   |
| Fertilisation group: Nov                     | 0.43     | 0.67 | 0.64    | 0.547   |

**Table S6:** Parameter estimates with associated standard errors (SE) for mixed model analysis testing the differences in the gill NKA activity (log transformed) and plasma chloride concentration of brown trout classed as smolts (migratory tactics), mature or unassigned (non-migratory tactics) after 24 saltwater immersion. Effects were contrasted against mature female fish from the Nov-Dec fertilisation group. Statistical significance was assessed at  $p < 0.05$ .

| Effect   | Estimate | SE   | t-value | p-value |
|--|----------|------|---------|---------|
| <i>Mixed model for gill NKA activity (log transformed)</i> |          |      |         |         |
| Intercept  | 0.12     | 0.16 | 0.77    | 0.445   |
| Life-history: Smolt  | 0.81     | 0.11 | 7.41    | < 0.001 |
| Life-history: Unassigned                                   | 0.02     | 0.12 | 0.15    | 0.879   |
| Sex: Male  | -0.09    | 0.08 | -1.12   | 0.266   |
| Fertilisation group: Nov                                   | 0.23     | 0.27 | 0.84    | 0.442   |
| <i>Mixed model for plasma chloride concentration</i>       |          |      |         |         |
| Intercept  | 161.61   | 3.59 | 45.00   | < 0.001 |
| Life-history: Smolt  | -16.24   | 2.92 | -5.56   | < 0.001 |
| Life-history: Unassigned                                   | 13.68    | 3.69 | 3.71    | < 0.001 |
| Sex: Male  | 3.84     | 2.36 | 1.63    | 0.105   |
| Fertilisation group: Nov                                   | -9.04    | 6.45 | -1.40   | 0.210   |



**Table S7:** Parameter estimates with associated standard errors (SE) for mixed model analysis testing the effects of fork length, food treatment, temperature treatment, fertilisation group, and sex on the gill NKA activity and plasma chloride concentration after 24 saltwater immersion of brown trout smolts after 18 months of experimental tank-rearing. Effects were contrasted against female fish from the Nov-Dec fertilisation group in the high food and cool temperature treatment. Responses are z-standardised and statistical significance was assessed at  $p < 0.05$ .

| Effect   | Estimate | SE   | t-value | p-value |
|--|----------|------|---------|---------|
| <i>Mixed model for gill NKA activity</i>             |          |      |         |         |
| Intercept  | 0.57     | 0.59 | 0.98    | 0.342   |
| Fork length  | -0.06    | 0.29 | -0.21   | 0.840   |
| Food: Low  | -0.09    | 0.62 | -0.15   | 0.889   |
| Temperature: Warm                                    | -0.42    | 0.52 | -0.80   | 0.435   |
| Sex: Male  | -0.64    | 0.47 | -1.37   | 0.189   |
| Fertilisation group: Nov                             | -1.12    | 0.69 | -1.64   | 0.200   |
| <i>Mixed model for plasma chloride concentration</i> |          |      |         |         |
| Intercept  | -0.88    | 0.47 | -1.90   | 0.072   |
| Fork length  | -0.43    | 0.20 | -2.18   | 0.041   |
| Food: Low  | 1.08     | 0.53 | 2.06    | 0.109   |
| Temperature: Warm                                    | 0.54     | 0.40 | 1.34    | 0.194   |
| Sex: Male  | -0.34    | 0.30 | -1.13   | 0.271   |
| Fertilisation group: Nov                             | 0.79     | 0.53 | 1.49    | 0.211   |

## Chapter 4

---

# Metabolic traits in brown trout (*Salmo trutta*) vary in response to food restriction and population background

Louise C. Archer<sup>\*1,2</sup>, Stephen A. Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, W. Russell Poole<sup>3</sup>, Patrick Gargan<sup>4</sup>, Philip McGinnity<sup>1,3</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> *School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.*

<sup>2</sup> *Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland.*

<sup>3</sup> *Marine Institute, Furnace, Newport, Co. Mayo, Ireland.*

<sup>4</sup> *Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.*

**\*Correspondence:** Louise Archer (l.archer@umail.ucc.ie)

**Key words:** climate change, partial migration, anadromy, aquatic, brown trout, metabolism, phenotypic flexibility, plasticity

**Contributions:** LA, TR, and PMcG conceived the study. LA, SH, and LH collected data and contributed to experimental design. LA conducted statistical analysis and wrote the manuscript. All authors contributed to interpretation of results and revisions of the manuscript.

## Abstract

Metabolic rates vary hugely within and between populations of the same species, yet we know relatively little about the factors causing this intraspecific variation. Given that metabolic rate determines the energetic cost of life, uncovering these sources of variation is important to understand and forecast population responses to environmental change. Moreover, few studies have examined factors causing intraspecific variation in metabolic flexibility. Here, we explore if, and how, extrinsic environmental conditions and population-specific factors contribute to variation in metabolic traits in brown trout, an iconic species that is capable of extreme variation in migratory life-history tactics. We measured metabolic traits in offspring from two wild populations of brown trout that naturally show divergent migratory tactics (one anadromous i.e. sea-migratory, and one non-anadromous) that had been reared under experimental conditions of long-term food restriction. Both populations showed decreased standard metabolic rates (SMR – baseline energy requirements) under low food conditions. The non-anadromous population had an overall lower SMR relative to the anadromous population, but maximum metabolic rate (MMR), and consequently, aerobic scope (AS), were both higher in the anadromous population, and were higher overall in males compared to females. Intriguingly, the anadromous population had a higher AS under low food conditions compared to optimal food conditions, indicating population-specific effects of food restriction on AS. Collectively, our results suggest the different components of metabolic rate can vary in their response to extrinsic environmental conditions, and can also vary according to intrinsic (i.e. population background/sex) effects. Moreover, populations can further differ in their flexibility of metabolic traits, potentially due to population-specific factors related to life history (*e.g.* migratory tactics). Overall, our study suggests that responses to environmental change may be population specific, but incorporating an understanding of variation in metabolic traits, and their flexibility, will improve our ability to conserve populations.

## Introduction

Metabolic rate represents the fundamental energetic cost of living that underpins organism performance in variable and changing environments. Since metabolism has profound implications for fitness (Pettersen *et al.* 2016, 2018), relatively higher or lower metabolic rates have in turn been linked to variation in fitness components such as growth rates (Auer *et al.* 2015c; Zeng *et al.* 2017), and survival (Bochdansky *et al.* 2005) in ways that often depend on environmental context (Burton *et al.* 2011; Auer *et al.* 2015b, c). The minimum energy expenditure required for tissue maintenance and homeostasis is termed standard metabolic rate (SMR) in ectotherms (basal metabolic rate (BMR) in endotherms within the thermoneutral range). SMR occurs when an organism is inactive, unstressed, and not digesting (Chabot *et al.* 2016). Maximum metabolic rate (MMR) sets the upper bounds of energy expenditure as the highest rate of aerobic metabolism (transport of oxygen and production of ATP) that can be achieved (Norin and Metcalfe 2019). Together, SMR and MMR define an organism's aerobic scope (AS), a trait that determines the amount of energy that can be directed towards key functions including digestion, movement, growth, and reproduction through increased metabolism, once baseline energy requirements (i.e. SMR) are met (Guderley and Pörtner 2010).

Large variation in both SMR/BMR and MMR (and consequently AS) exists among species, populations, and individuals (Burton *et al.* 2011; Konarzewski and Książek 2013; Hillman *et al.* 2013; Norin and Clark 2016), with variation linked to differences in lifestyle (Killen *et al.* 2010), geographic distribution (Angilletta 2001; Naya and Bozinovic 2012), thermal regime (Álvarez *et al.* 2006; Eliason *et al.* 2011; Sandblom *et al.* 2016), and behavioural differences (Metcalfe *et al.* 2016). In aquatic ectotherms, factors related to life-history tactics appear to underpin many inter-individual and intra-individual differences in metabolic traits. Along with a 16-fold variation in MMR reported across fish species that occupy different ecological niches (Norin and Clark 2016), metabolic rates can still show c. 3-fold inter-individual variation after accounting for age and size differences (Metcalfe *et al.* 2016). Such variation

likely arises because the optimal combination of the various components of metabolic phenotype is context-specific (Auer *et al.* 2015b, c), or because populations (or types of individuals within populations) experience different  
 65 selection pressures due to life-history differences, or extrinsic/intrinsic factors. For example, sockeye salmon *Oncorhynchus nerka* populations that undertake longer, or more challenging migrations have higher AS (Eliason *et al.* 2011), and higher metabolic rates have also been documented in males *versus* females, *e.g.* higher AS in male pink salmon *O. gorbuscha* (Clark *et al.* 2011).

70 On top of variation *per se*, patterns of covariation in metabolic phenotypes can also be different across and within species. SMR and MMR have been proposed to be tightly linked because of the “increased intake hypothesis”, whereby a high SMR requires investment in metabolic machinery that also facilitates a high MMR, with associated fitness benefits (Biro and Stamps 2010; Burton *et al.*  
 75 2011). While SMR and MMR generally do appear to be correlated within species (Auer *et al.* 2017), the traits can vary in their response to different environmental factors, and the coupling of metabolic traits can be context dependent (Killen *et al.* 2013; Norin *et al.* 2016). Moreover, a decoupling of SMR and MMR can occur over time because each is under individual selection  
 80 pressures (Norin and Metcalfe 2019), which often operate in parallel but may also act independently (*e.g.* Wone *et al.* 2015; Barceló *et al.* 2016). Thus, even if SMR and MMR are somewhat functionally linked, ecologically significant variation in overarching AS can arise due to differences in the sensitivities of each metabolic trait to environmental conditions. Thus, within-individual  
 85 variation in response to variation in the environment may account to some extent for intraspecific patterns of variation and covariation in metabolic traits.

The ability of a single genotype to display different physiological, morphological, or behavioural phenotypes in response to variation in environmental factors is called phenotypic plasticity. Phenotypic ‘flexibility’ has  
 90 been defined as a particular type of plasticity in which within-individual changes are reversible (Piersma and Drent 2003), as distinct from developmental plasticity, where phenotypic responses to early developmental

conditions remain relatively fixed for the rest of life (West-Eberhard 2003). Phenotypic flexibility is an important attribute that facilitates individuals in coping with changing conditions (Seebacher *et al.* 2015), with life-history consequences that may scale up to affect higher levels of organisation, for example, population persistence, community stability, and ecosystem processes (Bolnick *et al.* 2011). Flexibility in metabolic rate is likely to be an important component here, and indeed, there is a growing body of evidence supporting metabolic plasticity as a widespread response to environmental change (Hofmann and Todgham 2010). Factors including temperature (Seebacher *et al.* 2015; Sandblom *et al.* 2016), food availability (Auer *et al.* 2015c, 2016a; Zeng *et al.* 2018), food quality (Naya *et al.* 2007), oxygen availability (Hochachka *et al.* 1996; Norin *et al.* 2016) and salinity (Allan *et al.* 2006) have all been shown to induce short term and longer term (i.e. acclimation) changes in metabolic rates of organisms.

In ectotherms, SMR generally appears to be more flexible in the extent of its response to extrinsic factors than MMR (Norin and Metcalfe 2019). For example, increased temperatures were shown to cause reduction of SMR in European perch *Perca fluviatilis*, a thermal compensation response that was not apparent in MMR (Sandblom *et al.* 2016). Similar flexibility in BMR relative to MMR (or cold-induced maximum aerobic metabolism) has been demonstrated in endotherms in response to temperature (Nespolo *et al.* 2001; van de Ven *et al.* 2013; Dubois *et al.* 2016). Food availability has also been frequently shown to induce flexibility in SMR (and BMR) (Naya *et al.* 2007; Auer *et al.* 2015c, 2016a; Langer *et al.* 2018), suggesting reductions in baseline metabolism, rather than MMR, tends to underpin overall metabolic flexibility in response to food restriction (Zeng *et al.* 2018).

Although there is a significant number of studies detailing inter and intra specific variation in metabolic responses, we know considerably less about factors giving rise to differences in metabolic flexibility (Norin and Metcalfe 2019). Variation in metabolic rate flexibility between populations has been described primarily as changes in SMR or BMR, and particularly in response to

distribution or temperature factors *e.g.* cane toads *Rhinela marina* at high  
 125 latitudes show more plastic resting metabolic rates in response to temperature  
 than their counterparts at low latitudes (Winwood-Smith *et al.* 2015; McCann  
*et al.* 2018). Similarly, rufous-collared sparrow *Zonotrichia capensis* populations  
 from seasonally-variable or temperate environments show more flexible BMRs  
 in response to temperature than those from arid desert systems, though desert  
 130 populations conversely showed more BMR flexibility at low food conditions, a  
 finding that highlights the context-dependency of optimal metabolic  
 phenotypes (Cavieres and Sabat 2008; Maldonado *et al.* 2012). Given that the  
 optimal metabolic phenotype in a given context can show considerable  
 variability depending on the population background, incorporating population-  
 135 specific (or life history) factors into the investigation of metabolic variation and  
 flexibility is likely to have important implications for managing and conserving  
 species experiencing environmental change, yet few studies have addressed  
 this.

Salmonine fishes (salmons, trouts, and charrs) represent an excellent group to  
 140 study variation in metabolic phenotypes. As obligate freshwater spawning  
 species, salmonines display a multitude of life-history strategies that  
 incorporate a wide variety of migratory tactics (Klemetsen *et al.* 2003). Some  
 individuals remain resident in natal freshwaters for their entire life cycles,  
 whilst others migrate to more productive feeding grounds such as larger rivers  
 145 and lakes (termed “potomodromy”), or even undertake dramatic migrations to  
 the sea (termed “anadromy”) (Ferguson *et al.* 2019). Migration generally  
 facilitates high levels of growth in the new habitat, with migrants typically  
 returning to spawn in freshwater at larger sizes than non-migratory “residents”.  
 Facultative migration – where individuals can adopt either a migratory or a non-  
 150 migratory lifestyle - is common in salmonines, and populations can be primarily  
 resident, migratory, or comprise a mix of both tactics (Chapman *et al.* 2012).  
 Such alternative migratory phenotypes can be understood using the framework  
 of the “environmentally cued threshold model” (Tomkins and Hazel 2007;  
 Piche *et al.* 2008; Pulido 2011; Buoro *et al.* 2012), whereby tactic frequencies are

155 controlled by the relationship between an environmentally-sensitive trait (*e.g.*  
physiological condition or energetic status) and a genetically-variable  
threshold. Migration is triggered depending on whether or not an individual's  
"status" trait exceeds the threshold condition for residency. Energetic limitation  
in natal freshwaters is proposed to be a strong determinant of migration  
160 (Forseth *et al.* 1999). As such, variation in migratory tactics is likely to be linked  
to variation in metabolic rates *e.g.* steelhead trout *O. mykiss* that matured in  
freshwater *in lieu* of migrating tended to have lower SMR values (Sloat and  
Reeves 2014); and juvenile Atlantic salmon *Salmo salar* with higher SMR early  
in life were more likely to subsequently migrate (McCarthy 2000).

165 Moreover, once the migration decision has been made, different energetic  
demands are associated with the alternative tactics. For example, migrants must  
have sufficiently high aerobic capacity (*i.e.* MMR/ AS) to sustain swimming  
performance during the migration itself (which can cover distances of tens to  
several thousand kilometres), and to facilitate high growth in the new  
170 environment. In contrast, residents typically have lower energetic  
requirements, but must cope with access to fewer food resources in the  
freshwater environment (relative to lacustrine or marine resources) (Gross *et al.*  
1988). Populations that display predominantly one migratory phenotype over  
another are thus likely to experience different selection pressures on metabolic  
175 traits, whereby a migratory life-style might favour increases in the upper bounds  
of metabolisms (MMR), and residency might promote decreases in baseline  
energetic requirements (SMR), each with implications for overall AS and energy  
balance. This has some empirical support, for example, migratory three-spined  
stickleback *Gasterosteus aculeatus* had higher SMR, active metabolic rate, and  
180 AS than non-migratory morphs (Tudorache *et al.* 2007), and anadromous (sea-  
migratory) juvenile Atlantic salmon *S. salar* had higher SMR than non-migrants  
(Seppänen *et al.* 2010). Less is known about how differences in migratory  
lifestyle might interact with environmental conditions to cause variation in  
metabolic traits.



185 Here, we explore the effects of extrinsic (food supply) *versus* intrinsic (population/sex) factors on metabolic rates in experimental F1 offspring derived from two wild populations of brown trout that differ in their migratory tactics. Specifically, we aimed to (i) assess how a period of long-term food restriction alters SMR, MMR and AS (ii) test if populations that naturally vary in their life  
190 histories also vary in SMR, MMR, and AS and (iii) explore if populations show variation in their metabolic responses to conditions of food restriction. We expected that food restriction would overall have a greater effect on SMR compared to MMR. We also expected that offspring derived from the naturally anadromous (i.e. migratory) population would show relatively higher MMR or  
195 AS, and those from the non-anadromous population would show relatively lower SMR, but the populations might vary in their ability to adjust their metabolic traits.

## Methods

### *Study populations and fish rearing*

200 Brown trout brood stock from two wild populations were caught by seine netting in November 2015 in the Erriff (53° 37' 0.00" N: 09° 40' 17.10" W) and the Burrishoole (53° 57' N: 09° 35' W) catchments in the west of Ireland (Figure S1). Erriff brood stock were caught in the Tawnyard Lough, an upland lake of 56 ha which is fed primarily by the Glendavoch river and a number of smaller  
205 tributaries. The Tawnyard Lough population spawn mainly in the Glendavoch River, and move downstream as fry or parr to Tawnyard Lough (a distance of a few hundred metres to kilometres, depending on where spawning occurred). Tawnyard Lough produces a large run of out-migrating anadromous juveniles (smolts), with annual estimates of 500 to 3000 smolts enumerated at the  
210 outflow of the Lough over the last 30 years (Gargan *et al.* 2016). An unknown proportion of the population remain within the lake, and undergo several years of freshwater growth before returning to the natal stream to spawn, with local expertise indicating that the Tawnyard population in general has a strong anadromous component (broadly estimated as 50 – 60% of population  
215 expressing anadromy) (P. Gargan, *pers comm*).

Burrishoole brood stock were caught in Lough Bunaveela (46 ha) in the headwaters of the catchment. A population of non-anadromous trout remain resident in Lough Bunaveela for most of their lifecycle, undertaking only short-distance, directed movements (10 – 100s of metres) between the lake and inflowing/outflowing spawning streams. Although the anadromous life history is present in the larger Burrishoole catchment, the development of aquaculture in Clew Bay is believed to have caused the anadromous trout run to decline severely in Burrishoole in the late 1980s. Despite Bunaveela spawning streams being accessible to anadromous fish, there is no evidence that the Bunaveela population has ever produced anadromous fish, either historically, or recently (Poole *et al.* 2007; Magee 2017). In summary, we consider offspring derived from the Tawnyard brood stock to have a strong anadromous background (hereafter termed the “anadromous background population”), and offspring from the Bunaveela brood stock to have no recent anadromous background (termed the “non-anadromous background population”).

See Archer *et al.* (2019) for detailed description of crossing, fertilisation, and rearing procedures, which we describe here in brief. Each ripe female was mated to two males from the same source population. Fertilised eggs were incubated at a hatchery in the Burrishoole catchment. Post-hatching, fry were transferred to a rearing facility at University College Cork (Aquaculture and Fisheries Development Centre). Here, fry were initially held in two 100L growth tanks on a recirculating aquaculture system (RAS) maintained at natural temperature regime for the west of Ireland, and moved to 520L tanks on a larger RAS to facilitate growth in December 2016. Populations were reared separately for the entirety of the study to prevent emergence of dominance hierarchies. Fry were fed *ad libitum* with commercially available trout pellets (Skretting Ltd., Norway) until experimental food treatments (see below) began in September 2016. During the experimental phase, a programmed lighting system of LED lights above each tank mimicked the photoperiod of the source catchments. Water in the RAS was treated with mechanical filtration, bio filtration and UV skimming, and water quality (checked weekly) consistently remained within acceptable

levels for fish health. Great care was taken to ensure that all measured variables other than feeding regime (fish densities, temperature, photoperiod, lux, flow rates) were constant across the tanks.

#### 250 *Food restriction treatments*

Fish in this study experienced experimental food restriction treatments from September 2016 to June 2018. The study, and all associated procedures, were carried out with ethical approval from Health Products Regulatory Authority (HPRA) Ireland, under HPRA project license AE19130/P034, and individual  
255 licenses AE19130/1087, AE19130/1200, AE19130/1201 and AE19130/1202 with all fish humanely euthanized under licence in June 2018 as part of a parallel experiment (Archer *et al.* 2019).

To explore the effects of extrinsic environment (food restriction) and intrinsic factors on metabolism, juvenile brown trout from each population were  
260 randomly allocated one of four food treatments in September 2016 ( $n = 90$  per feeding treatment per population, at the beginning of the experimental phase). The food treatments were as follows: (i) High-High food: fish fed recommended daily pellet rations for optimal growth, calculated as a percentage of their body weight and adjusted for seasonally-changing temperatures (Skretting Ltd,  
265 Norway); (ii) Low-Low: fish fed 25% of recommended optimal rations. (iii) High-Low: fish switched from optimal rations to 25% optimal ration (i.e. from High to Low) in June 2017; and (iv) Low-High: fish switched from 25% of optimal rations to 100% optimal rations in June 2017 (i.e. from Low to High). The reductions to 25% of optimal food rations took place gradually over a four-  
270 week period to minimise stress. Within each tank, absolute rations were adjusted on a monthly basis to account for changes in body mass and temperature.

#### *Measurement of metabolic traits*

Eight to twelve individuals of each population in each food treatment were  
275 measured for SMR and MMR in February 2018 in a controlled-temperature (CT) chamber at 8 °C (mean temperature  $7.99 \text{ °C} \pm 0.26 \text{ SD}$ , matching the natural temperature in the wild for these populations).

*Measurement of MMR*

Whole-animal oxygen consumption ( $MO_2$ ) in animals operating at their maximum aerobic metabolic rate was used as proxy for MMR (Norin and Metcalfe 2019) following best practices outlined in Norin and Clark (2016). We used an exhaustive chase protocol following Norin and Clark (2016) to elicit MMR in the same individuals that we measured for SMR. Prior to SMR measurements, each individual fish due to be measured for SMR that day was first placed in an aerated 50L tank and manually chased by hand until exhaustion, determined to occur when the fish were unresponsive (i.e. did not elicit burst swimming) to tactile stimulus (typically after 2 to 3 minutes of sustained chasing). Once exhausted, the fish was immediately transferred to a respirometry chamber in the same system used to measure SMR, the chamber was sealed, and oxygen decline within the closed chamber loop (recirculation pump operational) was recorded for a 60s measurement period. The time taken to transfer fish to chambers and begin recording oxygen measurements never exceeded 20s, ensuring that minimal recovery from the exhaustive chase procedure occurred before recording oxygen consumption.

*Measurement of SMR*

The SMR of individual fish was determined overnight in a darkened CT chamber using intermittent-flow respirometry, following best practices outlined in Svendsen *et al.* (2016). The respirometry system consisted of four acrylic respirometry chambers (1200 ml) (Loligo Systems, Viborg, Denmark), submerged in a water bath, flushed with de-chlorinated water bubbled to 100% oxygen saturation by an air stone. PVC tubing (10mm diameter, non-permeable to oxygen) connected each individual chamber to two pumps (Eheim Ltd., Deizisau, Germany): the “flush” pump flushed fully oxygenated water through the chambers. A second “recirculation” pump recirculated water in a closed loop through the chamber, whereby water exiting the chamber was passed through a 10mm flow through oxygen cell (PreSens Ltd., Regensburg, Germany) that continually measured dissolved oxygen concentration, before being recirculated back to the same respirometry chamber via PVC tubing. Thus,

individual oxygen consumption ( $MO_2$ , used as a proxy for SMR in fasted, non-  
 310 active animals) was measured in repeated cycles that consisted of a flush period  
 (flush pump operational) and a measurement period (recirculation pump  
 operational) where oxygen level in each chamber was recorded at one second  
 intervals to estimate oxygen decline (i.e. oxygen uptake). Each cycle consisted  
 of 330s of flushing, and a measurement period of 200 - 300s (to ensure  
 315 sufficient  $O_2$  depletion for calculating  $MO_2$  in different-sized fish). We also  
 allowed a 30s buffer period before recording oxygen uptake once the flush  
 pump was switched off, to allow the chamber water and flush water to mix  
 completely and reach an equilibrium oxygen saturation .

Fish were fasted for 28 h prior to being placed into individual respirometry  
 320 chambers to ensure individuals were in a post absorptive state (Cutts *et al.*  
 2002). Fish entered the chambers between 11:00 and 12:00 each day, and were  
 left to acclimatise for five hours, with chambers continually flushed with  
 oxygen-saturated water during this acclimation period. SMR measurements  
 began between 16:00 and 17:00, and ended between 09:00 and 10:00 the  
 325 following morning, allowing for a minimum of 100 measurements of oxygen  
 uptake per individual. Fish were not disturbed during this ~20-hr SMR  
 measurement period. Once SMR measurements had finished, each fish was  
 removed from the chamber, lightly anaesthetised with MS-222, blotted dry, and  
 mass and fork length were recorded. Each fish was then given an individual  
 330 identifier tag using unique colour combinations of visible implant elastomer  
 tags (Northwest Marine Technology Ltd., USA). To limit bacterial growth in the  
 system, the entire respirometry set-up was rinsed with bleach after each  
 overnight SMR respirometry trial. We also measured background (i.e.,  
 bacterial) respiration rates in each chamber on a daily basis by recording oxygen  
 335 decline in empty chambers for one measurement cycle before fish entered the  
 respirometry system, and again for one measurement cycle the following  
 morning (once the respirometry measurements had ended and fish were  
 removed from the chambers).

*Determination of sex*

340 To determine sex, we euthanised fish that had been measured for SMR and MMR via an overdose of MS-222 in April 2018 (approximately 2 months after respirometry measurements due to involvement in an ongoing parallel study, see Archer *et al.* (2019)). If sex could not be determined anatomically, genotypic sex was later assigned using a microsatellite sex marker (P. Prodöhl, 345 *unpublished*). We were unable to re-identify six individuals due to tag loss, leaving  $n = 55$  fish successfully assigned for sex. Sex ratios were approximately similar across food treatment groups and across population backgrounds.

*Statistical analysis*

To estimate SMR ( $\text{mg O}_2 \text{ h}^{-1}$ ), we first calculated  $MO_2$  values for each repeated 350 measurement of oxygen uptake recorded during the overnight SMR respirometry trials.  $MO_2$  ( $\text{mg O}_2 \text{ h}^{-1}$ ) was calculated as the most consistent linear decline in oxygen recorded during each measurement cycle, estimated by rolling regression in the *respR* package in R (Harianto and Carey 2019). All measurements of  $MO_2$  were visually inspected to assess regression fit, and only 355  $MO_2$  values with an acceptable fit (associated  $R^2$  values  $> 0.90$ , unless a clear linear trend was determined upon visual inspection of fit) were included in subsequent SMR calculations. To account for any background respiration included in these  $MO_2$  values, we estimated background respiration by calculating  $MO_2$  values for the oxygen uptake measurements in empty 360 chambers, both before and after each overnight SMR respirometry trial (as described above). Because background  $MO_2$  rates were assumed to increase linearly through time over the course of the experiment (due to bacterial growth), we allowed for a dynamic background correction value (i.e. that increased overnight), calculated as:

$$365 \quad MO_{2\_bg} = bg_0 + (t \times bg)$$

Where  $MO_{2\_bg}$  is background  $MO_2$ , at a given measurement time point  $t$ , the time elapsed since initiating overnight SMR measurements,  $bg_0$  and  $bg$  are parameters (the intercept and slope respectively) estimated from the matrix

regression of the background oxygen uptake before, and background oxygen  
 370 uptake after, as a function of time elapsed. We then used  $MO_{2\_bg}$  to account for  
 background respiration by subtracting  $MO_{2\_bg}$  from each value of  $MO_2$  as  
 calculated for an individual fish at successive time points during the overnight  
 SMR respirometry trials.  $MO_{2\_bg}$  never exceeded more than 2% of total  $MO_2$  in  
 all cases, confirming that background respiration rates remained low  
 375 throughout the study.

SMR for each individual fish was calculated by taking the mean of the lowest  
 10<sup>th</sup> percentile of background-corrected  $MO_2$  values recorded over the 20-h  
 SMR measurement period, then excluding outliers (values more than two  
 standard deviations from this mean).

380 We estimated individual MMR ( $\text{mg O}_2 \text{ h}^{-1}$ ) using the *respR* package (Harianto  
 and Carey 2019) by calculating  $MO_2$  as the linear decline in oxygen in each  
 individual respirometry chamber in the 60s measurement period immediately  
 after the exhaustive chase protocol (i.e. extracting slopes from the linear  
 regression of oxygen concentration against time over a 60s period). Oxygen  
 385 sensor probe and equipment malfunctions resulted in respirometry  
 measurements for 6 fish being discarded, leaving a total of  $n = 61$  individuals  
 measured for SMR and MMR. Absolute aerobic scope (AS) for each fish was  
 calculated as the difference ( $\text{mg O}_2 \text{ h}^{-1}$ ) between MMR and SMR.

Since metabolic responses to food restriction are well-documented to be  
 390 reversible in salmonines (metabolic rates are restored to pre food restriction  
 levels once standard food rations are reinstated (O'Connor *et al.* 2000)), we first  
 assessed if any potential metabolic responses to food restriction had been  
 reversed/offset in the Low-High groups by February 2018 (when we measured  
 metabolic traits). No differences in SMR (ANOVA:  $\chi^2 = 0.23$ ,  $df = 1$ ,  $P = 0.633$ ),  
 395 MMR ( $\chi^2 = 0.40$ ,  $df = 1$ ,  $P = 0.528$ ), or AS ( $\chi^2 = 0.51$ ,  $df = 1$ ,  $P = 0.476$ ) existed  
 between the High-High food treatment and Low-High treatment. Similarly, we  
 tested whether potential metabolic responses to food restriction were affected  
 by the length of the food restriction period i.e. did Low-Low (17 months

restriction) differ from High-Low (7 months restriction). No differences existed  
 400 in SMR ( $\chi^2 = 0.04$ ,  $df = 1$ ,  $P = 0.836$ ), MMR ( $\chi^2 = 0.44$ ,  $df = 1$ ,  $P = 0.509$ ), or AS  
 ( $\chi^2 = 0.47$ ,  $df = 1$ ,  $P = 0.494$ ) between the Low-Low food treatment and the High-  
 Low treatment. Since our primary interest was simply in the overarching effects  
 of food restriction on metabolism (and not the effects of switching food  
 treatments *per se*), we combined the High-High and Low-High treatments into  
 405 a single “High Food” treatment group, and combined the Low-Low and the  
 High-Low treatments into a single “Low Food” treatment group. We present  
 analyses using the “High Food” and “Low Food” groups here, with the *caveat*  
 that “High” or “Low” refers specifically to the food treatment experienced in the  
 ~7-month period prior to metabolic measurements in year 2 of life (a timescale  
 410 over which metabolic rates have been shown to be consistent in salmonines  
 (Seppänen *et al.* 2010)). Moreover, pilot SMR measurements collected from our  
 populations in April and May 2017 (following similar respirometry protocols to  
 those described above) showed similar effects of High/Low food treatments as  
 the results described below (which focus solely on the 2018 measurements of  
 415 SMR, MMR, and AS). This indicates that responses to food treatments were: (i)  
 consistent though time (or at least between years); and (ii) most likely as a result  
 of phenotypic plasticity rather than random variation.

To avoid the pitfalls associated with solely using *P*-values (Halsey *et al.* 2015;  
 Halsey 2019), we first tested for factors influencing mass-independent measures  
 420 of SMR, MMR, and AS through estimation statistics (i.e. estimating effect sizes)  
 using the *dabestr* package (Ho *et al.* 2019). We used the residuals of the linear  
 relationships between  $\log_{10}$  body mass, and SMR, MMR, and AS (all  $\log_{10}$   
 transformed) to correct for body size in these analyses. Residuals generated  
 from these analyses (rSMR, rMMR, and rAS) gave mass-independent estimates  
 425 of metabolic rates (individuals with positive residuals have a higher than  
 expected metabolic rate for a given fish size, whereas negative residuals indicate  
 a lower than expected rate). Effect sizes for mean differences in rSMR, rMMR,  
 and rAS were computed for all pairwise comparisons between all levels of food  
 treatment (high or low) and population background (anadromous or non-



anadromous) factors, and 95% confidence intervals (CIs) were constructed by bootstrapped resampling for 5,000 resamples. An additional set of analyses tested for sex-based differences in metabolic traits, whereby we similarly estimated effect sizes for pairwise comparisons of rSMR, rMMR, and rAS between the two levels of sex. Similar analyses were also run using an alternative ANCOVA approach, which tested for variation in the relationships between body mass and SMR, MMR, and AS according to population, food treatment, and sex factors using general linear models (GLMs) (see Supporting Information). The results (shown in the SI) were qualitatively similar, suggesting that the findings based on estimation statistics that we present here are robust.

Finally, to explore whether population background and food treatments affected the size-independent relationships between different aspects of metabolism, we created three GLMs (normal errors). The first GLM included rMMR as a response variable, and rSMR, food treatment and population treatment as explanatory variables, interactions between rSMR and food treatment, and between rSMR and population, along with a three-way interaction term ( $\text{rSMR} \times \text{food} \times \text{population}$ ). The second GLM included rAS as the response variable, and similarly included rSMR, food treatment, population treatment, and interaction terms for  $\text{rSMR} \times \text{food}$ ,  $\text{rSMR} \times \text{population}$  and  $\text{rSMR} \times \text{food} \times \text{population}$ . We constructed a third GLM with rAS as the response variable, and rMMR, food treatment, population treatment as predictors, along with interaction terms for  $\text{rMMR} \times \text{food}$ ,  $\text{rMMR} \times \text{population}$  and  $\text{rMMR} \times \text{food} \times \text{population}$ .

For the estimation statistics approach, we considered an estimated difference in means between groups to exist (i.e. was significant) if the 95% CI of the effect size did not include zero. We used likelihood ratio tests (LRT) to assess statistical significance of predictor variables for all of the GLM models at a 5% alpha level, and non-significant interaction terms were excluded to interpret main effects. All analysis was carried out in R version 3.6.0 (R Core Team 2019) and all models were checked against assumptions of the given model

(independence, non-normality of residuals, heteroscedasticity and multicollinearity).

## Results


### *Effects of population and food restriction on metabolic rate*

Overall, whole-animal SMR, MMR and AS varied with food treatments and across populations, with higher mean SMR, MMR and AS in the anadromous population (see Table 1 for mean values and SD by population and treatment combinations).

**Table 1:** Mean values and associated standard deviations (SD) for the length (mm), mass (g), standard metabolic rate (SMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), maximum metabolic rate (MMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), and aerobic scope (AS) ( $\text{mg O}_2 \text{ hr}^{-1}$ ) of brown trout offspring derived from two wild populations (AB = anadromous background population, non-AB = non anadromous background population). Offspring experimentally reared under two food treatments (High = optimal rations, Low = 25% of optimal rations).

| Food,<br>Population | Length<br>(mean $\pm$ SD) | Mass<br>(mean $\pm$ SD) | SMR<br>(mean $\pm$ SD) | MMR<br>(mean $\pm$ SD) | AS<br>(mean $\pm$ SD) |
|---------------------|---------------------------|-------------------------|------------------------|------------------------|-----------------------|
| High, AB            | 201.6 $\pm$ 18.8          | 109.65 $\pm$ 30.40      | 5.39 $\pm$ 1.71        | 40.52 $\pm$ 14.47      | 35.13 $\pm$ 13.38     |
| Low, AB             | 200.6 $\pm$ 9.2           | 101.76 $\pm$ 16.38      | 4.51 $\pm$ 0.71        | 43.04 $\pm$ 8.94       | 38.52 $\pm$ 8.79      |
| High, Non-AB        | 199.7 $\pm$ 15.5          | 110.83 $\pm$ 27.59      | 5.18 $\pm$ 1.49        | 31.87 $\pm$ 10.79      | 26.69 $\pm$ 9.64      |
| Low, Non-AB         | 199.0 $\pm$ 17.7          | 101.37 $\pm$ 27.30      | 4.21 $\pm$ 1.54        | 30.31 $\pm$ 11.73      | 26.09 $\pm$ 11.13     |

Fish from the low food treatments had lower mass-independent SMR (lower rSMR) than those in the high food treatment (Figure 1A), and this difference in mean rSMR was evident in both populations (Figure 1A, Table 2). Fish from the anadromous background population had a marginally higher rSMR than those from the non-anadromous background population in both food treatments, however the 95% CIs for the mean difference in rSMR between populations overlapped zero (Figure 1B, Table 2).



**Figure 1:** (A) Residual standard metabolic rate (rSMR) values (body mass corrected) for brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental food treatments: optimal food rations (High); and 25% of optimal daily rations (Low). Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (B) Cumming estimation plots for each population background and food treatment combination, with effect sizes shown as black dots (i.e. the mean differences in rSMR among the groups), the distributions (shaded curves) and 95% confidence intervals (back bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).

**Table 2:** Effect sizes ( $\Delta$ ) and associated 95% confidence intervals (CIs) for differences in mean residual standard metabolic rate (rSMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), residual maximum metabolic rate (rMMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), and residual aerobic scope (rAS) ( $\text{mg O}_2 \text{ hr}^{-1}$ ) of brown trout offspring derived from two wild populations (AB = anadromous background population, non-AB = non anadromous background population), exposed to two food treatments (High = optimal rations, Low = 25% of optimal rations). CIs were constructed by non-parametric bootstrap resampling (5,000 resamples).

| Mean difference ( $\Delta$ )    | $\Delta$ rSMR<br>(95% CI)  | $\Delta$ rMMR<br>(95% CI)  | $\Delta$ rAS<br>(95% CI)   |
|---------------------------------|----------------------------|----------------------------|----------------------------|
| <i>Low AB – High AB</i>         | -0.039<br>(-0.067; -0.007) | 0.060<br>(-0.003; 0.129)   | 0.075<br>(0.004; 0.155)    |
| <i>Low non-AB – High non-AB</i> | -0.054<br>(-0.109; -0.002) | 0.006<br>(-0.066; 0.078)   | 0.013<br>(-0.071; 0.101)   |
| <i>High AB – High non-AB</i>    | 0.026<br>(-0.014; 0.061)   | 0.111<br>(0.039; 0.175)    | 0.125<br>(0.039; 0.200)    |
| <i>Low AB – Low non-AB</i>      | 0.040<br>(-0.006; 0.089)   | 0.165<br>(0.091; 0.236)    | 0.187<br>(0.101; 0.271)    |
| <i>Female – Male</i>            | -0.019<br>(-0.051; 0.020)  | -0.155<br>(-0.203; -0.104) | -0.178<br>(-0.236; -0.118) |


495

There was no effect of food on rMMR in either population (95% CIs for the mean difference in rMMR overlapped zero, Figure 2A, B). Fish from the anadromous background population had a higher rMMR than those from the non-anadromous background population in both food treatments (Figure 2A, B, Table 2).


500

Similarly, fish from the anadromous background population had a higher rAS than the non-anadromous background population under both food treatments (Figure 3A, B, Table 2). We also detected population-specific effects of food treatment on rAS, whereby fish in the anadromous population in the low food treatment had a marginally higher rAS than those in the high food treatment (Figure 3B, Table 2). This food treatment effect on rAS was not apparent in the non-anadromous population (Figure 3B).

505



**Figure 2:** (A) Residual maximum metabolic rate (rMMR) values (body mass corrected) for brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental food treatments: optimal food rations (High); and 25% of optimal daily rations (Low). Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (B) Cumming estimation plots for each population background and food treatment combination, with effect sizes shown as black dots (i.e. the mean differences in rMMR among the groups), the distributions (shaded curves) and 95% confidence intervals (back bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).



**Figure 3:** (A) Residual aerobic scope (rAS) values (body mass corrected) for brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental food treatments: optimal food rations (High); and 25% of optimal daily rations (Low). Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (B) Cumming estimation plots for each population background and food treatment combination, with effect sizes shown as black dots (i.e. the mean differences in rAS among the groups), the distributions (shaded curves) and 95% confidence intervals (black bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).

*Coupling of metabolic traits*


When considering size-independent effects of rSMR on rMMR, the interaction terms for rSMR  $\times$  food  $\times$  population ( $\chi^2 = 0.23$ ,  $df = 1$ ,  $P = 0.633$ ), rSMR  $\times$  food  
 515 ( $\chi^2 = 1.45$ ,  $df = 1$ ,  $P = 0.229$ ), rSMR  $\times$  population ( $\chi^2 = 0.90$ ,  $df = 1$ ,  $P = 0.344$ ),  
 and food  $\times$  population ( $\chi^2 = 2.66$ ,  $df = 1$ ,  $P = 0.103$ ) were all non-significant. The  
 main effects of rSMR ( $\chi^2 = 0.66$ ,  $df = 1$ ,  $P = 0.417$ ) and food ( $\chi^2 = 2.21$ ,  $df = 1$ ,  $P =$   
 0.137) were also non-significant. We detected a significant main effect of  
 population background ( $\chi^2 = 22.35$ ,  $df = 1$ ,  $P < 0.001$ ), whereby the anadromous  
 520 background population had a higher rMMR for a given rSMR (Figure 4A).

Effects of rSMR on rAS were similar, where we detected non-significant effects  
 of rSMR  $\times$  food  $\times$  population ( $\chi^2 = 0.24$ ,  $df = 1$ ,  $P = 0.624$ ), rSMR  $\times$  food ( $\chi^2 =$   
 1.39,  $df = 1$ ,  $P = 0.239$ ), rSMR  $\times$  population ( $\chi^2 = 0.92$ ,  $df = 1$ ,  $P = 0.337$ ), and food  
 $\times$  population ( $\chi^2 = 2.69$ ,  $df = 1$ ,  $P = 0.101$ ), and non-significant main effects of  
 525 rSMR ( $\chi^2 = 0.004$ ,  $df = 1$ ,  $P = 0.952$ ) and food ( $\chi^2 = 1.86$ ,  $df = 1$ ,  $P = 0.173$ ). The  
 anadromous population had a significantly higher rAS for a given rSMR (effect  
 of population background:  $\chi^2 = 21.98$ ,  $df = 1$ ,  $P < 0.001$ ; Figure 4B).

We detected a significant positive relationship between rMMR and rAS ( $\chi^2 =$   
 4689.8,  $df = 1$ ,  $P < 0.001$ ; Figure 4C), but interactions between rMMR  $\times$  food  $\times$   
 530 population ( $\chi^2 = 1.16$ ,  $df = 1$ ,  $P = 0.201$ ), rMMR  $\times$  food ( $\chi^2 = 0.2$ ,  $df = 1$ ,  $P = 0.673$ ),  
 rMMR  $\times$  population ( $\chi^2 = 2.3$ ,  $df = 1$ ,  $P = 0.1297$ ), and food  $\times$  population ( $\chi^2 = 0.1$ ,  
 $df = 1$ ,  $P = 0.768$ ) were all non-significant. The main effects of food ( $\chi^2 = 2.4$ ,  $df$   
 $= 1$ ,  $P = 0.123$ ) and population ( $\chi^2 = 1.9$ ,  $df = 1$ ,  $P = 0.163$ ) were also non-significant.

See Supporting Information for coefficient estimates for all of the above models.

535




**Figure 4:** Size-independent relationships between: (A) residual standard metabolic rate (rSMR) and residual maximum metabolic rate (rMMR); (B) rSMR and residual aerobic scope (rAS); and (C) rMMR and rAS for brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish experienced two food reduction treatments: optimal food rations (High) and 25% of optimal rations (Low).

#### *Effects of sex on metabolism*

There were no sex-based differences in rSMR between , with similar rSMR in males and females (Figure 5A, Table 2). However, male fish had higher rMMR than female fish (Figure 5B, Table 2), and similar sex-based differences were detected in rAS (Figure 5C, Table 2).





**Figure 5:** Gardner-Altman estimation plots for: (A) standard metabolic rate (SMR); (B) maximum metabolic rate (MMR); and (C) aerobic scope (AS) of brown trout classed as female or male after two years of experimental tank rearing. The Gardner-Altman estimation plots show the residual (body mass corrected) SMR/MMR/AS on the left axes and the effect size (mean difference between females and males) is represented by the black dot on the right axes, along with the distribution (shaded curve) and 95% confidence interval (black bars) of the effect size, obtained via non-parametric bootstrap resampling (5,000 resamples).

## 545 Discussion

Intra-specific variation in metabolic rates is widespread across species, yet there are still gaps in our understanding of how intrinsic and extrinsic environmental factors can interact to influence the various components of an individual's metabolism. Here, we exposed brown trout offspring from two population  
 550 backgrounds (one anadromous/sea-migratory, one non-anadromous) to long-term food restriction to determine if, and how, intrinsic factors (i.e. population/sex specific effects) *versus* extrinsic factors (food resources) affect metabolic rates. Fish from both populations had lower SMR under low food conditions, with slight differences in overall SMR between populations. Fish  
 555 from the anadromous population had higher MMR, and consequently, higher AS than the non-anadromous population under all food regimes. Intriguingly, fish from the anadromous-background also had a higher AS at low food compared to high food conditions, suggesting this population was more flexible in maximum metabolic traits than the non-anadromous population. We also  
 560 found differences in MMR and AS linked to sex, which were not apparent in SMR. Collectively, our results suggest the various components of metabolism are differentially affected by intrinsic and extrinsic factors. Moreover, populations may vary in their capacity to flexibly adjust metabolic traits in response to environmental conditions, with consequences for population  
 565 resilience to global change.

### *Effects of extrinsic environment on metabolic traits*

The lower SMR we observed in response to long-term food restriction (> 7 months) is in line with previous work showing SMR (or BMR) to be strongly sensitive to food availability, typically without corresponding changes in MMR  
 570 (Metcalf *et al.* 2016). SMR has been found to show similar flexible decreases in food-poor environments (Naya *et al.* 2007; Auer *et al.* 2015c, 2016a; Zeng *et al.* 2018; Langer *et al.* 2018) or, conversely, to increase at high food availability (Van Leeuwen *et al.* 2011, 2012). Reductions in SMR are assumed to be optimal when food is scarce because the overall energetic cost of living is similarly reduced,  
 575 thus facilitating higher growth (and consequently, fitness) (Auer *et al.* 2015c).

Such flexibility in SMR will likely have positive implications for species experiencing rapid environmental change, with temperature-induced plasticity in SMR linked to increased resilience to climate change (Magozzi and Calosi 2015). However, the adaptiveness of a given flexible response will depend upon  
 580 both the predictability of the environmental change (i.e. the pattern of fluctuations in the environment), and the speed at which organisms can flexibly adjust their phenotypes to match these changing conditions (Reed *et al.* 2010). Moreover, it is unclear whether SMR flexibility translates into overall fitness benefits in scenarios of multi-faceted environmental change. This is particularly  
 585 pertinent for aquatic ectotherms such as salmonines, which are likely to experience reductions in invertebrate prey size and abundance alongside warming (Durance and Ormerod 2007). It remains to be seen whether such populations have the capacity to sufficiently reduce SMR in response to combined stressors of food restriction and warming, though a study in common  
 590 carp *Cyprinus carpio* indicates that the benefits of food-induced SMR plasticity may be temperature dependent. (Zeng *et al.* 2018).

#### *Effects of intrinsic factors on metabolic traits*

We detected overall variation in SMR, MMR, and AS according to population factors, with higher metabolic traits observed for the anadromous population.  
 595 Population-level variation in metabolic traits could arise either through plasticity/flexibility/acclimation, or reflect genetic differences (which could include genetic variation in plasticity itself, *e.g.* variation among genotypes in their extent of flexibility). Metabolic rates are evolvable (Pettersen *et al.* 2018) and have been shown to respond to selection across relatively short time frames,  
 600 *e.g.* BMR increased within 11 generations of selection in bank voles *Myodes glareolus* (Sadowska *et al.* 2015). The differences in SMR (for a given fish size) between populations was in agreement with our expectations, with the non-anadromous population having a lower mass-adjusted SMR than the anadromous population. However, the effect size of population on SMR was  
 605 small in both food treatments, with 95% CIs that included zero, suggesting that differences between populations were marginal. Nevertheless, any differences

in SMR could be indicative of life-history differences between the two populations, where lower SMR in the non-anadromous population has arisen as an evolutionary or plastic response to lower levels of productivity in freshwaters relative to marine (Gross *et al.* 1988). Since we observed population-level differences in SMR in both high and low food treatments, it seems more likely that the differences in SMR between populations are due to inherited genetic differences rather than plastic responses (i.e. the non-anadromous population had lower SMR even at optimal food levels). Such inherited differences could arise from standard inherited allelic variation, or from inherited environmental influences (*e.g.* maternal effects) that could include epigenetic inheritance. Regardless of the inheritance mechanism, the resulting fixed phenotypic differences between the populations could be adaptive. For example, the non-anadromous population may have experienced stronger selection for reduced SMR in order to minimise their baseline energy requirements (Gross *et al.* 1988), whereas selection on SMR may have been in the opposite direction in the naturally anadromous population, whereby higher SMR (and indeed MMR or AS) could facilitate rapid somatic growth in order to reach target smolt sizes to successfully migrate (McCarthy 2000).

The relatively strong differences we observed between populations in MMR (and consequently AS) suggests the upper bounds of metabolism may be more affected by population-specific factors than the lower bounds. We had expected the anadromous population to show comparatively higher MMR than the non-anadromous population, and indeed this was the case, a finding that reflects higher MMR and AS previously described in migratory versus non-migratory ecotypes of three-spined sticklebacks (Tudorache *et al.* 2007). A genetic basis to MMR has been proposed to underpin metabolic variation between migratory forms of three-spined stickleback, where differences in MMR between anadromous and non-anadromous populations have been explained within the context of relaxed selection on swimming performance in stream-resident populations, mediated by reductions in MMR (Dalziel *et al.* 2012b). In contrast, anadromous populations that undertake more arduous/lengthy migrations

tend to have higher swimming/cardiac performances, and higher MMR (Lee *et al.* 2003; Eliason *et al.* 2011; Dalziel *et al.* 2012a), indicating migration effort may  
 640 further underpin consistent differences in MMR between migratory and non-migratory individuals or populations (Seppänen *et al.* 2010). A higher MMR in the anadromous population may also confer fitness benefits by facilitating high levels of growth, either through direct selection on MMR in the freshwater environment (where fast growth increased migration success), and indeed,  
 645 differences in intrinsic freshwater growth rates have previously been described for our study populations (Archer *et al.* 2019). Indirect selection on MMR in juveniles might also occur because of a positive genetic correlation with MMR expressed in the marine environment, where high growth rates are translated into increased fecundity, with rank-order MMR in fish generally repeatable  
 650 through time (Norin and Clark 2016).

Sex-based differences in MMR (and AS) that were not evident in SMR further suggests that MMR is more strongly influenced by intrinsic rather than environmental factors. We observed a higher MMR in males, suggesting that while males and females had similar basic energetic requirements, males had  
 655 more scope to increase their metabolism and divert resources into processes such as growth, or aggressive interactions underpinning competition. Salmonines generally show patterns of sex-specific aggression, with differences in aggression developing early, *e.g.* juvenile *O. mykiss* display more aggressive behaviour than females (Johnsson and Åkerman 1998), a trait likely genetically  
 660 correlated to sex-based differences in competitive ability as adults (Johnsson *et al.* 2001). On a broader scale, our finding corroborates evidence for sex-specific differences in AS described in pink salmon *O. gorbuscha* (Clark *et al.* 2011), and in cardiovascular performance of migrating sockeye salmon *O. nerka* (Sandblom *et al.* 2009). Collectively, these studies suggest that the relatively  
 665 lower AS of female salmonines could make them more susceptible to effects of global change.

*Differences in flexibility of metabolic traits*

That we detected stronger effects of food restriction on SMR compared to MMR or AS lends further support to the proposal that the “ceiling”, which constrains upper limits of metabolism, is less flexible than the metabolic “floor” (i.e. SMR) (Sandblom *et al.* 2016). Nonetheless, we did detect population-specific flexibility in AS, where the anadromous population had marginally higher AS at low rather than high food conditions, a difference that was not apparent in the non-anadromous population. The AS flexibility in the anadromous population appeared to be somewhat underpinned by decreased SMR at low food conditions (i.e. similar effect sizes for food treatment effects on SMR in both populations, but higher/positive effects sizes of low food on AS were only seen in the anadromous population). The few studies that have explored the effects of food restriction on MMR or AS have found little evidence for food-induced flexibility in these traits (Van Leeuwen *et al.* 2011; Killen 2014; Auer *et al.* 2016b; Zeng *et al.* 2018). The population-specific increase in AS in response to food restriction that we observed is initially counter-intuitive, but can be interpreted as further evidence for the optimal combination of metabolic traits being context-dependent. For example, context-dependency of flexibility in MMR and AS have previously been described in barramundi *Lates calcarifer* that showed variable plasticity to hypoxia, salinity, and temperature changes (Norin *et al.* 2016). It is less clear why a higher AS might be optimal in a low food environment. The ability to flexibly increase AS may perhaps be a consequence of the migratory background of this population, particularly if the conditions that promote a migratory life-history in this population also tend to promote flexibility in SMR, MMR, or AS (e.g. fluctuations in food resources/quality in the catchment-of-origin drive patterns of migration and also flexibility in AS).

Plasticity in MMR or AS in the anadromous population may be an adaptive response to conditions of low food, given that food restriction increases the frequency of migratory tactics in brown trout in general (Ferguson *et al.* 2019 and references therein) and has been shown to increase the prevalence of migrants in this population specifically (Archer *et al.* 2019). If low food

environments promote migration, individuals that can increase their MMR are likely to have higher fitness, since high aerobic capacity (i.e. MMR or AS) is required to fuel swimming performance of migrating fish (Claireaux *et al.* 2005; Eliason *et al.* 2011). A flexible MMR in low conditions could be thus be a population-level adaptive response to a migratory background, a response that only emerges at low food availability when it potentially facilitates improved migration performance. Such a response is comparable to studies documenting increased whole animal oxygen consumption of high latitude (i.e. cold-acclimated) killifish *Fundulus heteroclitus* compared to low-latitude fish, with differences only evident at cold extremes (Fangue *et al.* 2009; Dhillon and Schulte 2011).

Higher AS (but not SMR) have been previously associated with competitive performance (Killen *et al.* 2014), and increased food intake (Auer *et al.* 2015a). Selection for higher MMR in low food environments has recently been shown in juvenile Atlantic salmon *S. salar* (Auer *et al.* 2018a), with higher MMR explained in relation to increased competitive ability. As such, a flexible maximum metabolic rate may represent an alternative metabolic strategy to lowering SMR, in order to maintain food consumption rates (and growth) in response to long-term food restriction. The benefits of such an alternative metabolic strategy become clear in the context of long-term food restriction scenarios because SMR depression is associated with the accumulation of harmful mitochondrial reactive oxygen species (ROS) that can impose long-term costs on life-history traits (Salin *et al.* 2018). Alternatively, we cannot rule out that food reduction may have induced higher MMR as a by-product of increased numbers of aggressive interactions between individuals (Seebacher *et al.* 2013), with anadromous brown trout tending to show more aggressive behaviour than non-anadromous forms (Lahti *et al.* 2001).

#### Implications and considerations

Overall, the importance of population-specific factors evident here is consistent with a role for natural selection in the evolution of MMR and SMR, though SMR appears to also respond in a more flexible/plastic manner to environmental

conditions. Intriguingly, because we observed metabolic trait variation in  
 730 populations that naturally express different life histories (in terms of migratory  
 tactics) such population-level differences may be further influenced by  
 proximate and ultimate mechanisms underpinning alternative migration  
 (Hazel *et al.* 1990; Tomkins and Hazel 2007; Pulido 2011). For example, if  
 primarily non-migratory populations (or individuals) tend to have lower SMR,  
 735 this may increase the frequency of fish that meet their residency  
 threshold/“switch point”, meaning the population shifts further towards  
 residency and undergoes further selection for lower SMR up to some limit.  
 Conversely, if a migratory population (or individual) tends to have higher SMR,  
 this increases the likelihood that individuals become energetically constrained  
 740 in freshwater, (i.e. do not meet their residency threshold), and thus more fish  
 become migrants (and the population as a whole experiences positive selection  
 on SMR). Subsequently, migrants with a higher MMR will likely perform better  
 during migration and at sea, achieving higher levels of fitness, and thus  
 selection will favour a higher MMR in such populations. Environmental  
 745 conditions could also affect evolutionary processes, if factors such as low food  
 or inclement temperatures interact with metabolic traits and further alter the  
 costs and benefits of migration *versus* residency tactics. A tight evolutionary  
 coupling has recently been described for metabolic rate and pace of life history  
 in guppies *Poecilia reticulata* (Auer *et al.* 2018b), and similar mechanisms could  
 750 be at play in species with alternative migratory life histories. Because  
 metabolism and life-history traits are both influenced by environmental factors,  
 reciprocal translocation or “common garden” experiments in nature (*e.g.* Auer  
*et al.* (2018b)) can help to disentangle interactions between physiology,  
 environment, and genetics as both the causes and consequences of alternative  
 755 migratory tactics (i.e. whether covariation between metabolic traits and life  
 history is caused by phenotypic flexibility or fixed inherited differences).

It is important also to note that here we only considered metabolic variation at  
 a single life stage, and it is likely that metabolism (and its flexibility) can vary  
 though ontogeny (Pettersen *et al.* 2016), depending on energy requirements



760 associated with developmental stage (Beaman *et al.* 2016; Burggren 2018), or  
 seasonal changes (Versteegh *et al.* 2012; Petit *et al.* 2013). Expanding our  
 approach to incorporate repeated measurements on individuals throughout  
 their lives (or indeed on individuals at different stages of ontogeny) would  
 further illuminate when and how metabolic flexibility develops, and is most  
 765 beneficial. Nonetheless, our results indicate that metabolic traits can respond  
 differently to extrinsic and intrinsic factors, and metabolic responses can  
 further vary according to population-specific factors. That maximum  
 metabolism was more fixed than minimum metabolism suggests that trout may  
 be more constrained in their capacity to adjust AS in response to extrinsic  
 770 factors, with important conservation implications for a species that is already in  
 global decline due to anthropogenic activities (Limburg and Waldman 2009).  
 Moreover, the variation in metabolic traits that we observed in our two brown  
 trout populations indicates that responses to environmental change are unlikely  
 to be universal; and developing effective management strategies is not  
 775 necessarily a straightforward task. Nevertheless, greater plasticity is linked to  
 higher resilience (Magozzi and Calosi 2015; Seebacher *et al.* 2015) if  
 environmental changes are predictable (Reed *et al.* 2010) and understanding  
 the capacity of species and populations to flexibly adjust their metabolic traits  
 is essential for predicting and mitigating the effects of progressively changing  
 780 environmental conditions in natural systems.

### Acknowledgements

The authors would like to thank Brian Clarke, Deirdre Cotter, members of the  
 FishEyE team at UCC, and the staff of Inland Fisheries Ireland and the Marine  
 Institute for obtaining brood stock and for assistance in fish rearing, along with  
 785 Robert Wynne, Ronan O'Sullivan, Peter Moran and Adam Kane for assistance  
 in fish husbandry, and Jamie Coughlan for genotyping work. This research was  
 supported by an ERC Starting Grant (639192-ALH) and an SFI ERC Support  
 Award awarded to TER. PMcG was supported in part by grants from Science  
 Foundation Ireland (15/IA/3028 & 16/BBSRC/3316) and by grant-in-aid

790 (RESPI/FS/I6/01) from the Marine Institute (Ireland) as part of the Marine Research Programme by the Irish Government.

## References

- Allan EL, Froneman PW, and Hodgson AN. 2006. Effects of temperature and salinity on the standard metabolic rate (SMR) of the caridean shrimp *Palaemon peringueyi*. *Journal of Experimental Marine Biology and Ecology* **337**: 103–8.
- 795 Álvarez D, Cano JM, and Nicieza AG. 2006. Microgeographic variation in metabolic rate and energy storage of brown trout: Countergradient selection or thermal sensitivity? *Evolutionary Ecology* **20**: 345–63.
- Angilletta Jr Michael J. 2001. Variation in metabolic rate between populations of a geographically widespread lizard. *Physiological and Biochemical Zoology* **74**: 11–21.
- 800 Archer LC, Hutton SA, Harman L, O’Grady MN, Kerry JP, Poole WR, Gargan P, McGinnity P, and Reed TE. 2019. The interplay between extrinsic and intrinsic factors in determining migration decisions in brown trout (*Salmo trutta*): An experimental study. *Frontiers in Ecology and Evolution* **7**: 222.
- 805 Auer SK, Anderson GJ, McKelvey S, Bassar RD, McLennan D, Armstrong JD, Nislow KH, Downie HK, McKelvey L, Morgan TAJ, Salin K, Orrell DL, Gauthey A, Reid TC, and Metcalfe NB. 2018a. Nutrients from salmon parents alter selection pressures on their offspring. *Ecology Letters* **21**: 287–95.
- 810 Auer SK, Dick CA, Metcalfe NB, and Reznick DN. 2018b. Metabolic rate evolves rapidly and in parallel with the pace of life history. *Nature Communications* **9**: 14.
- Auer SK, Killen SS, and Rezende EL. 2017. Resting vs. active: A meta-analysis of the intra- and inter-specific associations between minimum, sustained, and maximum metabolic rates in vertebrates. *Functional Ecology* **31**: 1728–38.
- 815 Auer SK, Salin K, Anderson GJ, and Metcalfe NB. 2015a. Aerobic scope explains individual variation in feeding capacity. *Biology Letters* **11**: 20150793.
- Auer SK, Salin K, Anderson GJ, and Metcalfe NB. 2016a. Flexibility in metabolic rate and activity level determines individual variation in overwinter performance. *Oecologia* **182**: 703–12.
- 820 Auer SK, Salin Karine, Rudolf Agata M., Anderson Graeme J., and Metcalfe Neil B. 2016b. Differential effects of food availability on minimum and maximum rates of metabolism. *Biology Letters* **12**: 20160586.
- Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015b. The optimal combination of standard metabolic rate and aerobic scope for somatic growth depends on food availability. *Functional Ecology* **29**: 479–86.
- 825 Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015c. Flexibility in metabolic rate confers a growth advantage under changing food availability. *Journal of Animal Ecology* **84**: 1405–11.
- 830 Barceló G, Love OP, and Vézina F. 2016. Uncoupling basal and summit metabolic rates in white-throated sparrows: Digestive demand drives maintenance costs, but changes in muscle mass are not needed to improve thermogenic capacity. *Physiological and Biochemical Zoology* **90**: 153–65.

- 835 Beaman JE, White CR, and Seebacher F. 2016. Evolution of plasticity: Mechanistic link between development and reversible acclimation. *Trends in Ecology & Evolution* **31**: 237–49.
- Biro PA and Stamps JA. 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? *Trends in Ecology & Evolution* **25**: 653–9.
- 840 Bochdanský AB, Grønkjær P, Herra TP, and Leggett WC. 2005. Experimental evidence for selection against fish larvae with high metabolic rates in a food limited environment. *Marine Biology* **147**: 1413–7.
- Bolnick DI, Amarasekare P, Araújo MS, Bürger R, Levine JM, Novak M, Rudolf VHW, Schreiber SJ, Urban MC, and Vasseur DA. 2011. Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution* **26**: 183–92.
- 845 Buoro M, Gimenez O, and Prévost E. 2012. Assessing adaptive phenotypic plasticity by means of conditional strategies from empirical data: The latent environmental threshold model. *Evolution* **66**: 996–1009.
- Burggren W. 2018. Developmental phenotypic plasticity helps bridge stochastic weather events associated with climate change. *Journal of Experimental Biology* **221**: jeb161984.
- 850 Burton T, Killen SS, Armstrong JD, and Metcalfe NB. 2011. What causes intraspecific variation in resting metabolic rate and what are its ecological consequences? *Proceedings of the Royal Society B: Biological Sciences* **278**: 3465–73.
- Cavieres G and Sabat P. 2008. Geographic variation in the response to thermal acclimation in rufous-collared sparrows: Are physiological flexibility and environmental heterogeneity correlated? *Functional Ecology* **22**: 509–15.
- 855 Chabot D, Steffensen JF, and Farrell AP. 2016. The determination of standard metabolic rate in fishes. *Journal of Fish Biology* **88**: 81–121.
- 860 Chapman BB, Hulthén K, Brodersen J, Nilsson PA, Skov C, Hansson L-A, and Brönmark C. 2012. Partial migration in fishes: Causes and consequences. *Journal of Fish Biology* **81**: 456–78.
- Claireaux G, McKenzie DJ, Genge AG, Chatelier A, Aubin J, and Farrell AP. 2005. Linking swimming performance, cardiac pumping ability and cardiac anatomy in rainbow trout. *Journal of Experimental Biology* **208**: 1775–84.
- 865 Clark TD, Jeffries KM, Hinch SG, and Farrell AP. 2011. Exceptional aerobic scope and cardiovascular performance of pink salmon (*Oncorhynchus gorbuscha*) may underlie resilience in a warming climate. *Journal of Experimental Biology* **214**: 3074–81.
- Cutts CJ, Metcalfe NB, and Taylor AC. 2002. Juvenile Atlantic Salmon (*Salmo salar*) with relatively high standard metabolic rates have small metabolic scopes. *Functional Ecology* **16**: 73–8.
- 870 Dalziel AC, Ou M, and Schulte PM. 2012a. Mechanisms underlying parallel reductions in aerobic capacity in non-migratory threespine stickleback (*Gasterosteus aculeatus*) populations. *Journal of Experimental Biology* **215**: 746–59.
- 875 Dalziel AC, Vines TH, and Schulte PM. 2012b. Reductions in prolonged swimming capacity following freshwater colonization in multiple threespine stickleback populations. *Evolution* **66**: 1226–39.

- Dhillon RS and Schulte PM. 2011. Intraspecific variation in the thermal plasticity of mitochondria in killifish. *Journal of Experimental Biology* **214**: 3639–48.
- 880 Dubois K, Hallot F, and Vézina F. 2016. Basal and maximal metabolic rates differ in their response to rapid temperature change among avian species. *Journal of Comparative Physiology B* **186**: 919–35.
- Durance I and Ormerod SJ. 2007. Climate change effects on upland stream macroinvertebrates over a 25-year period. *Global Change Biology* **13**: 942–57.
- 885 Eliason EJ, Clark TD, Hague MJ, Hanson LM, Gallagher ZS, Jeffries KM, Gale MK, Patterson DA, Hinch SG, and Farrell AP. 2011. Differences in thermal tolerance among sockeye salmon populations. *Science* **332**: 109–12.
- Fangue NA, Richards JG, and Schulte PM. 2009. Do mitochondrial properties explain intraspecific variation in thermal tolerance? *The Journal of Experimental Biology*  
890 **212**: 514–22.
- Ferguson A, Reed TE, Cross TF, McGinnity P, and Prodöhl PA. 2019. Anadromy, potamodromy and residency in brown trout *Salmo trutta*: The role of genes and the environment. *Journal of Fish Biology* **95**: 692–718.
- Forseth T, Nesje TF, Jonsson B, and Hårsaker K. 1999. Juvenile migration in brown trout: A consequence of energetic state. *Journal of Animal Ecology* **68**: 783–93.
- 895 Gargan P, Kelly F, Shephard S, and Whelan K. 2016. Temporal variation in sea trout *Salmo trutta* life history traits in the Erriff River, western Ireland. *Aquaculture Environment Interactions* **8**: 675–89.
- Gross MR, Coleman RM, and McDowall RM. 1988. Aquatic productivity and the evolution of diadromous fish migration. *Science* **239**: 1291–3.
- 900 Guderley H and Pörtner HO. 2010. Metabolic power budgeting and adaptive strategies in zoology: Examples from scallops and fish. *Canadian Journal of Zoology* **88**: 753–63.
- Halsey LG. 2019. The reign of the p-value is over: What alternative analyses could we employ to fill the power vacuum? *Biology Letters* **15**: 20190174.
- 905 Halsey LG, Curran-Everett D, Vowler SL, and Drummond GB. 2015. The fickle P value generates irreproducible results. *Nature Methods* **12**: 179–85.
- Hariato J and Carey N. 2019. respR: Analyse, Convert, and Automate Respirometry-Related Data. R package version 1.0.5.1.  
910 <https://github.com/januarhariato/respR>.
- Hazel WN, Smock R., and Johnson M. D. 1990. A polygenic model for the evolution and maintenance of conditional strategies. *Proceedings of the Royal Society of London Series B: Biological Sciences* **242**: 181–7.
- Hillman SS, Hancock TV, and Hedrick MS. 2013. A comparative meta-analysis of maximal aerobic metabolism of vertebrates: Implications for respiratory and cardiovascular limits to gas exchange. *Journal of Comparative Physiology B* **183**:  
915 167–79.
- Ho J, Tumkaya T, Aryal S, Choi H, and Claridge-Chang A. 2019. Moving beyond P values: Data analysis with estimation graphics. *Nature Methods* **16**: 565.

- 920 Hochachka PW, Buck LT, Doll CJ, and Land SC. 1996. Unifying theory of hypoxia tolerance: Molecular/metabolic defense and rescue mechanisms for surviving oxygen lack. *Proceedings of the National Academy of Sciences* **93**: 9493–8.
- Hofmann GE and Todgham AE. 2010. Living in the now: Physiological mechanisms to tolerate a rapidly changing environment. *Annual Review of Physiology* **72**: 127–45.
- 925 Johnsson JI and Åkerman A. 1998. Intersexual differences in aggression in juvenile rainbow trout. *Journal of Fish Biology* **52**: 1292–4.
- Johnsson JI, Sernland E, and Blixt M. 2001. Sex-specific aggression and antipredator behaviour in young brown trout. *Ethology* **107**: 587–99.
- 930 Killen SS. 2014. Growth trajectory influences temperature preference in fish through an effect on metabolic rate. *Journal of Animal Ecology* **83**: 1513–22.
- Killen SS, Atkinson D, and Glazier DS. 2010. The intraspecific scaling of metabolic rate with body mass in fishes depends on lifestyle and temperature. *Ecology Letters* **13**: 184–93.
- 935 Killen SS, Marras S, Metcalfe NB, McKenzie DJ, and Domenici P. 2013. Environmental stressors alter relationships between physiology and behaviour. *Trends in Ecology & Evolution* **28**: 651–8.
- Killen SS, Mitchell MD, Rummer JL, Chivers DP, Ferrari MCO, Meekan MG, and McCormick MI. 2014. Aerobic scope predicts dominance during early life in a tropical damselfish. *Functional Ecology* **28**: 1367–76.
- 940 Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, and Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): A review of aspects of their life histories. *Ecology of Freshwater Fish* **12**: 1–59.
- 945 Konarzewski M and Książek A. 2013. Determinants of intra-specific variation in basal metabolic rate. *Journal of Comparative Physiology B* **183**: 27–41.
- Lahti K, Laurila A, Enberg K, and Piironen J. 2001. Variation in aggressive behaviour and growth rate between populations and migratory forms in the brown trout, *Salmo trutta*. *Animal Behaviour* **62**: 935–44.
- 950 Langer F, Havenstein N, and Fietz J. 2018. Flexibility is the key: Metabolic and thermoregulatory behaviour in a small endotherm. *Journal of Comparative Physiology B* **188**: 553–63.
- Lee CG, Farrell AP, Lotto A, Hinch SG, and Healey MC. 2003. Excess post-exercise oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*) salmon following critical speed swimming. *Journal of Experimental Biology* **206**: 3253–60.
- 955 Limburg KE and Waldman JR. 2009. Dramatic declines in north Atlantic diadromous fishes. *BioScience* **59**: 955–65.
- Magee J. 2017. A comparison of population structuring and genetic stock identification of brown trout (*Salmo trutta*) displaying distinct migratory strategies. In: PhD thesis, pp277. Queens University Belfast, Belfast, NI.
- 960 Magozzi S and Calosi P. 2015. Integrating metabolic performance, thermal tolerance, and plasticity enables for more accurate predictions on species vulnerability to acute and chronic effects of global warming. *Global Change Biology* **21**: 181–94.

- 965 Maldonado K, Bozinovic F, Cavieres G, Fuentes CA, Cortés A, and Sabat P. 2012. Phenotypic flexibility in basal metabolic rate is associated with rainfall variability among populations of rufous-collared sparrow. *Zoology* **115**: 128–33.
- McCann SM, Kosmala GK, Greenlees MJ, and Shine R. 2018. Physiological plasticity in a successful invader: Rapid acclimation to cold occurs only in cool-climate
- 970 populations of cane toads (*Rhinella marina*). *Conservation Physiology* **6**.
- McCarthy ID. 2000. Temporal repeatability of relative standard metabolic rate in juvenile Atlantic salmon and its relation to life history variation. *Journal of Fish Biology* **57**: 224–38.
- Metcalfe NB, Van Leeuwen TE, and Killen SS. 2016. Does individual variation in
- 975 metabolic phenotype predict fish behaviour and performance? *Journal of Fish Biology* **88**: 298–321.
- Naya DE and Bozinovic F. 2012. Metabolic scope of fish species increases with distributional range. *Evolutionary Ecology Research* **14**: 769–77.
- Naya DE, Lardies MA, and Bozinovic F. 2007. The effect of diet quality on physiological
- 980 and life-history traits in the harvestman *Pachylus paessleri*. *Journal of Insect Physiology* **53**: 132–8.
- Nespolo RF, Bacigalupe LD, Rezende EL, and Bozinovic F. 2001. When nonshivering thermogenesis equals maximum metabolic rate: Thermal acclimation and phenotypic plasticity of fossorial *Spalacopus cyanus* (Rodentia). *Physiological and Biochemical Zoology* **74**: 325–32.
- 985 Norin T and Clark TD. 2016. Measurement and relevance of maximum metabolic rate in fishes. *Journal of Fish Biology* **88**: 122–51.
- Norin T, Malte H, and Clark TD. 2016. Differential plasticity of metabolic rate phenotypes in a tropical fish facing environmental change. *Functional Ecology* **30**: 369–78.
- 990 Norin T and Metcalfe N. 2019. Ecological and evolutionary consequences of metabolic rate plasticity in response to environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**: 20180180.
- O'Connor K i., Taylor A c., and Metcalfe N b. 2000. The stability of standard metabolic rate during a period of food deprivation in juvenile Atlantic salmon. *Journal of Fish Biology* **57**: 41–51.
- 995 Petit M, Lewden A, and Vézina F. 2013. Intra-seasonal flexibility in avian metabolic performance highlights the uncoupling of basal metabolic rate and thermogenic capacity. *PLOS ONE* **8**: e68292.
- 1000 Pettersen AK, Marshall DJ, and White CR. 2018. Understanding variation in metabolic rate. *Journal of Experimental Biology* **221**: jeb166876.
- Pettersen AK, White CR, and Marshall DJ. 2016. Metabolic rate covaries with fitness and the pace of the life history in the field. *Proc R Soc B* **283**: 20160323.
- Piche J, Hutchings JA, and Blanchard W. 2008. Genetic variation in threshold reaction norms for alternative reproductive tactics in male Atlantic salmon, *Salmo salar*. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1571–5.
- 1005 Piersma T and Drent J. 2003. Phenotypic flexibility and the evolution of organismal design. *Trends in Ecology & Evolution* **18**: 228–33.

- Poole WR, Dillane M, DeEyto E, Rogan G, McGinnity P, and Whelan K. 2007. Characteristics of the Burrishoole sea trout population: Census, marine survival, enhancement and stock-recruitment relationship, 1971-2003. In: *Sea Trout: Biology, Conservation and Management* (Eds: G Harris and N Milner). Oxford, UK: Blackwells Scientific Publications.
- Pulido F. 2011. Evolutionary genetics of partial migration – the threshold model of migration revis(it)ed. *Oikos* **120**: 1776–83.
- R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Reed TE, Waples Robin S., Schindler Daniel E., Hard Jeffrey J., and Kinnison Michael T. 2010. Phenotypic plasticity and population viability: The importance of environmental predictability. *Proceedings of the Royal Society B: Biological Sciences* **277**: 3391–400.
- Sadowska ET, Stawski Clare, Rudolf Agata, Dheyongera Geoffrey, Chrzęścik Katarzyna M., Baliga-Klimczyk Katarzyna, and Koteja Paweł. 2015. Evolution of basal metabolic rate in bank voles from a multidirectional selection experiment. *Proceedings of the Royal Society B: Biological Sciences* **282**: 20150025.
- Salin K, Villasevil EM, Anderson GJ, Auer SK, Selman C, Hartley RC, Mullen W, Chinopoulos C, and Metcalfe NB. 2018. Decreased mitochondrial metabolic requirements in fasting animals carry an oxidative cost. *Functional Ecology* **32**: 2149–57.
- Sandblom E, Clark TD, Gräns A, Ekström A, Brijs J, Sundström LF, Odelström A, Adill A, Aho T, and Jutfelt F. 2016. Physiological constraints to climate warming in fish follow principles of plastic floors and concrete ceilings. *Nature Communications* **7**: 11447.
- Sandblom E, Clark TD, Hinch SG, and Farrell AP. 2009. Sex-specific differences in cardiac control and hematology of sockeye salmon (*Oncorhynchus nerka*) approaching their spawning grounds. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology* **297**: R1136–1143.
- Seebacher F, Ward AJW, and Wilson RS. 2013. Increased aggression during pregnancy comes at a higher metabolic cost. *Journal of Experimental Biology* **216**: 771–6.
- Seebacher F, White CR, and Franklin CE. 2015. Physiological plasticity increases resilience of ectothermic animals to climate change. *Nature Climate Change* **5**: 61–6.
- Seppänen E, Piironen J, and Huuskonen H. 2010. Consistency of standard metabolic rate in relation to life history strategy of juvenile Atlantic salmon *Salmo salar*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **156**: 278–84.
- Sloat MR and Reeves GH. 2014. Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences* **71**: 491–501.
- Svendsen MBS, Bushnell PG, and Steffensen JF. 2016. Design and setup of intermittent-flow respirometry system for aquatic organisms. *Journal of Fish Biology* **88**: 26–50.

- 1055 Tomkins JL and Hazel W. 2007. The status of the conditional evolutionarily stable strategy. *Trends in Ecology & Evolution* **22**: 522–8.
- Tudorache C, Blust R, and Boeck GD. 2007. Swimming capacity and energetics of migrating and non-migrating morphs of three-spined stickleback *Gasterosteus aculeatus* L. and their ecological implications. *Journal of Fish Biology* **71**: 1448–
- 1060 56.
- Van Leeuwen TE, Rosenfeld JS, and Richards JG. 2011. Adaptive trade-offs in juvenile salmonid metabolism associated with habitat partitioning between coho salmon and steelhead trout in coastal streams. *Journal of Animal Ecology* **80**: 1012–23.
- 1065 Van Leeuwen TE, Rosenfeld JS, and Richards JG. 2012. Effects of food ration on SMR: Influence of food consumption on individual variation in metabolic rate in juvenile coho salmon (*Oncorhynchus kisutch*). *Journal of Animal Ecology* **81**: 395–402.
- 1070 Ven TMFN van de, Mzilikazi N, and McKechnie AE. 2013. Phenotypic flexibility in body mass, basal metabolic rate and summit metabolism in southern red bishops (*Euplectes orix*): Responses to short term thermal acclimation. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **165**: 319–27.
- 1075 Versteegh MA, Helm B, Gwinner E, and Tieleman BI. 2012. Annual cycles of metabolic rate are genetically determined but can be shifted by phenotypic flexibility. *Journal of Experimental Biology* **215**: 3459–66.
- West-Eberhard MJ. 2003. Developmental plasticity and evolution. Oxford, UK: Oxford University Press.
- 1080 Winwood-Smith HS, Alton LA, Franklin CE, and White CR. 2015. Does greater thermal plasticity facilitate range expansion of an invasive terrestrial anuran into higher latitudes? *Conservation Physiology* **3**.
- 1085 Wone BWM, Madsen P, Donovan ER, Labocha MK, Sears MW, Downs CJ, Sorensen DA, and Hayes JP. 2015. A strong response to selection on mass-independent maximal metabolic rate without a correlated response in basal metabolic rate. *Heredity* **114**: 419–27.
- Zeng L-Q, Fu C, and Fu S-J. 2018. The effects of temperature and food availability on growth, flexibility in metabolic rates and their relationships in juvenile common carp. *Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology* **217**: 26–34.
- 1090 Zeng L-Q, Zhang A-J, Killen SS, Cao Z-D, Wang Y-X, and Fu S-J. 2017. Standard metabolic rate predicts growth trajectory of juvenile Chinese crucian carp (*Carassius auratus*) under changing food availability. *Biology Open* **6**: 1305–9.



## Supporting Information

---

# Metabolic traits in brown trout (*Salmo trutta*) vary in response to food restriction and population background


Louise C. Archer<sup>\*1,2</sup>, Stephen A. Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, W. Russell Poole<sup>3</sup>, Patrick Gargan<sup>4</sup>, Philip McGinnity<sup>1,3</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> *School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.*


<sup>2</sup> *Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland.*

<sup>3</sup> *Marine Institute, Furnace, Newport, Co. Mayo, Ireland.*

<sup>4</sup> *Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.*



**Figure S1:** (A) Bunaveela Lough in the Burrishoole catchment, where brown trout brood stock were collected by seine netting in November 2015, used to produce F1 offspring for the experimental study. The wild Bunaveela population does not express the anadromous life history, and experimental offspring from Bunaveela brood stock were considered to have non-anadromous population background. (B) Tawnyard Lough in the Erriff catchment, site of brown trout brood stock collections (via seine netting) in November 2015, used to produce F1 offspring for the experimental study. The wild Tawnyard Lough population has a strong anadromous component, and experimental offspring from Tawnyard brood stock were considered to have an anadromous population background.



**Figure S2:** (A) Body mass scaling of standard metabolic rate (SMR) measured in April and May 2017 in brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental food treatments: optimal food rations (High); and 25% of optimal daily rations (Low). (B) Residual SMR values (body mass corrected) for each population background and food treatment combination. Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (C) Cumming estimation plots with effect sizes shown as black dots (i.e. the mean differences in rSMR among the groups), the distributions (shaded curves) and 95% confidence intervals (back bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).

**Table S1:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from a mixed effect model (marginal  $R^2 = 0.92$ ) testing the effects of food treatment (high or low) and population background (anadromous or non-anadromous) on standard metabolic rate (SMR) measured in brown trout in April and May 2017. SMR was  $\log_{10}$ -transformed, and body mass ( $\log_{10}$ -transformed) was included as a covariate. Date of SMR measurement was included as a random effect. Significance was assessed at a 5% alpha level. Effects are contrasted against fish from the non-anadromous population background in the high food treatment.

| Response        | Parameter              | Estimate | SE   | <i>t</i> -value | <i>P</i> -value |
|-----------------|------------------------|----------|------|-----------------|-----------------|
| $\log_{10}$ SMR | Intercept              | -1.06    | 0.07 | -15.19          | < 0.001         |
|                 | Food: Low              | -0.08    | 0.02 | -3.69           | 0.005           |
|                 | Population: Anadromous | 0.03     | 0.02 | 1.40            | 0.196           |
|                 | $\log_{10}$ Body mass  | 0.99     | 0.05 | 21.11           | < 0.001         |

**Table S2:** Parameter estimates, with associated standard errors (SE), *t*-values, and *P*-values from the linear model describing  $\log_{10}$ -transformed metabolic rates ( $\text{mg O}_2 \text{ hr}^{-1}$ ) as a function of  $\log_{10}$ -transformed body mass in  $n = 61$  brown trout (SMR = standard metabolic rate, MMR = maximum metabolic rate, and AS = aerobic scope).

| Response        | Parameter             | Estimate | SE   | <i>t</i> -value | <i>P</i> -value |
|-----------------|-----------------------|----------|------|-----------------|-----------------|
| $\log_{10}$ SMR | Intercept             | -1.54    | 0.16 | -9.50           | < 0.001         |
|                 | $\log_{10}$ Body mass | 1.10     | 0.08 | 13.64           | < 0.001         |
| $\log_{10}$ MMR | Intercept             | -0.26    | 0.30 | -0.85           | 0.397           |
|                 | $\log_{10}$ Body mass | 0.89     | 0.15 | 5.99            | < 0.001         |
| $\log_{10}$ AS  | Intercept             | -0.24    | 0.35 | -0.68           | 0.497           |
|                 | $\log_{10}$ Body mass | 0.85     | 0.17 | 4.89            | < 0.001         |

**Table S3:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from the general linear models testing the effects of food treatment (high or low) and population background (anadromous or non-anadromous) on relationships between residual (i.e. body mass corrected) standard metabolic rate (rSMR) and residual maximum metabolic rate (rMMR), rSMR and residual aerobic scope (rAS), and rMMR and rAS in brown trout. Significance was assessed at a 5% alpha level. Effects are contrasted against fish from the non-anadromous population background in the high food treatment.

| Response | Parameter              | Estimate | SE    | <i>t</i> -value | <i>P</i> -value |
|----------|------------------------|----------|-------|-----------------|-----------------|
| rMMR     | Intercept              | -0.09    | 0.02  | -3.79           | < 0.001         |
|          | rSMR                   | 0.18     | 0.22  | 0.81            | 0.420           |
|          | Food: Low              | 0.04     | 0.03  | 1.49            | 0.142           |
|          | Population: Anadromous | 0.13     | 0.03  | 4.73            | < 0.001         |
| rAS      | Intercept              | -0.10    | 0.03  | -3.70           | < 0.001         |
|          | rSMR                   | 0.02     | 0.26  | 0.06            | 0.952           |
|          | Food: Low              | 0.05     | 0.03  | 1.36            | 0.178           |
|          | Population: Anadromous | 0.15     | 0.03  | 4.69            | < 0.001         |
| rAS      | Intercept              | 0.001    | 0.003 | 0.17            | 0.868           |
|          | rMMR                   | 1.172    | 0.017 | 68.48           | < 0.001         |
|          | Food: Low              | 0.005    | 0.004 | 1.54            | 0.128           |
|          | Population: Anadromous | -0.006   | 0.004 | -1.39           | 0.169           |

*Supplementary ANCOVA analyses*

As additional analyses to ascertain the robustness of our estimation statistics approach, we also tested for variation in the relationships between body mass and SMR, MMR, and AS according to population and food treatment factors using general linear models (GLMs). Body mass and metabolic rates were  $\log_{10}$  transformed to normalise and linearise the data. The GLMs included either  $\log_{10}$  SMR,  $\log_{10}$  MMR, or  $\log_{10}$  AS as response variables, and the models included  $\log_{10}$  body mass, food treatment (high or low) and population background (anadromous or non-anadromous) as explanatory variables. We also included an interaction between body mass and food treatment, an interaction between mass and population, and a food  $\times$  population interaction to test if responses to food reductions were similar for both populations. We created an additional set of GLMs to test for sex-based differences in  $\log_{10}$ -transformed SMR, MMR, and AS, where sex (male or female) and  $\log_{10}$  body mass were included as response variables. For all of the above GLMs, separate models were constructed for SMR, MMR and AS ( $\log_{10}$ -transformed) and we assumed normal errors in each case.

*Supplementary ANCOVA results*


$\log_{10}$  SMR increased with  $\log_{10}$  body mass ( $\chi^2 = 205.26$ ,  $df = 1$ ,  $P < 0.001$ ), and the food treatment  $\times$  population interaction term was non-significant (LRT for model with and without interaction term:  $\chi^2 = 0.22$ ,  $df = 1$ ,  $P = 0.643$ ).  $\log_{10}$  SMR was significantly affected by food treatment ( $\chi^2 = 8.72$ ,  $df = 1$ ,  $P = 0.003$ ) and population background ( $\chi^2 = 4.20$ ,  $df = 1$ ,  $P = 0.040$ ). For a given size, fish in the low food treatment had lower SMR values than those in the high food treatment, and fish from the non-anadromous population also had a lower SMR than those from the anadromous population (Figure S3A).

$\log_{10}$  MMR increased with  $\log_{10}$  body mass ( $\chi^2 = 51.70$ ,  $df = 1$ ,  $P < 0.001$ ), with a non-significant interaction between food treatment and population ( $\chi^2 = 0.99$ ,  $df = 1$ ,  $P = 0.320$ ), and no significant differences between food treatments ( $\chi^2 = 1.63$ ,  $df = 1$ ,  $P = 0.202$ ). There was a significant effect of population ( $\chi^2 = 25.88$ ,  $df = 1$ ,  $P < 0.001$ ), where for a given size, fish from the anadromous population


had a higher MMR than those from the non-anadromous population (Figure S3B).

$\log_{10}$  AS similarly varied with  $\log_{10}$  body mass ( $\chi^2 = 34.33$ ,  $df = 1$ ,  $P < 0.001$ ), with no significant interaction between food treatment and population ( $\chi^2 = 0.94$ ,  $df = 1$ ,  $P = 0.333$ ), and a non-significant effect of food treatment ( $\chi^2 = 2.08$ ,  $df = 1$ ,  $P = 0.149$ ). Fish from the anadromous population had a higher AS (for a given fish size) than those from the non-anadromous background (Figure S3C:  $\chi^2 = 23.74$ ,  $df = 1$ ,  $P < 0.001$ ).

The positive relationship between  $\log_{10}$  SMR and  $\log_{10}$  body mass ( $\chi^2 = 141.01$ ,  $df = 1$ ,  $P < 0.001$ ) was similar for female and male fish (Figure S4A:  $\chi^2 = 1.08$ ,  $df = 1$ ,  $P = 0.299$ ). However, males of a given size had a higher MMR than females (Figure S4B:  $\chi^2 = 36.28$ ,  $df = 1$ ,  $P < 0.001$ ) and sex-based differences were similarly detected in AS for a given fish size (Figure S4C:  $\chi^2 = 33.90$ ,  $df = 1$ ,  $P < 0.001$ ). See Table S4 and Table S5 for coefficients from all above GLMs.



**Figure S3:** Body mass scaling of (A) standard metabolic rate (SMR), (B) maximum metabolic rate (MMR), and (C) aerobic scope (AS) of brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental food treatments: optimal food rations (High); and 25% of optimal daily rations (Low).



**Figure S4:** Body mass scaling of (A) standard metabolic rate (SMR); (B) maximum metabolic rate (MMR); and (C) aerobic scope (AS) of brown trout classed as female or male after two years of experimental tank rearing.



**Table S4:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from the general linear models testing the effects of food treatment (high or low) and population background (anadromous or non-anadromous) on standard metabolic rate (SMR), maximum metabolic rate (MMR), and aerobic scope (AS) in brown trout. SMR, MMR, and AS were log<sub>10</sub>-transformed, and body mass (log<sub>10</sub>-transformed) was included as a covariate. Significance was assessed at a 5% alpha level. Effects are contrasted against fish from the non-anadromous population background in the high food treatment.

| Response              | Parameter                   | Estimate | SE   | <i>t</i> -value | <i>P</i> -value |
|-----------------------|-----------------------------|----------|------|-----------------|-----------------|
| log <sub>10</sub> SMR | Intercept                   | -1.47    | 0.15 | -9.73           | < 0.001         |
|                       | Food: Low                   | -0.05    | 0.02 | -2.95           | 0.005           |
|                       | Population: Anadromous      | 0.03     | 0.02 | 2.05            | 0.045           |
|                       | log <sub>10</sub> Body mass | 1.06     | 0.07 | 14.33           | < 0.001         |
| log <sub>10</sub> MMR | Intercept                   | -0.37    | 0.26 | -1.45           | 0.152           |
|                       | Food: Low                   | 0.03     | 0.03 | 1.28            | 0.207           |
|                       | Population: Anadromous      | 0.14     | 0.03 | 5.09            | < 0.001         |
|                       | log <sub>10</sub> Body mass | 0.90     | 0.13 | 7.19            | < 0.001         |
| log <sub>10</sub> AS  | Intercept                   | -0.38    | 0.30 | -1.27           | 0.211           |
|                       | Food: Low                   | 0.05     | 0.03 | 1.44            | 0.155           |
|                       | Population: Anadromous      | 0.15     | 0.03 | 4.87            | < 0.001         |
|                       | log <sub>10</sub> Body mass | 0.87     | 0.15 | 5.86            | < 0.001         |

**Table S5:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from the general linear models testing the effect of sex on standard metabolic rate (SMR), maximum metabolic rate (MMR), aerobic scope (AS) of brown trout (*n* = 55). SMR, MMR, and AS were log<sub>10</sub>-transformed, and log<sub>10</sub> body mass was included as a covariate in all models. Significance was assessed at a 5% alpha level. Effects are contrasted against female fish.

| Response              | Parameter                   | Estimate | SE   | <i>t</i> -value | <i>P</i> -value |
|-----------------------|-----------------------------|----------|------|-----------------|-----------------|
| log <sub>10</sub> RMR | Intercept                   | -1.58    | 0.19 | -8.36           | < 0.001         |
|                       | Sex: Male                   | 0.02     | 0.02 | 1.04            | 0.304           |
|                       | log <sub>10</sub> Body mass | 1.11     | 0.09 | 11.87           | < 0.001         |
| log <sub>10</sub> MMR | Intercept                   | -0.53    | 0.27 | -2.00           | 0.051           |
|                       | Sex: Male                   | 0.16     | 0.03 | 6.02            | < 0.001         |
|                       | log <sub>10</sub> Body mass | 0.99     | 0.13 | 7.56            | < 0.001         |
| log <sub>10</sub> AS  | Intercept                   | -0.56    | 0.32 | -1.76           | 0.084           |
|                       | Sex: Male                   | 0.18     | 0.03 | 5.82            | < 0.001         |
|                       | log <sub>10</sub> Body mass | 0.97     | 0.16 | 6.18            | < 0.001         |

## Chapter 5

---

# Developmental temperature and population factors affect variation and covariation of metabolic traits, with complex effects on growth rates in brown trout (*Salmo trutta*)

Louise C. Archer<sup>\*1,2</sup>, Stephen A. Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, W. Russell Poole<sup>3</sup>, Patrick Gargan<sup>4</sup>, Philip McGinnity<sup>1,3</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.

<sup>2</sup> Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland.

<sup>3</sup> Marine Institute, Furnace, Newport, Co. Mayo, Ireland.

<sup>4</sup> Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.

**Key words:** climate change, partial migration, anadromy, aquatic, brown trout, acclimation, *Salmo trutta*, metabolism

**\*Correspondence:** Louise Archer (l.archer@umail.ucc.ie)

**Contributions:** LA, TR, and PMG conceived the study. LA, SH, and LH collected data and contributed to experimental design. LA conducted statistical analysis and wrote the manuscript. All authors contributed to interpretation of results and revisions of the manuscript.

## Abstract

Metabolism defines the energetic cost of life, yet we still know relatively little about how intraspecific variation in standard metabolic rate (SMR), maximum metabolic rate (MMR), and aerobic scope (AS) arises and persists. Inconsistent associations between metabolic traits and fitness/performance metrics indicates that the optimal metabolic phenotype, or combination of metabolic traits, may be context-dependent. Here, we use a tank-rearing experiment to explore how environmental and population-specific factors influence the links between growth rates (a key performance trait) and metabolism in brown trout, a species that demonstrates considerable variation in migratory life histories. Growth rates and metabolic traits were measured in offspring from two populations that naturally vary in migratory tactics (one anadromous i.e. sea migratory, one non-anadromous), exposed to ~15 months of warming at 1.8 °C above natural temperatures. The anadromous population had overall higher SMR, MMR, and AS compared to the non-anadromous population, but both populations showed lower SMR in the warm treatment. We observed lower MMR and AS in the anadromous population in the warm temperature treatment, but not in the non-anadromous population. The variation in metabolic traits had complex implications for growth rates across the study that were dependent on population background and temperature. Lower SMR was associated with higher growth in the warm treatment, suggesting SMR reduction was an acclimation response to minimise maintenance costs under warm conditions. Higher MMR and AS were linked to higher growth rates in the anadromous population, potentially due to behavioural or life-history differences. A strong coupling between SMR and MMR or AS in this population further suggested that metabolic trait covariation was underpinned by the demands of fuelling a high-powered metabolism. Collectively, our study indicates that populations vary in their capacity to acclimate different metabolic traits in response to chronic warming and that the links between metabolic traits and growth rates are also population specific. In the wild, the effects of a given metabolic phenotype on growth rates and overall fitness are likely to depend on both environmental context and population background, and we

should consider both factors when forecasting population responses to warming.

## 35 **Introduction**

As the fundamental biological rate determining resource use and energy balance (Brown *et al.* 2016), metabolism underlies organism performance, and ultimately, fitness (Burton *et al.* 2011). Understanding the links between metabolism, performance, and environmental conditions is widely recognised  
 40 as being crucial to forecasting species' and population responses to global change (Seebacher *et al.* 2015), yet this is not necessarily straightforward. Despite considerable variation in metabolic traits apparent within and between populations, uncertainty still surrounds the drivers of such variation, or its consequences for fitness in different environmental contexts (Pettersen *et al.*  
 45 2018). It is imperative that we uncover the sources of metabolic variation, and assess how such variation can influence the performance of individuals, and hence the ability of populations to cope with global change.

The baseline energetic demands of ectotherms are defined by their standard metabolic rate (SMR), termed basal metabolic rate (BMR) in endotherms within  
 50 their thermoneutral zone. SMR represents the minimum energetic costs of maintaining tissues and homeostasis in an organism that is inactive, unstressed, and non-digestive (Chabot *et al.* 2016). The upper bounds of metabolism are set by the maximum metabolic rate (MMR), which is the highest rate of aerobic metabolism (i.e. oxygen transport and ATP production) that can be achieved  
 55 (Norin and Metcalfe 2019). Aerobic scope (AS) – the difference between an organism's SMR and MMR – determines the potential energy that can be allocated towards important functions including digestion, activity, growth and reproduction (Guderley and Pörtner 2010). SMR, the most frequently measured metabolic trait, shows significant inter-individual variation, with up to  
 60 threefold differences in SMR even among similarly sized and aged individuals from the same population (Burton *et al.* 2011; Konarzewski and Książek 2013). Although MMR (and consequently AS) is measured less often, the level of variation appears similar to that of SMR, once age and size are accounted for

(Hillman *et al.* 2013; Norin and Clark 2016). While SMR and MMR are often  
 65 correlated within species (Auer *et al.* 2017; Swanson *et al.* 2017), the relationship  
 between metabolic traits can also vary considerably (Cutts *et al.* 2002; Chappell  
*et al.* 2007; Wone *et al.* 2015; Norin *et al.* 2016; Barceló *et al.* 2016). This  
 substantial variation and covariation in metabolic phenotypes can be  
 understood in terms of ultimate mechanisms (evolved differences due to past  
 70 natural selection), or proximate mechanisms (differences due to developmental  
 variation or plastic responses to variable environments) (Tinbergen 1963;  
 Bateson and Laland 2013).

Given the likely importance of metabolism for fitness, the reason behind the  
 persistence of such variation is somewhat unclear, but one possibility is that the  
 75 optimum metabolic phenotype is context-dependent (Burton *et al.* 2011).  
 Variation in optimum phenotypes may occur across populations, *e.g.* due to  
 spatial variation in selection pressures such as temperature, food supply, or  
 habitat quality (Thomas *et al.* 2001; Angilletta *et al.* 2002). Within populations,  
 variation in optimal phenotypes might emerge through time via fluctuating  
 80 selection arising from temporal variation in environmental conditions (Schulte  
 2015). Moreover, at any given time, the optimum phenotype may vary between  
 different types of individuals within and among populations. For example,  
 within a population of three-spined sticklebacks *Gasterosteus aculeatus*,  
 migrants had higher standard and active metabolic rates than residents  
 85 (Tudorache *et al.* 2007), with similar differences among populations linked to  
 reduced selection on swimming performance in stream-resident populations  
 (Dalziel *et al.* 2012a, b). Variation in optimal phenotypes might also emerge  
 between different behavioural types *e.g.* shy *versus* bold individuals  
 (Dingemanse *et al.* 2004; Boon *et al.* 2007; Careau *et al.* 2008). Collectively, the  
 90 above ultimate mechanisms can lead to evolved (i.e. genetically based) variation  
 in metabolic traits being maintained within and across populations. Over time,  
 a decoupling of metabolic traits can also occur because each trait is subject to  
 subtly distinct selection pressures (Norin and Metcalfe 2019), which often act

in parallel but may also operate independently (Wone *et al.* 2015; Barceló *et al.* 95 2016).

Environmental factors are also instrumental in controlling metabolic rate variation from a proximate perspective. Dramatic changes in metabolic components have been reported in response to temperature (Fry 1971; Pörtner and Knust 2007; Clark *et al.* 2013), hypoxia (Dupont-Prinet *et al.* 2013; Norin *et al.* 100 *et al.* 2016), food availability (O'Connor *et al.* 2000; Van Leeuwen *et al.* 2011, 2012; Auer *et al.* 2015c), and habitat structure (Finstad *et al.* 2004; Millidine *et al.* 2006). Since aerobic metabolism is dependent on factors influencing oxygen demand and uptake (Fry 1971), temperature profoundly determines metabolic rates (Gillooly *et al.* 2001; Angilletta *et al.* 2002; Pörtner and Knust 2007; Brown 105 *et al.* 2016; Pettersen *et al.* 2018). However, the magnitude of any temperature-induced change in metabolism tends to decrease with exposure time as the animal becomes acclimated to the new temperature (Seebacher *et al.* 2015). Acute effects of temperature on metabolism are reasonably well researched, but we know less about the effects of chronic temperature exposure (i.e. shifts in 110 thermal regime lasting months to years) on different metabolic traits, despite such time scales being highly relevant in the context of climate change (Clark *et al.* 2013).

Predicting the effects of temperature is made more complicated because the various components of metabolism can show different sensitivities to 115 environmental conditions. There is some evidence for more plastic metabolic “floors” (SMR) than “ceilings” (MMR) in response to temperature, as documented in European perch *Perca fluviatilis* that reduced SMR after long-term warming with no compensatory adjustments seen in MMR (Sandblom *et al.* 2016). Indeed, many fish species appear to show minimal temperature-induced changes in AS (Lefevre 2016; Jutfelt *et al.* 2018), though this is not 120 always the case (Norin *et al.* 2014). Similar divergence in the relative responses of BMR and MMR (or cold-induced maximum metabolism) to temperature is apparent in endotherms (Nespolo *et al.* 2001; van de Ven *et al.* 2013; Dubois *et al.* 2016), indicating that decoupling of metabolic traits may be widespread.

125 While a positive relationship between SMR and MMR has been proposed under  
 the “increased intake” hypothesis – where a higher SMR maintains the  
 metabolic machinery that fuels high MMR (Biro and Stamps 2010; Burton *et al.*  
 2011) – covariation between metabolic components may be stronger, weaker, or  
 non-existent depending on environmental context (Killen *et al.* 2013; Norin *et*  
 130 *al.* 2016), and can vary between individuals (Norin and Malte 2011). It seems  
 that SMR and MMR might thus be subject to subtly different proximate (or  
 ultimate) constraints that might be revealed or masked by a given set of  
 environmental conditions (Killen *et al.* 2013; Norin and Metcalfe 2019).

While the fitness implications of metabolism are presumed to be considerable  
 135 because it shapes life histories (Stearns 1992), there is equivocal evidence for  
 effects of metabolic traits on various fitness metrics (Burton *et al.* 2011). Positive  
 relationships between SMR/RMR and growth (Yamamoto *et al.* 1998; McCarthy  
 2000), reproduction (Sadowska *et al.* 2013), and survival (Jackson *et al.* 2001)  
 imply fitness benefits of higher SMR that are in line with the “increased intake”  
 140 hypothesis. Yet SMR/BMR has also been negatively linked to growth (Álvarez  
 and Nicieza 2005; Norin and Malte 2011), reproduction (Blackmer *et al.* 2005)  
 and survival (Álvarez and Nicieza 2005; Larivée *et al.* 2010), supporting an  
 alternative “compensation” hypothesis, whereby a lower SMR is advantageous  
 for energy-saving purposes (Burton *et al.* 2011). Surprisingly little attention has  
 145 been paid to the associations between MMR or AS and fitness (Metcalf *et al.*  
 2016), even though AS has been proposed as the overarching physiological trait  
 governing fitness-related functions (Brown *et al.* 1993; Pörtner and Knust 2007;  
 Pörtner and Farrell 2008). Nonetheless, MMR appears to show similarly  
 inconsistent relationships with fitness (Hayes and O’Connor 1999; Boratyński  
 150 and Koteja 2009; Zub *et al.* 2014). Such inconsistencies are likely explained by  
 context-dependent fitness benefits of various metabolic trait phenotypes. For  
 example, higher SMR (or MMR) may only be beneficial when resources are  
 plentiful (Reid *et al.* 2011, 2012; Auer *et al.* 2015c; Killen *et al.* 2016) or  
 predictable (Hoogenboom *et al.* 2013), and have been found to have negative  
 155 (Bochdansky *et al.* 2005), or no effects (Álvarez and Nicieza 2005) when



resources are limited. Moreover, the fitness consequences of metabolic traits may depend on complex links between the larger metabolic phenotype (*e.g.* the coupling of SMR and MMR), and the environmental conditions encountered (Auer *et al.* 2015b).

160 Since the optimal metabolic phenotype (or combination of traits) varies across space/among populations as the environmental context changes, metabolic variation might be further associated with population-specific (*e.g.* evolved) differences in life history, such as migration tendency, yet relatively few studies have investigated these links. Salmonine fishes (salmon, trout, and charr) offer  
 165 great potential for exploration of the proximate and ultimate drivers of metabolic trait variation and covariation. In facilitating obligate freshwater spawning, salmonines exhibit diverse life-histories that encompass a continuum of migratory tactics, from individuals that remain resident in natal freshwater streams for their entire life cycle, to those that undertake spectacular migrations  
 170 to larger rivers, lakes, or the sea (Klemetsen *et al.* 2003). Facultative migration, where individuals in a population can display either migratory or non-migratory (resident) tactics (Chapman *et al.* 2011a), is proposed to be triggered by the relationship between an environmentally-sensitive trait and a genetically-variable threshold (Tomkins and Hazel 2007; Piche *et al.* 2008; Pulido 2011;  
 175 Buoro *et al.* 2012). While the exact trait controlling migration tactics remains unresolved (Ferguson *et al.* 2019), migration is often expressed as a consequence of energetic limitation in natal streams (Forseth *et al.* 1999). Variation in migratory life histories among individuals and populations is thus likely to be linked to variation in metabolic traits, *e.g.* anadromous (sea-migratory) Atlantic  
 180 salmon *Salmo salar* had higher SMR than non-migrants across three populations (Seppänen *et al.* 2010), and future migrants within a population had higher SMR than individuals that delayed migration (McCarthy 2000).

While variation in metabolic traits could have proximate/causal effects *i.e.* contribute to the emergence of migratory tactics among populations via the  
 185 threshold model, metabolic variation might persist as a compensatory adaptation in migratory or resident populations (via ultimate mechanisms) that

increases the benefits of migration/residency relative to costs (Hendry *et al.* 2004). For example, while larger body sizes and higher growth are associated with increased migration propensity (Jonsson 1985; Forseth *et al.* 1999; Morinville and Rasmussen 2003), survival-at-sea also increases with size (Armstrong *et al.* 2018). We might thus expect higher SMR, MMR, or AS to confer growth-mediated fitness benefits in migratory populations. In contrast, non-migratory populations must cope with unpredictable/patchy temperatures (Elliott 2000; Rutherford *et al.* 2004) and food availability typical of freshwater streams (Martin-Smith and Armstrong 2002), which might favour a lower SMR that minimises energy requirements.

Here, we explore the effects of temperature and population factors on metabolic traits and growth (a key fitness-associated trait) in experimentally reared FI offspring from two wild trout populations that naturally differ in migratory tactics. Specifically, we aimed to: (i) assess how long-term temperature elevation and population background affects metabolic traits (SMR, MMR, and AS); (ii) test if populations of variable migratory propensity show variation in metabolic responses to warming; (iii) test whether variation in metabolic traits influences growth performance; and (iv) investigate how relationships between metabolic traits and growth vary according to temperature and population background. We expected that increased temperature would result in compensatory responses in metabolic rates (lower SMR, and/or higher MMR and AS) and increased growth rates but that the migratory population would overall show higher metabolic rates and growth rates. We also expected that SMR, MMR, and AS would be positively related to growth, but the strength or direction of this relationship could vary between populations, and be magnified by warmer temperatures.

## Methods

### *Study populations and fish rearing*

We obtained brown trout brood stock from two wild populations in November 2015 by seine netting in the Erriff (53° 37' 0.00" N: 09° 40' 17.10" W) and the Burrishoole (53° 57' N: 09° 35' W) catchments in the west of Ireland (Figure S1).

Erriff broodstock were caught in the Tawnyard Lough (56 ha), an upland lake fed by the Glendavoch river and a number of smaller tributaries. Trout in the Tawnyard Lough primarily spawn in the Glenadavoch River, moving downstream as fry or parr to the Lough (covering several hundred metres to kilometres, depending on spawning location). A large run of out-migrating anadromous juveniles (smolts) are produced from Tawnyard Lough each year, with estimates of 500 to 3000 smolts enumerated annually at the outflow of the Lough (Gargan *et al.* 2016). An undetermined proportion of the population remain in the Lough, undergoing several years of freshwater growth before spawning in their natal stream, with local expertise indicating that the Tawnyard population overall has a strong anadromous component (broadly estimated as 50 – 60% of the wild population) (P. Gargan, *pers comm.*).

The Burrishoole brood stock were caught in the Srahrevagh River (*ca.* 7250 m<sup>2</sup> of salmonid habitat), a tributary of the Black River in the headwaters of the catchment that drains into Lough Feeagh, and ultimately Clew Bay. Although an anadromous life history occurred in the Burrishoole catchment up to 30 years ago (annual estimates of up to 2000 anadromous recruits), the anadromous trout run collapsed in the 1980s, coinciding with sea lice outbreaks from a salmon aquaculture facility established in the downstream estuary (Poole *et al.* 2007). Recent annual runs of up-migrating sea trout are typically in the low hundreds. Although the exact spawning locations of historic or contemporary anadromous individuals are uncertain, brood stock used in our study showed no signs of having undertaken a marine migration, i.e. were all non-anadromous fish. In summary, we consider offspring derived from the Tawnyard brood stock to have a strong anadromous background (hereafter termed the “anadromous background population”), and offspring from the Srahrevagh brood stock to have no recent anadromous background (termed the “non-anadromous background population”).

See Archer *et al.* (2019) and Chapter 3 for detailed description of crossing, fertilisation, and rearing procedures, described here in brief. Eggs from each female were fertilised by 1-2 males from the same source population and

incubated at the Burrishoole hatchery facility. Post-hatching, fry were transferred to a rearing facility at University College Cork (Aquaculture and Fisheries Development Centre), Ireland. Here, they were held in two 100L growth tanks (one per population) on a recirculating aquaculture system (RAS), maintained at a natural temperature regime typical of the west of Ireland under a constant photoperiod (12:12 hours of light: dark), until experimental treatments began. During this initial rearing period, fish were fed *ad libitum* with commercially available trout pellets (Skretting Ltd., Norway). In December 2016, 140 fry were allocated to one of four 203L capacity tanks in a larger experimental RAS (two tanks per population, initial  $n = 35$  per tank). Within the experimental RAS, LED lights above each tank mimicked the natural photoperiod of the source catchments. Water flowed continually through tanks, was treated with mechanical filtration, biofiltration, protein and UV skimming, and was mixed in a central sump tank. Mixed water from the sump was passed through one of two conditioning units, one that maintained a seasonal temperature regime typical of the west of Ireland, or one that was elevated to 1.8 °C ( $\pm 0.55$  SD) above the natural temperature regime. Thus, each tank received water from the same recirculating source, heated to one of two temperatures. Fish were fed recommended daily pellet rations for optimal growth calculated as a percentage of their body mass, with absolute rations adjusted monthly to account for changes in temperatures and body mass (Skretting Ltd., Norway). Water quality in the RAS (checked weekly) was consistently within acceptable levels for fish health, and great care was taken to ensure that all measured variables other than temperature regime (fish densities, feeding, photoperiod, lux, flow rates) were constant across tanks. Mortality was negligible during the experimental period, but fish were haphazardly culled ( $n = 20$ ) over the course of tank rearing for inclusion in parallel studies.

The study and all associated procedures were carried out with ethical approval from Health Products Regulatory Authority (HPRA) Ireland, under HPRA project license AEI9130/P034, and individual licenses AEI9130/1087,

280 AEI9I30/I200, AEI9I30/I201 and AEI9I30/I202 with all fish humanely euthanized under licence in April 2018.

#### *Temperature treatments*

To explore the effects of temperature and population factors on growth and metabolism, juvenile brown trout from each population were randomly  
 285 allocated to one of two temperature regimes in January 2016, with the populations reared separately to prevent emergence of dominance hierarchies. The two temperature regimes were: (i) cool temperature treatment: temperatures matched the natural, seasonally-varying temperatures of the source catchments; and (ii) warm temperature treatment: temperature elevated  
 290 to  $1.8\text{ }^{\circ}\text{C} \pm 0.55\text{ (SD)}$  above the cool treatment. The cool treatment ranged from  $5.9 - 16.4\text{ }^{\circ}\text{C}$  (mean temperature =  $10.8\text{ }^{\circ}\text{C} \pm 3.3\text{ SD}$ ) and the warm treatment ranged from  $7.5 - 18.2\text{ }^{\circ}\text{C}$  (mean temperature =  $12.6\text{ }^{\circ}\text{C} \pm 3.4\text{ SD}$ ). The  $1.8\text{ }^{\circ}\text{C}$  elevation in the warm treatment was chosen to reflect increases of  $1 - 3\text{ }^{\circ}\text{C}$  projected under climate change scenarios (IPCC 2014), but is within sub-lethal  
 295 ranges for brown trout (Forseth *et al.* 2009; Jonsson and Jonsson 2009). Temperature was increased by  $0.5\text{ }^{\circ}\text{C}$  per week when initiating treatments to minimise stress. Within each temperature treatment, 24-26 fish of each population were lightly anaesthetised with MS-222, and marked with a unique colour combination of visible implant elastomer (VIE) tags (Northwest Marine  
 300 Technology, USA) to allow for re-identification. Six individuals lost VIE tags over the course of the experiment, leaving  $n = 96$  individually identifiable fish.

#### *Data collection*

To calculate growth rates of VIE tagged individuals across the study period, the fork length (mm) and mass (g) of lightly anaesthetised fish was recorded in April,  
 305 June, July, August, September, and December 2017. In February 2018 (approximately 12 months after temperature treatments were established), eight fish from each population and temperature treatment combination underwent respirometry trials to measure metabolic traits ( $n = 32$  individuals in total).

### 310 *Measurement of MMR*

Whole-animal oxygen consumption ( $MO_2$ ) in animals operating at their maximum aerobic metabolic rate was used as proxy for MMR (Norin and Metcalfe 2019), following best practices outlined in Norin and Clark (2016). We used an exhaustive chase protocol following Norin and Clark (2016) to elicit

315 MMR in the same individuals that we measured for SMR. Prior to SMR measurements, each individual fish due to be measured for SMR later that day was first placed in an aerated 50L tank in a controlled-temperature (CT) chamber at  $7.9\text{ }^{\circ}\text{C} \pm 0.1\text{ SD}$ , and manually chased by hand until exhaustion. We determined exhaustion to occur when the fish were unresponsive (i.e. did not

320 exhibit burst swimming) to tactile stimulus (typically after 2 to 3 minutes of sustained chasing). Once exhausted, the fish was immediately transferred to a respirometry chamber in the same system used to measure SMR, the chamber was sealed, and oxygen decline within the closed chamber loop (recirculation pump operational) was recorded for a 60s measurement period. The time taken

325 to transfer fish to the chamber and begin recording oxygen measurements never exceeded 20s, ensuring that minimal recovery from the exhaustive chase procedure occurred before recording oxygen consumption.

### *Measurement of SMR*

The SMR of individual fish was determined using intermittent-flow

330 respirometry, following best practices outlined in Svendsen *et al.* (2016). SMR measurements took place overnight in a darkened CT chamber maintained at  $7.9\text{ }^{\circ}\text{C} \pm 0.1\text{ SD}$  (the mid-point between the cool and warm temperature treatments at the time of measurements). The respirometry system consisted of four acrylic respirometry chambers (1200 ml) (Loligo Systems, Viborg,

335 Denmark), submerged in a water bath, flushed with de-chlorinated water bubbled to 100% oxygen saturation by an air stone. PVC tubing (10mm diameter, non-permeable to oxygen) connected each individual chamber to two pumps (Eheim Ltd., Deizisau, Germany). A “flush” pump flushed fully oxygenated water through the chambers. A second “recirculation” pump

340 recirculated water in a closed loop through the chamber, whereby water exiting

the chamber was passed through a 10mm flow through oxygen cell (PreSens Ltd., Regensburg, Germany) that continually measured dissolved oxygen concentration, before being recirculated back to the same respirometry chamber via PVC tubing. Thus, individual oxygen consumption ( $MO_2$ , used as  
345 a proxy for SMR in fasted, inactive animals) was measured in repeated cycles that consisted of a flush period (flush pump operational) and a measurement period (recirculation pump operational) where oxygen level in each chamber was recorded at one second intervals to estimate oxygen decline (i.e. oxygen uptake). Each cycle consisted of 330s of flushing, and a measurement period of  
350 200 - 300s (to ensure sufficient  $O_2$  depletion for calculating  $MO_2$  in different-sized fish). We also allowed a 30s buffer period before recording oxygen uptake once the flush pump was switched off, to allow the chamber water and flush water to mix completely and reach an equilibrium oxygen saturation.

Fish were fasted for 28h prior to respirometry measurements to ensure  
355 individuals were in a post absorptive state (Cutts *et al.* 2002). Fish were placed in individual respirometry chambers between 11:00 and 12:00 each day, and were left to acclimatise for five hours, with chambers continually flushed with oxygen-saturated water during this acclimation period. SMR measurements began between 16:00 and 17:00, and ended between 09:00 and 10:00 the  
360 following morning, allowing for a minimum of 100 measurements of oxygen uptake per individual over a ~20-h period, during which time fish were not disturbed. Once SMR measurements had finished, fish were returned to their rearing tanks. To limit bacterial growth in the system, the entire respirometry set-up was washed with bleach after each overnight SMR respirometry trial. We  
365 also measured background (i.e. bacterial) respiration rates in each chamber on a daily basis by recording oxygen decline in empty chambers for one measurement cycle before fish entered the respirometry system, and again for one measurement cycle once the respirometry measurements had ended and fish were removed from the chambers.

### 370 *Measurement of MMR*

Whole-animal oxygen consumption ( $MO_2$ ) in animals operating at their maximum aerobic metabolic rate was used as proxy for MMR (Norin and Metcalfe 2019), following best practices outlined in Norin and Clark (2016). We used an exhaustive chase protocol following Norin and Clark (2016) to elicit

375 MMR in the same individuals that we measured for SMR. Prior to SMR measurements, each individual fish due to be measured for SMR later that day was first placed in an aerated 50L tank and manually chased by hand until exhaustion. We determined exhaustion to occur when the fish were unresponsive (i.e. did not exhibit burst swimming) to tactile stimulus (typically

380 after 2 to 3 minutes of sustained chasing). Once exhausted, the fish was immediately transferred to a respirometry chamber in the same system used to measure SMR, the chamber was sealed, and oxygen decline within the closed chamber loop (recirculation pump operational) was recorded for a 60s measurement period. The time taken to transfer fish to the chamber and begin

385 recording oxygen measurements never exceeded 20s, ensuring that minimal recovery from the exhaustive chase procedure occurred before recording oxygen consumption.

### *Statistical Analysis*

We estimated SMR ( $\text{mg O}_2 \text{ h}^{-1}$ ) by first calculating an individual's  $MO_2$  values

390 ( $\text{mg O}_2 \text{ h}^{-1}$ ) for each repeated measurement of oxygen uptake recorded during the overnight SMR respirometry trials.  $MO_2$  was calculated as the most consistent linear decline in oxygen recorded during each measurement loop, estimated by rolling regression using the *respR* package (Harianto and Carey 2019). We inspected all  $MO_2$  measurements to assess regression fit, and only

395 included  $MO_2$  values with an acceptable fit ( $R^2$  values  $> 0.90$ ) in subsequent SMR calculations, unless a clear linear trend was determined upon visual inspection of fit. Any background respiration in  $MO_2$  values was accounted for by subtracting the  $MO_2$  values calculated in empty chambers before and after each trial. We assumed background  $MO_2$  rates increased linearly over the

400 course of each trial (due to bacterial growth), and therefore allowed for a



dynamic background correction value (i.e. background  $MO_2$  increased overnight), calculated as:

$$MO_{2_{bg}} = bg_0 + (t \times bg)$$

where  $MO_{2_{bg}}$  is background  $MO_2$ , at a given measurement time point  $t$ , the  
 405 time elapsed since initiating overnight SMR measurements,  $bg_0$  and  $bg$  are  
 parameters (the intercept and slope respectively) estimated from the matrix  
 regression of the background oxygen uptake before and after experiments, as a  
 function of time elapsed. We then subtracted  $MO_{2_{bg}}$  from each value of  $MO_2$ ,  
 as calculated for an individual fish at successive time points during the  
 410 overnight SMR respirometry trials.  $MO_{2_{bg}}$  never exceeded more than 2% of total  
 $MO_2$  in all cases, confirming that background respiration rates remained low  
 throughout the study. Individual SMR was then calculated by taking the mean  
 of the lowest 10<sup>th</sup> percentile of background-corrected  $MO_2$  values recorded over  
 the 20-h SMR measurement period, then excluding outliers (values more than  
 415 two SD from this mean).

We estimated individual MMR ( $mg\ O_2\ h^{-1}$ ) using the *respR* package (Harianto  
 and Carey 2019) by calculating  $MO_2$  as the linear decline in oxygen in each  
 individual respirometry chamber in the 60s measurement period immediately  
 after the exhaustive chase protocol (i.e. extracting slopes from the linear  
 420 regression of oxygen concentration against time over a 60s period). We  
 calculated individual absolute aerobic scope (AS) ( $mg\ O_2\ h^{-1}$ ) as the difference  
 between MMR and SMR.

We explored variation in metabolic traits by first testing if the relationships  
 between body mass and SMR, MMR, or AS were affected by temperature and  
 425 population factors (Aims 1 and 2) using three general linear models (GLMs).  
 Body mass and metabolic rates were  $\log_{10}$  transformed to normalise and  
 linearise the data. Each GLM (normal errors) included either  $\log_{10}$  SMR,  $\log_{10}$   
 MMR, or  $\log_{10}$  AS as the response variable, and  $\log_{10}$  body mass, temperature  
 treatment and population background as explanatory variables. To test for  
 430 variation in the scaling of metabolic rates, each model included interactions

between  $\log_{10}$  body mass and temperature, and  $\log_{10}$  body mass and population, along with an interaction between temperature and population, and a three-way interaction term ( $\log_{10}$  body mass  $\times$  temperature  $\times$  population).

To explore how factors affected metabolic rate independent of mass, and to  
 435 avoid the pitfalls associated with solely using *P*-values (Halsey *et al.* 2015; Halsey 2019), we next tested for differences in mass-independent measures of SMR, MMR, and AS by estimation statistics (i.e. calculating effect sizes) using the *dabestr* package (Ho *et al.* 2019). The residuals of the linear relationship between  $\log_{10}$  body mass and SMR, MMR, and AS (all  $\log_{10}$  transformed) were  
 440 used to correct for body mass in these analyses (see Table SI for model summaries). Residual values of metabolic rates (rSMR, rMMR, rAS) gave mass-independent estimates of metabolic rates (individuals with positive residuals have a higher than expected metabolic rate for a given fish size, whereas negative residuals indicate a lower than expected rate). Effect sizes for mean  
 445 differences in rSMR, rMMR, and rAS were computed for all pairwise comparisons between all levels of temperature treatment and population background, and 95% confidence intervals (CIs) were constructed by bootstrapped resampling for 5,000 resamples.

To assess whether temperature treatment and population background  
 450 influenced the relationship between metabolic traits, we constructed three GLMs (normal errors). We used residual metabolic rates in these analyses to give mass-independent estimates of metabolic rates. The first GLM included rMMR as a response variable, with rSMR, temperature treatment, and population background included as explanatory variables. A second GLM  
 455 included rAS as the response, and rSMR, temperature treatment, and population background were similarly included as predictors. We included two-way interaction terms for rSMR  $\times$  temperature, and rSMR  $\times$  population, along with a three-way rSMR  $\times$  temperature  $\times$  population interaction to test for differences in covariation between MMR or AS with SMR. The third GLM  
 460 modelled rAS as function as rMMR, temperature treatment, and population

background, and similarly included the interaction terms  $rMMR \times \text{temperature}$ ,  $rMMR \times \text{population}$ , and  $rMMR \times \text{temperature} \times \text{population}$ .

Finally, we explored how metabolic rates, population and temperature treatments influence growth rates across the study period (Aims 3 and 4) within  
 465 a mixed effects modelling framework using the *nlme* package (Pinheiro *et al.* 2019). We estimated growth rates as the specific growth rate (% day<sup>-1</sup>) in terms of fork length ( $G_L$ ) between measurement periods according to:

$$G_L = 100 \times (\ln S_t - \ln S_i) / d$$

where  $S_t$  is the fork length at time  $t$ ,  $S_i$  is the initial fork length, and  $d$  is the time  
 470 elapsed (in days) between  $S_i$  and  $S_t$  (Hopkins 1992). We built three mixed effects models (normal errors) to examine how  $rSMR$ ,  $rMMR$  and  $rAS$  influenced growth rates in fish that underwent respirometry trials. Each model included time (continuous variable corresponding to months since start of experiment),  
 475 a third order polynomial term for time (to account for non-linearity through time), temperature treatment, population background, and metabolic rate (either  $rSMR$ ,  $rMMR$ , or  $rAS$ ) as fixed effects, and individual identity as a random effect to account for multiple growth rate measurements of individuals. We included two-way interactions between metabolic rate and temperature, metabolic rate and population, and temperature and population, and a three-  
 480 way interaction (metabolic rate  $\times$  temperature  $\times$  population). Since growth rate is size dependent (Nicieza and Álvarez 2009; Killen 2014), we included initial fork length as a covariate in the models, and we accounted for temporal autocorrelation of growth rates by modelling an autoregressive error structure as a first order lag function of time. To test if metabolic rate effects were  
 485 consistent across the experiment, we constructed three additional mixed effect models as described above, but included an additional interaction between metabolic rate and time.

We used likelihood ratio tests (LRT) to assess statistical significance of predictor variables for all models at a 5% alpha level. Non-significant interaction terms  
 490 were excluded to interpret main effects. For the estimation statistics approach,

we considered an estimated difference in means between groups to exist (i.e. was significant) if the 95% CI of the effect size did not include zero. Marginal  $R^2$  values for mixed effect models were calculated using the *MuMIn* package (Barton 2019). Analysis was carried out in R version 3.6.0 (R Core Team 2019) and all models were checked against assumptions of the given model (independence, non-normality of residuals, heteroscedasticity and multicollinearity).

## Results

### *Variation in metabolism*

Whole-animal SMR, MMR, and AS varied between temperature treatments and population background (see Table 1 for a summary of mean values of metabolic rates for each temperature and population combination).

As expected,  $\log_{10}$  SMR increased significantly with  $\log_{10}$  body mass ( $\chi^2 = 31.79$ ,  $df = 1$ ,  $P < 0.001$ , Table S2), but we did not find any significant interactions between body mass, temperature, and population ( $\chi^2 = 0.31$ ,  $df = 1$ ,  $P = 0.580$ ), between body mass and temperature ( $\chi^2 = 0.11$ ,  $df = 1$ ,  $P = 0.735$ ), or between body mass and population ( $\chi^2 = 0.13$ ,  $df = 1$ ,  $P = 0.716$ ). The temperature  $\times$  population interaction was also non-significant ( $\chi^2 = 0.01$ ,  $df = 1$ ,  $P = 0.985$ ), but the main effects of temperature treatment ( $\chi^2 = 9.55$ ,  $df = 1$ ,  $P = 0.002$ ), and population background ( $\chi^2 = 5.05$ ,  $df = 1$ ,  $P = 0.025$ ) were significant. For a given size, fish in the warm temperature treatment had lower SMR values, as did fish from the non-anadromous population (Figure 1A, Table S2). Similarly, estimation statistics showed fish in the warm temperature treatment had lower mass-independent SMR (lower rSMR) than those from the cool treatment (Figure 1B, Table 2). The non-anadromous population also tended to have lower rSMR at both temperatures, although this difference was less apparent in the warm temperature treatment, where the 95% CIs for the mean difference in rSMR between populations overlapped zero (Figure 1C, Table 2).

**Table 1:** Mean values and associated standard deviations (SD) for the length (mm), mass (g), standard metabolic rate (SMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), maximum

metabolic rate (MMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), and aerobic scope (AS) ( $\text{mg O}_2 \text{ hr}^{-1}$ ) of brown trout ( $n = 32$ ) offspring derived from two wild populations (AB = Anadromous background, Non-AB = Non-anadromous background). Fish were measured for metabolic traits in February 2018 after ~12 months of experimental rearing at one of two temperature treatments (Cool = natural temperature regime, Warm =  $1.8^\circ\text{C}$  above natural temperature regime).

| Temperature,<br>Population | Length<br>(mean $\pm$ SD) | Mass<br>(mean $\pm$ SD) | SMR<br>(mean $\pm$ SD) | MMR<br>(mean $\pm$ SD) | AS<br>(mean $\pm$ SD) |
|----------------------------|---------------------------|-------------------------|------------------------|------------------------|-----------------------|
| Cool, AB                   | 218.50 $\pm$ 11.30        | 119.00 $\pm$ 23.00      | 5.97 $\pm$ 1.32        | 54.00 $\pm$ 9.60       | 48.04 $\pm$ 8.65      |
| Warm, AB                   | 222.88 $\pm$ 6.90         | 133.59 $\pm$ 17.13      | 5.57 $\pm$ 1.34        | 45.57 $\pm$ 10.89      | 40.00 $\pm$ 10.29     |
| Cool, Non-AB               | 201.75 $\pm$ 8.22         | 108.58 $\pm$ 14.66      | 4.80 $\pm$ 1.02        | 36.83 $\pm$ 8.32       | 32.04 $\pm$ 8.01      |
| Warm, Non-AB               | 202.75 $\pm$ 19.95        | 108.10 $\pm$ 29.88      | 3.95 $\pm$ 1.03        | 35.16 $\pm$ 8.79       | 31.21 $\pm$ 8.08      |


After accounting for the effects of individual variation in  $\log_{10}$  body mass on  $\log_{10}$  MMR ( $\chi^2 = 11.13$ ,  $df = 1$ ,  $P = 0.001$ , Table S2), there were no significant interactions between body mass, temperature, and population ( $\chi^2 = 0.01$ ,  $df = 1$ ,  $P = 0.992$ ), between body mass and temperature ( $\chi^2 = 0.06$ ,  $df = 1$ ,  $P = 0.810$ ), between body mass and population ( $\chi^2 = 0.87$ ,  $df = 1$ ,  $P = 0.352$ ), or between temperature and population ( $\chi^2 = 1.15$ ,  $df = 1$ ,  $P = 0.284$ ). We detected significant main effects of temperature treatment ( $\chi^2 = 4.10$ ,  $df = 1$ ,  $P = 0.043$ ) and population background ( $\chi^2 = 10.59$ ,  $df = 1$ ,  $P = 0.001$ ) on  $\log_{10}$  MMR. Overall, MMR (for a given size) was lower in the warm treatment, and in the non-anadromous background population (Figure 2A, Table S2). When considering size-independent MMR (rMMR), a lower rMMR in the warm treatment was evident only fish from the anadromous background population (Figure 2B, C, Table 2), and population differences in rMMR were seen in the cool treatment only (Figure 2B, C).

**Table 2:** Effect sizes ( $\Delta$ ) and associated 95% confidence intervals (CIs) for differences in mean residual standard metabolic rate (rSMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), residual maximum metabolic rate (rMMR) ( $\text{mg O}_2 \text{ hr}^{-1}$ ), and residual aerobic scope (rAS) ( $\text{mg O}_2 \text{ hr}^{-1}$ ) of brown trout ( $n = 32$ ) offspring derived from two wild populations (AB = anadromous background population, non-AB = non


anadromous background population), exposed to two temperature treatments (Cool = natural temperature regime, Warm = 1.8 °C above natural temperature regime). CIs were constructed by non-parametric bootstrap resampling (5,000  
550 resamples).

| Mean difference ( $\Delta$ )     | $\Delta$ rSMR<br>(95% CI)  | $\Delta$ rMMR<br>(95% CI)  | $\Delta$ rAS<br>(95% CI)   |
|----------------------------------|----------------------------|----------------------------|----------------------------|
| <i>Warm AB – Cool AB</i>         | -0.084<br>(-0.153; -0.020) | -0.116<br>(-0.192; -0.029) | -0.121<br>(-0.206; -0.031) |
| <i>Warm non-AB – Cool non-AB</i> | -0.075<br>(-0.134; -0.005) | -0.012<br>(-0.073; 0.057)  | -0.002<br>(-0.073; 0.076)  |
| <i>Cool non-AB – Cool AB</i>     | -0.060<br>(-0.118; -0.005) | -0.144<br>(-0.217; -0.068) | -0.157<br>(-0.241; -0.073) |
| <i>Warm non-AB – Warm AB</i>     | -0.050<br>(-0.125; 0.023)  | -0.040<br>(-0.122; 0.026)  | -0.038<br>(-0.124; 0.037)  |

We detected similar trends in AS, with non-significant interactions between body mass, temperature, and population ( $\chi^2 = 0.01$ ,  $df = 1$ ,  $P = 0.942$ ), between body mass and temperature ( $\chi^2 = 0.04$ ,  $df = 1$ ,  $P = 0.834$ ), between body mass  
555 and population ( $\chi^2 = 0.93$ ,  $df = 1$ ,  $P = 0.336$ ), and between temperature and population ( $\chi^2 = 1.27$ ,  $df = 1$ ,  $P = 0.260$ ). After accounting for the effects of  $\log_{10}$  body mass on  $\log_{10}$  AS ( $\chi^2 = 7.81$ ,  $df = 1$ ,  $P = 0.005$ , Table S2), the main effect of temperature treatment was also non-significant ( $\chi^2 = 2.98$ ,  $df = 1$ ,  $P = 0.084$ ). Population background significantly affected  $\log_{10}$  AS ( $\chi^2 = 9.51$ ,  $df = 1$ ,  $P =$   
560 0.002), whereby AS (for a given size) was lower in the non-anadromous background population (Figure 3A, Table S2). However, rAS tended to be lower in the warm treatment in the anadromous background population (Figure 3B, C, Table 2), but not in the non-anadromous population. Population differences in rAS were strongest in the cool treatment (Figure 3C).




**Figure 1:** (A) Body mass scaling of standard metabolic rate (SMR) of brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental temperature treatments: a natural temperature regime (Cool); and temperature elevated 1.8°C above the cool regime (Warm). (B) Residual SMR values (body mass corrected) for each population background and temperature treatment combination. Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (C) Cumming estimation plots with effect sizes shown as black dots (i.e. the mean differences in rSMR among the groups), the distributions (shaded curves) and 95% confidence intervals (back bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).



**Figure 2:** (A) Body mass scaling of maximum metabolic rate (MMR) of brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental temperature treatments: a natural temperature regime (Cool); and temperatures elevated 1.8°C above the Cool regime (Warm). (B) Residual MMR values (body mass corrected) for each population background and temperature treatment combination. Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (C) Cumming estimation plots with effect sizes shown as black dots (i.e. the mean differences in rMMR among the groups), the distributions (shaded curves) and 95% confidence intervals (back bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).






**Figure 3:** (A) Body mass scaling of aerobic scope (AS) of brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish were reared under two experimental temperature treatments: a natural temperature regime (Cool); and temperatures elevated 1.8°C above the cool regime (Warm). (B) Residual AS values (body mass corrected) for each population background and temperature treatment combination. Black vertical bars represent the standard deviation around the mean (shown as a gap in the bars), and sample size is shown as “N =”. (C) Cumming estimation plots with effect sizes shown as black dots (i.e. the mean differences in rAS among the groups), the distributions (shaded curves) and 95% confidence intervals (back bars) of the effect sizes obtained from non-parametric bootstrap resampling (5,000 resamples).

*Relationships between metabolic traits*

570 After adjusting for body mass, rMMR was positively related to rSMR, with a significant interaction between rSMR and population ( $\chi^2 = 5.16$ ,  $df = 1$ ,  $P = 0.023$ ) whereby the anadromous population had a higher rMMR for a given rSMR (Figure 4A, Table S3). The two-way interactions between temperature and rSMR ( $\chi^2 = 0.63$ ,  $df = 1$ ,  $P = 0.429$ ), and between temperature and population  
 575 ( $\chi^2 = 0.25$ ,  $df = 1$ ,  $P = 0.618$ ), and the three-way interaction between temperature, population, and rSMR ( $\chi^2 = 0.64$ ,  $df = 1$ ,  $P = 0.425$ ) were all non-significant. We did not detect a significant main effect of temperature ( $\chi^2 = 1.22$ ,  $df = 1$ ,  $P = 0.270$ ).

The positive relationship between rAS and rSMR varied according to a  
 580 significant rSMR  $\times$  population interaction ( $\chi^2 = 5.02$ ,  $df = 1$ ,  $P = 0.025$ ), whereby the anadromous population had a higher rAS for a given rSMR (Figure 4B, Table S3). The rAS  $\times$  temperature interaction ( $\chi^2 = 0.66$ ,  $df = 1$ ,  $P = 0.420$ ), the temperature  $\times$  population interaction ( $\chi^2 = 0.28$ ,  $df = 1$ ,  $P = 0.597$ ), and the rSMR  $\times$  temperature  $\times$  population interaction ( $\chi^2 = 0.66$ ,  $df = 1$ ,  $P = 0.418$ ) were all  
 585 non-significant. The main effect of temperature was also non-significant ( $\chi^2 = 1.17$ ,  $df = 1$ ,  $P = 0.276$ ).

There was a strong positive relationship between rAS and rMMR (Figure 4C,  $\chi^2 = 2490.9$ ,  $df = 1$ ,  $P < 0.001$ ). We did not detect significant interactions between rMMR and temperature ( $\chi^2 = 0.50$ ,  $df = 1$ ,  $P = 0.479$ ), between rMMR and  
 590 population ( $\chi^2 = 1.41$ ,  $df = 1$ ,  $P = 0.234$ ), or between rMMR, temperature, and population ( $\chi^2 = 0.26$ ,  $df = 1$ ,  $P = 0.613$ ). The temperature  $\times$  population interaction was also non-significant ( $\chi^2 = 0.47$ ,  $df = 1$ ,  $P = 0.493$ ). While the main effect of population was non-significant ( $\chi^2 = 1.46$ ,  $df = 1$ ,  $P = 0.227$ ), the main effect of temperature was significant ( $\chi^2 = 6.23$ ,  $df = 1$ ,  $P = 0.013$ ), whereby fish  
 595 in the warm treatment had a marginally higher rAS for a given rMMR (Figure 4C, Table S3).



**Figure 4:** Relationships between: (A) Residual standard metabolic rate (rSMR) and residual maximum metabolic rate (rMMR); (B) rSMR and residual aerobic scope (rAS); and (C) rMMR and rAS for brown trout offspring derived from an anadromous background population (AB) and a non-anadromous background population (non-AB). Fish experienced two temperature treatments: a natural temperature regime (Cool), or temperatures elevated 1.8°C above the cool regime (Warm). Residual (size-independent) values for metabolic rates were estimated from the linear relationship between each metabolic rate (SMR, MMR, or AS) as a function of body mass (both log<sub>10</sub>-transformed).

#### *Metabolic traits and growth*

Specific growth rates of fish included in respirometry trials varied non-linearly through time (polynomial term for time:  $\chi^2 = 134.32$ ,  $df = 3$ ,  $P < 0.001$ ; Figure 5A). After controlling for significant negative effects of initial body size on growth ( $\chi^2 = 14.23$ ,  $df = 1$ ,  $P < 0.001$ ; Table S4), the mixed effects model describing the effects of rSMR on specific growth rate (marginal  $R^2 = 0.69$ ) retained significant two-way interactions between rSMR and temperature, and between temperature and population (Table 3). Overall, the negative rSMR  $\times$  temperature term indicated that in the warm treatment, higher rSMR were associated with lower growth rates (Figure 5B, Table S4). The negative interaction between temperature and population indicated that non-anadromous population had marginally lower growth rates in the warm treatment (Figure 5A, B). Neither the interaction term for rSMR  $\times$  temperature


× population, nor the term for rSMR × population was significant in the model (Table 3).

The mixed effects model describing the effects of rMMR on specific growth rate (marginal  $R^2 = 0.67$ ) retained a significant two-way interaction between rMMR and population (Table 3), once the effects of initial size and time were accounted for (effect of time:  $\chi^2 = 133.15$ ,  $df = 3$ ,  $P < 0.001$ ; and of initial size:  $\chi^2 = 14.50$ ,  $df = 1$ ,  $P < 0.001$ , Table S4). The anadromous background population showed a weak, positive effect of rMMR on growth, but the relationship was negative in the non-anadromous population (Figure 5A, C, and Table S4). Interaction terms for rMMR × temperature × population, for rMMR × temperature, and for temperature × population were not significant in the model describing growth rate as a function of rMMR (Table 3).

The model describing the effects of rAS on specific growth rate (marginal  $R^2 = 0.69$ ) similarly retained a significant two-way interaction between rAS and population (Table 3), once the effects of initial size and time were accounted for (effect of time:  $\chi^2 = 132.99$ ,  $df = 3$ ,  $P < 0.001$ ; and of initial size:  $\chi^2 = 12.52$ ,  $df = 1$ ,  $P < 0.001$  Table S4). The effect of rAS on growth was positive in the anadromous population but negative in the non-anadromous population (Figure 5A, D, Table S4). The model did not include significant interaction terms for rAS × temperature × population, for rAS × temperature, or for temperature × population (Table 3).

The additional set of mixed effect model analyses indicated that relationships between metabolic rates and growth were variable through time (Figure S2, Table S5) (LRT: rSMR × polynomial term for time:  $\chi^2 = 38.60$ ,  $df = 3$ ,  $P < 0.001$ ; rMMR × polynomial term for time:  $\chi^2 = 10.45$ ,  $df = 3$ ,  $P = 0.015$ ; rAS × polynomial term for time:  $\chi^2 = 8.80$ ,  $df = 3$ ,  $P = 0.032$ ).

See Supporting Information for summaries of all GLM and mixed effects model coefficients.



**Figure 5:** (A) Coefficient estimates ( $\pm$  95% confidence intervals) from the mixed effect models describing the effects of residual metabolic rate components (SMR = standard metabolic rate, MMR = maximum metabolic rate, and AS = aerobic scope, temperature treatment (Cool and Warm), and population background (AB = anadromous background, Non-AB = non-anadromous background) on specific growth rates of brown trout obtained from repeated measurements across the study (“Time” = months since initiating treatments). The mixed models were used to predict specific growth rates of brown trout in response to marginal effects of (B) rSMR, temperature treatment, and population background, (C) marginal effects of rMMR, temperature treatment, and population background and (D) marginal effects of rAS, temperature treatment, and population background. Shaded regions in (B), (C), and (D) show the 95% confidence intervals for the predictions. Growth rates were predicted at mean values for the remaining explanatory variables in the models.

640 **Table 3:** Results of the mixed effect model analysis for specific growth rate trajectories (% day<sup>-1</sup>) of brown trout that were measured for standard metabolic rate (SMR), maximum metabolic rate (MMR) and aerobic scope (AS). The results of the model selection procedure on interaction terms are given (significance assessed at  $P < 0.05$ ), and the selected model for specific growth rate as a function of a given metabolic component is highlighted in bold. The models included a random effect of individual identity and a first-order autoregressive correlation structure with respect to time was modelled (“Time” = months since beginning of treatment).

| Model terms  | df | AIC     | logLik | L-ratio | P-value |
|--|----|---------|--------|---------|---------|
| rSMR*Temperature*Population + poly(Time, 3) + Length                                 | 15 | -487.46 | 258.73 |         |         |
| rSMR*Temperature + rSMR*Population + Temperature*Population + poly(Time, 3) + Length | 14 | -486.41 | 257.20 | 3.05    | 0.081   |
| rSMR*Population + Temperature*Population + poly(Time, 3) + Length                    | 13 | -483.75 | 254.88 | 4.66    | 0.031   |
| <b>rSMR*Temperature + Temperature*Population + poly(Time, 3) + Length</b>            | 13 | -487.76 | 256.88 | 0.64    | 0.422   |
| rSMR*Temperature + rSMR*Population + poly(Time, 3) + Length                          | 13 | -482.70 | 254.35 | 5.71    | 0.017   |
| rMMR*Temperature*Population + poly(Time, 3) + Length                                 | 15 | -486.17 | 258.09 |         |         |
| rMMR*Temperature + rMMR*Population + Temperature*Population + poly(Time, 3) + Length | 14 | -488.12 | 258.06 | 0.05    | 0.818   |
| rMMR*Population + Temperature*Population + poly(Time, 3) + Length                    | 13 | -489.95 | 257.97 | 0.17    | 0.678   |
| rMMR*Temperature + Temperature*Population + poly(Time, 3) + Length                   | 13 | -484.62 | 255.31 | 5.50    | 0.019   |
| rMMR*Temperature + rMMR*Population + poly(Time, 3) + Length                          | 13 | -488.36 | 257.18 | 1.76    | 0.184   |
| <b>rMMR*Population + Temperature + poly(Time, 3) + Length</b>                        | 12 | -488.00 | 256.00 | 4.13    | 0.127   |
| rAS*Temperature*Population + poly(Time, 3) + Length                                  | 15 | -487.10 | 258.55 |         |         |
| rAS*Temperature + rAS*Population + Temperature*Population + poly(Time, 3) + Length   | 14 | -489.10 | 258.55 | 0.01    | 0.982   |
| rAS*Population + Temperature*Population + poly(Time, 3) + Length                     | 13 | -490.67 | 258.33 | 0.43    | 0.513   |
| rAS*Temperature + Temperature*Population + poly(Time, 3) + Length                    | 13 | -485.32 | 255.66 | 5.78    | 0.016   |
| rAS*Temperature + rAS*Population + poly(Time, 3) + Length                            | 13 | -489.73 | 257.87 | 1.36    | 0.243   |
| <b>rAS*Population + Temperature + poly(Time, 3) + Length</b>                         | 12 | -488.70 | 256.35 | 4.40    | 0.111   |

645

## Discussion

Metabolism defines the energetic balance of organisms, with considerable implications for performance, and ultimately, fitness. Yet despite widespread variation in metabolic traits, we know surprisingly little about how or why such variation persists, or how metabolic variation affects individual performance. Here, we explored the intrinsic and extrinsic causes of variation in metabolism in facultatively migratory brown trout by experimentally rearing two populations (that naturally differ in anadromy tactics) under a temperature regime that simulated a 1.8°C increase above natural temperatures, in line with climate change projections. Both populations had lower SMR in the warm temperature treatment, with lower MMR and AS also seen in the anadromous population in the warm temperature treatment, a difference that was not evident in the non-anadromous population. The variation in metabolic traits had complex implications for growth rates across the study that varied by population background and temperature. Lower SMR were associated with higher growth in the warm treatment, whereas lower MMR and AS were linked to higher growth rates in the non-anadromous population. Overall, our study indicates that populations may show variable responses to changing environmental conditions in terms of metabolic traits, with potential fitness consequences in a warming world.

### *Effects of temperature on metabolism*

The lower SMR displayed by both populations in the warm treatment suggests that adjustment of this key physiological trait is a plastic, or acclimation, response to chronic warming. While acute warming is well known to cause an initial increase in ectotherm metabolic rates (Clarke and Johnston 1999; Angilletta *et al.* 2002), exposure over longer time scales (i.e. those comparable to the 15 months of warming in our study) tends to reduce the magnitude of the response because acclimation occurs (Seebacher *et al.* 2015). The reduction in SMR we observed supports the potential for thermal compensation by way of the “plastic floors” hypothesis, where a lower SMR at warmer temperatures is beneficial because it reduces maintenance costs (Sandblom *et al.* 2016). Such

acclimation responses in fish can be considerable, *e.g.* Atlantic halibut (*Hippoglossus hippoglossus*) acclimated at warmer temperatures (ranging from 5-18°C) for 14 weeks showed stable SMR from 10-16°C (Gräns *et al.* 2014).  
 680 Acclimation capacity has been generally linked to increased resilience to environmental change (Magozzi and Calosi 2015; Seebacher *et al.* 2015). However, while temperatures are broadly projected to increase, more extreme and frequent warming events are also forecast (Meehl and Tebaldi 2004; Seneviratne *et al.* 2014). The adaptiveness of a given flexible response will thus  
 685 greatly depend on both the pattern of fluctuations in temperatures, and the speed at which individuals can alter their phenotype (Reed *et al.* 2010).

#### *Population effects on metabolism*

In addition to temperature-induced changes, we observed variation in SMR between our study populations that matched our expectations of lower baseline  
 690 energetic requirements in the non-anadromous population. Metabolic traits are evolvable (Rønning *et al.* 2007; Nilsson *et al.* 2009; Pettersen *et al.* 2018) and can respond to selection over relatively short periods (Sadowska *et al.* 2015). As such, population-level variation could reflect genetic differences or plastic/acclimation responses related to life-history differences between  
 695 populations. For example, a lower SMR could be optimal for the non-anadromous population because freshwater systems have relatively low productivity (Gross *et al.* 1988), whereas higher SMR in the anadromous population might facilitate rapid growth to reach sufficient smolt size to successfully migrate (Ward and Slaney 1988), particularly if SMR and MMR are  
 700 linked under the “increased intake” hypothesis (Burton *et al.* 2011). The higher MMR and AS we observed in the anadromous population might similarly reflect evolutionary change or plasticity driven by the demands of migration. MMR and AS is higher in anadromous sockeye salmon *Oncorhynchus nerka* populations that undertake more challenging migrations (Lee *et al.* 2003; Eliason *et al.* 2011),  
 705 whereas relatively lower MMR and AS has been reported in resident *versus* migratory three-spine stickleback populations, linked to weak selection on swimming performance (Dalziel *et al.* 2012a, b). Higher SMR and MMR might



subsequently be favoured in the marine environment by facilitating predator avoidance (*e.g.* locomotor capacity) (Plaut 2001; Killen *et al.* 2015; Eliason and  
 710 Farrell 2016). We also cannot rule out that metabolic traits may be reflective of environmental constraints specific to each population, *e.g.* counter gradient selection driven by poor/fluctuating resources experienced by the anadromous population, whereby a higher AS maximises consumption and growth when opportunities arise (Álvarez *et al.* 2006).

715 Intriguingly, while both of our study populations showed similar reductions in SMR in the warm temperature treatment, we detected variable responses in size-independent measures of MMR and AS with warming. Similar MMR and AS in the non-anadromous populations in both warm and cool treatments suggested that aerobic performance in this population was either unaffected by  
 720 warming, or else showed a heightened acclimation response to the temperature increase than the anadromous population (which had lower MMR and AS in the warm *versus* cool treatment). Relatively little is known about the response of MMR or AS to chronic temperature increases (Schulte *et al.* 2011; Clark *et al.* 2013), but the population-level variation we observed supports mounting  
 725 evidence that effects of long-term warming may vary considerably between and potentially within species. For example, thermal compensation in AS by way of acclimation in MMR has previously been seen in Atlantic halibut (Gräns *et al.* 2014). However, tropical barramundi *Lates calcarifer* showed reduced MMR and AS after acclimation to high temperatures, despite observed increases in MMR  
 730 after acute exposure to warming (Norin *et al.* 2014). Similarly, a small decline in MMR was observed in shorthorn sculpin *Myoxocephalus scorpius* after eight weeks of warming, even though a reduction in SMR partially restored AS by the end of the acclimation period (Sandblom *et al.* 2014).

Since a high AS is generally considered to increase individual performance and  
 735 fitness (Pörtner and Farrell 2008; Biro and Stamps 2010), and is a trait linked with migration effort (Eliason *et al.* 2011; Clark *et al.* 2011) it is initially unclear why the anadromous population did not appear capable of maintaining MMR. This implies there might be costs associated with maintaining a high MMR that

are specific to the anadromous population, in other words, temperature-induced flexibility in MMR may not be adaptive in this population *e.g.* if environmental fluctuations are rapid/extreme and the speed of acclimation is slow, the benefits of plasticity may be limited (Reed *et al.* 2010). Alternatively, the reduction in SMR may have caused an inevitable reduction in MMR, which is in line with the “increased intake” hypothesis underpinning energetic capacity in the anadromous population (Burton *et al.* 2011). This seems likely given the strong coupling we observed between SMR and MMR, which was absent in the non-anadromous population.

#### *Consequences of metabolic rate variation for growth*

The variation in metabolic traits we observed had consequences for growth, which depended largely on temperature treatment in the case of SMR. Lower SMR were associated with higher growth in warm, but not in cool treatments, supporting a context-dependency to the fitness consequences of a given SMR (Burton *et al.* 2011). However, it is important to note that growth rate is just one component of fitness, and might not always map positively or linearly onto fitness (Arendt 1997; Blanckenhorn 1998). Nonetheless, in this case, a lower SMR under warm conditions was likely beneficial for growth because maintenance energy costs were reduced, analogous to reductions in SMR facilitating higher growth or energy storage when food is limited (Naya *et al.* 2007; Auer *et al.* 2015c, 2016; Zeng *et al.* 2017). However, we also detected a complex interaction between SMR and time, indicating that the relationship between SMR and growth can further vary according to temporal factors. For example, the negative association between warm temperatures and SMR on growth may have been strongest during periods when temperatures exceeded the optimal ranges for growth in trout *e.g.* during the summer months (Elliott *et al.* 1995; Ojanguren *et al.* 2001). Moreover, while relative metabolic rates of salmonines tend to be stable through time (Nespolo and Franco 2007; Seppänen *et al.* 2010), absolute metabolic rates can vary considerably depending on a suite of environmental factors (Metcalf *et al.* 2016; Norin and Clark 2016). As such, the link between SMR, growth, and temperature may also

770 depend on additional factors, *e.g.* food supply (Zeng *et al.* 2018), which in turn show fluctuations in time and space (Álvarez and Nicieza 2005; Robertsen *et al.* 2014).

A similarly complex mapping emerged between both MMR and AS and growth rate, which depended on population background and varied through time. 775 Higher MMR and AS were associated with slightly higher growth rates in the anadromous population (independent of temperature), but this relationship was reversed in the non-anadromous population. Food intake (Auer *et al.* 2015a), and digestive/assimilation capacity (Alsop and Wood 1997) increase with MMR and AS, which presumably underlies the positive correlation 780 between MMR, AS, and growth in the anadromous population. Higher growth (relative to a non-anadromous population) has previously been described in fish originating from the same source catchment as the anadromous-background population in the present study (Archer *et al.* 2019). Fast juvenile growth has been associated with the anadromy lifestyle in general (Forseth *et al.* 1999), 785 appearing to be partially underpinned here by a higher metabolic scope. This implies that while a high metabolic scope is optimal in migratory populations, (perhaps because MMR and AS are linked to swimming performance (Dalziel *et al.* 2012b), and migration effort/performance (Eliason *et al.* 2011; Clark *et al.* 2011)), growth rates in non-anadromous populations are not limited by aerobic 790 capacity.

Observed differences in the strength of the association between both MMR and AS and growth, and its variability through time, reflect inconsistent outcomes among studies that have previously assessed the relationship between metabolism and fitness-related traits (Metcalf *et al.* 2016). In this case, 795 behavioural differences between populations may have altered the magnitude of the association between MMR and AS and growth. For example, migratory fish populations (and individuals) typically show higher levels of activity, boldness, and aggression (Metcalf *et al.* 1995; Lahti *et al.* 2001; Chapman *et al.* 2011b), which may in turn cause fish from anadromous populations to operate 800 closer to MMR more frequently. Thus, a higher MMR and AS is likely be more

advantageous in terms of maximising growth (Burton *et al.* 2011), but the association may be weaker, or absent in the non-anadromous population, which potentially approaches MMR infrequently. There is a dearth of knowledge regarding how often individuals approach their MMR, and whether this varies  
805 among individuals or populations, though existing studies suggest considerable variation is likely (Murchie *et al.* 2011; Seebacher *et al.* 2013; Killen *et al.* 2014). The negative association between MMR and growth in the non-anadromous population implies a stronger influence of SMR on growth, supporting the “compensation” hypothesis as a factor underpinning metabolic variation in this  
810 population (Burton *et al.* 2011) (particularly when considered alongside the weaker correlation between SMR and MMR in the non-anadromous population). Alternatively, variability in the relationship between MMR and AS and growth between populations, might reflect differences in the allocation of surplus resources (fuelled by AS) towards other competing functions besides  
815 somatic growth, *e.g.* lipid accumulation and gonadal development (Jonsson *et al.* 2013), or competitive/aggressive interactions (Killen *et al.* 2014). This could also explain temporal variability in the MMR and AS – growth relationship, where the benefits of a higher MMR and AS may be enhanced when growing conditions are optimal *e.g.* at times of high food availability, and growth  
820 benefits reduce when food is scarce (Auer *et al.* 2015b).

#### *Implications and considerations*

Variation in the relationship between growth, population, and MMR and AS underscores that considering additional biotic and abiotic factors that vary through time or space might further disentangle the links between  
825 environment, metabolism, and fitness components (Pettersen *et al.* 2016). Here, we measured metabolic traits only after a long-term period of temperature acclimation and growth. Although relative metabolic rates in salmonines are consistent over this time scale (Seppänen *et al.* 2010), repeated measurements of metabolic traits could further illuminate how metabolic phenotype (or  
830 associations between metabolic traits and fitness) can vary according to fluctuating extrinsic and intrinsic conditions (Versteegh *et al.* 2012; Biro *et al.*

2018), or developmental stage (Beaman *et al.* 2016; Burggren 2018). Moreover, metabolic measurements at a finer temporal scale would allow exploration of variation in acclimation time, a trait that has received relatively little attention but likely has important implications for ectotherms experiencing both chronic and variable temperature changes (Sandblom *et al.* 2014). Additionally, we did not consider acute responses to warming, which may preclude any acclimation response. For example, if acute warming initially increases SMR before thermal compensation occurs, the resulting reduction in AS may limit capacity for essential functions such as feeding or predator avoidance (Clark *et al.* 2011; Sandblom *et al.* 2014). Extending this study to include more realistic/natural conditions (*e.g.* co-occurring abiotic or biotic stressors), coupled with tracking of individual reproductive success would give further insight into how optimal combinations of metabolic traits and life history are shaped by environmental context.

Nonetheless, the variation (and covariation) we detected in metabolic traits arising from long-term temperature exposure and population factors has implications for the performance of populations experiencing rapid climate change (Parmesan 2006). Studies have most often focused on the link between SMR or BMR and fitness components, but we add to a growing body of research suggesting that MMR (and consequently, AS) can influence fitness components (Boratyński and Koteja 2009; Clark *et al.* 2013; Zub *et al.* 2014; Auer *et al.* 2015b). In our case, links between MMR and growth rates – a key fitness-related trait – were mediated by population-specific factors. Moreover, population-based variation in the correlation between SMR and MMR and AS hints at the intriguing possibility of fundamental links between these traits that may vary according to extrinsic or intrinsic factors, thus limiting or enhancing capacity to respond to environmental change (Metcalf *et al.* 2016; Norin *et al.* 2016). Though metabolic acclimation capacity is linked to increased resilience (Seebacher *et al.* 2015) – at least when environmental change is predictable (Reed *et al.* 2010) – such population-level variation underscores that responses to change are likely to be highly variable, even within species. Understanding

the context-dependency of associations between metabolic phenotypes and fitness, and how these can vary depending on population-specific factors, is  
 865 required for successful management and conservation of aquatic species that are already in widespread decline due to progressive warming and global change (Pörtner and Knust 2007; Limburg and Waldman 2009; Martins *et al.* 2011).

### Acknowledgements

The authors would like to thank Brian Clarke, Deirdre Cotter, members of the  
 870 FishEyE team at UCC, and the staff of Inland Fisheries Ireland and the Marine Institute for obtaining brood stock and for assistance in fish rearing, along with Robert Wynne, Ronan O’Sullivan, Peter Moran and Adam Kane for assistance in fish husbandry, and Jamie Coughlan for genotyping work. This research was supported by an ERC Starting Grant (639192-ALH) and an SFI ERC Support  
 875 Award awarded to TER. PMcG was supported in part by grants from Science Foundation Ireland (15/IA/3028 & 16/BBSRC/3316) and by grant-in-aid (RESPI/FS/16/01) from the Marine Institute (Ireland) as part of the Marine Research Programme by the Irish Government.

### References

- 880 Alsop D and Wood C. 1997. The interactive effects of feeding and exercise on oxygen consumption, swimming performance and protein usage in juvenile rainbow trout (*Oncorhynchus mykiss*). *Journal of Experimental Biology* **200**: 2337–46.
- Álvarez D, Cano JM, and Nicieza AG. 2006. Microgeographic variation in metabolic rate and energy storage of brown trout: Countergradient selection or thermal  
 885 sensitivity? *Evolutionary Ecology* **20**: 345–63.
- Álvarez D and Nicieza AG. 2005. Is metabolic rate a reliable predictor of growth and survival of brown trout (*Salmo trutta*) in the wild? *Canadian Journal of Fisheries and Aquatic Sciences* **62**: 643–9.
- 890 Angilletta MJ, Niewiarowski PH, and Navas CA. 2002. The evolution of thermal physiology in ectotherms. *Journal of Thermal Biology* **27**: 249–68.
- Archer LC, Hutton SA, Harman L, O’Grady MN, Kerry JP, Poole WR, Gargan P, McGinnity P, and Reed TE. 2019. The interplay between extrinsic and intrinsic factors in determining migration decisions in brown trout (*Salmo trutta*): An experimental study. *Frontiers in Ecology and Evolution* **7**: 222.
- 895 Arendt JD. 1997. Adaptive intrinsic growth rates: An integration across taxa. *The Quarterly Review of Biology* **72**: 149–77.
- Armstrong JD, McKelvey S, Smith GW, Rycroft P, and Fryer RJ. 2018. Effects of individual variation in length, condition and run-time on return rates of wild-reared Atlantic salmon *Salmo salar* smolts. *Journal of Fish Biology* **92**: 569–78.

- 900 Auer SK, Killen SS, and Rezende EL. 2017. Resting vs. active: A meta-analysis of the intra- and inter-specific associations between minimum, sustained, and maximum metabolic rates in vertebrates. *Functional Ecology* **31**: 1728–38.
- Auer SK, Salin K, Anderson GJ, and Metcalfe NB. 2015a. Aerobic scope explains individual variation in feeding capacity. *Biology Letters* **11**: 20150793.
- 905 Auer SK, Salin K, Anderson GJ, and Metcalfe NB. 2016. Flexibility in metabolic rate and activity level determines individual variation in overwinter performance. *Oecologia* **182**: 703–12.
- Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015b. The optimal combination of standard metabolic rate and aerobic scope for somatic growth depends on food availability. *Functional Ecology* **29**: 479–86.
- 910 Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015c. Flexibility in metabolic rate confers a growth advantage under changing food availability. *Journal of Animal Ecology* **84**: 1405–11.
- Barceló G, Love OP, and Vézina F. 2016. Uncoupling basal and summit metabolic rates in white-throated sparrows: Digestive demand drives maintenance costs, but changes in muscle mass are not needed to improve thermogenic capacity. *Physiological and Biochemical Zoology* **90**: 153–65.
- 915 Barceló G, Love OP, and Vézina F. 2016. Uncoupling basal and summit metabolic rates in white-throated sparrows: Digestive demand drives maintenance costs, but changes in muscle mass are not needed to improve thermogenic capacity. *Physiological and Biochemical Zoology* **90**: 153–65.
- Barton K. 2019. MuMIn: Multi-Model Inference. R package version 1.43.6. <https://CRAN.R-project.org/package=MuMIn>.
- 920 Bateson P and Laland KN. 2013. Tinbergen's four questions: An appreciation and an update. *Trends in Ecology & Evolution* **28**: 712–8.
- Beaman JE, White CR, and Seebacher F. 2016. Evolution of plasticity: Mechanistic link between development and reversible acclimation. *Trends in Ecology & Evolution* **31**: 237–49.
- 925 Biro PA, Garland T, Beckmann C, Ujvari B, Thomas F, and Post JR. 2018. Metabolic scope as a proximate constraint on individual behavioral variation: Effects on personality, plasticity, and predictability. *The American Naturalist* **192**: 142–54.
- Biro PA and Stamps JA. 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? *Trends in Ecology & Evolution* **25**: 653–9.
- 930 Blackmer AL, Mauck RA, Ackerman JT, Huntington CE, Nevitt GA, and Williams JB. 2005. Exploring individual quality: basal metabolic rate and reproductive performance in storm-petrels. *Behavioral Ecology* **16**: 906–13.
- Blanckenhorn WU. 1998. Adaptive Phenotypic Plasticity in Growth, Development, and Body Size in the Yellow Dung Fly. *Evolution* **52**: 1394–407.
- 935 Bochdansky AB, Grønkjær P, Herra TP, and Leggett WC. 2005. Experimental evidence for selection against fish larvae with high metabolic rates in a food limited environment. *Marine Biology* **147**: 1413–7.
- Boon AK, Réale D, and Boutin S. 2007. The interaction between personality, offspring fitness and food abundance in North American red squirrels. *Ecology Letters* **10**: 1094–104.
- 940 Boratyński Z and Koteja P. 2009. The association between body mass, metabolic rates and survival of bank voles. *Functional Ecology* **23**: 330–9.

- 945 Brown JH, Gillooly JF, Allen AP, Savage VM, and West GB. 2016. Toward a metabolic theory of ecology. *Ecology*: 1771–89.
- Brown JH, Marquet PA, and Taper ML. 1993. Evolution of body size: Consequences of an energetic definition of fitness. *The American Naturalist* **142**: 573–84.
- Buoro M, Gimenez O, and Prévost E. 2012. Assessing adaptive phenotypic plasticity by means of conditional strategies from empirical data: The latent environmental threshold model. *Evolution* **66**: 996–1009.
- 950 Burggren W. 2018. Developmental phenotypic plasticity helps bridge stochastic weather events associated with climate change. *Journal of Experimental Biology* **221**: jeb161984.
- Burton T, Killen SS, Armstrong JD, and Metcalfe NB. 2011. What causes intraspecific variation in resting metabolic rate and what are its ecological consequences? *Proceedings of the Royal Society B: Biological Sciences* **278**: 3465–73.
- 955 Careau V, Thomas D, Humphries MM, and Réale D. 2008. Energy metabolism and animal personality. *Oikos* **117**: 641–53.
- Chabot D, Steffensen JF, and Farrell AP. 2016. The determination of standard metabolic rate in fishes. *Journal of Fish Biology* **88**: 81–121.
- 960 Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011a. Partial migration: An introduction. *Oikos* **120**: 1761–3.
- Chapman BB, Hulthén K, Blomqvist DR, Hansson L-A, Nilsson J-Å, Brodersen J, Nilsson PA, Skov C, and Brönmark C. 2011b. To boldly go: Individual differences in boldness influence migratory tendency. *Ecology Letters* **14**: 871–6.
- 965 Chappell MA, Garland T, Robertson GF, and Saltzman W. 2007. Relationships among running performance, aerobic physiology and organ mass in male Mongolian gerbils. *Journal of Experimental Biology* **210**: 4179–97.
- Clark TD, Jeffries KM, Hinch SG, and Farrell AP. 2011. Exceptional aerobic scope and cardiovascular performance of pink salmon (*Oncorhynchus gorbuscha*) may underlie resilience in a warming climate. *Journal of Experimental Biology* **214**: 3074–81.
- 970 Clark TD, Sandblom E, and Jutfelt F. 2013. Aerobic scope measurements of fishes in an era of climate change: Respirometry, relevance and recommendations. *Journal of Experimental Biology* **216**: 2771–82.
- 975 Clarke A and Johnston NM. 1999. Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology* **68**: 893–905.
- Cutts CJ, Metcalfe NB, and Taylor AC. 2002. Juvenile Atlantic Salmon (*Salmo salar*) with relatively high standard metabolic rates have small metabolic scopes.
- 980 *Functional Ecology* **16**: 73–8.
- Dalziel AC, Ou M, and Schulte PM. 2012a. Mechanisms underlying parallel reductions in aerobic capacity in non-migratory threespine stickleback (*Gasterosteus aculeatus*) populations. *Journal of Experimental Biology* **215**: 746–59.
- 985 Dalziel AC, Vines TH, and Schulte PM. 2012b. Reductions in prolonged swimming capacity following freshwater colonization in multiple threespine stickleback populations. *Evolution* **66**: 1226–39.



- Dingemanse NJ, Both C, Drent PJ, and Tinbergen JM. 2004. Fitness consequences of avian personalities in a fluctuating environment. *Proceedings of the Royal Society of London Series B: Biological Sciences* **271**: 847–52.
- 990 Dubois K, Hallot F, and Vézina F. 2016. Basal and maximal metabolic rates differ in their response to rapid temperature change among avian species. *Journal of Comparative Physiology B* **186**: 919–35.
- Dupont-Prinet A, Vagner M, Chabot D, and Audet C. 2013. Impact of hypoxia on the metabolism of Greenland halibut (*Reinhardtius hippoglossoides*). *Canadian*  
 995 *Journal of Fisheries and Aquatic Sciences* **70**: 461–9.
- Eliason EJ, Clark TD, Hague MJ, Hanson LM, Gallagher ZS, Jeffries KM, Gale MK, Patterson DA, Hinch SG, and Farrell AP. 2011. Differences in thermal tolerance among sockeye salmon populations. *Science* **332**: 109–12.
- Eliason EJ and Farrell AP. 2016. Oxygen uptake in Pacific salmon *Oncorhynchus* spp.:  
 1000 When ecology and physiology meet. *Journal of Fish Biology* **88**: 359–88.
- Elliott JM. 2000. Pools as refugia for brown trout during two summer droughts: Trout responses to thermal and oxygen stress. *Journal of Fish Biology* **56**: 938–48.
- Elliott JM, Hurley MA, and Fryer RJ. 1995. A new, improved growth model for brown trout, *Salmo trutta*. *Functional Ecology* **9**: 290–8.
- 1005 Ferguson A, Reed TE, Cross TF, McGinnity P, and Prodöhl PA. 2019. Anadromy, potamodromy and residency in brown trout *Salmo trutta*: The role of genes and the environment. *Journal of Fish Biology* **95**: 692–718.
- Finstad AG, Forseth T, Næsje TF, and Ugedal O. 2004. The importance of ice cover for energy turnover in juvenile Atlantic salmon. *Journal of Animal Ecology* **73**: 959–  
 1010 66.
- Forseth T, Larsson S, Jensen AJ, Jonsson B, Näslund I, and Berglund I. 2009. Thermal growth performance of juvenile brown trout *Salmo trutta*: No support for thermal adaptation hypotheses. *Journal of Fish Biology* **74**: 133–49.
- Forseth T, Nesje TF, Jonsson B, and Hårsaker K. 1999. Juvenile migration in brown  
 1015 trout: A consequence of energetic state. *Journal of Animal Ecology* **68**: 783–93.
- Fry FE. 1971. The effect of environmental factors on the physiology of fish. In: *Fish physiology* (eds WS Hoar & DJ Randall). New York, NY: Academic Press.
- Gargan P, Kelly F, Shephard S, and Whelan K. 2016. Temporal variation in sea trout *Salmo trutta* life history traits in the Erriff River, western Ireland. *Aquaculture*  
 1020 *Environment Interactions* **8**: 675–89.
- Gillooly JF, Brown JH, West GB, Savage VM, and Charnov EL. 2001. Effects of size and temperature on metabolic rate. *Science* **293**: 2248–51.
- Gräns A, Jutfelt F, Sandblom E, Jönsson E, Wiklander K, Seth H, Olsson C, Dupont S, Ortega-Martinez O, Einarsdottir I, Björnsson BT, Sundell K, and Axelsson M.  
 1025 2014. Aerobic scope fails to explain the detrimental effects on growth resulting from warming and elevated CO<sub>2</sub> in Atlantic halibut. *Journal of Experimental Biology* **217**: 711–7.
- Gross MR, Coleman RM, and McDowall RM. 1988. Aquatic productivity and the evolution of diadromous fish migration. *Science* **239**: 1291–3.

- 1030 Guderley H and Pörtner HO. 2010. Metabolic power budgeting and adaptive strategies in zoology: Examples from scallops and fish. *Canadian Journal of Zoology* **88**: 753–63.
- Halsey LG. 2019. The reign of the p-value is over: What alternative analyses could we employ to fill the power vacuum? *Biology Letters* **15**: 20190174.
- 1035 Halsey LG, Curran-Everett D, Vowler SL, and Drummond GB. 2015. The fickle P value generates irreproducible results. *Nature Methods* **12**: 179–85.
- Hariato J and Carey N. 2019. respR: Analyse, Convert, and Automate Respirometry-Related Data. R package version 1.0.5.1. <https://github.com/januarhariato/respr>.
- 1040 Hayes JP and O'Connor CS. 1999. Natural selection on thermogenic capacity of high-altitude deer mice. *Evolution* **53**: 1280–7.
- Hendry AP, Bohlin T, Jonsson B, and Berg OK. 2004. To sea or not to sea? Anadromy v. non-anadromy in salmonids. In: *Evolution illuminated - Salmon and their relatives*. Edited by A.P. Hendry & S. C. Sterns. New York, NY: Oxford University Press.
- 1045 Hillman SS, Hancock TV, and Hedrick MS. 2013. A comparative meta-analysis of maximal aerobic metabolism of vertebrates: Implications for respiratory and cardiovascular limits to gas exchange. *Journal of Comparative Physiology B* **183**: 167–79.
- 1050 Ho J, Tumkaya T, Aryal S, Choi H, and Claridge-Chang A. 2019. Moving beyond P values: Data analysis with estimation graphics. *Nature Methods* **16**: 565.
- Hoogenboom MO, Armstrong JD, Groothuis TGG, and Metcalfe NB. 2013. The growth benefits of aggressive behavior vary with individual metabolism and resource predictability. *Behavioral Ecology* **24**: 253–61.
- 1055 Hopkins KD. 1992. Reporting fish growth: A review of the basics. *Journal of the World Aquaculture Society* **23**: 173–9.
- IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. In: Geneva, Switzerland.
- 1060 Jackson DM, Trayhurn P, and Speakman JR. 2001. Associations between energetics and over-winter survival in the short-tailed field vole *Microtus agrestis*. *Journal of Animal Ecology* **70**: 633–40.
- Jonsson B. 1985. Life history patterns of freshwater resident and sea-run migrant brown trout in Norway. *Transactions of the American Fisheries Society* **114**: 182–94.
- 1065 Jonsson B and Jonsson N. 2009. A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of Fish Biology* **75**: 2381–447.
- 1070 Jonsson B, Jonsson N, and Finstad AG. 2013. Effects of temperature and food quality on age and size at maturity in ectotherms: An experimental test with Atlantic salmon. *Journal of Animal Ecology* **82**: 201–10.
- Jutfelt F, Norin T, Ern R, Overgaard J, Wang T, McKenzie DJ, Lefevre S, Nilsson GE, Metcalfe NB, Hickey AJR, Brijs J, Speers-Roesch B, Roche DG, Gamperl AK,

- 1075 Raby GD, Morgan R, Esbaugh AJ, Gräns A, Axelsson M, Ekström A, Sandblom E, Binning SA, Hicks JW, Seebacher F, Jørgensen C, Killen SS, Schulte PM, and Clark TD. 2018. Oxygen- and capacity-limited thermal tolerance: Blurring ecology and physiology. *Journal of Experimental Biology* **221**: jeb169615.
- 1080 Killen SS. 2014. Growth trajectory influences temperature preference in fish through an effect on metabolic rate. *Journal of Animal Ecology* **83**: 1513–22.
- Killen SS, Fu C, Wu Q, Wang Y-X, and Fu S-J. 2016. The relationship between metabolic rate and sociability is altered by food deprivation. *Functional Ecology* **30**: 1358–65.
- 1085 Killen SS, Marras S, Metcalfe NB, McKenzie DJ, and Domenici P. 2013. Environmental stressors alter relationships between physiology and behaviour. *Trends in Ecology & Evolution* **28**: 651–8.
- Killen SS, Mitchell MD, Rummer JL, Chivers DP, Ferrari MCO, Meekan MG, and McCormick MI. 2014. Aerobic scope predicts dominance during early life in a tropical damselfish. *Functional Ecology* **28**: 1367–76.
- 1090 Killen SS, Reid D, Marras S, and Domenici P. 2015. The interplay between aerobic metabolism and antipredator performance: Vigilance is related to recovery rate after exercise. *Frontiers in Physiology* **6**.
- 1095 Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, and Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): A review of aspects of their life histories. *Ecology of Freshwater Fish* **12**: 1–59.
- Konarzewski M and Książek A. 2013. Determinants of intra-specific variation in basal metabolic rate. *Journal of Comparative Physiology B* **183**: 27–41.
- 1100 Lahti K, Laurila A, Enberg K, and Piironen J. 2001. Variation in aggressive behaviour and growth rate between populations and migratory forms in the brown trout, *Salmo trutta*. *Animal Behaviour* **62**: 935–44.
- Larivée ML, Boutin S, Speakman JR, McAdam AG, and Humphries MM. 2010. Associations between over-winter survival and resting metabolic rate in juvenile North American red squirrels. *Functional Ecology* **24**: 597–607.
- 1105 Lee CG, Farrell AP, Lotto A, Hinch SG, and Healey MC. 2003. Excess post-exercise oxygen consumption in adult sockeye (*Oncorhynchus nerka*) and coho (*O. kisutch*) salmon following critical speed swimming. *Journal of Experimental Biology* **206**: 3253–60.
- 1110 Lefevre S. 2016. Are global warming and ocean acidification conspiring against marine ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO<sub>2</sub> and their interaction. *Conservation Physiology* **4**.
- Limburg KE and Waldman JR. 2009. Dramatic declines in north Atlantic diadromous fishes. *BioScience* **59**: 955–65.
- 1115 Magozzi S and Calosi P. 2015. Integrating metabolic performance, thermal tolerance, and plasticity enables for more accurate predictions on species vulnerability to acute and chronic effects of global warming. *Global Change Biology* **21**: 181–94.
- Martins EG, Hinch SG, Patterson DA, Hague MJ, Cooke SJ, Miller KM, Lapointe MF, English KK, and Farrell AP. 2011. Effects of river temperature and climate

- warming on stock-specific survival of adult migrating Fraser River sockeye salmon (*Oncorhynchus nerka*). *Global Change Biology* **17**: 99–114.
- 1120
- Martin-Smith KM and Armstrong JD. 2002. Growth rates of wild stream-dwelling Atlantic salmon correlate with activity and sex but not dominance. *Journal of Animal Ecology* **71**: 413–23.
- McCarthy ID. 2000. Temporal repeatability of relative standard metabolic rate in juvenile Atlantic salmon and its relation to life history variation. *Journal of Fish Biology* **57**: 224–38.
- 1125
- Meehl GA and Tebaldi C. 2004. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* **305**: 994–7.
- Metcalfe NB, Taylor AC, and Thorpe JE. 1995. Metabolic rate, social status and life-history strategies in Atlantic salmon. *Animal Behaviour* **49**: 431–6.
- 1130
- Metcalfe NB, Van Leeuwen TE, and Killen SS. 2016. Does individual variation in metabolic phenotype predict fish behaviour and performance? *Journal of Fish Biology* **88**: 298–321.
- Millidine KJ, Armstrong JD, and Metcalfe NB. 2006. Presence of shelter reduces maintenance metabolism of juvenile salmon. *Functional Ecology* **20**: 839–45.
- 1135
- Morinville GR and Rasmussen JB. 2003. Early juvenile bioenergetic differences between anadromous and resident brook trout (*Salvelinus fontinalis*). *Canadian Journal of Fisheries and Aquatic Sciences* **60**: 401–10.
- Murchie KJ, Cooke SJ, Danylchuk AJ, and Suski CD. 2011. Estimates of field activity and metabolic rates of bonefish (*Albula vulpes*) in coastal marine habitats using acoustic tri-axial accelerometer transmitters and intermittent-flow respirometry. *Journal of Experimental Marine Biology and Ecology* **396**: 147–55.
- 1140
- Naya DE, Lardies MA, and Bozinovic F. 2007. The effect of diet quality on physiological and life-history traits in the harvestman *Pachylus paessleri*. *Journal of Insect Physiology* **53**: 132–8.
- 1145
- Nespolo RF, Bacigalupe LD, Rezende EL, and Bozinovic F. 2001. When nonshivering thermogenesis equals maximum metabolic rate: Thermal acclimation and phenotypic plasticity of fossorial *Spalacopus cyanus* (Rodentia). *Physiological and Biochemical Zoology* **74**: 325–32.
- Nespolo RF and Franco M. 2007. Whole-animal metabolic rate is a repeatable trait: A meta-analysis. *The Journal of Experimental Biology* **210**: 2000–5.
- 1150
- Nicieza AG and Álvarez D. 2009. Statistical analysis of structural compensatory growth: How can we reduce the rate of false detection? *Oecologia* **159**: 27–39.
- Nilsson J-Å, Åkesson M, and Nilsson JF. 2009. Heritability of resting metabolic rate in a wild population of blue tits. *Journal of Evolutionary Biology* **22**: 1867–74.
- 1155
- Norin T and Clark TD. 2016. Measurement and relevance of maximum metabolic rate in fishes. *Journal of Fish Biology* **88**: 122–51.
- Norin T and Malte H. 2011. Repeatability of standard metabolic rate, active metabolic rate and aerobic scope in young brown trout during a period of moderate food availability. *Journal of Experimental Biology* **214**: 1668–75.
- 1160

- Norin T, Malte H, and Clark TD. 2014. Aerobic scope does not predict the performance of a tropical eurythermal fish at elevated temperatures. *Journal of Experimental Biology* **217**: 244–51.
- 1165 Norin T, Malte H, and Clark TD. 2016. Differential plasticity of metabolic rate phenotypes in a tropical fish facing environmental change. *Functional Ecology* **30**: 369–78.
- Norin T and Metcalfe N. 2019. Ecological and evolutionary consequences of metabolic rate plasticity in response to environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**: 20180180.
- 1170 O'Connor K i., Taylor A c., and Metcalfe N b. 2000. The stability of standard metabolic rate during a period of food deprivation in juvenile Atlantic salmon. *Journal of Fish Biology* **57**: 41–51.
- Ojanguren AF, Reyes-Gavilán FG, and Braña F. 2001. Thermal sensitivity of growth, food intake and activity of juvenile brown trout. *Journal of Thermal Biology* **26**:  
1175 165–70.
- Parmesan C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics* **37**: 637–69.
- Pettersen AK, Marshall DJ, and White CR. 2018. Understanding variation in metabolic rate. *Journal of Experimental Biology* **221**: jeb166876.
- 1180 Pettersen AK, White CR, and Marshall DJ. 2016. Metabolic rate covaries with fitness and the pace of the life history in the field. *Proc R Soc B* **283**: 20160323.
- Piche J, Hutchings JA, and Blanchard W. 2008. Genetic variation in threshold reaction norms for alternative reproductive tactics in male Atlantic salmon, *Salmo salar*. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1571–5.
- 1185 Pinheiro J, Bates D, DebRoy S, Sarkar D, and R Core Team. 2019. nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-139:. URL <https://CRAN.R-project.org/package=nlme>.
- Plaut I. 2001. Critical swimming speed: Its ecological relevance. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **131**: 41–50.  
1190
- Poole WR, Dillane M, DeEyto E, Rogan G, McGinnity P, and Whelan K. 2007. Characteristics of the Burrishoole sea trout population: Census, marine survival, enhancement and stock-recruitment relationship, 1971-2003. In: *Sea Trout: Biology, Conservation and Management* (Eds: G Harris and N Milner).  
1195 Oxford, UK: Blackwells Scientific Publications.
- Pörtner HO and Farrell AP. 2008. Physiology and climate change. *Science* **322**: 690–2.
- Pörtner HO and Knust R. 2007. Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* **315**: 95–7.
- Pulido F. 2011. Evolutionary genetics of partial migration – the threshold model of migration revis(it)ed. *Oikos* **120**: 1776–83.  
1200
- R Core Team. 2019. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

- 1205 Reed TE, Waples Robin S., Schindler Daniel E., Hard Jeffrey J., and Kinnison Michael T. 2010. Phenotypic plasticity and population viability: The importance of environmental predictability. *Proceedings of the Royal Society B: Biological Sciences* **277**: 3391–400.
- 1210 Reid D, Armstrong JD, and Metcalfe NB. 2011. Estimated standard metabolic rate interacts with territory quality and density to determine the growth rates of juvenile Atlantic salmon. *Functional Ecology* **25**: 1360–7.
- Reid D, Armstrong JD, and Metcalfe NB. 2012. The performance advantage of a high resting metabolic rate in juvenile salmon is habitat dependent. *Journal of Animal Ecology* **81**: 868–75.
- 1215 Robertsen G, Armstrong JD, Nislow KH, Herfindal I, McKelvey S, and Einum S. 2014. Spatial variation in the relationship between performance and metabolic rate in wild juvenile Atlantic salmon. *Journal of Animal Ecology* **83**: 791–9.
- Rønning B, Jensen H, Moe B, and Bech C. 2007. Basal metabolic rate: Heritability and genetic correlations with morphological traits in the zebra finch. *Journal of Evolutionary Biology* **20**: 1815–22.
- 1220 Rutherford JC, Marsh NA, Davies PM, and Bunn SE. 2004. Effects of patchy shade on stream water temperature: How quickly do small streams heat and cool? *Marine and Freshwater Research* **55**: 737–48.
- 1225 Sadowska J, Gębczyński Andrzej K., and Konarzewski Marek. 2013. Basal metabolic rate is positively correlated with parental investment in laboratory mice. *Proceedings of the Royal Society B: Biological Sciences* **280**: 20122576.
- Sadowska ET, Stawski Clare, Rudolf Agata, Dheyongera Geoffrey, Chrzęścik Katarzyna M., Baliga-Klimczyk Katarzyna, and Koteja Paweł. 2015. Evolution of basal metabolic rate in bank voles from a multidirectional selection experiment. *Proceedings of the Royal Society B: Biological Sciences* **282**: 20150025.
- 1230 Sandblom E, Clark TD, Gräns A, Ekström A, Brijs J, Sundström LF, Odelström A, Adill A, Aho T, and Jutfelt F. 2016. Physiological constraints to climate warming in fish follow principles of plastic floors and concrete ceilings. *Nature Communications* **7**: 11447.
- 1235 Sandblom E, Gräns Albin, Axelsson Michael, and Seth Henrik. 2014. Temperature acclimation rate of aerobic scope and feeding metabolism in fishes: Implications in a thermally extreme future. *Proceedings of the Royal Society B: Biological Sciences* **281**: 20141490.
- 1240 Schulte PM. 2015. The effects of temperature on aerobic metabolism: Towards a mechanistic understanding of the responses of ectotherms to a changing environment. *Journal of Experimental Biology* **218**: 1856–66.
- Schulte PM, Healy TM, and Fangue NA. 2011. Thermal performance curves, phenotypic plasticity, and the time scales of temperature exposure. *Integrative and Comparative Biology* **51**: 691–702.
- 1245 Seebacher F, Ward AJW, and Wilson RS. 2013. Increased aggression during pregnancy comes at a higher metabolic cost. *Journal of Experimental Biology* **216**: 771–6.
- Seebacher F, White CR, and Franklin CE. 2015. Physiological plasticity increases resilience of ectothermic animals to climate change. *Nature Climate Change* **5**: 61–6.

- 1250 Seneviratne SI, Donat MG, Mueller B, and Alexander LV. 2014. No pause in the increase of hot temperature extremes. *Nature Climate Change* **4**: 161–3.
- Seppänen E, Piironen J, and Huuskonen H. 2010. Consistency of standard metabolic rate in relation to life history strategy of juvenile Atlantic salmon *Salmo salar*. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **156**: 278–84.
- 1255 Stearns SC. 1992. The evolution of life histories. Oxford, UK: Oxford University Press.
- Svendsen MBS, Bushnell PG, and Steffensen JF. 2016. Design and setup of intermittent-flow respirometry system for aquatic organisms. *Journal of Fish Biology* **88**: 26–50.
- 1260 Swanson DL, McKechnie AE, and Vézina F. 2017. How low can you go? An adaptive energetic framework for interpreting basal metabolic rate variation in endotherms. *Journal of Comparative Physiology B* **187**: 1039–56.
- Thomas DW, Blondel J, Perret P, Lambrechts MM, and Speakman JR. 2001. Energetic and fitness costs of mismatching resource supply and demand in seasonally breeding birds. *Science* **291**: 2598–600.
- 1265 Tinbergen N. 1963. On aims and methods of Ethology. *Zeitschrift für Tierpsychologie* **20**: 410–33.
- Tomkins JL and Hazel W. 2007. The status of the conditional evolutionarily stable strategy. *Trends in Ecology & Evolution* **22**: 522–8.
- 1270 Tudorache C, Blust R, and Boeck GD. 2007. Swimming capacity and energetics of migrating and non-migrating morphs of three-spined stickleback *Gasterosteus aculeatus* L. and their ecological implications. *Journal of Fish Biology* **71**: 1448–56.
- 1275 Van Leeuwen TE, Rosenfeld JS, and Richards JG. 2011. Adaptive trade-offs in juvenile salmonid metabolism associated with habitat partitioning between coho salmon and steelhead trout in coastal streams. *Journal of Animal Ecology* **80**: 1012–23.
- 1280 Van Leeuwen TE, Rosenfeld JS, and Richards JG. 2012. Effects of food ration on SMR: Influence of food consumption on individual variation in metabolic rate in juvenile coho salmon (*Oncorhynchus kisutch*). *Journal of Animal Ecology* **81**: 395–402.
- 1285 Ven TMFN van de, Mzilikazi N, and McKechnie AE. 2013. Phenotypic flexibility in body mass, basal metabolic rate and summit metabolism in southern red bishops (*Euplectes orix*): Responses to short term thermal acclimation. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology* **165**: 319–27.
- Versteegh MA, Helm B, Gwinner E, and Tieleman BI. 2012. Annual cycles of metabolic rate are genetically determined but can be shifted by phenotypic flexibility. *Journal of Experimental Biology* **215**: 3459–66.
- 1290 Ward BR and Slaney PA. 1988. Life history and smolt-to-adult survival of Keogh River steelhead trout (*Salmo gairdneri*) and the relationship to smolt size. *Canadian Journal of Fisheries and Aquatic Sciences* **45**: 1110–22.
- Wone BWM, Madsen P, Donovan ER, Labocha MK, Sears MW, Downs CJ, Sorensen DA, and Hayes JP. 2015. A strong response to selection on mass-independent

- maximal metabolic rate without a correlated response in basal metabolic rate.  
 1295 *Heredity* **114**: 419–27.
- Yamamoto T, Ueda H, and Higashi S. 1998. Correlation among dominance status, metabolic rate and otolith size in masu salmon. *Journal of Fish Biology* **52**: 281–90.
- 1300 Zeng L-Q, Fu C, and Fu S-J. 2018. The effects of temperature and food availability on growth, flexibility in metabolic rates and their relationships in juvenile common carp. *Comparative Biochemistry and Physiology Part A, Molecular & Integrative Physiology* **217**: 26–34.
- Zeng L-Q, Zhang A-J, Killen SS, Cao Z-D, Wang Y-X, and Fu S-J. 2017. Standard metabolic rate predicts growth trajectory of juvenile Chinese crucian carp (1305 *Carassius auratus*) under changing food availability. *Biology Open* **6**: 1305–9.
- Zub K, Borowski Z, Szafrńska PA, Wieczorek M, and Konarzewski M. 2014. Lower body mass and higher metabolic rate enhance winter survival in root voles, *Microtus oeconomus*. *Biological Journal of the Linnean Society* **113**: 297–309.



## Supporting Information

---

### **Developmental temperature and population factors affect variation and covariation of metabolic traits, with complex effects on growth rates in brown trout (*Salmo trutta*)**


Louise C. Archer<sup>\*1,2</sup>, Stephen A. Hutton<sup>1,2</sup>, Luke Harman<sup>1,2</sup>, W. Russell Poole<sup>3</sup>, Patrick Gargan<sup>4</sup>, Philip McGinnity<sup>1,3</sup>, Thomas E. Reed<sup>1,2</sup>

<sup>1</sup> *School of Biological, Earth and Environmental Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland.*


<sup>2</sup> *Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland.*

<sup>3</sup> *Marine Institute, Furnace, Newport, Co. Mayo, Ireland.*

<sup>4</sup> *Inland Fisheries Ireland, 3044 Lake Drive, Citywest Business Campus, Dublin, D24 Y265, Ireland.*



**Figure S1:** (A) Srahrevagh River in the Burrishoole catchment, where brown trout brood stock were collected by seine netting in November 2015, used to produce F1 offspring for the experimental study. The wild Srahrevagh population does not express the anadromous life history, and experimental offspring from Srahrevagh brood stock were considered to have non-anadromous population background. (B) Tawnyard Lough in the Erriff catchment, site of brown trout brood stock collections (via seine netting) in November 2015, used to produce F1 offspring for the experimental study. The wild Tawnyard Lough population has a strong anadromous component, and experimental offspring from Tawnyard brood stock were considered to have an anadromous population background.



**Figure S2:** Mean specific growth rate trajectories (residual values corrected for initial body size) (with associated standard errors) of brown trout classed as having: (A) High or low standard metabolic rate (SMR); (B) High or low maximum metabolic rate (MMR); and (C) High or low aerobic scope (AS) according to residual SMR/MMR/AS values after accounting for variation in body mass (“High” metabolic phenotype =  $rSMR/rMMR/rAS > 0$  and “Low” =  $rSMR/rMMR/rAS < 0$ ). Fish were offspring from two population backgrounds (AB = Anadromous background, Non-AB = Non-Anadromous background), reared under two temperature treatments (Cool = natural temperature regime, Warm = elevated 1.8°C above Cool). (D) Coefficient estimates (± 95% confidence intervals) from mixed effect models testing interactive effects of time (months since experiment start) and metabolic rate ( $rSMR/rMMR/rAS$ ), along with effects of temperature treatment, and population background on specific growth rate trajectories of brown trout.

**Table S1:** Parameter estimates, with associated standard errors (SE), *t*-values, and *P*-values from the linear model describing  $\log_{10}$ -transformed metabolic rates ( $\text{mg O}_2 \text{ hr}^{-1}$ ) as a function of  $\log_{10}$ -transformed body mass in  $n = 32$  brown trout (SMR = standard metabolic rate, MMR = maximum metabolic rate, and AS = aerobic scope).

| <b>Response</b> | <b>Parameter</b>      | <b>Estimate</b> | <b>SE</b> | <b>t-value</b> | <b>P-value</b> |
|-----------------|-----------------------|-----------------|-----------|----------------|----------------|
| $\log_{10}$ SMR | Intercept             | -1.28           | 0.33      | -3.89          | 0.001          |
|                 | $\log_{10}$ Body mass | 0.95            | 0.16      | 6.00           | < 0.001        |
| $\log_{10}$ MMR | Intercept             | 0.16            | 0.39      | 0.42           | 0.676          |
|                 | $\log_{10}$ Body mass | 0.71            | 0.19      | 3.74           | 0.001          |
| $\log_{10}$ AS  | Intercept             | 0.18            | 0.42      | 0.42           | 0.680          |
|                 | $\log_{10}$ Body mass | 0.67            | 0.21      | 3.27           | 0.003          |

**Table S2:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from the general linear models testing the effects of temperature treatment (cool or warm) and population background (anadromous or non-anadromous) on standard metabolic rate (SMR), maximum metabolic rate (MMR), and aerobic scope (AS) in brown trout. SMR, MMR, and AS were log<sub>10</sub>-transformed, and body mass (log<sub>10</sub>-transformed) was included as a covariate. Significance was assessed at  $P < 0.05$ . Effects are contrasted against fish from the anadromous population background in the cool temperature treatment.

| Response              | Parameter                   | Estimate | SE    | t-value | P-value |
|-----------------------|-----------------------------|----------|-------|---------|---------|
| log <sub>10</sub> SMR | Intercept                   | -1.045   | 0.300 | -3.479  | 0.002   |
|                       | Population: Non-anadromous  | -0.061   | 0.027 | -2.246  | 0.033   |
|                       | Temperature: Warm           | -0.078   | 0.025 | -3.090  | 0.004   |
|                       | log <sub>10</sub> Body mass | 0.875    | 0.144 | 6.092   | < 0.001 |
| log <sub>10</sub> MMR | Intercept                   | 0.605    | 0.356 | 1.698   | 0.101   |
|                       | Population: Non-anadromous  | -0.104   | 0.032 | -3.255  | 0.003   |
|                       | Temperature: Warm           | -0.061   | 0.030 | -2.024  | 0.053   |
|                       | log <sub>10</sub> Body mass | 0.531    | 0.170 | 3.117   | 0.004   |
| log <sub>10</sub> AS  | Intercept                   | 0.649    | 0.399 | 1.626   | 0.115   |
|                       | Population: Non-anadromous  | -0.111   | 0.036 | -3.084  | 0.005   |
|                       | Temperature: Warm           | -0.058   | 0.034 | -1.727  | 0.095   |
|                       | log <sub>10</sub> Body mass | 0.483    | 0.191 | 2.532   | 0.017   |

**Table S3:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from the general linear models testing the effects of temperature treatment (cool or warm) and population background (anadromous or non-anadromous) on relationships between size-corrected standard metabolic rate (SMR) and maximum metabolic rate (MMR), SMR and aerobic scope (AS), and MMR and AS in brown trout. SMR, MMR, and AS were corrected for body mass by taking the residuals of the linear relationship between metabolic rate and body mass to give rSMR, rMMR, and rAS. Significance was assessed at  $P < 0.05$ . Effects are contrasted against fish from the anadromous population background in the cool temperature treatment.

| Response | Parameter                  | Estimate | SE    | t-value | P-value |
|----------|----------------------------|----------|-------|---------|---------|
| rMMR     | Intercept                  | 0.043    | 0.027 | 1.572   | 0.128   |
|          | rSMR                       | 0.731    | 0.261 | 2.798   | 0.009   |
|          | Population: Non-anadromous | -0.073   | 0.029 | -2.499  | 0.019   |
|          | Temperature: Warm          | -0.035   | 0.031 | -1.103  | 0.280   |
|          | rSMR × Non-anadromous      | -0.774   | 0.349 | -2.221  | 0.035   |
| rAS      | Intercept                  | 0.049    | 0.032 | 1.561   | 0.130   |
|          | rSMR                       | 0.690    | 0.301 | 2.292   | 0.030   |
|          | Population: Non-anadromous | -0.084   | 0.034 | -2.482  | 0.020   |
|          | Temperature: Warm          | -0.039   | 0.036 | -1.089  | 0.286   |
|          | rSMR × Non-anadromous      | -0.878   | 0.402 | -2.187  | 0.038   |
| rAS      | Intercept                  | -0.007   | 0.004 | -2.053  | 0.050   |
|          | rMMR                       | 1.113    | 0.022 | 49.909  | < 0.001 |
|          | Population: Non-anadromous | 0.005    | 0.004 | 1.208   | 0.237   |
|          | Temperature: Warm          | 0.010    | 0.004 | 2.495   | 0.019   |

**Table S4:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from three mixed effects models testing the effects of standard metabolic rate (SMR), maximum metabolic rate (MMR), aerobic scope (AS) on specific growth rate trajectories of brown trout from two population backgrounds. Specific growth rate was calculated from repeated fork length measurements in fish experiencing either natural (Cool) or elevated (Warm) temperature regimes. SMR, MMR, and AS were corrected for body mass by taking the residuals of the relationship between body mass and metabolic rate to give rSMR, rMMR, and rAS. Initial fork length was included as a covariate in all models. Significance was assessed at  $P < 0.05$ . Effects are contrasted against fish from the anadromous population in the cool treatment.

| Response             | Effect                 | Estimate | SE    | <i>t</i> -value | <i>P</i> -value |
|----------------------|------------------------|----------|-------|-----------------|-----------------|
| Specific growth rate | Intercept              | 0.332    | 0.073 | 4.559           | < 0.001         |
|                      | Non-Anadromous         | 0.019    | 0.019 | 0.996           | 0.332           |
|                      | Time                   | -0.136   | 0.114 | -1.200          | 0.233           |
|                      | Time <sup>2</sup>      | -0.148   | 0.051 | -2.939          | 0.004           |
|                      | Time <sup>3</sup>      | 0.466    | 0.046 | 10.081          | < 0.001         |
|                      | Initial length         | -0.013   | 0.004 | -3.644          | < 0.001         |
|                      | rSMR                   | 0.149    | 0.115 | 1.294           | 0.211           |
|                      | Temperature: Warm      | 0.033    | 0.014 | 2.335           | 0.031           |
|                      | rSMR × Warm            | -0.278   | 0.134 | -2.071          | 0.052           |
|                      | Warm × Non- Anadromous | -0.044   | 0.018 | -2.433          | 0.025           |
| Specific growth rate | Intercept              | 0.356    | 0.072 | 4.955           | < 0.001         |
|                      | Non-Anadromous         | -0.024   | 0.016 | -1.513          | 0.146           |
|                      | Time                   | -0.133   | 0.115 | -1.152          | 0.252           |
|                      | Time <sup>2</sup>      | -0.143   | 0.051 | -2.794          | 0.006           |
|                      | Time <sup>3</sup>      | 0.458    | 0.046 | 9.997           | < 0.001         |
|                      | Initial length         | -0.013   | 0.004 | -3.692          | < 0.001         |
|                      | rMMR                   | 0.003    | 0.067 | 0.046           | 0.964           |
|                      | Temperature: Warm      | 0.011    | 0.010 | 1.125           | 0.274           |
|                      | rMMR × Non-Anadromous  | -0.267   | 0.128 | -2.094          | 0.049           |
| Specific growth rate | Intercept              | 0.331    | 0.071 | 4.695           | < 0.001         |
|                      | Non-Anadromous         | -0.020   | 0.015 | -1.335          | 0.197           |
|                      | Time                   | -0.166   | 0.114 | -1.465          | 0.146           |
|                      | Time <sup>2</sup>      | -0.135   | 0.051 | -2.633          | 0.010           |
|                      | Time <sup>3</sup>      | 0.462    | 0.046 | 10.083          | < 0.001         |
|                      | Initial length         | -0.012   | 0.004 | -3.431          | 0.001           |
|                      | rAS                    | 0.018    | 0.060 | 0.303           | 0.765           |
|                      | Temperature: Warm      | 0.013    | 0.009 | 1.375           | 0.184           |
|                      | rAS × Non-Anadromous   | -0.259   | 0.111 | -2.345          | 0.029           |

**Table S5:** Parameter estimates and associated standard errors (SE), *t*-values, and *P*-values from the mixed effects models testing the for interactive effects of time (weeks since start of experiment) and standard metabolic rate (SMR), maximum metabolic rate (MMR), and aerobic scope (AS) on specific growth rate trajectories of brown trout. Fish were from two population backgrounds (AB = anadromous and non-AB = non-anadromous). Specific growth rate was calculated from repeated fork length measurements in fish experiencing either natural (Cool) or elevated (Warm) temperature regimes. SMR, MMR, and AS were corrected for body mass by taking the residuals of the relationship between body mass and metabolic rate to give rSMR, rMMR, and rAS. Initial fork length was included as a covariate in all models. Significance was assessed at  $P < 0.05$ . Effects are contrasted against fish from the anadromous population in the cool treatment.

| Response             | Effect                   | Estimate | SE    | <i>t</i> -value | <i>P</i> -value |
|----------------------|--------------------------|----------|-------|-----------------|-----------------|
| Specific growth rate | Intercept                | 0.316    | 0.071 | 4.456           | < 0.001         |
|                      | Non-Anadromous           | 0.020    | 0.019 | 1.049           | 0.307           |
|                      | Time                     | -0.155   | 0.108 | -1.437          | 0.153           |
|                      | Time <sup>2</sup>        | -0.162   | 0.047 | -3.479          | 0.001           |
|                      | Time <sup>3</sup>        | 0.487    | 0.045 | 10.924          | < 0.001         |
|                      | Initial length           | -0.012   | 0.003 | -3.549          | 0.001           |
|                      | rSMR                     | 0.128    | 0.116 | 1.106           | 0.283           |
|                      | Temperature: Warm        | 0.036    | 0.014 | 2.513           | 0.021           |
|                      | rSMR × Warm              | -0.263   | 0.134 | -1.955          | 0.065           |
|                      | Warm × Non-AB            | -0.046   | 0.018 | -2.512          | 0.021           |
|                      | rSMR × Time              | 2.459    | 0.469 | 5.244           | < 0.001         |
|                      | rSMR × Time <sup>2</sup> | -0.708   | 0.506 | -1.401          | 0.164           |
|                      | rSMR × Time <sup>3</sup> | 1.181    | 0.521 | 2.268           | 0.025           |
| Specific growth rate | Intercept                | 0.331    | 0.072 | 4.585           | < 0.001         |
|                      | Non-Anadromous           | -0.021   | 0.016 | -1.317          | 0.203           |
|                      | Time                     | -0.164   | 0.115 | -1.424          | 0.157           |
|                      | Time <sup>2</sup>        | -0.136   | 0.051 | -2.654          | 0.009           |
|                      | Time <sup>3</sup>        | 0.469    | 0.045 | 10.421          | < 0.001         |
|                      | Initial length           | -0.012   | 0.004 | -3.324          | 0.001           |
|                      | rMMR                     | 0.004    | 0.066 | 0.057           | 0.955           |
|                      | Temperature: Warm        | 0.011    | 0.009 | 1.145           | 0.266           |
|                      | rMMR × Non-AB            | -0.280   | 0.127 | -2.209          | 0.039           |
|                      | rMMR × Time              | 0.686    | 0.479 | 1.432           | 0.155           |
|                      | rMMR × Time <sup>2</sup> | -0.409   | 0.455 | -0.899          | 0.370           |
|                      | rMMR × Time <sup>3</sup> | 1.023    | 0.421 | 2.428           | 0.017           |



|                      |                         |        |       |        |         |
|----------------------|-------------------------|--------|-------|--------|---------|
| Specific growth rate | Intercept               | 0.314  | 0.071 | 4.399  | < 0.001 |
|                      | Non-Anadromous          | -0.018 | 0.015 | -1.215 | 0.239   |
|                      | Time                    | -0.189 | 0.114 | -1.658 | 0.100   |
|                      | Time <sup>2</sup>       | -0.129 | 0.052 | -2.501 | 0.014   |
|                      | Time <sup>3</sup>       | 0.472  | 0.045 | 10.423 | < 0.001 |
|                      | Initial length          | -0.011 | 0.004 | -3.142 | 0.002   |
|                      | rAS                     | 0.019  | 0.060 | 0.320  | 0.752   |
|                      | Temperature: Warm       | 0.013  | 0.009 | 1.399  | 0.177   |
|                      | rAS × Non-AB            | -0.272 | 0.110 | -2.472 | 0.023   |
|                      | rAS × Time              | 0.480  | 0.445 | 1.077  | 0.284   |
|                      | rAS × Time <sup>2</sup> | -0.362 | 0.422 | -0.858 | 0.392   |
|                      | rAS × Time <sup>3</sup> | 0.911  | 0.388 | 2.350  | 0.020   |

## Chapter 6

---

### General Discussion

Enormous phenotypic diversity is evident between populations and among individuals within populations (Roff 1996), with pivotal roles in ecosystem structure and function (Bolnick *et al.* 2011; Des Roches *et al.* 2018). Understanding why, and how, such intraspecific diversity arises and persists is necessary to inform management and conservation of species in a changing world (Naish and Hard 2008), not least because different phenotypes may show variable responses to environmental change. Mechanisms underpinning intraspecific diversity can be understood from proximate and ultimate perspectives (Tinbergen 1963), where proximate mechanisms relate to the environmental or ontogenetic factors influencing phenotypic expression, and ultimate mechanisms concern evolutionary function and phylogenetic control of phenotypes (Laland *et al.* 2011; Bateson and Laland 2013). With this in mind, the overarching objective of this thesis was to advance the understanding of how proximate and ultimate factors (with more emphasis on the former) contribute to shaping phenotypic diversity in brown trout, a culturally and economically important species that is iconic for the impressive array of life histories it displays (Klemetsen *et al.* 2003; Ferguson *et al.* 2019; Nevoux *et al.* 2019).

#### Overview of each chapter

My overall approach has been to address the causes and consequences of life history diversity in brown trout by exploring the interplays between physiology, environment, and phenotypic diversity, focusing on alternative migratory tactics. In Chapter 2 I aimed to explore how extrinsic (i.e. environmental) and intrinsic (i.e. population) factors interactively determine expression of migratory *versus* non-migratory tactics. In experimental offspring from two populations that naturally differ in migration frequency, I measured rates of anadromy, maturation, and various traits associated with physiological

condition, in response to long-term food restriction treatments. To clarify when in early life the migratory decision might occur I also investigated how the timing of food restriction influences migration, which proved to be variable between the two populations. Although anadromy rates tended to be higher in the offspring from the anadromous population, anadromous phenotypes emerged at lower frequencies among offspring from the non-migratory population, with implications for conservation and restoration of the anadromous life history among brown trout populations.

In Chapter 3, I aimed to extend the study of the proximate drivers of life history variation by exploring how co-occurring environmental factors collectively influence migratory tactics. I reared offspring from an anadromous-background population under conditions of low food and warm temperature treatments that simulated potential climate change scenarios. The food and temperature stressor treatments had additive, but opposing effects on migration tactics, whereby food restriction increased migration rates, but warm temperatures increased maturation rates *in lieu* of migration. The combined stressors had antagonistic effects on size-associated traits and underlying physiological condition, suggesting that responses to cumulative effects of global change will not be straightforward, and may depend upon the response considered.

Chapter 4 explored how metabolic traits vary according to intrinsic and extrinsic factors. Specifically, I measured SMR, MMR, and AS in brown trout offspring from two populations that differ in migration tendency, after exposure to long-term conditions of food restriction. Variation in SMR and MMR was related to population background and food restriction treatments, and I detected population-specific variation in the ability to adjust AS in response to food restriction. The results in Chapter 4 contribute to our understanding of how populations can respond to fluctuating food resources via phenotypic plasticity in metabolic rate, with potential consequences for population persistence if food resources are limited or temporally variable.

I further explored causes and consequences of metabolic rate variation in Chapter 5 by assessing how temperature influences variation in metabolic traits

among populations, and the consequences of metabolic variation for a fitness-related trait. I measured SMR, MMR, AS, and individual growth rates in offspring from two populations with divergent migratory tactics that were reared under long-term warming conditions. Metabolic traits showed both  
 65 population-level variation and plastic responses to warming. The relationship between metabolic traits and growth depended on either temperature (for SMR) or population background (for MMR and AS), which suggests that future climate warming effects on performance in wild populations are likely to be context dependent.

70 In this chapter, I discuss how these studies contribute to our understanding of the interaction between genes and environment underpinning phenotypic diversity in terms of: (i) life-history tactics, and (ii) physiology. In synthesising the results of this thesis with existing studies, I highlight the broader implications of the findings presented here. I also address some considerations  
 75 faced while undertaking this research, and identify areas for future research to build upon this body of work.

### **Proximate drivers of facultative migration**

#### *Support for the threshold model*

Much of our understanding of facultative migration has been aided by applying  
 80 the environmentally cued threshold model (Tomkins and Hazel 2007). Under this framework, alternative tactics are conditional upon the relationship between a “status” trait (which is cued by the environment, but also probably influenced by genetic factors), and an inherited threshold for the status trait (Tomkins and Hazel 2007; Piche *et al.* 2008; Pulido 2011; Buoro *et al.* 2012). It  
 85 follows that divergent phenotypes can emerge under similar environmental conditions, but also that similar phenotypes might be produced from genetic dissimilarity. Based on the threshold reaction norm concept, I had expected that manipulations of environmental conditions (i.e. food treatments in Chapter 2, along with food and temperature treatments in Chapter 3) would result in  
 90 different frequencies of migratory phenotypes emerging in the various treatments. The relatively higher rates of migratory phenotypes observed in

food restriction treatments in Chapters 2 and 3 were compatible with the threshold model, as was the higher frequency of the mature phenotype in warm treatments in Chapter 3 (although the expression of migratory tactics in this case was in the opposite direction to expectations). Moreover, the emergence of anadromous phenotypes in fish derived from the Bunaveela population (that naturally does not show any anadromy in the wild) suggests that experimental food restriction prevented some individuals within this population from meeting their threshold for residency. Nonetheless, despite the apparent re-expression of anadromous phenotypes within the Bunaveela under low food conditions, anadromy was comparatively more frequent among fish from the naturally anadromous Erriff population, indicating a genetic component to anadromy. This offers some insight into ultimate mechanisms underlying facultative migration, whereby the mean threshold values have potentially evolved to differ between the two populations (Piche *et al.* 2008).

An important *caveat* here is that smolting was measured in year two of life, meaning total anadromy rates across all potential smolt ages were not quantified. Outcomes could potentially differ if the probability of smolting in year three varies by population. Within the Erriff catchment, the majority of smolts migrate in the second (2+) or third (3+) year of life, with approximately equal frequencies of 2+ and 3+ smolts (Gargan *et al.* 2016). Since the Bunaveela population does not express anadromy in nature, we have no definitive information on the distribution of smolts among different age classes. However, smolts historically produced in the wider Burrishoole catchment (pre-collapse of the anadromous stocks) exhibited approximately equal ratios of 2+ to 3+ smolts (Poole *et al.* 1996), and more recent records indicate a bias towards 2+ smolts (Poole *et al.* 2007). We thus have no compelling reason to suspect that the population-level patterns of anadromy might vary if the study had been extended for an additional year.

#### 120 *Evidence for potential status traits*

The environmentally cued threshold model proposes that individuals undertaking migratory decisions assess a “status” trait, yet studies have not

conclusively identified any one trait that controls migratory decisions in salmonines (Kendall *et al.* 2014; Ferguson *et al.* 2019). The results described in Chapters 2 and 3 add to a growing body of evidence supporting a multitude of potential status traits as being instrumental in migratory decisions. In light of previous studies describing both positive and negative relationships between migration tactics and proposed status traits *e.g.* size (Morinville and Rasmussen 2003; Acolas *et al.* 2012), mass (Winter *et al.* 2016), condition (Boel *et al.* 2014; Hecht *et al.* 2015), growth (Jonsson 1985; Morinville and Rasmussen 2003; Acolas *et al.* 2012), energetic demands (Forseth *et al.* 1999), and lipid deposition (McMillan *et al.* 2012; Sloat and Reeves 2014), it seem increasingly apparent that it is more likely that a suite of interlinked physiological (or morphological) traits combine to influence migration tactics. In Chapters 2 and 3, future migrants and non-migrants showed divergent trajectories in traits such as condition, body size, and mass, far in advance of the natural migration period. Though we cannot definitively say if these traits had causal effects in the migratory decision-making process, or were consequences of adopting a chosen tactic, nonetheless, the results in Chapters 2 and 3 do suggest that a number of size/physiologically-associated traits are integral to the migratory phenotype. Such traits may reflect some underlying energetic state that is assessed, or their association with migration tactics might arise from genetic or environmental covariation between these traits and other physiological traits that influence migration (*e.g.* rate of change in energy balance or lipid deposition) (Doctor *et al.* 2014).

Relationships between migratory tactics and different physiological components or potential “status” traits may be further mediated by intrinsic or extrinsic conditions, making it more difficult to characterise clear associations, particularly if traits vary in environmental sensitivities. For example, antagonistic effects of environmental stressors on traits in Chapter 3 did not translate at the level of migratory tactics, perhaps due to variable effects of co-occurring environmental factors on key underlying traits (Galic *et al.* 2018). Moreover, links between the underlying traits themselves may be variable, for

example, associations between metabolic traits and growth (two strong  
 155 candidates for status traits) varied according to environmental factors (*i.e.*  
 temperature) and population background in Chapter 5, in addition to showing  
 temporal sensitivity. When considered alongside population-level variation in  
 metabolic traits in offspring from populations of different migratory  
 backgrounds (Chapters 4 and 5), the divergence in physiological and  
 160 morphological traits implies fundamental differences in energy uptake and  
 expenditure between migrants and residents (Forseth *et al.* 1999; Sloat and  
 Reeves 2014), or at least between predominantly migratory and non-migratory  
 populations. Individual-level tracking of relationships among metabolic traits  
 and other surrogate status traits (size/physiologically-associated traits) across  
 165 an extended period during development would help to clarify how these traits  
 collectively form an “energetic phenotype” that influences migratory tactics.

#### *Environmental factors influencing migration*

Understanding how environmental factors mediate the expression of life-  
 history tactics via underlying traits was a key goal of this thesis, and was  
 170 specifically addressed in Chapters 2 and 3. The higher rates of anadromy in  
 response to food limitation (Chapters 2 and 3) is in agreement with studies that  
 have found direct and indirect manipulations of food availability to alter  
 patterns of adfluvial/marine migration and freshwater maturation (Olsson *et al.*  
 2006; Wysujack *et al.* 2009; O’Neal and Stanford 2011; Jonsson *et al.* 2012; Jones  
 175 *et al.* 2015). An interesting result emerging from Chapter 2 was that the timing  
 of food limitation can alter patterns of migration *versus* residency in a  
 population-specific manner. This could be explained by variable “decision  
 windows” between populations, or differential sensitivity between populations  
 to annually recurring decision windows. While we could not distinguish in  
 180 Chapter 2 if populations showed temporally distinct decisions windows, or if  
 the migration decision is revisited annually, threshold-type models used to  
 understand life-history trajectories in Atlantic salmon *S. salar* (Thorpe *et al.*  
 1998) and steelhead *O. mykiss* (Satterthwaite *et al.* 2009) propose existence of  
 multiple distinct decision windows. Moreover, each window might relate to

185 separate maturation or emigration decisions. A similar series of decisions (or “switches”) could exist for brown trout, each of which is independently controlled by a combination of environmental and genetic factors (Ferguson *et al.* 2019).

The prevalence of individuals that could not be assigned a life-history phenotype in Chapters 2 and 3 further supports the existence of additional life-  
 190 history decisions that may be more related to the timing of phenotypic expression. Further exploration is required to determine whether similar threshold mechanisms are at play here. Existence of additional decision windows could perhaps explain why associations between potential status traits  
 195 and migration tactics have generally proved to be inconsistent (Kendall *et al.* 2014; Ferguson *et al.* 2019), particularly if different traits hold varying degrees of importance for various decisions related to migration and residency. Condition (and/or other traits related to energy status) might form the basis for the initial migration *versus* residency decision, whereas size, lipid stores, or  
 200 growth might be more important in decisions regarding *when* to migrate or mature, respectively (Thorpe *et al.* 1998; Thorpe and Metcalfe 1998). Similarly, migration destination might be influenced by factors related to, but nonetheless distinct from, the initial migratory decision. For example, metabolic rate has been implicated in the decision of brown trout to migrate from natal streams,  
 205 but subsequent growth rates (Cucherousset *et al.* 2005), or rate of lipid depletion (Boel *et al.* 2014) might dictate whether trout terminate migration in the larger river stem or lake, or continue to the sea. It seems increasingly obvious that we should reject the notion of life-history tactics as a simple dichotomy of migration *versus* residency, which belies the complex mosaic of  
 210 life-history decisions in trout (Birnie-Gauvin *et al.* 2019). Instead, consideration of myriad trajectories – encompassing freshwater maturation at a variety of ages and migration/anadromy at a variety of ages – is required, with each trajectory controlled by a combination of inherited and environmental factors mediated by interlinked physiological traits.



215 While the overarching effects of food availability in Chapters 2 and 3 were in line with expectations, the negative relationship between temperature and migratory tactics in Chapter 3 was somewhat unexpected given the fundamental role of temperature in defining energetic balance in ectotherms (Fry 1971; Clarke and Johnston 1999; Gillooly *et al.* 2001). Since cooler water temperatures are associated with increased lipid deposition (Kammerer and Heppell 2013), which is linked to maturation in salmonines (McMillan *et al.* 2012; Sloat and Reeves 2014), higher temperatures have generally been postulated to increase the frequency of migratory phenotypes (Sogard *et al.* 2012; Sloat and Reeves 2014; Sloat *et al.* 2014; Kendall *et al.* 2014). However, the results in Chapter 3 suggest that warming acted in this case to increase early freshwater maturation. While the mechanisms behind warming-induced maturation are not immediately obvious, it seems that an acclimation response to temperature (*e.g.* in minimum energy requirements, similar to SMR reductions in Chapter 5) resulted in more individuals reaching their threshold for residency. Any such reduction in baseline energy costs would thus allow for accumulation of energy stores/body condition necessary for maturation. Temperature effects on maturation might thus have been mediated via effects on condition (or energy allocation) at key times (even though effects on size/mass were negative, we detected a neutral or positive relationship between temperature and condition that varied through time in Chapter 3). Positive relationships between body condition and temperatures up to 20 °C have been observed for Atlantic salmon within aquaculture settings (Tromp *et al.* 2018), and have been linked to more frequent and earlier maturation in the species (Good and Davidson 2016; Debes *et al.* 2019).

240 The unexpected results in Chapter 3 underscore that temperature changes, or indeed, any environmental changes, are likely to depend on the pre-existing conditions in a given environment. Changes in thermal regimes (or other freshwater conditions) that remain within the optimal range for growth may promote residency if physiological condition can be maintained, a scenario particularly likely in relatively cool systems that undergo warming (Benjamin *et*

245

al. 2013). This is in agreement with predictions from state-dependent models indicating that dramatic temperature changes ( $> 3^{\circ}\text{C}$ ) would be required to alter the balance of life-history expression in *O. mykiss* inhabiting cool streams (Satterthwaite *et al.* 2010). Nevertheless, the results in Chapters 2 and 3 collectively indicate that changes in the freshwater environment may still alter the frequency of the anadromous and resident life-history tactics relatively quickly. As implied by this thesis, such changes could occur at the population level via changes in individual energy uptake/allocation in response to abiotic conditions or fluctuations in local patterns of resource availability. However, from these results we can infer changes that may occur on a broader scale. The prevalence of anadromy generally increases with latitude, reflecting the gradient in the balance between freshwater and marine productivity (Gross *et al.* 1988). As such, observed and predicted decreases in macroinvertebrate prey (*e.g.* Durance and Ormerod 2007) might drive more marine migration in some areas, whereas increased eutrophication may promote freshwater residency in others (Gross 1987). Predicted increases in temperature (IPCC 2014) might overwhelmingly act to tip the balance from anadromy to residency across much of the species' distribution in northern Europe, if effects of warming on growth rates are largely positive and within thermal optima (*e.g.* Chapters 3 and 5; Benjamin *et al.* 2013). Thus, changes in the balance of anadromy *versus* residency may vary depending on geographic location, and proximity to physiological optima, either enhancing or reducing life history diversity. A shift towards uniformity in life histories will have negative implications for the "portfolio effect" in salmonines, where high levels of life-history diversity can buffer species as a whole from environmental change (Schindler *et al.* 2010, 2015). Changes in the relative frequency of alternative life histories might also have consequences for the broader ecosystem. For example, changes in marine nutrients supplied by returning anadromous fish can alter freshwater community structure and ecosystem function (Naiman *et al.* 2002; Janetski *et al.* 2009; Doughty *et al.* 2016), or even result in eco-evolutionary feedback dynamics (Auer *et al.* 2018).

### Proximate and ultimate factors influencing metabolic traits

While the importance of proximate factors in driving migratory tactics was the primary aim of Chapters 2 and 3, in Chapters 4 and 5 I explored how interactions  
 280 between environmental and intrinsic (population) factors mediate physiology at a more fundamental level: metabolic rate variation. Despite considerable intraspecific variation evident in minimum metabolic rates (Biro and Stamps 2010; Burton *et al.* 2011), the reasons underpinning the persistence of such variation are somewhat unclear. Lower SMR in offspring from non-anadromous  
 285 populations (Chapters 4 and 5), and in conditions of food restriction (Chapter 4) and a long-term temperature increase (Chapter 5) collectively support SMR variation arising from both environmental and population factors (Norin and Metcalfe 2019). Variation in maximum aerobic metabolism is less-often studied but purportedly similar, and likely to be of great ecological relevance (Metcalfe  
 290 *et al.* 2016; Norin and Clark 2016). That MMR and AS tended overall to be higher in the anadromous-background populations hints at a genetic component to these traits. Moreover, population-specific effects of food restriction (Chapter 4) and temperature (Chapter 5) point towards potential genotype-by-environment effects on MMR and AS.

295 Reductions in SMR in warm temperatures/low food conditions presumably occurred as plastic or acclimation responses to energetically challenging environments. Such responses are in line with the principle of “plastic metabolic floors” (a reduction in baseline energetic demands) as a key compensation mechanism in coping with inclement conditions (Sandblom *et al.* 2016).  
 300 However, MMR, and consequently, AS, appear to be somewhat more immutable and to be largely determined by population-specific factors. While capacity for physiological plasticity can improve resilience to predictable effects of warming (Seebacher *et al.* 2015), constraints in the flexibility of MMR and AS may mean that compensation responses to environmental change are insufficient  
 305 (Sandblom *et al.* 2016).

The population-level differences observed in SMR, MMR, and AS might be indicative of divergent plastic responses by populations, and also raises the

intriguing possibility of ultimate mechanisms underlying metabolic trait variation, *e.g.* selection within the anadromous background population favouring higher SMR, MMR, or AS facilitating fast growth (McCarthy 2000) or increased swimming performance (Eliason *et al.* 2011). Interestingly, though AS was marginally higher in food-restricted fish from the anadromous-background population (Chapter 4), AS was lower in fish originating from the same population after long-term warming (Chapter 5). While these variable responses may simply be related to food/temperature induced shifts in the patterns of future life-history trajectories (as seen in Chapter 2 and 3), it nonetheless raises questions about the future performance of this population with climate warming (Pörtner and Farrell 2008; Sandblom *et al.* 2016). Will constraints in the upper boundaries of aerobic metabolism limit the capacity of migratory populations to respond to change? Some population-specific responses to warming have been predicted in salmonines (Martins *et al.* 2011), but the viability of many migratory populations is in doubt in a rapidly warming world (Crozier *et al.* 2008; Jonsson and Jonsson 2009; Farrell 2009).

#### *Implications of metabolic trait variation for fitness*

Collectively, results from Chapters 4 and 5 indicate that while metabolism is generally assumed pivotal in shaping life histories though effects on fitness (Brown *et al.* 1993; Ricklefs and Wikelski 2002; Guderley and Pörtner 2010), this association is mediated by environmental and genetic factors. The negative relationship between SMR and growth under warm conditions in Chapter 5 adds further support to studies describing context-dependent fitness benefits of a given metabolic phenotype (Norin and Malte 2011; Auer *et al.* 2015a, b). Although growth is a key fitness-related trait, it is important to note here that improved growth performance may not necessarily translate into improved fitness (Mangel and Stamps 2001), and growth is itself a trait likely to show context-dependent fitness benefits. For example, faster growth in populations that tend to migrate might confer additional fitness benefits in terms of survival-at-sea (Kendall *et al.* 2014; Armstrong *et al.* 2018) but could be less important for non-migratory populations, and may thus have contributed to the

population-specific variation in the relationship between MMR and AS and  
 340 growth observed in Chapter 5. Examination of additional fitness-related traits  
 (e.g. lipid stores) may reveal population-specific differences in how surplus  
 energy (fuelled by MMR and AS) is allocated and traded off between competing  
 functions.

On a broader scale, the ability to adjust physiological components in response  
 345 to environmental factors has been positively associated with species' resilience  
 to global change (Seebacher *et al.* 2015). However, the advantages of phenotypic  
 plasticity for population persistence will depend upon both the speed at which  
 the phenotype can change, but also the nature of the environmental change, i.e.  
 whether fluctuations are predictable, and how accurately the current  
 350 environment predicts future conditions (Reed *et al.* 2010). It is also important  
 to note that in Chapters 4 and 5, fish were exposed to long-term changes in food  
 and temperature but acute responses/short-term changes in environmental  
 conditions were not considered, which could have altered the relationship  
 between metabolic and fitness-related traits. This is particularly relevant since  
 355 more frequent extreme weather events are forecasted with a changing climate  
 (Meehl and Tebaldi 2004; Seneviratne *et al.* 2014), particularly if some acute  
 effects (e.g. those affecting locomotion/survival) may be more important for  
 fitness than longer term ones (e.g. growth). Nevertheless, understanding how  
 facultatively migratory species such as brown trout respond to global change in  
 360 terms of energetics may potentially contribute some mechanistic insights into  
 changes that might scale up and partially underpin any changes in life histories.  
 For example, though I did not directly test the association between metabolic  
 rates and life-history expression at the individual level, the capacity of fish to  
 lower SMR over a long timescale may shed some light on why relatively high  
 365 rates of maturation were still possible in the low food conditions and warm  
 temperature treatments in Chapters 2 and 3. Building on the knowledge of how  
 various metabolic components relate to life history, and how this nexus is  
 mediated by environmental factors will provide further insight into the  
 ecological significance of metabolic variation.

## 370 **Considerations and limitations of the research**

### *Experimental approach*

The results presented here help to reveal how environmental factors can combine with intrinsic factors to influence phenotypic diversity, but it is important to consider them within the context of the experimental tank-based  
 375 approach that formed the basis for this research. While experimental tank rearing is a powerful approach that allows clean manipulation of key environmental variables of interest, while holding other factors constant, these advantages must be traded off against the inevitable loss of natural complexity in an artificial setting. Inferring the applicability of findings from controlled  
 380 experiments to wild populations is somewhat difficult, and effects described here may vary in nature depending on the location or population considered. For example, while I explored combined environmental factors in Chapter 2, responses in wild populations may be swamped by additional stressors or other confounding factors (Merilä and Hendry 2014; de Eyto *et al.* 2016; Galic *et al.*  
 385 2018). Extending this common-garden approach to more variable natural systems, or reciprocal transplant experiments in the wild between populations of divergent migratory tactics would complement the tank-based manipulations described here and give insight on whether these findings hold up in more realistic and ecologically relevant contexts.

390 An additional drawback of this experimental approach is the considerable spatial and logistical constraints associated with the housing and husbandry of large numbers of fish under artificial conditions. A recurring limitation arising from spatial constraints is replication of experimental manipulations at the tank-level, which would allow any potential inter-tank variation/noise to be  
 395 estimated and accounted for. Although I attempted to minimise any such nuisance variation related to tank effects (*e.g.* by standardising all conditions besides manipulated variables, analysing the data at the individual level and fitting random effects where appropriate), additional tank-level replication would make any future experimental studies more robust. Nonetheless, despite  
 400 any potential nuisance variation related to tank effects, the results I describe

here are broadly in line with theoretical predictions, suggesting that conclusions drawn with respect to population and environmental effects are more parsimonious than simple tank-related effects. Likewise, logistics of tank rearing also limited our capacity for replicating treatments at the population level. While the use of two populations in Chapters 2, 4, and 5 is an advance on a single population approach - a common limitation among studies of facultatively migratory species (Olsson *et al.* 2006; Wysujack *et al.* 2009; Chapman *et al.* 2011) - inclusion of additional populations would determine the general applicability and robustness of my findings described here. This is particularly desirable given population-level variation is a key outcome of this thesis (Chapters 2, 4, and 5).

The importance of population background for traits related to both migration and metabolism throughout much of this thesis is clear. However, less obvious is whether such population-background effects can be attributed to life-history differences between populations, or are simply indicative of the different catchments of origin (i.e. the various populations having evolved in distinct river systems, possibly originating from different lineages). For example, brood stock from each population were of unknown life history, but were assumed to represent the naturally occurring balance of migration *versus* residency for each population (ranging from non-anadromous, to strongly anadromous). Using parents of a known life history (*e.g.* via stable isotope analysis) would help to parse out factors related specifically to life history from those related to various other population-level differences. This would be particularly powerful if representatives of each possible life-history phenotype could be obtained as brood stock from populations, however, this is not always possible (*e.g.* in wholly resident populations). The presence of reciprocal hybrids from each population/life history would also be desirable to control for parental effects, which can be considerable (Taborsky 2006; Burton *et al.* 2013; Moore *et al.* 2019). Nevertheless, life history is intricately linked to population-specific factors, being both proximately, and ultimately (via selective forces) determined by such factors (*e.g.* growth opportunity in the local environment), and is thus

likely to be representative of any major differences between populations. On a more philosophical note, disentangling whether divergent phenotypic responses among populations to environmental manipulations represents an adaptive genetic basis to the trait of interest, or implies other population-specific explanations (*e.g.* genetic drift or gene flow) is a fundamental challenge of the common-garden style approach employed in this thesis (Merilä and Hendry 2014). While the results in this thesis are largely consistent with a genetic basis to migration and metabolic traits, other potential population-specific explanations cannot be excluded completely.

### Conclusions and future directions

A recurring theme throughout this thesis is the decomposition of drivers underpinning phenotypic diversity into genetic and environmental components. The results described here offer some insight into how the culturally and economically important brown trout (and other salmonine species) will respond to changing environmental conditions, with implications for conservation and management strategies. The potential for (re-)emergence of migration in resident populations (Chapter 2; Thrower *et al.* 2004) offers some positive perspective on the restoration of anadromous salmonine populations that have drastically declined by as much as 80% in recent years (Limburg and Waldman 2009). Similar experiments using salmonine populations where migration has ceased due to impassable barriers/ dams would help to clarify further whether freshwater resident populations maintain the capacity to produce migratory individuals.

The general flexibility in life-history expression displayed by my focal populations offers some encouraging evidence that brown trout have the capacity to maintain a diverse suite of life histories among populations, which may help to cushion the entire species from global change (Schindler *et al.* 2015). However, the ecological and evolutionary impacts of any environmental changes on physiology and life-history expression will depend on a multitude of factors, not least geographic location and population-specific proximity to optimal growth. Some populations may show increased diversity, whereas



others may become more uniform. The consequences of altered intraspecific diversity require further investigation (Bálint *et al.* 2011; Ceballos *et al.* 2017),  
 465 but are somewhat analogous to biotic homogenisation at the community level in response to climate change, with negative impacts on community stability (O’Gorman *et al.* 2012, 2019). The logical next step is to consider environmental change across the broader range of brown trout distribution, in populations that may be closer to their thermal limits or those experiencing additional pressures  
 470 from interspecific dynamics or disease. Consideration of multiple, co-occurring environmental factors (*e.g.* Chapter 2) is essential in future studies in order to make realistic predictions regarding environmental change (Côté *et al.* 2016). A novel approach would be to take advantage of natural environmental gradients, which allow for *in situ* manipulation of environmental conditions. So-called  
 475 “natural laboratories” such as geothermally-heated systems (Woodward *et al.* 2010; O’Gorman *et al.* 2014), or areas warmed as a by-product of nuclear energy production (Sandblom *et al.* 2016; Huss *et al.* 2019) offer unique opportunities to explore the effects of warming/biotic interactions on metabolic traits and life-histories (*e.g.* O’Gorman *et al.* 2016; Pilakouta *et al.* 2019). By combining natural  
 480 complexity with semi-controlled conditions, these could prove to be an exciting extension to the common-garden or translocation experiments that are invaluable to disentangling environmentally-induced/plastic responses from long-term evolutionary change.

While much of this thesis has focused on direct effects of proximate factors on  
 485 phenotypic diversity, the existence of an underlying genetic component to migration-associated traits is clear. Consideration of the evolutionary effects of environmental forces, or how these may indirectly alter life histories via ultimate mechanisms is equally important. For example, directly testing how changes in freshwater conditions can alter the future fitness of various  
 490 phenotypes (*e.g.* reduced salinity tolerance in Chapter 3 and growth in Chapter 5) will indicate how ultimate mechanisms give rise to phenotypic change within, and divergence across, populations. Beyond effects in the natal freshwaters (explored in this thesis), environmental changes on the migration

journey, and in the destination environment, will alter the costs and benefits of  
 495 each tactic, and ultimately the relative fitness of migrants *versus* residents.  
 Direct stress from various factors – changes in pH, oxygen levels, temperatures  
 – on the migration route (Eliason *et al.* 2011; Peiman *et al.* 2017), or large-scale  
 shifts in freshwater or marine food webs will likely impose strong selection on  
 facultatively migratory species (Crozier and Hutchings 2014). Such  
 500 environmental changes could act as forces of selection on underlying migration  
 thresholds, or on the genes influencing physiological condition or  
 energetic/metabolic traits (Phillis *et al.* 2016) with consequences for migration  
 propensity, or perhaps, migration destination. Future studies on (i) the  
 heritability of metabolic phenotypes, energy budgets, and migratory tactics, and  
 505 (ii) their relative fitness in different contexts are fundamental to understanding  
 the evolutionary basis to alternative life histories, along with the physiological  
 underpinnings.

In an era of rapid global change and biodiversity loss, there is a pressing need  
 to understand how species with strikingly diverse and complex life histories  
 510 might respond. Can facultatively migratory species display sufficient flexibility  
 to cope with changes to key elements of the environment? The results of this  
 thesis suggest that brown trout have considerable capacity to respond in terms  
 of physiology and life history, but outstanding questions still remain on the  
 future of migratory species (Lennox *et al.* 2019). Developing further knowledge  
 515 of how genetic and environmental forces interactively shape physiology, and  
 ultimately, life histories is essential to anticipating and managing the  
 consequences of anthropogenic change for declining salmonine populations.

## References

- 520 Acolas ML, Labonne J, Baglinière JL, and Roussel JM. 2012. The role of body size versus  
 growth on the decision to migrate: A case study with *Salmo trutta*.  
*Naturwissenschaften* **99**: 11–21.
- Archer LC, Hutton SA, Harman L, O’Grady MN, Kerry JP, Poole WR, Gargan P,  
 McGinnity P, and Reed TE. 2019. The interplay between extrinsic and intrinsic  
 525 factors in determining migration decisions in brown trout (*Salmo trutta*): An  
 experimental study. *Frontiers in Ecology and Evolution* **7**: 222.

- Armstrong JD, McKelvey S, Smith GW, Rycroft P, and Fryer RJ. 2018. Effects of individual variation in length, condition and run-time on return rates of wild-reared Atlantic salmon *Salmo salar* smolts. *Journal of Fish Biology* **92**: 569–78.
- 530 Auer SK, Anderson GJ, McKelvey S, Bassar RD, McLennan D, Armstrong JD, Nislow KH, Downie HK, McKelvey L, Morgan TAJ, Salin K, Orrell DL, Gauthey A, Reid TC, and Metcalfe NB. 2018. Nutrients from salmon parents alter selection pressures on their offspring. *Ecology Letters* **21**: 287–95.
- Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015a. The optimal combination of standard metabolic rate and aerobic scope for somatic growth depends on food availability. *Functional Ecology* **29**: 479–86.
- 535 Auer SK, Salin K, Rudolf AM, Anderson GJ, and Metcalfe NB. 2015b. Flexibility in metabolic rate confers a growth advantage under changing food availability. *Journal of Animal Ecology* **84**: 1405–11.
- Bálint M, Domisch S, Engelhardt CHM, Haase P, Lehrian S, Sauer J, Theissinger K, Pauls SU, and Nowak C. 2011. Cryptic biodiversity loss linked to global climate change. *Nature Climate Change* **1**: 313–8.
- 540 Bateson P and Laland KN. 2013. Tinbergen's four questions: An appreciation and an update. *Trends in Ecology & Evolution* **28**: 712–8.
- Benjamin JR, Connolly PJ, Romine JG, and Perry RW. 2013. Potential effects of changes in temperature and food resources on life history trajectories of juvenile *Oncorhynchus mykiss*. *Transactions of the American Fisheries Society* **142**: 208–20.
- 545 Birnie-Gauvin K, Thorstad EB, and Aarestrup K. 2019. Overlooked aspects of the *Salmo salar* and *Salmo trutta* lifecycles. *Reviews in Fish Biology and Fisheries*.
- Biro PA and Stamps JA. 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? *Trends in Ecology & Evolution* **25**: 653–9.
- 550 Boel M, Aarestrup K, Baktoft H, Larsen T, Søndergaard Madsen S, Malte H, Skov C, Svendsen JC, and Koed A. 2014. The physiological basis of the migration continuum in brown trout (*Salmo trutta*). *Physiological and Biochemical Zoology* **87**: 334–45.
- 555 Bolnick DI, Amarasekare P, Araújo MS, Bürger R, Levine JM, Novak M, Rudolf VHW, Schreiber SJ, Urban MC, and Vasseur DA. 2011. Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution* **26**: 183–92.
- 560 Brown JH, Marquet PA, and Taper ML. 1993. Evolution of body size: Consequences of an energetic definition of fitness. *The American Naturalist* **142**: 573–84.
- Buoro M, Gimenez O, and Prévost E. 2012. Assessing adaptive phenotypic plasticity by means of conditional strategies from empirical data: The latent environmental threshold model. *Evolution* **66**: 996–1009.
- 565 Burton T, Killen SS, Armstrong JD, and Metcalfe NB. 2011. What causes intraspecific variation in resting metabolic rate and what are its ecological consequences? *Proceedings of the Royal Society B: Biological Sciences* **278**: 3465–73.
- Burton T, McKelvey S, Stewart DC, Armstrong JD, and Metcalfe NB. 2013. Early maternal experience shapes offspring performance in the wild. *Ecology* **94**: 618–26.
- 570

- Ceballos G, Ehrlich PR, and Dirzo R. 2017. Biological annihilation via the ongoing sixth mass extinction signaled by vertebrate population losses and declines. *Proceedings of the National Academy of Sciences* **114**: E6089–96.
- 575 Chapman BB, Brönmark C, Nilsson J-Å, and Hansson L-A. 2011. The ecology and evolution of partial migration. *Oikos* **120**: 1764–75.
- Clarke A and Johnston NM. 1999. Scaling of metabolic rate with body mass and temperature in teleost fish. *Journal of Animal Ecology* **68**: 893–905.
- 580 Côté IM, Darling ES, and Brown CJ. 2016. Interactions among ecosystem stressors and their importance in conservation. *Proceedings of the Royal Society B: Biological Sciences* **283**: 20152592.
- Crozier LG, Hendry AP, Lawson PW, Quinn TP, Mantua NJ, Battin J, Shaw RG, and Huey RB. 2008. Potential responses to climate change in organisms with complex life histories: Evolution and plasticity in Pacific salmon. *Evolutionary Applications* **1**: 252–70.
- 585 Crozier LG and Hutchings JA. 2014. Plastic and evolutionary responses to climate change in fish. *Evolutionary Applications* **7**: 68–87.
- Cucherousset J, Ombredane D, Charles K, Marchand F, and Baglinière J-L. 2005. A continuum of life history tactics in a brown trout (*Salmo trutta*) population. *Canadian Journal of Fisheries and Aquatic Sciences* **62**: 1600–10.
- 590 Debes PV, Piavachenko N, Ruokolainen A, Ovaskainen O, Moustakas-Verho JE, Parre N, Aykanat T, Erkinaro J, and Primmer CR. 2019. Large single-locus effects for maturation timing are mediated via condition variation in Atlantic salmon. *bioRxiv* doi/10.1101/780437.
- 595 Des Roches S, Post DM, Turley NE, Bailey JK, Hendry AP, Kinnison MT, Schweitzer JA, and Palkovacs EP. 2018. The ecological importance of intraspecific variation. *Nature Ecology & Evolution* **2**: 57–64.
- Doctor K, Berejikian B, Hard JJ, and VanDoornik D. 2014. Growth-mediated life history traits of steelhead reveal phenotypic divergence and plastic response to temperature. *Transactions of the American Fisheries Society* **143**: 317–33.
- 600 Doughty CE, Roman J, Faurby S, Wolf A, Haque A, Bakker ES, Malhi Y, Dunning JB, and Svenning J-C. 2016. Global nutrient transport in a world of giants. *Proceedings of the National Academy of Sciences* **113**: 868–73.
- Durance I and Ormerod SJ. 2007. Climate change effects on upland stream macroinvertebrates over a 25-year period. *Global Change Biology* **13**: 942–57.
- 605 Eliason EJ, Clark TD, Hague MJ, Hanson LM, Gallagher ZS, Jeffries KM, Gale MK, Patterson DA, Hinch SG, and Farrell AP. 2011. Differences in thermal tolerance among sockeye salmon populations. *Science* **332**: 109–12.
- 610 Eyto E de, Dalton C, Dillane M, Jennings E, McGinnity P, O'Dwyer B, Poole R, Rogan G, and Taylor D. 2016. The response of North Atlantic diadromous fish to multiple stressors, including land use change: A multidecadal study. *Canadian Journal of Fisheries and Aquatic Sciences* **73**: 1759–69.
- Farrell AP. 2009. Environment, antecedents and climate change: Lessons from the study of temperature physiology and river migration of salmonids. *Journal of Experimental Biology* **212**: 3771–80.

- 615 Ferguson A, Reed TE, Cross TF, McGinnity P, and Prodöhl PA. 2019. Anadromy, potamodromy and residency in brown trout *Salmo trutta*: The role of genes and the environment. *Journal of Fish Biology* **95**: 692–718.
- Forseth T, Nesje TF, Jonsson B, and Hårsaker K. 1999. Juvenile migration in brown trout: A consequence of energetic state. *Journal of Animal Ecology* **68**: 783–93.
- 620 Fry FE. 1971. The effect of environmental factors on the physiology of fish. In: Fish physiology (eds WS Hoar & DJ Randall). New York, NY: Academic Press.
- Galic N, Sullivan LL, Grimm V, and Forbes VE. 2018. When things don't add up: Quantifying impacts of multiple stressors from individual metabolism to ecosystem processing. *Ecology Letters* **21**: 568–77.
- 625 Gargan P, Kelly F, Shephard S, and Whelan K. 2016. Temporal variation in sea trout *Salmo trutta* life history traits in the Erriff River, western Ireland. *Aquaculture Environment Interactions* **8**: 675–89.
- Gillooly JF, Brown JH, West GB, Savage VM, and Charnov EL. 2001. Effects of size and temperature on metabolic rate. *Science* **293**: 2248–51.
- 630 Good C and Davidson J. 2016. A review of factors influencing maturation of Atlantic salmon, *Salmo salar*, with focus on water recirculation aquaculture system environments. *Journal of the World Aquaculture Society* **47**: 605–32.
- Gross MR. 1987. Evolution of diadromy in fishes. *Common Strategies of Anadromous and Catadromous Fishes*: 14–25.
- 635 Gross MR, Coleman RM, and McDowall RM. 1988. Aquatic productivity and the evolution of diadromous fish migration. *Science* **239**: 1291–3.
- Guderley H and Pörtner HO. 2010. Metabolic power budgeting and adaptive strategies in zoology: Examples from scallops and fish. *Canadian Journal of Zoology* **88**: 753–63.
- 640 Hecht BC, Hard JJ, Thrower FP, and Nichols KM. 2015. Quantitative genetics of migration-related traits in rainbow and steelhead trout. *G3: Genes, Genomes, Genetics* **5**: 873–89.
- Huss M, Lindmark M, Jacobson P, Dorst RM van, and Gårdmark A. 2019. Experimental evidence of gradual size-dependent shifts in body size and growth of fish in response to warming. *Global Change Biology* **25**: 2285–95.
- 645 IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]. In: Geneva, Switzerland.
- 650 Janetski DJ, Chaloner DT, Tiegs SD, and Lamberti GA. 2009. Pacific salmon effects on stream ecosystems: A quantitative synthesis. *Oecologia* **159**: 583–95.
- Jones DA, Bergman E, and Greenberg L. 2015. Food availability in spring affects smolting in brown trout (*Salmo trutta*). *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 1694–9.
- 655 Jonsson B. 1985. Life history patterns of freshwater resident and sea-run migrant brown trout in Norway. *Transactions of the American Fisheries Society* **114**: 182–94.

- Jonsson B, Finstad AG, and Jonsson N. 2012. Winter temperature and food quality affect age at maturity: An experimental test with Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences* **69**: 1817–26.
- 660 Jonsson B and Jonsson N. 2009. A review of the likely effects of climate change on anadromous Atlantic salmon *Salmo salar* and brown trout *Salmo trutta*, with particular reference to water temperature and flow. *Journal of Fish Biology* **75**: 2381–447.
- 665 Kammerer BD and Heppell SA. 2013. Individual condition indicators of thermal habitat quality in field populations of redband trout (*Oncorhynchus mykiss gairdneri*). *Environmental Biology of Fishes* **96**: 823–35.
- Kendall NW, McMillan JR, Sloat MR, Buehrens TW, Quinn TP, Pess GR, Kuzishchin KV, McClure MM, and Zabel RW. 2014. Anadromy and residency in steelhead and rainbow trout (*Oncorhynchus mykiss*): A review of the processes and patterns. *Canadian Journal of Fisheries and Aquatic Sciences* **72**: 319–42.
- 670 Klemetsen A, Amundsen P-A, Dempson JB, Jonsson B, Jonsson N, O'Connell MF, and Mortensen E. 2003. Atlantic salmon *Salmo salar* L., brown trout *Salmo trutta* L. and Arctic charr *Salvelinus alpinus* (L.): A review of aspects of their life histories. *Ecology of Freshwater Fish* **12**: 1–59.
- 675 Laland KN, Sterelny K, Odling-Smee J, Hoppitt W, and Uller T. 2011. Cause and effect in biology revisited: Is Mayr's proximate-ultimate dichotomy still useful? *Science* **334**: 1512–6.
- Lennox RJ, Paukert CP, Aarestrup K, Auger-Méthé M, Baumgartner L, Birnie-Gauvin K, Bøe K, Brink K, Brownscombe JW, Chen Y, Davidsen JG, Eliason EJ, Filous A, Gillanders BM, Helland IP, Horodysky AZ, Januchowski-Hartley SR, Lowerre-Barbieri SK, Lucas MC, Martins EG, Murchie KJ, Pompeu PS, Power M, Raghavan R, Rahel FJ, Secor D, Thiem JD, Thorstad EB, Ueda H, Whoriskey FG, and Cooke SJ. 2019. One hundred pressing questions on the future of global fish migration science, conservation, and policy. *Frontiers in Ecology and*
- 680 *Evolution* **7**.
- 685 Limburg KE and Waldman JR. 2009. Dramatic declines in north Atlantic diadromous fishes. *BioScience* **59**: 955–65.
- Mangel M and Stamps JA. 2001. Trade-offs between growth and mortality and the maintenance of individual variation in growth. *Evolutionary Ecology Research*
- 690 **3**: 611–32.
- Martins EG, Hinch SG, Patterson DA, Hague MJ, Cooke SJ, Miller KM, Lapointe MF, English KK, and Farrell AP. 2011. Effects of river temperature and climate warming on stock-specific survival of adult migrating Fraser River sockeye salmon (*Oncorhynchus nerka*). *Global Change Biology* **17**: 99–114.
- 695 McCarthy ID. 2000. Temporal repeatability of relative standard metabolic rate in juvenile Atlantic salmon and its relation to life history variation. *Journal of Fish Biology* **57**: 224–38.
- McMillan JR, Dunham JB, Reeves GH, Mills JS, and Jordan CE. 2012. Individual condition and stream temperature influence early maturation of rainbow and steelhead trout, *Oncorhynchus mykiss*. *Environmental Biology of Fishes* **93**: 343–55.
- 700

- Meehl GA and Tebaldi C. 2004. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* **305**: 994–7.
- 705 Merilä J and Hendry AP. 2014. Climate change, adaptation, and phenotypic plasticity: the problem and the evidence. *Evolutionary Applications* **7**: 1–14.
- Metcalfe NB, Van Leeuwen TE, and Killen SS. 2016. Does individual variation in metabolic phenotype predict fish behaviour and performance? *Journal of Fish Biology* **88**: 298–321.
- 710 Moore MP, Whiteman HH, and Martin RA. 2019. A mother's legacy: The strength of maternal effects in animal populations. *Ecology Letters* **22**: 1620–8.
- Morinville GR and Rasmussen JB. 2003. Early juvenile bioenergetic differences between anadromous and resident brook trout (*Salvelinus fontinalis*). *Canadian Journal of Fisheries and Aquatic Sciences* **60**: 401–10.
- 715 Naiman RJ, Bilby RE, Schindler DE, and Helfield JM. 2002. Pacific salmon, nutrients, and the dynamics of freshwater and riparian ecosystems. *Ecosystems* **5**: 399–417.
- Naish KA and Hard JJ. 2008. Bridging the gap between the genotype and the phenotype: Linking genetic variation, selection and adaptation in fishes. *Fish and Fisheries* **9**: 396–422.
- 720 Nevoux M, Finstad B, Davidsen JG, Finlay R, Josset Q, Poole R, Höjesjö J, Aarestrup K, Persson L, Tolvanen O, and Jonsson B. 2019. Environmental influences on life history strategies in partially anadromous brown trout (*Salmo trutta*, Salmonidae). *Fish and Fisheries* **In Press**.
- 725 Norin T and Clark TD. 2016. Measurement and relevance of maximum metabolic rate in fishes. *Journal of Fish Biology* **88**: 122–51.
- Norin T and Malte H. 2011. Repeatability of standard metabolic rate, active metabolic rate and aerobic scope in young brown trout during a period of moderate food availability. *Journal of Experimental Biology* **214**: 1668–75.
- 730 Norin T and Metcalfe N. 2019. Ecological and evolutionary consequences of metabolic rate plasticity in response to environmental change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **374**: 20180180.
- O'Gorman EJ, Benstead JP, Cross WF, Friberg N, Hood JM, Johnson PW, Sigurdsson BD, and Woodward G. 2014. Climate change and geothermal ecosystems: Natural laboratories, sentinel systems, and future refugia. *Global Change Biology* **20**: 3291–9.
- 735 O'Gorman EJ, Ólafsson ÓP, Demars BOL, Friberg N, Guðbergsson G, Hannesdóttir ER, Jackson MC, Johansson LS, McLaughlin ÓB, Ólafsson JS, Woodward G, and Gíslason GM. 2016. Temperature effects on fish production across a natural thermal gradient. *Global Change Biology* **22**: 3206–20.
- 740 O'Gorman EJ, Petchey OL, Faulkner KJ, Gallo B, Gordon TAC, Neto-Cerejeira J, Ólafsson JS, Pichler DE, Thompson MSA, and Woodward G. 2019. A simple model predicts how warming simplifies wild food webs. *Nature Climate Change* **9**: 611–6.
- 745 O'Gorman EJ, Pichler DE, Adams G, Benstead JP, Cohen H, Craig N, Cross WF, Demars BOL, Friberg N, Gíslason GM, Gudmundsdóttir R, Hawczak A, Hood JM, Hudson LN, Johansson L, Johansson MP, Junker JR, Laurila A, Manson JR,

- Mavromati E, Nelson D, Ólafsson JS, Perkins DM, Petchey OL, Plebani M, Reuman DC, Rall BC, Stewart R, Thompson MSA, and Woodward G. 2012. Chapter 2 - Impacts of warming on the structure and functioning of aquatic communities: Individual- to ecosystem-level responses. In: Woodward G, Jacob U, O’Gorman EJ (Eds). *Advances in Ecological Research*. Cambridge, Massachusetts: Academic Press.
- 750 Olsson IC, Greenberg LA, Bergman E, and Wysujack K. 2006. Environmentally induced migration: The importance of food. *Ecology Letters* **9**: 645–51.
- 755 O’Neal SL and Stanford JA. 2011. Partial migration in a robust brown trout population of a Patagonian river. *Transactions of the American Fisheries Society* **140**: 623–35.
- 760 Peiman KS, Birnie-Gauvin K, Midwood JD, Larsen MH, Wilson ADM, Aarestrup K, and Cooke SJ. 2017. If and when: Intrinsic differences and environmental stressors influence migration in brown trout (*Salmo trutta*). *Oecologia* **184**: 375–84.
- Phillis CC, Moore JW, Buoro M, Hayes SA, Garza JC, and Pearse DE. 2016. Shifting thresholds: Rapid evolution of migratory life histories in steelhead/rainbow trout, *Oncorhynchus mykiss*. *Journal of Heredity* **107**: 51–60.
- 765 Piche J, Hutchings JA, and Blanchard W. 2008. Genetic variation in threshold reaction norms for alternative reproductive tactics in male Atlantic salmon, *Salmo salar*. *Proceedings of the Royal Society B-Biological Sciences* **275**: 1571–5.
- Pilakouta N, Killen SS, Kristjánsson BK, Skúlason S, Lindström J, Metcalfe NB, and Parsons KJ. 2019. Reduction in standard metabolic rate after multigenerational exposure to elevated temperatures in the wild. *bioRxiv* doi/10.1101/749986.
- 770 Poole WR, Dillane M, DeEyto E, Rogan G, McGinnity P, and Whelan K. 2007. Characteristics of the Burrishoole sea trout population: Census, marine survival, enhancement and stock-recruitment relationship, 1971-2003. In: Sea Trout: Biology, Conservation and Management (Eds: G Harris and N Milner). Oxford, UK: Blackwells Scientific Publications.
- 775 Poole WR, Whelan KF, Dillane MG, Cooke DJ, and Matthews M. 1996. The performance of sea trout, *Salmo trutta* L., stocks from the Burrishoole system western Ireland, 1970–1994. *Fisheries Management and Ecology* **3**: 73–92.
- Pörtner HO and Farrell AP. 2008. Physiology and climate change. *Science* **322**: 690–2.
- 780 Pulido F. 2011. Evolutionary genetics of partial migration – the threshold model of migration revis(it)ed. *Oikos* **120**: 1776–83.
- Reed TE, Waples Robin S., Schindler Daniel E., Hard Jeffrey J., and Kinnison Michael T. 2010. Phenotypic plasticity and population viability: The importance of environmental predictability. *Proceedings of the Royal Society B: Biological Sciences* **277**: 3391–400.
- 785 Ricklefs RE and Wikelski M. 2002. The physiology/life-history nexus. *Trends in Ecology & Evolution* **17**: 462–8.
- Roff DA. 1996. The evolution of threshold traits in animals. *The Quarterly Review of Biology* **71**: 3–35.
- 790 Sandblom E, Clark TD, Gräns A, Ekström A, Brijs J, Sundström LF, Odelström A, Adill A, Aho T, and Jutfelt F. 2016. Physiological constraints to climate warming in



- fish follow principles of plastic floors and concrete ceilings. *Nature Communications* **7**: 11447.
- 795 Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, and Mangel M. 2009. Steelhead life history on California's Central Coast: Insights from a state-dependent model. *Transactions of the American Fisheries Society* **138**: 532–48.
- 800 Satterthwaite WH, Beakes MP, Collins EM, Swank DR, Merz JE, Titus RG, Sogard SM, and Mangel M. 2010. State-dependent life history models in a changing (and regulated) environment: Steelhead in the California Central Valley. *Evolutionary Applications* **3**: 221–43.
- Schindler DE, Armstrong JB, and Reed TE. 2015. The portfolio concept in ecology and evolution. *Frontiers in Ecology and the Environment* **13**: 257–63.
- 805 Schindler DE, Hilborn R, Chasco B, Boatright CP, Quinn TP, Rogers LA, and Webster MS. 2010. Population diversity and the portfolio effect in an exploited species. *Nature* **465**: 609–12.
- Seebacher F, White CR, and Franklin CE. 2015. Physiological plasticity increases resilience of ectothermic animals to climate change. *Nature Climate Change* **5**: 61–6.
- 810 Seneviratne SI, Donat MG, Mueller B, and Alexander LV. 2014. No pause in the increase of hot temperature extremes. *Nature Climate Change* **4**: 161–3.
- Sloat MR, Fraser DJ, Dunham JB, Falke JA, Jordan CE, McMillan JR, and Ohms HA. 2014. Ecological and evolutionary patterns of freshwater maturation in Pacific and Atlantic salmonines. *Reviews in Fish Biology and Fisheries* **24**: 689–707.
- 815 Sloat MR and Reeves GH. 2014. Individual condition, standard metabolic rate, and rearing temperature influence steelhead and rainbow trout (*Oncorhynchus mykiss*) life histories. *Canadian Journal of Fisheries and Aquatic Sciences* **71**: 491–501.
- 820 Sogard SM, Merz JE, Satterthwaite WH, Beakes MP, Swank DR, Collins EM, Titus RG, and Mangel M. 2012. Contrasts in habitat characteristics and life history patterns of *Oncorhynchus mykiss* in California's Central Coast and Central Valley. *Transactions of the American Fisheries Society* **141**: 747–60.
- Taborsky B. 2006. Mothers determine offspring size in response to own juvenile growth conditions. *Biology Letters* **2**: 225–8.
- 825 Thorpe JE, Mangel M, Metcalfe NB, and Huntingford FA. 1998. Modelling the proximate basis of salmonid life-history variation, with application to Atlantic salmon, *Salmo salar* L. *Evolutionary Ecology* **12**: 581–99.
- Thorpe JE and Metcalfe NB. 1998. Is smolting a positive or a negative developmental decision? *Aquaculture* **168**: 95–103.
- 830 Thrower FP, Hard JJ, and Joyce JE. 2004. Genetic architecture of growth and early life-history transitions in anadromous and derived freshwater populations of steelhead. *Journal of Fish Biology* **65**: 286–307.
- Tinbergen N. 1963. On aims and methods of Ethology. *Zeitschrift für Tierpsychologie* **20**: 410–33.
- 835 Tomkins JL and Hazel W. 2007. The status of the conditional evolutionarily stable strategy. *Trends in Ecology & Evolution* **22**: 522–8.

- Tromp JJ, Jones PL, Brown MS, Donald JA, Biro PA, and Afonso LOB. 2018. Chronic exposure to increased water temperature reveals few impacts on stress physiology and growth responses in juvenile Atlantic salmon. *Aquaculture* **495**: 196–204.
- 840 Winter ER, Tummers JS, Aarestrup K, Baktoft H, and Lucas MC. 2016. Investigating the phenology of seaward migration of juvenile brown trout (*Salmo trutta*) in two European populations. *Hydrobiologia* **775**: 139–51.
- Woodward G, Dybkjær JB, Ólafsson JS, Gíslason GM, Hannesdóttir ER, and Friberg N. 2010. Sentinel systems on the razor's edge: Effects of warming on Arctic geothermal stream ecosystems. *Global Change Biology* **16**: 1979–91.
- 845 Wysujack K, Greenberg LA, Bergman E, and Olsson IC. 2009. The role of the environment in partial migration: Food availability affects the adoption of a migratory tactic in brown trout *Salmo trutta*. *Ecology of Freshwater Fish* **18**: 52–9