CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	The provenance of the Devonian Old Red Sandstone of the Dingle Peninsula, SW Ireland; the earliest record of Laurentian and peri-Gondwanan sediment mixing in Ireland
Authors	Fairey, Brenton J.;Kerrison, Aidan;Meere, Patrick A.;Mulchrone, Kieran F.;Hofmann, Mandy;Gärtner, Andreas;Sonntag, Benita-Lisette;Linnemann, Ulf;Kuiper, Klaudia F.;Ennis, Meg;Mark, Chris;Cogné, Nathan;Chew, David
Publication date	2018-01-17
Original Citation	Fairey, B. J., Kerrison, A., Meere, P. A., Mulchrone, K. F., Hofmann, M., Gärtner, A., Sonntag, B.-L., Linnemann, U., Kuiper, K. F., Ennis, M., Mark, C., Cogné, N. and Chew, D. (2018) 'The provenance of the Devonian Old Red Sandstone of the Dingle Peninsula, SW Ireland; the earliest record of Laurentian and peri-Gondwanan sediment mixing in Ireland', Journal of the Geological Society, 175(3), pp. 411-424. doi:10.1144/jgs2017-099
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://doi.org/10.6084/m9.figshare.c.3971862 - 10.1144/jgs2017-099
Rights	© 2018, the Authors. Published by The Geological Society of London. All rights reserved. This document is the Accepted Manuscript version of a Published Work that appeared in final form in Journal of the Geological Society. To access the final edited and published work see https://doi.org/10.1144/jgs2017-099
Download date	2026-10-31 11:43:22
Item downloaded from	https://hdl.handle.net/10468/6494



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

The provenance of the Devonian Old Red Sandstone of the Dingle Peninsula, SW Ireland – the earliest record of Laurentian and peri-Gondwanan sediment mixing in Ireland.

Brenton J. Fairey[1], Aidan Kerrison[1], Patrick A. Meere[1], Kieran F. Mulchrone[2], Mandy Hofmann[3], Andreas Gärtner[3], Benita-Lisette Sonntag[3], Ulf Linnemann[3], Klaudia F. Kuiper[4], Meg Ennis[1], Chris Mark[5], Nathan Cogné[5] and David Chew[5].

[1]School of Biological, Earth and Environmental Sciences, University College Cork, Cork, Ireland.

[2]School of Applied Mathematics, University College Cork, Cork, Ireland

[3]Senckenberg Naturhistorische Sammlungen Dresden, Museum für Mineralogie und Geologie, Königsbrücker Landstraße 159, D-01109 Dresden, Germany.

[4]Faculty of Earth Sciences Vrije Universiteit Amsterdam, De Boelelaan 1085, NL-1081 HV Amsterdam, The Netherlands.

[5]Department of Geology, Museum Building, Trinity College Dublin, Dublin 2, Ireland.

Abstract

The Lower Old Red Sandstone (LORS) in southern Ireland is hosted in the Lower Devonian Dingle Basin which lies immediately south of the Iapetus Suture on the Dingle Peninsula, County Kerry. The basin developed as a post-Caledonian pull-apart structure prior to Acadian deformation which in turn was followed by end-Carboniferous Variscan deformation. Detrital zircon U-Th-Pb geochronology is complimented by mica Ar-Ar and apatite U-Pb geochronology to gain a

comprehensive understanding of the provenance of the Lower Devonian LORS of the Dingle Basin and assess contributions of major tectonic components (e.g. Laurentia, Ganderia). Sedimentary rocks in the LORS have similar detrital zircon age distributions which are dominated by ca. 1.2 Ga zircons as well as late Neoproterozoic grains. This indicates a dominant contribution of detritus of Laurentian affinity as well as contributions from westerly and southerly derived Ganderian detritus. Caledonian uplift of the area north of the Iapetus Suture would have facilitated a large contribution of (peri-)Laurentian material. The Upper Old Red Sandstone on the Dingle Peninsula has a distinctly different detrital zircon character including few late Neoproterozoic zircons and abundant zircons of ca. 1.05 Ga age, indicating sediment derivation only from Laurentia and no recycling from the LORS.

Supplementary material: the full detrital U-Pb zircon (Table-1) and apatite (Table-2) analytical dataset as well as revised detrital mica age dataset (Table-3) is available at xxxxx.

The Dingle Basin in County Kerry, southwest Ireland represents the only record of Early Devonian sedimentation south of the Iapetus Suture in Ireland. Its structure records a critical period in the tectonic history and development of Ireland, having been affected by both the Acadian and Variscan Orogenies (Meere & Mulchrone 2006). Furthermore, the basin's sedimentary rocks offer an opportunity to understand the palaeogeography of the basin, possibly recording Grampian to Late Caledonian (430-420 Ma) detrital input following the final Silurian accretion of Ganderia to the margin of Laurentia.

Existing provenance studies (e.g. Todd 2000; Ennis *et al.* 2015) have greatly improved our understanding of the development and palaeogeography of the Dingle Basin. However, a comprehensive investigation utilising detrital single-grain techniques has yet to be undertaken. Such a study would serve to elucidate regional sediment source contributions in a basin which is intimately associated with the Iapetus Suture (Todd 2000) – arguably the most important structural entity for in the Phanerozoic tectonic history of Ireland.

This study provides the first multiproxy (zircon, apatite, mica) single-grain datasets from the area, with the aim of determining the provenance of the Lower Old Red Sandstone (LORS) in the Dingle Group of the Lower Devonian Dingle Basin and assessing the roles of Laurentian and peri-Gondwanan domains in contributing detritus. The study also considers the detrital zircon provenance of the Upper Old Red Sandstone (UORS) on the Dingle Peninsula.

Regional Geology and Review of Terranes in the British Isles

The oldest potential source of detrital zircons lies to the north of the Iapetus Suture Zone (ISZ) in the form of the long-lived Laurentian craton (Fig. 1). A compilation of existing detrital zircon data of known Laurentian sources (Fig. 2) shows three broad peaks that correspond to major crust-forming events which contributed to the formation of Laurentia (Cawood *et al.* 2007). These peaks occur in the Archaean, Palaeoproterozoic and Mesoproterozoic recording essentially uninterrupted zircon production from 1.9 Ga to 0.9 Ga. A major characteristic of detrital zircon distributions from this sector of the Laurentian continent is the absence of zircons of late Neoproterozoic age due to the absence of an active margin on the Laurentian continent at this time (Pointon *et al.* 2012). The Rhinns Complex on the island of Inishtrahull off the north coast of County Donegal and the Annagh Gneiss

74 Complex in County Mayo comprise the exposed Proterozoic Laurentian basement in
75 Ireland, and are also the oldest rocks exposed in Ireland. The Grenville Orogeny,
76 represented by the large peak in late Mesoproterozoic zircon ages in Figure 2, is
77 recorded in the Annagh Gneiss Complex by the 1.17 Ga Doolough gneiss, the 1.01
78 Ga Doolough Granite and by 0.99 to 0.96 Ga late orogenic pegmatites and
79 migmatitic leucosomes (Daly 1996). A variety of marine sedimentary (e.g. Clew Bay
80 Complex), ocean-arc-volcanic (e.g. Lough Nafooey Arc – see (Chew *et al.* 2007) and
81 ophiolitic rocks (e.g. Deer Park Complex) make up the material accreted to the
82 margin of Laurentia during the Grampian Orogeny (465-475 Ma) in Ireland which
83 represents the early stage of the Caledonian orogenic cycle (Chew & Stillman 2009).
84 This was followed by sinistral transpressive docking of a peri-Gondwanan terrane to
85 the newly-accreted Laurentian margin (Dewey & Strachan 2003).

86 The term `peri-Gondwanide' was first proposed by Van Der Voo (1988) to
87 describe a number of tectonostratigraphic elements which existed as terranes in the
88 Iapetus Ocean during the Ordovician period. Using palaeomagnetic evidence, Van
89 Der Voo (1988) suggested that these terranes were proximal to northwest African
90 Gondwana, rather than Laurentia. This has since been supported by other studies
91 (see Nance *et al.* 2008, and references therein). We use the term *domain* to refer to
92 tectonostratigraphic units consisting of one or more terranes (Hibbard *et al.* 2007).
93 Peri-Gondwanan domains include Avalonia, Ganderia, Megumia, Carolinia in
94 present-day North America, and Avalonia and Cadomia (including Iberia, Bohemia
95 and Armorica) in Europe (Nance *et al.* 2008). Those domains that had the potential,
96 given their Devonian positions relative to southern Ireland, to provide sediment to the
97 basin being investigated include Avalonia, Ganderia and Megumia. For a
98 comprehensive review of peri-Gondwanan terranes, the reader is referred to Nance

et al. (2008). The correlation of the Meguma terrane to the Harlech Dome is discussed by Waldron *et al.* (2011), White *et al.* (2012) and Nance *et al.* (2015).

The main rock exposures of pre-Devonian basement south of the Iapetus Suture Zone in Ireland are found in the Leinster Massif in the southeastern part of the island. The massif hosts a number of Cambrian to Silurian volcanic and sedimentary units intruded by Caledonian-Acadian granites. There has been wide acceptance that pre-Silurian southern Ireland represents part of Avalonia (e.g. Van Der Voo 1983; Livermore *et al.* 1985; Van Der Voo 1988; Pickering *et al.* 1988; Ford *et al.* 1992; Cocks *et al.* 1997; McConnell & Morris 1997; Keppie *et al.* 2003; Tyrrell *et al.* 2007; Woodcock & Strachan 2012; Fullea *et al.* 2014; Todd 2015), but others have suggested linkages to different peri-Gondwanan domains. One school of thought contends that southern Ireland, Wales and southern England formed part of a number of terranes that collectively formed Cadomia (e.g. Soper & Hutton 1984; Max *et al.* 1990).

Kennedy (1979) first implied that southern Ireland is a trans-Atlantic extension of Ganderia. Van Staal *et al.* (1996) (and references therein) extend the Gander Zone of Newfoundland, where Ganderia was first described, into Ireland, Wales, England and the Isle of Man. As evidence of this they cited a correlation of Cambrian and Ordovician clastic successions, a similarity of overstepping successions, juxtaposition of mafic to ultramafic, possibly ophiolitic rocks (e.g. Rosslare Complex) and a similarity in fossil fauna. This extension of Ganderia into southern Ireland is further supported by Van Staal *et al.* (1998) who suggest that Avalonia and Ganderia may have become juxtaposed at least during the late Cambrian. If this is the case then it was an amalgamated Avalonia-Ganderia microcontinent that collided with Laurentia during the Caledonian Orogeny. More recently, Waldron *et*

al. (2014) have drawn similarities between detrital zircon ages from Monian Composite Terrane (county Wexford) and Leinster-Lakesman samples and samples analysed by Fyffe *et al.* (2009) for Ganderia in New Brunswick and Maine. The difference between Ganderian detrital zircon samples from Cambrian sedimentary rocks and West Avalonian samples is that the latter are lacking in Mesoproterozoic and Palaeoproterozoic zircons (Waldron *et al.* 2014). Waldron *et al.* (2014) attribute these Mesoproterozoic and Palaeoproterozoic zircons in Ganderian sediments to a possible Amazonian source in West Gondwana.

Distinguishing between Avalonia and Ganderia by detrital zircon populations alone is difficult (Fig. 2). Stratigraphic, faunal and other isotopic evidence is required, as exemplified in a review of the East Avalonian terranes by Schofield *et al.* (2016). They use geochronological data as well as magmatic whole-rock Sm-Nd and O-isotopes to show that what is currently viewed as East Avalonian basement in England has closer isotopic and age affinities to Ganderia than to West Avalonia. Elimination of East Avalonia in the British Isles simplifies the interpretation of the provenance of Neoproterozoic detrital zircons in Devonian sedimentary rocks in the region. Although Ganderia is the most proximal potential source of Neoproterozoic zircons, one cannot eliminate the possible detrital influence from other peri-Gondwanan sources.

Waldron *et al.* (2011) proposed that the Cambrian successions of the Meguma terrane of Nova Scotia and the Harlech Dome of Wales be considered a single palaeogeographical domain which they named ‘Megumia’. They show that detrital zircon age distributions from the Harlech Dome have greater similarity to those in the Meguma terrane than those in Avalonia. The major difference in detrital zircon age distributions between Megumia and Avalonia is the significant presence of 1.95

149 to 2.1 Ga zircons in Megumia and the lack thereof in Avalonia. These zircon ages
150 are believed to be associated with Eburnean orogenic magmatic activity (Waldron *et*
151 *al.* 2011). For the purpose of this study, a significant contribution of Neoproterozoic
152 zircon grains to the sediments under investigation is used predominantly as a tool to
153 distinguish between peri-Gondwana-derived and Laurentia-derived zircons, due to
154 the fact that Laurentia is not known to have any major source of Neoproterozoic
155 zircons (Fig. 2). Furthermore, given the Devonian age and tectonic setting of the
156 sediments, it is likely that they represent heterogeneous source areas. However, the
157 reader is urged to bear in mind the complexities associated with sources of
158 Neoproterozoic zircons in the British Isles, as described above.

159 The Grampian Orogeny, which occurred in the Ordovician period, is considered
160 to be the first stage of the Caledonian orogenic cycle (Chew & Stillman 2009) and
161 records the collision of an oceanic arc (recognised in Ireland as the Lough Nafooey
162 Arc) with the Laurentian margin (Chew & Strachan 2014). Ordovician sediments
163 from the South Mayo Trough record dominant input of zircons in the age range ca.
164 490 to ca. 467 Ma (McConnell *et al.* 2009). However, this range bears two
165 populations, one around 487 Ma and one between ca. 474 and ca. 467 Ma,
166 interpreted by McConnell *et al.* (2009) to be sourced from the Lough Nafooey
167 Arc/Clew Bay Complex and the Connemara orthogneiss suite respectively. The
168 Tyrone Igneous Complex is also of similar age to the Connemara suite (Cooper *et*
169 *al.* 2011). The Grangegeeth volcanic terrane in eastern Ireland has a maximum age
170 of ca. 465 Ma and inherited zircon within it indicate that it is of Laurentian origin,
171 perhaps being related to the Tyrone Igneous Complex (McConnell *et al.* 2010). Its
172 anomalous position south of the Southern Uplands – Longford Down terrane is
173 attributed to transpressive strike-slip deformation in Middle Silurian times

(McConnell *et al.* 2010). To the south of the Grangegeeth terrane lies the Bellewstown terrane (Fig. 1. Regional map of the British Isles showing the major terranes of Ireland. The rectangle indicates the study area shown in Figure 3). This terrane is considered part of the Ganderian margin and zircons dated from a sandstone within a volcanogenic breccia place the age of volcanism in the terrane at ca. 474 Ma (McConnell *et al.* 2015).

Plagiogranite boulders from Silurian conglomerates which lie unconformably upon the Lough Nafooey Group are considered to be sourced from the Lough Nafooey Arc (Chew *et al.* 2007). U-Pb zircon ages of around 490 Ma from these boulders, supported by Nd-isotope data, led Chew *et al.* (2007) to conclude that the arc had encountered Laurentian margin sediments by this time. This represents a source of Early Ordovician detrital zircons. The Southern Uplands – Longford Down terrane predominantly consists of Ordovician to Silurian metasedimentary rocks that were originally deposited in the Iapetus Ocean on the margin of Laurentia and were subsequently accreted to this margin (McConnell *et al.* 2016; Waldron *et al.* 2008). On the Irish side, McConnell *et al.* (2016) found that these sediments contained Proterozoic zircons indicative of a Laurentian origin. In addition, their samples contained an abundance of Early to Middle Ordovician zircons which they interpret as representing a volcanic arc source (e.g. Tryrone Igneous Complex, Lough Nafooey Arc). On the Scottish side, Waldron *et al.* (2008) and Waldron *et al.* (2014) found that the majority of detrital zircons are of Proterozoic age and sourced from Laurentia. Generally, samples from these studies produced fewer Early to Middle Ordovician zircons relative to those from the Irish side of the terrane.

Another potentially important Ordovician zircon source occurs in the Duncannon and Ribband Groups in the Leinster Massif. The minor calc-alkaline volcanic rocks

in the Ribband Group represent arc development during initial stages of subduction of the Iapetus Ocean crust beneath Ganderia (East Avalonia) in Tremadocian times (Woodcock 2012). The Duncannon Group volcanic suite consists of basalts and rhyolites, extruded in Caradoc times (Sandbian-Katian) which is considered indicative of a back-arc region (Woodcock 2012). These volcanic suites have not yet been isotopically dated and their ages are largely constrained by faunal evidence in associated sedimentary successions (e.g. Owen & Parkes 2000).

The final welding of Ganderia to the margin of Laurentia, which included an accreted ocean arc following the Grampian Orogeny, was achieved during the Caledonian Orogeny, by about 430-425 Ma (Mac Niocaill 2000; Waldron *et al.* 2014). However, abundant evidence of Late Caledonian (including Acadian, *sensu* Chew & Stillman 2009) magmatism exists in the form of widely distributed intrusions in Ireland and Britain. These range in age from ca. 430 Ma to 380 Ma (Fig. 2). An example of such an intrusion, which has been proposed as a proximal source of detritus to the Munster Basin (e.g. Penney 1980) and which represents the age of the majority of these intrusions (Fig. 2), is the Leinster granite batholith. O'Connor *et al.* (1989) obtained a U-Pb monazite age of 405 ± 2 Ma for the batholith. However, recent work by Fritschle *et al.* (2017) shows that it was emplaced over an extended period from ca. 417 Ma to 405 Ma. Vermeulen *et al*. (2000) have shown, by seismic analysis of southern Ireland, that the UORS south of the Killarney-Mallow Fault Zone is likely to be concealing a granitic body. Such an interpretation has been suggested by other studies (e.g. Ford *et al*. 1991; Meere 1995; Masson *et al*. 1998; Vermeulen *et al*. 2000; Todd 2000). Todd (2000) suggested that, based on clast analysis of the Trabeg Conglomerate Formation, a granite body similar to the Leinster Batholith was exposed in the southern hinterlands of the Dingle Basin

during its development and that the Leinster terrane therefore extends westward following Caledonian trends. Other examples of intrusives of Late Caledonian age in Ireland include the Donegal (418-388 Ma), Galway (412-380 Ma) and Newry Granites (403-387 Ma) and the Carnsore and Saltees Granites (436-428 Ma). The age ranges presented above are, in most cases, the result of dating by multiple geochronogical techniques which yield different ages. The studies from which these ages are obtained are reviewed in Chew & Stillman (2009). The Newry Igneous Complex has recently been redated by Cooper *et al.* (2016) who showed that, like the Leinster Batholith, the complex was emplaced over a similarly extended period from 414 Ma to 407 Ma.

Following the closure of the Iapetus Ocean, a Silurian to Early Devonian period of sinistral transtension accommodated deposition of the LORS in the Dingle Basin and elsewhere in the southern British Isles (Todd 1989; Soper & Woodcock 2003). This Emsian transtension across the ISZ in Ireland and Britain possibly initiated emplacement of granites of similar age on either side of the ISZ (Brown *et al.* 2008; Cooper *et al.* 2016). The rocks of the Dingle Basin record an Emsian deformational event which is considered to be part of the Acadian orogenic episode (Todd 1989; Todd 2000; Meere & Mulchrone 2006; Todd 2015). Deformation occurred in a transpressive regime but its kinematic character in the Dingle Basin is debated (e.g. Todd 1989; Meere & Mulchrone 2006; Todd 2015) because the structural fabrics within the LORS are complicated by Carboniferous Variscan overprinting.

Local Geology and Sample Location

The majority of the rocks that crop out on the Dingle Peninsula, southwest Ireland, form part of the Lower to Middle Devonian Lower Old Red Sandstone of the Dingle Basin. Basement to this basin is also exposed on the peninsula in the form of

the Ordovician Annascaul Formation and the Silurian Dunquin Group (Todd *et al.* 2000) (Fig. 3). These have been correlated with Ordovician and Silurian rocks in the Leinster Massif and therefore have a peri-Gondwanan affinity (Todd *et al.* 2000). The axis of the Dingle Basin approximates the regional northeast-southwest Caledonian trend in Ireland. The basin's sedimentary fill is complicated by a series of unconformities which separate five lithostratigraphic groups (Boyd & Sloan 2000). Todd *et al.* (1988) consider the main mechanism of subsidence to be in the form of a sinistral pull-apart structure. Meere and Mulchrone (2006) recognise two broad phases of extension: an Early Devonian phase that accommodated Dingle Basin sediments, and a Late Devonian to Carboniferous phase that accommodated the sediments of the Munster Basin. An intervening Middle Devonian transpression, likely recording the Acadian orogenic event (Dewey & Strachan 2003; Soper & Woodcock 2003), led Ennis *et al.* (2015) to consider the possibility of sedimentary recycling from the Dingle Basin into the Munster Basin. Such recycling of the Dingle Basin LORS into the UORS of the Munster Basin has also been suggested by Todd (2015).

Within the Dingle Basin, the Dingle Group, which is the most voluminous, represents a fluvial/alluvial environment in which axial (flowing from southwest to northeast along the basin axis) braid-plains and flood sheets, represented by the Eask, Coumeenoole, Slea Head and Ballymore Formations, were deposited by generally perennial axial river systems that flowed from the southwest (Todd 2000). These were abutted by alluvial fans, mainly represented by the Glashabeg Conglomerate Formation in the north and the Trabeg Conglomerate Formation in the south, that drained transversely into the basin (Todd 2000). Spores from the Eask and Slea Head Formation have yielded ages of Early Pragian and Early Emsian

274 respectively (Higgs 1999). Two samples from the LORS of the Dingle Basin were
275 taken from the Coumeenoole Formation and the overlying Slea Head Formation from
276 the Dingle Group (Fig. 3) in close proximity to the locations of equivalent samples
277 from Ennis *et al.* (2015).

278 Todd (2000) carried out an extensive study of clasts from the correlative
279 Glashabeg Conglomerate, Slea Head and Trabeg Conglomerate formations in the
280 Lower to Middle Devonian Dingle Basin. Based on the large number of mafic to
281 intermediate volcanic clasts and the presence of limestone clasts in the Glashabeg
282 Conglomerate Formation, he concluded that the source area was dominated by
283 Ordovician volcanic rocks lying in close proximity to the north of the basin and likely
284 intersecting the Iapetus Suture Zone. Rivers feeding the Slea Head Formation (and
285 the conformably underlying Coumeenoole Formation, which has a similar palaeoflow
286 direction) flowed axially through the basin toward the northeast, draining an area to
287 the southwest (Todd 2000). Pebble clasts indicate sediment derivation from Silurian
288 volcanic rocks and from an extension of the Leinster Massif basement (Todd 2000).
289 The Trabeg Conglomerate Formation formed the southern flank of the Slea Head
290 system and drained an area to the south and southwest of the basin which was
291 made up of rocks similar to those observed in the Leinster Massif (Todd 2000).

292 Unconformably overlying the Dingle Group is the Smerwick Group, which consists
293 of sedimentary rocks of aeolian and fluvial origin (Todd *et al.* 1988). The two groups
294 are truncated by an unconformity that developed due to erosion during Late Emsian
295 Acadian basin inversion (Meere & Mulchrone 2006). This was followed by deposition
296 of the Pointagare Group, exposed only on the northern coast of the peninsula, and
297 the Caherbla Group to the south which contains the Inch Conglomerate Formation.
298 These formations are thought to have been deposited in Middle Devonian (Eifelian)

299 times (Todd 2015, and references therein). Finally, the overstepping sandstone and
300 conglomerate successions of the Slieve Mish Group were deposited during the Late
301 Devonian and are considered by Williams (2000) to be a correlative of the
302 Ballinskelligs Formation and equivalents in the Munster Basin. Sample AK17 was
303 obtained from a pebbly sandstone of the fluvial to lacustrine (Todd 2015) Cappagh
304 Sandstone Formation of the Slieve Mish Group.

305 Analytical procedures and sampling

306 The apatite U-Pb data were originally generated as part of a bedrock thermal
307 history study utilising the apatite fission track (AFT) low-temperature
308 thermochronometer (Cogné *et al*. 2014), because the laser ablation-inductively
309 coupled plasma-mass spectrometry (LA-ICP-MS) approach to AFT analysis permits
310 U-Pb and AFT ages to be determined on the same grains during a single analytical
311 session (Chew & Donelick 2012).The sampling and separation process is therefore
312 different to the zircon separation process.

313 Zircon U-Pb

314 Data to produce the source 'signals' shown in Figure 2 were derived from a
315 number of sources. Data from samples interpreted to be of Laurentian origin were
316 sourced from Cawood *et al*. (2003), Friend *et al*. (2003), Cawood *et al*. (2007),
317 Kirkland *et al*. (2008), Waldron *et al*. (2008), McAteer *et al*. (2010), Cawood *et al*.
318 (2012), Strachan *et al*. (2013), Waldron *et al*. (2014) and Johnson *et al*. (2016). Data
319 from samples interpreted from Cadomia-Armorica are from Fernandez-Suarez *et al*.
320 (2002), Samson *et al*. (2005), Linnemann *et al*. (2008) and Strachan *et al*. (2008).
321 Data from sample of Ganderian association are from Fyffe *et al*. (2009), Waldron *et*
322 *al*. (2014) and Willner *et al*. (2014). Data from samples interpreted to be of
323 Megumian affinity are from Krogh & Keppie (1990), Waldron *et al*. (2009, 2011) and

Pothier *et al*. (2015). `East' Avalonia data were taken from Collins and Buchan (2004), Murphy *et al*. (2004b), Strachan *et al*. (2007), Linnemann *et al*. (2012) and Willner *et al*. (2013). Data from samples interpreted to be sourced from `West' Avalonia are from Keppie *et al*. (1998), Thompson & Bowring (2000), Barr *et al*. (2003), Murphy *et al*. (2004a, 2004b), Pollock (2007), Satkoski *et al*. (2010), Thompson *et al*. (2012), Dorais *et al*. (2012); Barr *et al*. (2012), Force & Barr (2012), Pollock *et al*. (2012), Willner *et al*. (2013) and Henderson *et al*. (2016). Caledonian granite ages are from Ireland only and include various isotopic geochronological techniques. The granite data were sourced from a compilation by Chew & Stillman (2009).

Sample separation was undertaken at Vrije Universiteit Amsterdam. Detrital zircons were liberated from samples using a jaw crusher and disc mill. Density separation was achieved using diiodomethane in a centrifuge as per Ijlst (1973) and magnetic separation using a Frantz magnetic separator. Zircons of all morphologies and colours, between 60 and 250 µm, were hand-picked under binocular microscope. Typically, between 120 and 180 zircons per sample were mounted in epoxy disks and ground and polished to expose the approximate centre of the grains. Cathodolumenescence (CL) imaging was undertaken at the University of St. Andrews and at Trinity College Dublin in order to identify optimal positions for laser ablation.

Uranium, thorium and lead isotopes were measured by laser ablation-sector field-inductively coupled plasma-mass spectrometry (LA-SF-ICP-MS) at the Museum für Mineralogie und Geologie (Geoplasma Lab, Senckenberg Naturhistorische Sammlungen Dresden) using a Thermo-Scientific Element 2 XR sector field ICP-MS coupled to a New Wave UP-193 Excimer Laser System. Each

analysis consisted of approximately 15 s background acquisition followed by 30 s data acquisition. A common-Pb correction based on the interference- and background-corrected ^{204}Pb signal and a model Pb composition (Stacey & Kramers 1975) was carried out if necessary. The necessity of the correction is judged on whether the corrected ^{207}Pb/^{206}Pb lies outside of the internal errors of the measured ratios. Raw data were corrected for background signal, common Pb, laser induced elemental fractionation, instrumental mass discrimination, and time-dependant elemental fractionation of Pb/Th and Pb/U using a Microsoft Excel spreadsheet program developed by Axel Gerdes (Institute of Geosciences, Johann Wolfgang Geothe-University Frankfurt, Frankfurt am Main, Germany). Concordia diagrams and concordia ages were produced using Isoplot/Ex 3.7 of Ludwig (2012). Frequency and kernel density estimation (KDE) curves were plotted using DensityPlotter (Vermeesch 2012). Frequency plots were assigned a binwidth of 25 Ma. KDEs were plotted using a bandwidth of 20 Ma and a Gaussian kernel. This bandwidth was chosen by trial and error (by comparison to histograms and probability density plots) because in many cases the 'optimal bandwidth' calculated in DensityPlotter caused severe oversmoothing. More importantly, the same bandwidth was applied to all samples and to detrital zircons from potential source areas so that comparisons were like-for-like. Multi-dimensional scaling (MDS) analysis was performed using the R package `provenance` by Vermeesch *et al*. (2016).

White mica Ar-Ar

Detrital white mica Ar-Ar ages for the Coumeenoole Formation in the Dingle Group from Ennis *et al.* (2015) are recalculated in this study using the 28.201 ± 0.046 Ma age of Kuiper *et al.* (2008) for the Fish Canyon sanidine standard,

generally increasing the age of individual grains by ca. 1 % . This was done to facilitate comparison of future detrital white mica analyses which will be conducted using this age for the standard. Details of analytical procedures can be found in Ennis *et al.* (2015).

Apatite U-Pb

At each outcrop, ~10 kg of material was collected across several adjacent beds to reduce any bias arising during deposition from localized heavy mineral concentrating processes. Subsequent sample preparation and analysis were conducted at Trinity College Dublin. The sub–300 μm nonmagnetic heavy mineral fraction was obtained by standard jaw crushing, sieving, magnetic, and heavy liquid separation techniques. Grains were mounted in epoxy resin, ground to expose internal surfaces, and polished. To avoid sample bias, no attempt was made to exclude anhedral or inclusion-bearing grains; the LA-ICP-MS technique permits identification and exclusion of U-rich inclusions (e.g., zircon) from the time-resolved (i.e., downhole) ablation signal of the appropriate isotopes.

Analyses were conducted using a Photon Machines Analyte Excite 193 nm ArF Excimer laser ablation system coupled to a Thermo Scientific iCAP Qc ICPMS, employing laser spots of 30 μm, a fluence of 4.5 J cm^{-2}, a repetition rate of 5 Hz, and an ablation time for each spot of 45 s followed by a 25 s background measurement. Repeated measurements of the primary Madagascar apatite mineral standard (Thomson *et al.* 2012) were used to correct for downhole U-Pb fractionation, mass bias, and intrasession instrument drift using the “VizualAge_UcomPbine” data reduction scheme for IOLITE (Chew *et al.* 2014; Paton *et al.* 2011), while the secondary McClure Mountain and Durango apatite standards were analysed as unknowns (Schoene & Bowring 2006; McDowell *et al.* 2005). Unlike phases that

399 exclude common (initial or nonradiogenic) Pb during crystallization, such as zircon,
400 the often high common-Pb content in apatite typically renders apatite grains
401 discordant in the U-Pb system. Common-Pb in the Madagascar apatite primary
402 standard was corrected for using a ^{207}Pb-based correction method using a known
403 initial ^{207}Pb/^{206}Pb ratio (Chew *et al.* 2014). Variable common-Pb content in the
404 detrital apatite unknowns was corrected using an initial common-Pb composition
405 derived from a terrestrial Pb evolution model (Stacey & Kramers 1975) applied to an
406 initial estimate for the age of the apatite, and then by adopting an iterative approach
407 based on a ^{207}Pb correction (Chew *et al.* 2011). The ^{207}Pb-based correction assumes
408 U-Pb* (radiogenic Pb) concordance – a reasonable assumption in the case of
409 standards and magmatic grains, but one which may not be the case for detrital
410 grains that have experienced partial Pb loss. As a result, independent geological
411 evidence is required to discriminate between partially and wholly reset detrital U-Pb
412 ages, similar to partially reset AFT ages (Mark *et al.* 2016).
413 Due to the ^{207}Pb-based correction, no apatite U-Pb age data can be excluded
414 based on discordance criteria. However, the relatively low U content of apatite
415 (sometimes <1 ppm) and consequent near-zero radiogenic Pb content of some
416 grains can result in undesirably large analytical uncertainties. We therefore excluded
417 grains with 2σ errors >25%, similar to the approach of Zattin *et al.* (2012).
418 As post-deposition temperatures exceeded the thermal sensitivity of the AFT
419 technique (ca. 120-60 °C; e.g., Gallagher *et al.* 1998), the resultant AFT ages
420 typically defined a single population for each sample. Because only a single AFT age
421 population was defined for each sample, it was only necessary to analyse ca. 20-30
422 grains for each sample.

Detrital zircon U-Th-Pb results

Two hundred and seventy three zircons were analysed from the Lower Devonian Dingle Group on the Dingle Peninsula. Another 119 detrital zircon grains were analysed from the Upper Devonian Slieve Mish Group on the Dingle Peninsula. The number of concordant ages obtained from each sample ensures an extremely low probability of missing an age component that contributes 10 % of the detrital zircon age population of the analysed sediment (Vermeesch *et al*. 2016). For all three samples, where age populations contribute over six percent, interpretations can be made at the 95 % confidence level.

Age uncertainties reported in the text are at the 2σ level unless otherwise stated. Core-rim analyses were undertaken where these features could be identified in CL images and where grains were large enough. Only rim ages are used for provenance interpretations in order to determine the most recent source of sediment. The results are presented as Wetherill concordia (Wetherill 1956), frequency and KDE plots (Figs 4 and Fig. 5). The $^{207}Pb/^{206}Pb$ age is reported where the $^{206}Pb/^{238}U$ age is greater than 1.0 Ga because the $^{207}Pb/^{206}Pb$ age is more precise in older zircon grains (e.g. Gehrels *et al*. 2008).

Coumeenoole Formation - Sample AK19

Sample AK19, collected at Connor Pass, consists of grey, moderately sorted, medium- to coarse-grained sandstone and is the stratigraphically oldest sample in the study area. One hundred and fourteen zircon grains were analysed yielding a total of 86 concordant analyses ranging from 394 ± 9 Ma to 2142 ± 16 Ma (Fig. 4a). Thirty seven grains (43 %) yielded Mesoproterozoic ages which form a major peak at around 1.20 Ga (Fig. 5a). Neoproterozoic zircons make up 26 % (n = 22) of the sample, the second largest peak in the sample occurring at around 550 Ma. The rest

448 of the sample is composed of 14 (16 %) Palaeoproterozoic zircons and 13 (15 %)
449 Palaeozoic zircons. Of the Palaeozoic grains, four are Cambrian, four are
450 Ordovician, two are Silurian and three are Devonian. Two of the Devonian zircons,
451 the youngest in the sample, have ages of 394 ± 9 Ma and 402 ± 10 Ma and have
452 length:width ratios of 3:1 and 5:1 respectively. Kostov (1973) suggests that high
453 length-width ratios may indicate rapid cooling rates. However, such ratios may
454 simply represent first order detrital zircons that have not been reworked (Poldervaart
455 1955).

456 **Slea Head Formation - Sample AK21**

457 This sample was taken from Cooleen Pier in Ventry Bay and consists of dark
458 grey, medium- to coarse-grained, pebbly sandstones. Ninety eight concordant ages
459 were obtained from measurement of 159 zircon grains. Ages range from 395 ± 8 Ma
460 to 2765 ± 29 Ma and the largest peak is developed at 1.18 Ga (Figs Fig**. 4**b and Fig**.**
461 **5**b). Mesoproterozoic zircons form the bulk of the sample at 40 % (n = 39). Twenty
462 six zircon grains (27 %) are Neoproterozoic in age, forming a major peak at ca. 630
463 Ma. Palaeoproterozoic zircons contribute 13 % (n = 13) of the sample. Eighteen
464 grains are Palaeozoic in age, including one Cambrian zircon, eight Ordovician
465 zircons, three Silurian zircons and six Devonian zircons. This sample, unlike AK19,
466 also contains Archaean grains (n = 2). A concordia age of 405 ± 4 Ma ($MSWD_{conc}$ =
467 0.18; p_{conc} = 0.67) was calculated using the six Devonian zircons. Two of these
468 zircons display length:width ratios of greater than 3:1.

469 **Cappagh Sandstone Formation - Sample AK17**

470 This sample, consisting of grey to pink, poorly sorted medium- to coarse-grained,
471 pebbly sandstone, was taken from an outcropping ridge at Aughils, north of the
472 R561. One hundred and nineteen zircons were analysed, producing 84 concordant

ages ranging from 403 ± 9 Ma to 3677 ± 21 Ma (Fig. 4c). The majority of these grains are Mesoproterozoic in age (43 grains representing 51 % of the sample) and the largest peak in the sample is Mesoproterozoic at ca. 1.05 Ga (Fig. 5c). Sixteen grains (19 %) are Palaeoproterozoic in age and represent the second largest population in the sample. Neoproterozoic zircons represent 12 % (n = 10) of the sample. Palaeozoic zircons represent 10 %, including one Cambrian zircon, three Ordovician zircons, three Silurian zircons and one Devonian zircon. Seven Archaean zircon grains are present in the sample. There is a paucity of zircon ages (n = 3) between ca. 550 Ma and ca. 980 Ma.

Apatite U-Pb and Mica $^{40}Ar/^{39}Ar$ Geochronology

Apatite U-Pb

These data were generated concurrently with apatite fission track analyses that were undertaken for bedrock thermal history studies by Cogné *et al.* (2014), but were not previously published. Due to the low number of detrital apatite grains analysed, the data must be interpreted with caution. In some cases, samples from the same formation are combined to yield the minimum statistical requirement of 60 analyses, as per Dodson *et al.* (1988). The limited variability in detrital apatite ages in these larger samples suggests that it is unlikely that multiple apatite sources of significantly different age and thermal history were being sampled by the sediments under investigation. The apatite U-Pb ages were not utilised in the aforementioned study. Therefore the data have been included in this study (**Fig. 6**) to provide a geochronological proxy which has an intermediate closure temperature (ca. 375-550 °C; Cochrane *et al.* 2014) between the mica $^{40}Ar/^{39}Ar$ and zircon U-Pb systems. The sampling transect was collected from Mount Brandon on the Dingle Peninsula (Fig. 3). This transect intersected two formations and includes four samples taken from

the Ballymore Sandstone Formation and one sample from the Farran Sandstone Formation.

Combined analyses from four samples (Mb-1, Mb-4, Mb-5 and Mb-7) from the Ballymore Sandstone Formation, the uppermost formation in the Dingle Group, yields a KDE spectrum composed of 70 ages forming a single KDE peak at ca. 420 Ma. Detrital apatite ages in this formation range from 356 ± 80 to 896 ± 30 Ma (Fig. 6a). However, 63 % of grains have ages between 380 and 440 Ma and 10 % have ages of less than 393 Ma (end-Emsian). Eighteen apatite grains were analysed from the Farran Sandstone Formation (sample Mb-9), the lowest formation in the Smerwick Group. These grains range in age from 347 ± 58 to 1356 ± 227 Ma. The KDE spectrum (Fig. 6b) shows the highest age concentration at ca. 420 Ma.

Mica $^{40}Ar/^{39}Ar$

Ennis *et al.* (2015) acquired detrital white mica ages for a sample from the Coumeenoole Formation, at a similar location (Connor Pass) to detrital zircon sample AK19. These ages have been recalculated here using the revised Fish Canyon standard age of Kuiper *et al.* (2008). Detrital white mica ages in the sample range from 308 ± 10.3 Ma to 440 ± 7 Ma, including two main age groups forming KDE peaks at ca. 414 Ma and ca. 382 Ma (**Fig. 7**).

Discussion

The overall age distribution of detrital zircons in samples from the Coumeenoole and Slea Head formations (similar Proterozoic distributions, a gap in ages between 730 and 940 Ma, mostly continuous distributions between 390 and 730 Ma and major KDE peaks at around 1.2 Ga) suggests a common source throughout their deposition (Kolmogorov-Smirnov test p-value of 0.871). Palaeozoic peaks in both samples, at around 430 Ma, likely correspond to a Caledonian (430-420 Ma) source.

A mean $^{207}Pb/^{206}Pb$ age of 432 ± 3 Ma as well as a whole-rock Rb-Sr isochron age of 428 ± 11 Ma was obtained by O'Connor *et al.* (1988) for the Carnsore granite. A Rb-Sr whole-rock isochron age of 436 ± 7 Ma for the Saltees granite by Max *et al.* (1979) led O'Connor *et al.* (1988) to conclude that the two intrusions are genetically related. These intrusions are the only ones of this age in Ireland and the 430 Ma age of detrital zircons from the LORS, as well as the northerly directed palaeoflow of the Trabeg Conglomerate Formation (Todd 2000), suggests that the intrusions extend westward beneath the Munster Basin along the strike of Caledonian lineaments. The idea of such a buried granite is not new; it was first proposed by Murphy (1960) and has since been expanded upon by a number of authors (e.g. Ford *et al.* 1991; Meere 1995; Masson *et al.* 1998; Vermeulen *et al.* 2000; Todd 2000). Vermeulen *et al.* (2000) interpreted seismic profiles across southern Ireland as showing a shallow, granitic body buried beneath the UORS just south of the Killarney-Mallow Fault Zone. Todd (2000) suggested that granite clasts within the Trabeg Conglomerate Formation might represent the unroofing of a presently buried granitic intrusion which forms part of an extension of the Leinster Massif.

Detrital apatite ages from the uppermost part of the Dingle Group, in the Ballymore Formation, yield a single peak at ca. 420 Ma (**Fig. 6**a) which is compatible with a Late Caledonian granitic source. Williams *et al.* (1999) obtained an age of 411 Ma for the Cooscrawn Tuff Bed in the Ballymore Formation which is older than 22 of the 70 detrital apatites analysed in this formation. A concordia age from six zircons in the underlying Slea Head Formation provides a minimum depositional age of 405 ± 4 Ma ($MSWD_{conc}$ = 0.18; p_{conc} = 0.67) and suggests that the Cooscrawn Tuff age is likely erroneous. Palynological evidence places the deposition of the Slea Head Formation in early to possibly middle Emsian times and is compatible with the

548 youngest group of detrital zircons in this formation. The high length to width ratio of
549 some of the youngest zircons in both the Coumeenoole and Slea Head formations
550 may suggest an igneous source that had cooled rapidly (Kostov 1973) – possibly a
551 syn-sedimentary volcanic source. Although U-Pb ages of apatite grains from the
552 Ballymore Formation are younger, they are nonetheless indistinguishable from the
553 zircon ages of the Coumeenoole and Slea Head formations at the 2-sigma level. A
554 dominant age peak of ca. 420 Ma (**Fig. 6**b) is also observed for 17 detrital apatite
555 grains in the Farran Sandstone Formation (Smerwick Group) which unconformably
556 overlies the Ballymore Formation.

557 A lower Palaeozoic source of detrital mica is reflected in the oldest age peak (ca.
558 414 Ma, **Fig. 7**) from a sample from the Coumeenoole Formation. However, the
559 dominant age peak is much younger, at ca. 382 Ma. This peak is much younger than
560 the Emsian depositional age of the host sedimentary rock and suggests that the
561 LORS experienced a thermal resetting event. A review of the potential causes of
562 resetting is provided in Ennis *et al.* (2015) and the extension associated with Lough
563 Guitane Volcanism in the Munster Basin is briefly mentioned. Given that Williams *et*
564 *al.* (2000) obtained a U-Pb zircon age of 378.5 Ma for the Horses Glen Volcanic
565 Centre and 384.5 Ma for the Killeen Volcanic Centre (which lie to the southeast of
566 the Dingle Basin), it is likely that the younger detrital mica ages represent resetting
567 by high heat flow associated with extension and volcanism which may have been
568 caused by emplacement of a granitic body at depth below the Munster Basin (Avison
569 1984). Alternatively, these younger ages could represent partial resetting due to low-
570 grade Variscan metamorphism at the end of the Carboniferous Period.

571 The low proportion of Palaeozoic zircons in the Dingle Group does not reflect the
572 high proportion of volcanic clasts reported by Todd (2000) and interpreted to have

been derived from rocks of Ordovician to Silurian age. This, however, may simply be a function of low zircon fertility in the volcanic source due to the dominant mafic to intermediate compositions (Todd 2000).

Determination of the ultimate source of late Neoproterozoic detrital zircons in these samples is relatively straightforward because of the lack of abundant known sources of this age on this part of the Laurentian craton (Fig. 2). The LORS samples also contain some 1.9 to 2.1 Ga zircons, considered to be indicative of the Gondwanan Eburnean Orogeny (Nance *et al.* 2008). Late Neoproterozoic zircons are ubiquitous in peri-Gondwanan terranes as a result of arc magmatism at that time (Nance *et al.* 2008) and therefore represent the most likely source. It should be noted, however, that Cawood & Nemchin (2001) report extensional magmatism associated with rifting in Laurentia between 520 and 555 Ma. But sedimentary rocks interpreted to be of Laurentian affinity do not record extensive Neoproterozoic zircon production (Fig.Fig. 2). Combined detrital zircon ages from various Cambrian formations in the Leinster Massif (Waldron *et al.* 2014) yield a high proportion in the range 500 to 770 Ma forming a peak at around 590 Ma (**Fig. 8**d). Waldron *et al.* (2014) proposed that these zircon ages represent a Ganderian source for the host sediments. If the source of granitic material for the Dingle Group was indeed Late Caledonian granites as suggested above, then, by extension, the Cambrian to Ordovician rocks into which the granites intrude are also a viable source of detritus. Support for derivation of late Neoproterozoic zircons from recycling of peri-Gondwanan sediments can be found in the study by Todd (2000) who found coticule and tourmalinite clasts akin to rocks found in the Ordovician Ribband Group. Todd (2000) also suggests that quartzite clasts in the Trabeg Conglomerate Formation could be related to quartzites of the Cambrian Bray Group. The similarity in the

598 ranges of detrital zircon ages younger than 800 Ma in both the Coumeenoole and
599 Slea Head formations suggests that this southerly source remained available
600 throughout their deposition.

601 Zircons with ages in the range of 1.1 to 1.25 Ga, the most abundant in both
602 samples, are likely ultimately sourced from Laurentia. Cawood and Nemchin (2001)
603 consider grains in this age range to be representative of magmatic and metamorphic
604 activity associated with the Grenville Orogeny. Zircons of similar age are present in
605 the Ganderia-derived Cambrian sediments in the Leinster Massif. However, the far
606 greater abundance of zircons of this age range relative to zircons of late
607 Neoproterozoic age in the Dingle Group samples requires an additional source from
608 which the older zircons can be derived.

609 Constructing a palaeodrainage pattern for the Coumeenoole and Slea Head
610 formations requires a source (or sources) of detritus that meet the following criteria:

611 a) Abundant zircons yielding ca. 1.2 Ga U-Pb ages (likely of ultimate Laurentian
612 affinity).
613 b) Presence of late Neoproterozoic and 1.9 to 2.1 Ga zircons (of peri-
614 Gondwanan affinity)
615 c) Paucity of zircons between 730 and 940 Ma
616 d) Low proportion of Palaeozoic zircons
617 e) Late Caledonian apatite U-Pb and white mica $^{40}Ar/^{39}Ar$ ages (ca. 420 Ma and
618 414 Ma, respectively).

619 Transverse drainage, as indicated by palaeocurrent directions (Todd 2000),
620 particularly from the south but possibly also from the north, was the likely means by
621 which Late Caledonian and peri-Gondwanan material was supplied to the Dingle
622 Basin (**Fig. 9**). Evidence of Late Caledonian detrital input to the basin is given by 420

623 Ma apatite and 414 Ma mica ages. Although the ultimate source of Mesoproterozoic
624 zircons in the Coumeenoole and Slea Head formations was probably the Grenville
625 Orogeny, defining the immediate source area is more problematic because a source
626 of abundant 1.2 Ga zircons has not been found in Ireland. Ordovician and Silurian
627 sediments from the Southern Uplands – Longford Down terrane show an abundance
628 of 1.1 Ga (**Fig. 8**a and **Fig. 8**c) and Ordovician zircons (**Fig. 8**b). This accretionary
629 wedge material is therefore a poor candidate unless the source character of these
630 sedimentary rocks changes westward, along strike. The apparent dissimilarity,
631 revealed by multi-dimensional scaling analysis, between the Dingle Group samples
632 and samples of Laurentian derivation in the Southern Uplands – Longford Down
633 terrane (**Fig. 8**g) reflects the lack of 1.2 Ga zircons in the available source data and
634 presence of late Neoproterozoic zircons of peri-Gondwanan affinity in the Dingle
635 Group samples.

636 Given the bulk northeastward palaeoflow in the Coumeenoole and Slea Head
637 Formations (Todd *et al.* 1988), it is likely that most of the detritus was deposited as a
638 result of this flow. A source to the southwest of the Dingle Basin must account for
639 the high proportion of Laurentian zircons in the analysed samples. Therefore,
640 although the main river course likely flowed from the west and southwest, we
641 suggest that Caledonian highlands in the north produced a large number of
642 tributaries flowing from north to south, into the main Coumeenoole/Slea Head
643 system (Fig. 9). Alternatively, it is also possible that the Iapetus Suture forms a
644 regional S-shape and that the strike of the suture swings from roughly east-west
645 onshore to southwest-northeast off the southwest coast of Ireland. This is highly
646 speculative however, as there is no other supporting evidence for such a hypothesis.

Unlike the LORS on the Dingle Peninsula, the UORS, represented by sample AK17 from the Cappagh White Sandstone Formation, contains very few late Neoproterozoic zircons but is instead dominated by Mesoproterozoic grains. Also unlike the LORS, the UORS contains a dominant 1.05 Ga peak as opposed to the 1.2 Ga peak of the LORS. The lack of late Neoproterozoic grains, dominant 1.05 Ga KDE peak and similarity to detrital zircon age spectra of Laurentian affinity (**Fig. 8**a, **Fig. 8**c and **Fig. 8**g) indicates a northerly-derived source area and no input from sources south of the Iapetus Suture. It would also indicate that recycling of the LORS into the UORS in southern Ireland, as previously suggested, is not likely to have occurred. It is more conceivable that this sediment was recycled from material similar to that found in the Longford Down terrane.

It is becoming widely accepted that a period of regional sinistral transtension occurred across the Iapetus Suture Zone in the Early Devonian Period (Todd 1989; Phillips *et al.* 1995; Dewey & Strachan 2003; Brown *et al.* 2008; Cooper *et al.* 2016). In the present study, the discrepancy between 1.2 Ga Laurentian zircons found in the Dingle Group and 1.1 Ga Laurentian zircons found in the Longford Down and Southern Uplands terranes as well as the abundance of 1.05 Ga zircons in the UORS of the Dingle Peninsula is supportive of large-scale sinistral transtenional displacement along the Iapetus Suture, removing the 1.2 Ga source of detrital zircon and introducing the 1.05-1.1 Ga source to the UORS in the Late Devonian Period. Of course, this discrepancy may not have tectonic significance and may simply be representative of topographic separation of sources or complete denudation of the 1.2 Ga source and subsequent exhumation of the 1.1 Ga source. High resolution detrital zircon sampling of younger formations within and overlying the Dingle Group would serve to elucidate the nature and timing of the change in source character.

672 Finally, a minimum depositional age of ca. 405 Ma for the Coumeenoole Formation
673 adds to growing evidence (Soper & Woodcock 2003, and references therein) of a
674 discrete Emsian orogenic event in the British Isles.

675 ## Conclusions

676 This paper presents the first multi-proxy, single-grain detrital geochronological
677 study of the LORS in the Lower Devonian Dingle Basin in southwestern Ireland
678 which suggests the following:

679 1) The Dingle Basin sedimentary fill is dominated by Laurentian detritus but
680 also includes a mixture of peri-Gondwanan and Late Caledonian (430-420
681 Ma) to Acadian (410-390 Ma) source areas.

682 2) A dominant drainage area lay to the west but the basin must have received
683 much detritus from transverse drainage to the north which intersected
684 Laurentian material.

685 3) Detrital zircons from the UORS Cappagh White Sandstone Formation
686 show very different characteristics to the LORS where sediment was being
687 supplied solely from detritus of Laurentian affinity.

688 4) The paucity of late Neoproterozoic zircons in the UORS compared to the
689 high proportion in the LORS suggests that the UORS was not derived from
690 recycling of LORS sediments as previously suggested.

691 5) Well documented regional sinistral transtension during the Early Devonian
692 Period was likely responsible for a switch in Laurentian source character
693 from the LORS (prevalence of 1.2 Ga detrital zircons) to the UORS
694 (prevalence of 1.1 Ga detrital zircons).

695

B.J.F, A.K, P.A.M and K.F.M acknowledge the financial support of the Irish Shelf Petroleum Studies Group (ISPSG) of the Irish Petroleum Infrastructure Programme (PIP) Group 4 (project code IS 12/05 UCC). The ISPSG comprises: Atlantic Petroleum (Ireland) Ltd, Cairn Energy Plc, Chrysaor E&P Ireland Ltd, Chevron North Sea Limited, ENI Ireland BV, Europa Oil & Gas, ExxonMobil E&P Ireland (Offshore) Ltd., Husky Energy, Kosmos Energy LLC, Maersk Oil North Sea UK Ltd, Petroleum Affairs Division of the Department of Communications, Energy and Natural Resources, Providence Resources Plc, Repsol Exploración SA, San Leon Energy Plc, Serica Energy Plc, Shell E&P Ireland Ltd, Sosina Exploration Ltd, Tullow Oil Plc and Woodside Energy (Ireland) Pty Ltd. D.C., C.M. and N.C. acknowledge support from Science Foundation Ireland grant 12/IP/1663 and 13/RC/2092 (iCRAG Research Centre, project HC4.2PD6a). iCRAG is funded under the SFI Research Centres Programme and is co-funded under the European Regional Development Fund. D. Pastor-Galan is thanked for his geochronology and sampling advice. C. Reid and R. van Elsas are thanked for their technical assistance for SEM and sample preparation respectively. J.W.F Waldron and J.F. Dewey are thanked for their constructive reviews which significantly improved the manuscript.

References

AVISON, M. 1984. The Lough Guitane Volcanic Complex, County Kerry: A Preliminary Survey. *Irish Journal of Earth Sciences*, **6**, 127–136.

BARR, S.M., DAVIS, D.W., KAMO, S. & WHITE, C.E. 2003. Significance of U–Pb detrital zircon ages in quartzite from peri-Gondwanan terranes, New Brunswick and Nova Scotia, Canada. *Precambrian Research*, **126**, 123–145, doi: 10.1016/S0301-9268(03)00192-X.

BARR, S.M., HAMILTON, M.A., SAMSON, S.D., SATKOSKI, A.M. & WHITE, C.E. 2012. Provenance variations in northern Appalachian Avalonia based on detrital zircon age patterns in Ediacaran and Cambrian sedimentary rocks, New Brunswick and Nova Scotia, Canada. *Canadian Journal of Earth Sciences*, **49**, 533–546, doi: 10.1139/e11-070.

BOYD, J.D. & SLOAN, R.J. 2000. Initiation and early development of the Dingle Basin, SW Ireland, in the context of the closure of the Iapetus Ocean. *Geological Society, London, Special Publications*, **180**, 123–145, doi: 10.1144/GSL.SP.2000.180.01.08.

BROWN, P.E., RYAN, P.D., SOPER, N.J. & WOODCOCK, N.H. 2008. The Newer Granite problem revisited: a transtensional origin for the Early Devonian Trans-Suture Suite. *Geological Magazine*, **145**, 235–256, doi: 10.1017/S0016756807004219.

CAWOOD, P.A. & NEMCHIN, A.A. 2001. Paleogeographic development of the east Laurentian margin: Constraints from U-Pb dating of detrital zircons in the Newfoundland Appalachians. *Geological Society of America Bulletin*, **113**, 1234–1246, doi: 10.1130/0016-7606(2001)113<1234:PDOTEL>2.0.CO;2.

CAWOOD, P.A., NEMCHIN, A.A., SMITH, M. & LOEWY, S. 2003. Source of the Dalradian Supergroup constrained by U-Pb dating of detrital zircon and implications for the East Laurentian margin. *Journal of the Geological Society*, **160**, 231–246, doi: 10.1144/0016-764902-039.

CAWOOD, P.A., NEMCHIN, A.A., STRACHAN, R., PRAVE, T. & KRABBENDAM, M. 2007. Sedimentary basin and detrital zircon record along East Laurentia and Baltica during assembly and breakup of Rodinia. *Journal Of The Geological Society*, **164**, 257–275, doi: 10.1144/0016-76492006-115.

CAWOOD, P.A., MERLE, R.E., STRACHAN, R.A. & TANNER, P.W.G. 2012. Provenance of the Highland Border Complex: constraints on Laurentian margin accretion in the Scottish Caledonides. *Journal of the Geological Society*, **169**, 575–586, doi: 10.1144/0016-76492011-076.

CHEW, D.M. & DONELICK, R.A. 2012. Combined apatite fission track and U-Pb dating by LA-ICP-MS and its application in apatite provenance analysis. *In*: Sylvester,

P. J. (ed.) *Quantitative Mineralogy and Microanalysis of Sediments and Sedimentary Rocks*. Mineralogical Association of Canada, 219–247.

CHEW, D.M. & STILLMAN, C.J. 2009. Late Caledonian orogeny and magmatism. *In*: *The Geology of Ireland*. 143–174.

CHEW, D.M. & STRACHAN, R.A. 2014. The Laurentian Caledonides of Scotland and Ireland. *Geological Society, London, Special Publications*, **390**, 45–91, doi: 10.1144/SP390.16.

CHEW, D.M., GRAHAM, J.R. & WHITEHOUSE, M.J. 2007. U-Pb zircon geochronology of plagiogranites from the Lough Nafooey (= Midland Valley) arc in western Ireland: constraints on the onset of the Grampian orogeny. *Journal of the Geological Society*, **164**, 747–750, doi: 10.1144/0016-76492007-025.

CHEW, D.M., SYLVESTER, P.J. & TUBRETT, M.N. 2011. U-Pb and Th-Pb dating of apatite by LA-ICPMS. *Chemical Geology*, **280**, 200–216, doi: 10.1016/j.chemgeo.2010.11.010.

CHEW, D.M., PETRUS, J.A. & KAMBER, B.S. 2014. U–Pb LA–ICPMS dating using accessory mineral standards with variable common Pb. *Chemical Geology*, **363**, 185–199, doi: 10.1016/j.chemgeo.2013.11.006.

COCHRANE, R., SPIKINGS, R.A., ET AL. 2014. High temperature (>350°C) thermochronology and mechanisms of Pb loss in apatite. *Geochimica et Cosmochimica Acta*, **127**, 39–56, doi: 10.1016/j.gca.2013.11.028.

COCKS, L.R.M., MCKERROW, W.S. & VAN STAAL, C.R. 1997. The margins of Avalonia. *Geological Magazine*, **134**, 627–636.

COGNÉ, N., CHEW, D. & STUART, F.M. 2014. The thermal history of the western Irish onshore. *Journal of the Geological Society*, **171**, 779–792, doi: 10.1144/jgs2014-026.

COLLINS, A.S. & BUCHAN, C. 2004. Provenance and age constraints of the South Stack Group, Anglesey, UK: U–Pb SIMS detrital zircon data. *Journal of the Geological Society*, **161**, 743–746, doi: 10.1144/0016-764904-036.

COOPER, M.R., CROWLEY, Q.G., ET AL. 2011. Age constraints and geochemistry of the Ordovician Tyrone Igneous Complex, Northern Ireland: implications for the Grampian orogeny. *Journal of the Geological Society*, **168**, 837–850, doi: 10.1144/0016-76492010-164.

COOPER, M.R., ANDERSON, P., CONDON, D.J., STEVENSON, C.T.E., ELLAM, R.M., MEIGHAN, I.G. & CROWLEY, Q.G. 2016. Shape and intrusion history of the Late Caledonian, Newry Igneous Complex, Northern Ireland. *In*: Young, M. E. (ed.) *Unearthed: Impacts of the Tellus Surveys of Ireland*. Dublin, Royal Irish Academy, 145–155., doi: DOI:10.3318/ 978-1-908996-88-6.ch11.

DALY, J.S. 1996. Pre-Caledonian history of the Annagh Gneiss Complex, North-western Ireland, and correlation with Laurentia-Baltica. *Irish Journal of Earth Sciences*, **15**, 5–18.

DEWEY, J.F. & STRACHAN, R.A. 2003. Changing Silurian-Devonian relative plate motion in the Caledonides: sinistral transpression to sinistral transtension. *Journal of the Geological Society*, **160**, 219–229, doi: 10.1144/0016-764902-085.

DODSON, M.H., COMPSTON, W., WILLIAMS, I.S. & WILSON, J.F. 1988. A search for ancient detrital zircons in Zimbabwean sediments. *Journal of the Geological Society*, **145**, 977–983, doi: 10.1144/gsjgs.145.6.0977.

DORAIS, M.J., WINTSCH, R.P., KUNK, M.J., ALEINIKOFF, J., BURTON, W., UNDERDOWN, C. & KERWIN, C.M. 2012. P-T-t conditions, Nd and Pb isotopic compositions and detrital zircon geochronology of the Massabesic Gneiss Complex, New

798 Hampshire: Isotopic and metamorphic evidence for the identification of Gander
799 basement, central New England. *American Journal of Science*, **312**, 1049–1097,
800 doi: 10.2475/10.2012.01.
801 ENNIS, M., MEERE, P.A., TIMMERMAN, M.J. & SUDO, M. 2015. Post-Acadian sediment
802 recycling in the Devonian Old Red Sandstone of Southern Ireland. *Gondwana*
803 *Research*, **28**, 1415–1433, doi: 10.1016/j.gr.2014.10.007.
804 FERNANDEZ-SUAREZ, J., GUTIERREZ ALONSO, G. & JEFFRIES, T.E. 2002. The
805 importance of along-margin terrane transport in northern Gondwana: Insights
806 from detrital zircon parentage in Neoproterozoic rocks from Iberia and Brittany.
807 *Earth and Planetary Science Letters*, **204**, 75–88, doi: 10.1016/S0012-
808 821X(02)00963-9.
809 FORCE, E.R. & BARR, S.M. 2012. Provenance of the Lower Carboniferous Horton
810 Group, Petit-de-Grat Island, Nova Scotia, as revealed by detrital zircon ages.
811 *Atlantic Geology*, **48**, 137–145, doi: 10.4138/atlgeol.2012.007.
812 FORD, M., BROWN, C. & READMAN, P. 1991. Analysis and tectonic interpretation of
813 gravity data over the Variscides of southwest Ireland. *Journal of the Geological*
814 *Society*, **148**, 137–148, doi: 10.1144/gsjgs.148.1.0137.
815 FORD, M., KLEMPERER, S.L. & RYAN, P.D. 1992. Deep structure of southern Ireland: a
816 new geological synthesis using BIRPS deep reflection profiling. *Journal of the*
817 *Geological Society*, **149**, 915–922.
818 FRIEND, C.R.L., STRACHAN, R.A., KINNY, P.D. & WATT, G.R. 2003. Provenance of the
819 Moine Supergroup of NW Scotland: evidence from geochronology of detrital and
820 inherited zircons from (meta) sedimentary rocks , granites and migmatites.
821 *Journal of the Geological Society, London*, **160**, 247–257.
822 FRITSCHLE, T., DALY, J.S., WHITEHOUSE, M.J., MCCONNELL, B. & BUHRE, S. 2017.

Multiple intrusive phases in the Leinster Batholith , Ireland : geochronology , isotope geochemistry and constraints on the deformation history.

FULLEA, J., MULLER, M.R., JONES, A.G. & AFONSO, J.C. 2014. The lithosphere-asthenosphere system beneath Ireland from integrated geophysical-petrological modeling II: 3D thermal and compositional structure. *Lithos*, **189**, 49–64, doi: 10.1016/j.lithos.2013.09.014.

FYFFE, L.R., BARR, S.M., ET AL. 2009. Detrital zircon ages from Neoproterozoic and Early Paleozoic conglomerate and sandstone units of New Brunswick and coastal Maine: implications for the tectonic evolution of Ganderia. *Atlantic Geology*, **45**, 110–144, doi: 10.4138/atlgeol.2009.006.

GALLAGHER, K., BROWN, R. & JOHNSON, C. 1998. Fission track analysis and its applications to geological problems. *Annual Review of Earth and Planetary Sciences*, **26**, 519–572, doi: 10.1146/annurev.earth.26.1.519.

GEHRELS, G.E., VALENCIA, V.A. & RUIZ, J. 2008. Enhanced precision, accuracy, efficiency, and spatial resolution of U-Pb ages by laser ablation-multicollector-inductively coupled plasma-mass spectrometry. *Geochemistry, Geophysics, Geosystems*, **9**, 1–13, doi: 10.1029/2007GC001805.

HENDERSON, B.J., COLLINS, W.J., MURPHY, J.B., GUTIERREZ-ALONSO, G. & HAND, M. 2016. Gondwanan basement terranes of the Variscan–Appalachian orogen: Baltican, Saharan and West African hafnium isotopic fingerprints in Avalonia, Iberia and the Armorican Terranes. *Tectonophysics*, **681**, 278–304, doi: 10.1016/j.tecto.2015.11.020.

HIBBARD, J.P., VAN STAAL, C.R. & RANKIN, D.W. 2007. A comparative analysis of pre-Silurian crustal building blocks of the northern and the southern Appalachian orogen. *American Journal of Science*, **307**, 23–45, doi: 10.2475/01.2007.02.

HIGGS, K.T. 1999. Early Devonian spore assemblages from the Dingle Group, County Kerry, Ireland. *Bollettino della Societa Paleontologica Italiana*, **38**, 187–196.

IJLST, L. 1973. A Laboratory Overflow-Centrifuge for Heavy Liquid Mineral Separation. *American Mineralogist*, **58**, 1088–1093.

JOHNSON, T.E., KIRKLAND, C.L., REDDY, S.M., EVANS, N.J. & MCDONALD, B.J. 2016. The source of Dalradian detritus in the Buchan Block, NE Scotland: application of new tools to detrital datasets. *Journal of the Geological Society*, **173**, 773–782, doi: 10.1144/jgs2016-019.

KENNEDY, M.J. 1979. The continuation of the Canadian Appalachians into the Caledonides of Britain and Ireland. *Geological Society, London, Special Publications*, **8**, 33–64, doi: 10.1144/GSL.SP.1979.008.01.03.

KEPPIE, J.D., DAVIS, D.W. & KROGH, T.E. 1998. U-Pb geochronological constraints on Precambrian stratified units in the Avalon Composite Terrane of Nova Scotia, Canada: tectonic implications. *Canadian Journal Of Earth Sciences*, **35**, 222–236, doi: 10.1139/e97-109.

KEPPIE, J.D., NANCE, R.D., MURPHY, J.B. & DOSTAL, J. 2003. Tethyan, Mediterranean, and Pacific analogues for the Neoproterozoic–Paleozoic birth and development of peri-Gondwanan terranes and their transfer to Laurentia and Laurussia. *Tectonophysics*, **365**, 195–219, doi: 10.1016/S0040-1951(03)00037-4.

KIRKLAND, C.L., STRACHAN, R.A. & PRAVE, A.R. 2008. Detrital zircon signature of the Moine Supergroup, Scotland: Contrasts and comparisons with other Neoproterozoic successions within the circum-North Atlantic region. *Precambrian Research*, **163**, 332–350, doi: 10.1016/j.precamres.2008.01.003.

873 KOSTOV, I. 1973. Zircon morphology as a Crystallogenetic indicator. *Kristall und*
874 *Technik*, **8**, 11–19, doi: 10.1002/crat.19730080103.
875 KROGH, T.E. & KEPPIE, J.D. 1990. Age of detrital zircon and titanite in the Meguma
876 Group, southern Nova Scotia, Canada: Clues to the origin of the Meguma
877 Terrane. *Tectonophysics*, **177**, 307–323, doi: 10.1016/0040-1951(90)90287-I.
878 KUIPER, K.F., DEINO, A., HILGEN, F.J., KRIJGSMAN, W., RENNE, P.R. & WIJBRANS, J.R.
879 2008. Synchronizing Rock Clocks of Earth History. *Science*, **320**, 500–504, doi:
880 10.1126/science.1154339.
881 LINNEMANN, U., GERDES, A., DROST, K. & BUSCHMANN, B. 2007. The continuum
882 between Cadomian orogenesis and opening of the Rheic Ocean: Constraints
883 from LA-ICP-MS U-Pb zircon dating and analysis of plate-tectonic setting (Saxo-
884 Thuringian zone, northeastern Bohemian Massif, Germany). *Geological Society*
885 *of America Special Papers*, **423**, 61–96, doi: 10.1130/2007.2423(03).
886 LINNEMANN, U., PEREIRA, F., JEFFRIES, T.E., DROST, K. & GERDES, A. 2008. The
887 Cadomian Orogeny and the opening of the Rheic Ocean: The diacrony of
888 geotectonic processes constrained by LA-ICP-MS U-Pb zircon dating (Ossa-
889 Morena and Saxo-Thuringian Zones, Iberian and Bohemian Massifs).
890 *Tectonophysics*, **461**, 21–43, doi: 10.1016/j.tecto.2008.05.002.
891 LINNEMANN, U., HERBOSCH, A., LIÉGEOIS, J.P., PIN, C., GÄRTNER, A. & HOFMANN, M.
892 2012. The Cambrian to Devonian odyssey of the Brabant Massif within
893 Avalonia: A review with new zircon ages, geochemistry, Sm-Nd isotopes,
894 stratigraphy and palaeogeography. *Earth-Science Reviews*, **112**, 126–154, doi:
895 10.1016/j.earscirev.2012.02.007.
896 LIVERMORE, R.A., SMITH, A.G. & BRIDEN, J.C. 1985. Palaeomagnetic Constraints on
897 the Distribution of Continents in the Late Silurian and Early Devonian.

Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, **309**, 29–56.

LUDWIG, K.R. 2012. User's Manual for Isoplot 3.75. *Berkeley Geochronology Centre Special Publication*, 1–72.

MAC NIOCAILL, C. 2000. A new Silurian palaeolatitude for eastern Avalonia and evidence for crustal rotations in the Avalonian margin of southwestern Ireland. *Geophysical Journal International*, **141**, 661–671, doi: 10.1046/j.1365-246X.2000.00101.x.

MARK, C., COGNÉ, N. & CHEW, D. 2016. Tracking exhumation and drainage divide migration of the Western Alps: A test of the apatite U-Pb thermochronometer as a detrital provenance tool. *Bulletin of the Geological Society of America*, **128**, 1439–1460, doi: 10.1130/B31351.1.

MASSON, F., JACOB, A.W.B., PRODEHL, C., READMAN, P.W., SHANNON, P.M., SCHULZE, A. & ENDERLE, U. 1998. A wide-angle seismic traverse through the Variscan of southwest Ireland. *Geophysical Journal International*, **134**, 689–705, doi: 10.1046/j.1365-246X.1998.00572.x.

MAX, M.D., PLOQUIN, A. & SONET, J. 1979. The age of the Saltees granite in the Rosslare Complex. *In*: *The Caledonides of the British Isles Reviewed*. Geological Society of London, 723–725., doi: 10.1144/GSL.SP.1979.008.01.88.

MAX, M.D., BARBER, A.J. & MARTINEZ, J. 1990. Terrane assemblage of the Leinster Massif , SE Ireland , during the Lower Palaeozoic. *Journal of the Geological Society, London*, **147**, 1035–1050.

MCATEER, C.A., STEPHEN DALY, J., FLOWERDEW, M.J., CONNELLY, J.N., HOUSH, T.B. & WHITEHOUSE, M.J. 2010. Detrital zircon, detrital titanite and igneous clast U-Pb geochronology and basement-cover relationships of the Colonsay Group, SW

Scotland: Laurentian provenance and correlation with the Neoproterozoic Dalradian Supergroup. *Precambrian Research*, **181**, 21–42, doi: 10.1016/j.precamres.2010.05.013.

McConnell, B. & Morris, J. 1997. Initiation of Iapetus subduction under Irish Avalonia. *Geological Magazine*, **134**, 213–218.

McConnell, B., Riggs, N. & Crowley, Q.G. 2009. Detrital zircon provenance and Ordovician terrane amalgamation, western Ireland. *Journal of the Geological Society, London*, **166**, 473–484, doi: 10.1144/0016-76492008-081.

McConnell, B., Crowley, Q.G. & Riggs, N. 2010. Laurentian origin of the Ordovician Grangegeeth volcanic arc terrane, Ireland. *Journal of the Geological Society*, **167**, 469–474, doi: 10.1144/0016-76492009-139.

McConnell, B., Parkes, M., Crowley, Q. & Rushton, A. 2015. No Exploits back-arc basin in the Iapetus suture zone of Ireland. *Journal of the Geological Society*, **172**, 740–747, doi: 10.1144/jgs2015-044.

McConnell, B., Rogers, R. & Crowley, Q. 2016. Sediment provenance and tectonics on the Laurentian margin: implications of detrital zircons ages from the Central Belt of the Southern Uplands–Down–Longford Terrane in Co. Monaghan, Ireland. *Scottish Journal of Geology*, **52**, 11–17, doi: 10.1144/sjg2015-013.

McDowell, F.W., McIntosh, W.C. & Farley, K.A. 2005. A precise 40Ar–39Ar reference age for the Durango apatite (U–Th)/He and fission-track dating standard. *Chemical Geology*, **214**, 249–263, doi: 10.1016/j.chemgeo.2004.10.002.

McIlroy, D. & Horák, J.M. 2006. Neoproterozoic: the late Precambrian terranes that formed Eastern Avalonia. *In*: Brenchley, P. J. & Rawson, P. F. (eds) *The*

948 *Geology of England and Wales*. Geological Society of London, 9–24., doi:
949 10.1144/GOEWP.2.
950 MEERE, P. 1995. The structural evolution of the western Irish Variscides: an example
951 of obstacle tectonics? *Tectonophysics*, **246**, 97–112.
952 MEERE, P.A. & MULCHRONE, K.F. 2006. Timing of deformation within Old Red
953 Sandstone lithologies from the Dingle Peninsula, SW Ireland. *Journal of the*
954 *Geological Society*, **163**, 461–469, doi: 10.1144/0016-764905-099.
955 MURPHY, J.B., FERNÁNDEZ-SUÁREZ, J. & JEFFRIES, T.E. 2004a. Lithogeochemical and
956 Sm-Nd and U-Pb isotope data from the Silurian–Lower Devonian Arisaig Group
957 clastic rocks, Avalon terrane, Nova Scotia: A record of terrane accretion in the
958 Appalachian-Caledonide orogen. *Geological Society of America Bulletin*, **116**,
959 1183, doi: 10.1130/B25423.1.
960 MURPHY, J.B., FERNÁNDEZ-SUÁREZ, J., JEFFRIES, T.E. & STRACHAN, R.A. 2004b. U-Pb
961 (LA-ICP-MS) dating of detrital zircons from Cambrian clastic rocks in Avalonia:
962 erosion of a Neoproterozoic arc along the northern Gondwanan margin. *Journal*
963 *of the Geological Society*, **161**, 243–254, doi: 10.1144/0016-764903-064.
964 MURPHY, T. 1960. Gravity map in Ireland - sheet 5 - south west. *Geophysical*
965 *Bulletin*, **18**.
966 NANCE, R., NEACE, E., BRAID, J., MURPHY, J., DUPUIS, N. & SHAIL, R. 2015. Does the
967 Meguma Terrane Extend into SW England? *Geoscience Canada*, **42**, 61–76,
968 doi: 10.12789/geocanj.2014.41.056.
969 NANCE, R.D., MURPHY, J.B., ET AL. 2008. Neoproterozoic-early Palaeozoic
970 tectonostratigraphy and palaeogeography of the peri-Gondwanan terranes:
971 Amazonian v. West African connections. *In*: Ennih, N. & Liégeois, J.-P. (eds)
972 *The Boundaries of the West African Craton*. The Geological Society of London,

345–383., doi: 10.1144/SP297.17.

NANCE, R.D., GUTIÉRREZ-ALONSO, G., ET AL. 2012. A brief history of the Rheic Ocean. *Geoscience Frontiers*, **3**, 125–135, doi: 10.1016/j.gsf.2011.11.008.

O'CONNOR, P.J., KENNAN, P.S. & AFTALION, M. 1988. New Rb-Sr and U-Pb Ages for the Carnsore Granite and their bearing on the antiquity of the Rosslare Complex, southeastern Ireland. *Geological Magazine*, **125**, 25–29, doi: 10.1017/S0016756800009341.

O'CONNOR, P.J., AFTALION, M. & KENNAN, P.S. 1989. Isotopic U–Pb ages of zircon and monazite from the Leinster Granite, southeast Ireland. *Geological Magazine*, **126**, 725, doi: 10.1017/S0016756800007044.

OWEN, A.W. & PARKES, M.A. 2000. Trilobite faunas of the Duncannon Group: Caradoc stratigraphy, environments and palaeobiogeography of the Leinster terrane, Ireland. *Palaeontology*, **43**, 219–269, doi: 10.1111/1475-4983.00125.

PATON, C., HELLSTROM, J., PAUL, B., WOODHEAD, J. & HERGT, J. 2011. Iolite: Freeware for the visualisation and processing of mass spectrometric data. *Journal of Analytical Atomic Spectrometry*, **26**, 2508, doi: 10.1039/c1ja10172b.

PENNEY, S. 1980. A new look at the Old Red Sandstone succession of the Comeragh Mountains, County Waterford. *Journal of Earth Sciences*, **3**, 155–178.

PHILLIPS, E.R., BARNES, R.P., BOLAND, M.P., FORTEY, N.J. & MCMILLAN, A.A. 1995. The Moniaive Shear Zone: a major zone of sinistral strike-slip deformation in the Southern Uplands of Scotland. *Scottish Journal of Geology*, **31**, 139–149, doi: 10.1144/sjg31020139.

PICKERING, K.T., BASSETT, M.G. & SIVETER, D.J. 1988. Late Ordovician-early Silurian destruction of the Iapetus Ocean : Newfoundland , British Isles and Scandinavia — a discussion. *Transactions of the Royal Society of Edinburgh: Earth*

998 *Sciences*, **79**, 361–382.
999 POINTON, M.A., CLIFF, R.A. & CHEW, D.M. 2012. The provenance of Western Irish
1000 Namurian Basin sedimentary strata inferred using detrital zircon U-Pb LA-ICP-
1001 MS geochronology. *Geological Journal*, **47**, 77–98, doi: 10.1002/gj.1335.
1002 POLDERVAART, A. 1955. Zircons in rocks; Part 1, Sedimentary rocks; Part 2, Igneous
1003 rocks. *American Journal of Science*, **253**, 433–461, doi: 10.2475/ajs.253.8.433.
1004 POLLOCK, J.C. 2007. *The Neoproterozoic-Early Paleozoic Tectonic Evolution of the*
1005 *Peri-Gondwanan Margin of the Appalachian Orogen - an Integrated*
1006 *Geochronological, Geochemical and Isotopic Study from North Carolina and*
1007 *Newfoundland*. North Carolina State University.
1008 POLLOCK, J.C., HIBBARD, J.P. & VAN STAAL, C.R. 2012. A paleogeographical review
1009 of the peri-Gondwanan realm of the Appalachian orogen. *Canadian Journal of*
1010 *Earth Sciences*, **49**, 259–288, doi: 10.1139/E11-049.
1011 POTHIER, H.D., WALDRON, J.W.F., SCHOFIELD, D.I. & DUFRANE, S.A. 2015. Peri-
1012 Gondwanan terrane interactions recorded in the Cambrian–Ordovician detrital
1013 zircon geochronology of North Wales. *Gondwana Research*, **28**, 987–1001, doi:
1014 10.1016/j.gr.2014.08.009.
1015 SAMSON, S.D., D'LEMOS, R.S., MILLER, B. V. & HAMILTON, M.A. 2005. Neoproterozoic
1016 palaeogeography of the Cadomia and Avalon terranes: constraints from detrital
1017 zircon U-Pb ages. *Journal of the Geological Society*, **162**, 65–71, doi:
1018 10.1144/0016-764904-003.
1019 SATKOSKI, A.M., BARR, S.M. & SAMSON, S.D. 2010. Provenance of Late
1020 Neoproterozoic and Cambrian Sediments in Avalonia: Constraints from Detrital
1021 Zircon Ages and Sm-Nd Isotopic Compositions in Southern New Brunswick,
1022 Canada. *Journal Of Geology*, **118**, 187–200, doi: 10.1086/649818.

1023 SCHOENE, B. & BOWRING, S.A. 2006. U–Pb systematics of the McClure Mountain
1024 syenite: thermochronological constraints on the age of the 40Ar/39Ar standard
1025 MMhb. *Contributions to Mineralogy and Petrology*, **151**, 615–630, doi:
1026 10.1007/s00410-006-0077-4.
1027 SCHOFIELD, D., POTTER, J., BARR, S.M., HORAK, J., MILLAR, I.L. & LONGSTAFFE, F.
1028 2016. Reappraising the Neoproterozoic 'East Avalonian' terranes of southern
1029 Great Britain. *Gondwana Research*, **35**, 257–271, doi: 10.1016/j.gr.2015.06.001.
1030 SOPER, N.J. & HUTTON, D.H.W. 1984. Late Caledonian sinistral displacements in
1031 Britain: Implication for a three-plate model. *Tectonics*, **3**, 781–794.
1032 SOPER, N.J. & WOODCOCK, N.H. 2003. The lost Lower Old Red Sandstone of
1033 England and Wales : a record of post-Iapetan flexure or Early Devonian
1034 transtension? *Geological Magazine*, **140**, 627–647, doi:
1035 10.1017/S0016756803008380.
1036 STACEY, J.S. & KRAMERS, J.D. 1975. Approximation of terrestrial lead isotope
1037 evolution by a two-stage model. *Earth and Planetary Science Letters*, **26**, 207–
1038 221, doi: 10.1016/0012-821X(75)90088-6.
1039 STRACHAN, R.. 2008. Early Earth history and the development of the Archaean crust.
1040 *In*: Woodcock, N. & Strachan, R. (eds) *Geological History of Britain and Ireland*.
1041 Oxford, Blackwell Science Ltd., 41–51.
1042 STRACHAN, R.A., COLLINS, A.S., BUCHAN, C., NANCE, R.D., MURPHY, J.B. & D'LEMOS,
1043 R.S. 2007. Terrane analysis along a Neoproterozoic active margin of
1044 Gondwana: insights from U-Pb zircon geochronology. *Journal of the Geological*
1045 *Society, London*, **164**, 57–60.
1046 STRACHAN, R.A., PRAVE, A.R., KIRKLAND, C.L. & STOREY, C.D. 2013. U–Pb detrital
1047 zircon geochronology of the Dalradian Supergroup, Shetland Islands, Scotland:

implications for regional correlations and Neoproterozoic–Palaeozoic basin development. *Journal of the Geological Society*, **170**, 905–916, doi: 10.1144/jgs2013-057.

THOMPSON, M.D. & BOWRING, S.A. 2000. Age of the Squantum 'Tillite,' Boston Basin, Massachusetts: U-Pb zircon constraints on terminal Neoproterozoic glaciation. *American Journal of Science*, **300**, 630–655, doi: 10.2475/ajs.300.8.630.

THOMPSON, M.D., BARR, S.M. & GRUNOW, A.M. 2012. Avalonian perspectives on Neoproterozoic paleogeography: Evidence from Sm-Nd isotope geochemistry and detrital zircon geochronology in SE New England, USA. *Bulletin of the Geological Society of America*, **124**, 517–531, doi: 10.1130/B30529.1.

THOMSON, S.N., GEHRELS, G.E., RUIZ, J. & BUCHWALDT, R. 2012. Routine low-damage apatite U-Pb dating using laser ablation-multicollector-ICPMS. *Geochemistry, Geophysics, Geosystems*, **13**, doi: 10.1029/2011GC003928.

TODD, S. 1989. Role of the Dingle Bay Lineament in the evolution of the Old Red Sandstone of southwest Ireland. *In*: Arthurton, R., Gutteridge, P. & Nolan, S. (eds) *The Role of Tectonics in Devonian and Carboniferous Sedimentation in the British Isles*. Bradford, The Yorkshire Geological Society, 35–54.

TODD, S.P. 2000. Taking the roof off a suture zone: basin setting and provenance of conglomerates in the ORS Dingle Basin of SW Ireland. *Geological Society, London, Special Publications*, **180**, 185–222, doi: 10.1144/GSL.SP.2000.180.01.10.

TODD, S.P. 2015. Structure of the Dingle Peninsula, SW Ireland: evidence for the nature and timing of Caledonian, Acadian and Variscan tectonics. *Geological Magazine*, **152**, 242–268, doi: 10.1017/S0016756814000260.

TODD, S.P., WILLIAMS, B.P.J. & HANCOCK, P.L. 1988. Lithostratigraphy and structure

of the Old Red Sandstone of the Northern Dingle Peninsula, Co. Kerry, southwest Ireland. *Geological Journal*, **23**, 107–120.

TODD, S.P., MURPHY, F.C. & KENNAN, P.S. 1991. On the trace of the Iapetus suture in Ireland and Britain. *Journal of the Geological Society, London*, **148**, 869–880.

TODD, S.P., CONNERY, C., HIGGS, K.T. & MURPHY, F.C. 2000. An Early Ordovician age for the Annascaul Formation of the SE Dingle Peninsula, SW Ireland. *Journal of the Geological Society*, **157**, 823–833, doi: 10.1144/jgs.157.4.823.

TYRRELL, S., HAUGHTON, P.D.W. & DALY, J.S. 2007. Drainage reorganization during breakup of Pangea revealed by in-situ Pb isotopic analysis of detrital K-feldspar. *Geology*, **35**, 971–974, doi: 10.1130/G4123A.1.

VAN DER VOO, R. 1983. Paleomagnetic constraints on the assembly of the Old Red Continent. *Tectonophysics*, **91**, 271–283.

VAN DER VOO, R. 1988. Paleozoic paleogeography of North America, Gondwana, and intervening displaced terranes: Comparisons of paleomagnetism with paleoclimatology and biogeographical patterns. *Geological Society of America Bulletin*, **100**, 311–324.

VAN STAAL, C.R., SULLIVAN, R.W. & WHALEN, J.B. 1996. Provenance and tectonic history of the Gander Zone in the Caledonian/Appalachian orogen: Implications for the origin and assembly of Avalon. *Geological Society of American Special Paper*, **304**, 347–367.

VAN STAAL, C.R., DEWEY, J.F., NIOCAILL, C.M. & MCKERROW, W.S. 1998. The Cambrian-Silurian tectonic evolution of the northern Appalachians and British Caledonides: history of a complex, west and southwest Pacific-type segment of Iapetus. *Geological Society, London, Special Publications*, **143**, 197–242, doi: 10.1144/GSL.SP.1998.143.01.17.

1098 VERMEESCH, P. 2012. On the visualisation of detrital age distributions. *Chemical*
1099 *Geology*, **312**–**313**, 190–194, doi: 10.1016/j.chemgeo.2012.04.021.
1100 VERMEESCH, P. 2013. Multi-sample comparison detrital age distributions. *Chemical*
1101 *Geology*, **341**, 140–146, doi: 10.1016/j.chemgeo.2013.01.010.
1102 VERMEESCH, P., RESENTINI, A. & GARZANTI, E. 2016. An R package for statistical
1103 provenance analysis. *Sedimentary Geology*, **336**, 14–25, doi:
1104 10.1016/j.sedgeo.2016.01.009.
1105 VERMEULEN, N.J., SHANNON, P.M., MASSON, F. & LANDES, M. 2000. Wide-angle
1106 seismic control on the development of the Munster Basin, SW Ireland.
1107 *Geological Society, London, Special Publications*, **180**, 223–237, doi:
1108 10.1144/GSL.SP.2000.180.01.11.
1109 WALDRON, J.W.F., FLOYD, J.D., SIMONETTI, A. & HEAMAN, L.M. 2008. Ancient
1110 Laurentian detrital zircon in the closing Iapetus ocean, Southern Uplands
1111 terrane, Scotland. *Geology*, **36**, 527–530, doi: 10.1130/G24763A.1.
1112 WALDRON, J.W.F., WHITE, C.E., BARR, S.M., SIMONETTI, A. & HEAMAN, L.M. 2009.
1113 Provenance of the Meguma terrane, Nova Scotia: rifted margin of early
1114 Paleozoic Gondwana. *Canadian Journal Of Earth Sciences*, **46**, 1–8, doi:
1115 10.1139/E09-004.
1116 WALDRON, J.W.F., SCHOFIELD, D.I., WHITE, C.E. & BARR, S.M. 2011. Cambrian
1117 successions of the Meguma Terrane, Nova Scotia, and Harlech Dome, North
1118 Wales : dispersed fragments of a peri-Gondwanan basin? *Journal of the*
1119 *Geological Society*, **168**, 83–98, doi: 10.1144/0016-76492010-068.Cambrian.
1120 WALDRON, J.W.F., SCHOFIELD, D.I., ET AL. 2014. Ganderia-Laurentia collision in the
1121 Caledonides of Great Britain and Ireland. *Journal of the Geological Society*, **171**,
1122 555–569, doi: 10.1144/jgs2013-131.

WETHERILL, G. 1956. Discordant Uraniam-Lead Ages, I. *Transactions, American Geophysical Union*, **37**, 320–326.

WHITE, C.E., PALACIOS, T., JENSEN, S. & BARR, S.M. 2012. Cambrian-Ordovician acritarchs in the Meguma terrane, Nova Scotia, Canada: Resolution of early Paleozoic stratigraphy and implications for paleogeography. *Bulletin of the Geological Society of America*, **124**, 1773–1792, doi: 10.1130/B30638.1.

WILLIAMS, E.A. 2000. Flexural cantilever models of extensional subsidence in the Munster Basin (SW Ireland) and Old Red Sandstone fluvial dispersal systems. *Geological Society, London, Special Publications*, **180**, 239–268, doi: 10.1144/GSL.SP.2000.180.01.12.

WILLIAMS, E.A., SERGEEV, S.A., STOSSEL, I., FORD, M. & HIGGS, K.T. 1999. U-Pb Zircon Geochronology of Silicic Tuffs from the Old Red Sandstone of the Dingle and Munster Basins (SW Ireland): Refining the Devonian Time Scale. *EUG Journal of Conference Abstracts*, **4**.

WILLIAMS, E.A., SERGEEV, S.A., STOSSEL, I., FORD, M. & HIGGS, K.T. 2000. U-Pb zircon geochronology of silicic tuffs and chronostratigraphy of the earliest Old Red Sandstone in the Munster Basin, SW Ireland. *Geological Society, London, Special Publications*, **180**, 269–302, doi: 10.1144/GSL.SP.2000.180.01.13.

WILLNER, A.P., BARR, S.M., GERDES, A., MASSONNE, H.-J. & WHITE, C.E. 2013. Origin and evolution of Avalonia: evidence from U–Pb and Lu–Hf isotopes in zircon from the Mira terrane, Canada, and the Stavelot–Venn Massif, Belgium. *Journal of the Geological Society* , **170**, 769–784, doi: 10.1144/jgs2012-152.

WILLNER, A.P., GERDES, A., MASSONNE, H.-J., VAN STAAL, C.R. & ZAGOREVSKI, A. 2014. Crustal evolution of the Northeast Laurentian Margin and the Peri-Gondwanan Microcontinent Ganderia Prior to and During Closure of the Iapetus

Ocean: Detrital Zircon U-Pb and Hf Isotope Evidence from Newfoundland. *Geoscience Canada*, **41**, 345–364.

WOODCOCK, N.H. 2012. Ordovician Volcanism and Sedimentation on Eastern Avalonia. *In*: Woodcock, N. H. & Strachan, R. A. (eds) *Geological History of Britain and Ireland*. Blackwell Publishing Ltd., 162–176.

ZATTIN, M., ANDREUCCI, B., THOMSON, S.N., REINERS, P.W. & TALARICO, F.M. 2012. New constraints on the provenance of the ANDRILL AND-2A succession (western Ross Sea, Antarctica) from apatite triple dating. *Geochemistry, Geophysics, Geosystems*, **13**, doi: 10.1029/2012GC004357.

Figure Captions

Fig. 1. Regional map of the British Isles showing the major terranes of Ireland. The rectangle indicates the study area shown in Figure 3 (modified after McIlroy & Horák 2006; Waldron *et al.* 2014; McConnell *et al.* 2016; alternative ISZ after Todd *et al.* 1991). Inset: regional map showing broad tectonic domains (Linnemann *et al.* 2007; Nance *et al.* 2012; Waldron *et al.* 2014; Waldron *et al.* 2011; Waldron *et al.* 2009) Key: BT, Bellewstown Terrane; CG, Carnsore Granite; DB, Dingle Basin; GG, Galway Granite; GT, Grangegeeth Terrane; ISZ, Iapetus Suture Zone; NIC, Newry Igneous Complex; SG; Saltees Granite; TIC, Tyrone Igneous Complex.

Fig. 2. Kernel density plots (bandwidth = 20 Ma) of published detrital zircon ages from various potential source terranes (modified and expanded after Pointon *et al.* 2012). Laurentia data from Cawood *et al.* (2003); Friend *et al.* (2003); Cawood *et al.* (2007); Kirkland *et al.* (2008); Waldron *et al.* (2008); McAteer *et al.* (2010); Cawood *et al.* (2012); Strachan *et al.* (2013); Waldron *et al.* (2014); Johnson *et al.* (2016).

Cadomia-Armorica data from Fernandez-Suarez *et al.* (2002); Samson *et al.* (2005); Linnemann *et al.* (2008); Strachan *et al.* (2008). Ganderia data from Fyffe *et al.* (2009); Waldron *et al.* (2014); Willner *et al.* (2014). Megumia data from Krogh & Keppie (1990); Waldron *et al.* (2009, 2011); Pothier *et al.* (2015). `East' Avalonia data from Collins & Buchan (2004); Murphy *et al.* (2004); Strachan *et al.* (2007); Linnemann *et al.* (2012); Willner *et al.* (2013). `West' Avalonia data from Keppie *et al.* (1998); Thompson & Bowring (2000); Barr *et al.* (2003); Murphy *et al.* (2004a, 2004b); Pollock (2007); Satkoski *et al.* (2010); Thompson *et al.* (2012); Dorais *et al.* (2012); Barr *et al.* (2012); Force & Barr (2012); Pollock *et al.* (2012); Willner *et al.* (2013); Henderson *et al.* (2016). Caledonian granite ages are from Ireland only and include various isotopic geochronological techniques. The Caledonian data are taken from Chew & Stillman (2009).

Fig. 3. Geological map and generalised north-south geological cross section of the Dingle Peninsula showing sample locations (modified after Todd 1989; Ennis *et al.* 2015). Detrital apatite and detrital zircon sample location marked by bold asterisks and white stars respectively.

Fig. 4. Wetherill concordia plots for all detrital zircon samples analysed in this study (youngest ages inset). (a) Coumeenoole Formation (sample AK19). (b) Slea Head Formation (sample AK21). (c) Cappagh Sandstone Formation (sample AK17).


Fig. 5. Detrital zircon age KDE, histogram plots and age percentage distribution plots from Dingle Peninsula samples. (a) Coumeenoole Formation (sample AK19). (b)

1197 Slea Head Formation (sample AK21). (c) Cappagh White Sandstone Formation
1198 (sample AK17).
1199
1200 **Fig. 6**. KDE and histogram plots for detrital apatite U-Pb ages in (a) the Ballymore
1201 Formation and (b) the Farran Sandstone Formation.
1202
1203 **Fig. 7**. Histogram and KDE plots of revised detrital white mica ages from a sample
1204 from the Coumeenoole Formation at Connor Pass on the Dingle Peninsula (original
1205 data from Ennis *et al.* 2015). Bold arrow represents approximate depositional age
1206 (see text for explanation of 382 Ma peak).
1207
1208 **Fig. 8**. KDE plots of detrital zircon ages from various potential local sediment
1209 sources as well as age distributions for detrital zircons analysed in this study. (a)
1210 Upper Ordovician to Llandovery sedimentary rocks of the Southern Uplands terrane
1211 (Waldron *et al.* 2008, 2014) with dominant Laurentian provenance. (b) Upper
1212 Ordovician to Llandovery sedimentary rocks of the Longford Down terrane showing
1213 peri-Laurentian arc provenance (McConnell *et al.* 2016). (c) A single sample from the
1214 Llandovery Lough Avaghon Formation in the Longford Down terrane showing
1215 dominant Laurentian provenance (McConnell *et al.* 2016). (d) Ganderian provenance
1216 of Cambrian sedimentary rocks in the Leinster terrane (Waldron *et al.* 2014). (e)
1217 UORS on the Dingle Peninsula. (f) Composite of two samples from the LORS Dingle
1218 Group. (g) Multi-dimensional scaling map of individual samples used in plots (a)-(f).
1219 Labels indicate broad provenance interpretations from original studies. Axes values
1220 are dimensionless K-S distances (Vermeesch 2013). The most closely related

samples are joined by a solid line and the second closest are marked with a dashed line. N, number of samples; n, number of single zircon grain ages.

Fig. 9. Possible palaeodrainage pattern in the Dingle basin during deposition of the Coumeenoole and Slea Head Formations (modified after Todd 2000). CG, Carnsore Granite; DBL, Dingle Bay Lineament; GG, Galway Granite Batholith; LB, Leinster Batholith; NKL, North Kerry Lineament.

Figure 1



Main map:
- Cenozoic rocks
- Mesozoic rocks
- Carboniferous + Permian rocks
- Devonian + Pridoli rocks
- Caledonian intrusions
- Cambrian-Silurian rocks
- Precambrian

Inset:
- Cadomia
- Avalonia
- Ganderia
- Megumia
- Peri-Laurentia
- Laurentia
- Baltica

Figure 2



Laurentia (n=2744)
Cadomia-Armorica (n=276)
Ganderia (n=1046)
Megumia (n=635)
'East' Avalonia (n=1049)
'West' Avalonia (n=2119)
Caledonian granites (n=52)
0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
4.0
Age (Ga)

Figure 3



N
S
LK
CA
LS
AK17
KM
BD
SC
BM
400 m
FN
TC
IC
SH
AK21
GB
AK19
BF
CO
EK
BH
Unconformity
Acadian unconformity
Fault
Dingle Succession
N
Tralee Bay
AK19
AK17
AK21
0 5 10 km
Upper Devonian
Lower/ Middle Devonian
Silurian
Ordovician
Slieve Mish Group: LS - Lough Slat Fm; CA - Cappagh Sandstone Fm; LK - Lack Sandstone Fm
Pointagare Group
Caherbla Group: IC - Inch Conglomerate Fm; KM - Killmurray Sandstone Fm
Smerwick Group: FN - Farran Sandstone Fm; BD - Ballydavid Sandstone Fm; SC - Sause Creek Fm
Dingle Group: BH - Bulls Head Fm; BF - Ballyferriter Fm; EK - Eask Sandstone Fm; CO - Coumeenoole Sandstone Fm; GB - Glashabeg Conglomerate Fm; Trabeg Conglomerate Fm; SH - Slea Head Fm; BM - Ballymore Sandstone Fm
Dunquin Group
Annascaul Fm

Figure 4



(c) AK17
data-point error ellipses are 2σ
3800
3400
3000
2600
2200
1800
800
700
600
500
400
300
(b) AK21
data-point error ellipses are 2σ
3000
2600
2200
1800
1400
800
700
600
500
400
300
(a) AK19
data-point error ellipses are 2σ
3000
2600
2200
1800
1400
800
700
600
500
400
300
$^{206}Pb/^{238}U$
$^{207}Pb/^{235}U$

Figure 5



Palaeozoic
Mesoproterozoic
Neoproterozoic
Palaeoproterozoic
Archaean
(c) AK17 (n=84)
(b) AK21 (n=98)
(a) AK19 (n=86)
Zircon count
Age (Ma)
0
500
1000
1500
2000
2500
3000
3500
4000
2
4
6
8

Figure 6



(b)
420 Ma
Farran Sandstone Fm. (n=17)
376 Ma
458 Ma
(a)
419 Ma
Ballymore Fm. (n=70)
Apatite count
Age (Ma)
0
2
4
6
8
10
12
1
2
3
300
400
500
600
700
800
900

Figure 7



Coumeenoole Formation
(n=43)
~382 Ma
~414 Ma
Mica count
Age (Ma)
0
1
2
3
4
5
200
250
300
350
400
450
500

Figure 8




A.
Southern Uplands terrane
Waldron et al. (2014)
Waldron et al. (2008)
N=7 ; n=566
B.
Longford-Down terrane
McConnell et al. (2016)
N=5; n=491
C.
Longford-Down terrane
McConnell et al. (2016)
N=1; n=100
D.
Leinster terrane
Waldron et al. (2014)
N=4; n=315
E.
AK17 (UORS)
This study
N=1; n=84
F.
LORS
This study
N=2; n=184
0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
4.0
Age (Ga)
G. MDS plot of study samples and potential source areas
Grampian-volcanic-arc-derived
peri-Gondwana-derived
Laurentia-derived
nearest neighbour
second nearest
0.2
0.0
−0.2
−0.4
−0.6
Dimension 2
−1.0
−0.5
0.0
0.5
Dimension 1

Figure 9


N
Coumeenoole/
Slea Head
drainage basin
GG
~400 Ma
Glashabeg
drainage
basin
LB
~405 Ma
??
NKL
DBL
Trabeg
drainage
basin
SG+CG
~430 Ma
Old Red Sandstone
Caledonian intrusions
Lower Palaeozoic
0
50
100 km
Peri-Laurentia
Ganderia
Iapetus Suture Zone