


Title	Cholestasis induced by bile duct ligation promotes changes in the intestinal microbiome in mice
Authors	Cabrera-Rubio, Raul;Patterson, Angela M.;Cotter, Paul D.;Beraza, Naiara
Publication date	2019-08-23
Original Citation	Cabrera-Rubio, R., Patterson, A. M., Cotter, P. D. and Beraza, N. (2019) 'Cholestasis induced by bile duct ligation promotes changes in the intestinal microbiome in mice', Scientific Reports, 9(1), 12324. (10pp.) doi: 10.1038/s41598-019-48784-z
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1038/s41598-019-48784-z
Rights	© The Author(s) 2019. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/ http://creativecommons.org/licenses/by/4.0/.
Download date	2025-06-01 11:24:43
Item downloaded from	https://hdl.handle.net/10468/9335




Cholestasis induced by bile duct ligation promotes changes in the intestinal microbiome in mice

Raul Cabrera-Rubio^{1, 2}, Angela M Patterson³, Paul D Cotter^{1, 2}, Naiara Beraza³

Supplemental material for Cabrera-Rubio et al.



В



Supplemental Figure legend

Supplemental Figure 1: Effects of BDL treatment on bacterial composition. (A) Phylum level bacterial composition in individual samples obtained from mice (control, BDL 3d and BDL 7d). (B) Family level bacterial composition in individual samples obtained from mice (control, BDL 3d and BDL 7d).