

Title	Life is like a box of chocolates: meeting the periodontal challenge
Titte	Life is the a box of endedtates. Meeting the periodomat enditering
Authors	Roberts, Anthony
Publication date	2014-06
Original Citation	Roberts, A. (2014) Life is like a box of chocolates: meeting the periodontal challenge, The Probe, 14 June, pp. 52-53.
Type of publication	Report
Link to publisher's version	http://www.dentalrepublic.co.uk/the-probe
Rights	© 2014, Dental Republic.
Download date	2024-04-20 11:30:32
Item downloaded from	https://hdl.handle.net/10468/2850



Life is like a box of chocolates: meeting the periodontal challenge

PROFESSOR ANTHONY ROBERTS on why managing patients with perio disease is akin to delving into a box of chocolates - you never know what you're going to get...

HEN considering periodontal disease, it seems prudent, initially, to consider the magnitude of the challenge in dentistry. The Adult Dental Health (ADH) Survey 2009 indicates that the number of people aged 65 years and over who are edentulous has

reduced from circa 75 per cent in 1978 to less than 25 per cent, hopefully due, in no small part, to improvements in the quality care we provide our patients. Clearly, not all of the teeth lost are solely due to periodontal disease, but the ADH Survey tells us that approximately 50 per cent

sport with these fantastic

treatment package offers

PER MONTH

Purchase an A-dec 400 chai packase and receive a Free

upgrade to Fibre optics and

the A-dec LED chair mounted light

Cal H-dec today (READER ENQUIRY NO: 40) Freephone DBOO 233 285 a sole c

KICK OFF the summer of

affecting the periodontal tissues. In addition, overall, 45 per cent of adults had periodontal (gum) pocketing exceeding 4mm and, no doubt, unfortunately, some of these teeth will be lost. The conclusions we can draw from the ADH Survey in periodontal terms are:

- Perio disease remains common at a low level
- Overall, a reduction in mild disease.
- There is a slight increase in the prevalence of more severe disease.
- The impact of severe disease is concentrated

• There are associations with a range of health behaviours, including smoking, infrequent toothbrushing and social gradient (albeit minor).

Clearly, when this information is considered as a whole, the prevalence of periodontal conditions is significant.

Gingivitis and periodontitis

This was the focus of the fifth European Workshop on Periodontology in 2005, where the panel considered whether gingivitis and periodontitis are a continuum of the same inflammatory disease. To summarise their conclusions:

- · Gingivitis and periodontitis are a continuum of the same inflammatory disease
- Not all patients with gingivitis will go on to develop periodontitis.
- Chronic inflammatory periodontal disease encompasses gingivitis and periodontitis as tangible clinical forms

So, it would appear that there are two patient types. Some patients will retain their gingivitis state in perpetuity; they have chronic gingivitis, poor oral health and high plaque scores, and yet they seem resistant to attachment loss. However, others will go on to develop periodontitis, which, of course, may, unfortunately, lead to tooth loss.

Importantly, given the evidence reviewed at the European Workshop, the precursor to periodontitis is always gingivitis, and to extrapolate from this, perhaps greater focus should be directed by dental care professionals on the management of gingivitis so that initiation of periodontitis is prevented.

Who's going to get periodontitis?

Epidemiological data indicates that a significant number of patients with gingivitis will develop periodontitis, but how do we know who's at risk?

There are a variety of elements in what we do on a daily basis in our clinics that can help us try to understand who will develop periodontitis For example, there is the Basic Periodontal Examination (BPE) screening tool to help identify patients developing periodontal problems. Readers should note that there have been some recent changes to the BPE system and the latest quidance document from the British Society of Periodontology can be found at www.bsnerio. org.uk/publications/downloads/39 143748 bpe2011.pdf, and www.bsperio.org.uk/ publications/downloads/53_085556_executivesummary-bsp_bspd-perio-guidelines-for-theunder-18s.pdf.

Of course, the BPE is just one assessment tool that identifies individual patients where the disease process has already begun. Significant efforts have also been made to try to identify those individuals at risk of developing disease, but that may not have the disease already. Lang & Tonetti's (1996) Bernese spider web/hexagor offers a further tool for patient assessment, this time from a risk factor perspective. Versions are available at www.perio-tools.com/pra/en/.

As the various criteria are entered for each patient (for example, age, smoking status, etc.), the schematic diagram pictorially represents the level of risk that the patient exhibits. with the anticipation that any modifiable risk factors can be addressed to minimise the risk of developing periodontitis. Other excellent examples of periodontal risk assessments/ calculators are changing the way we bring these factors together in a form that is understandable preventing it before it starts: www.previser.co.uk.

How will periodontitis progress?

There are many studies that have aimed to identify the natural history of periodontal diseases and, broadly, there are two patterns of disease progression

- 1. The linear model where a patient loses attachment in a linear manner, which is relatively predictable.
- 2. The discontinuous model where the disease activity increases sporadically over time. There will be periods of increased and decreased disease activity and progressive loss of attachment over time in an episodic manner. This pattern of disease is much less predictable.

Ultimately, however, there are diverse groups of patients whose oral health behaves differently and it is very difficult – almost impossible – to predict how periodontitis will progress in individual natients

Types of treatment

Currently, oral plaque biofilm disruption is the most effective way to treat and prevent both conditions, through mechanical and, where appropriate, chemical intervention. On a dayto-day basis, the emphasis is on the patient to affect their plaque biofilm (removal or disruption) using the variety of homecare products available. From a clinician's perspective, in terms of scheduling treatment for patients with chronic periodontitis, there are three main approaches that have been implemented over the years: Conventional Staged Debridement (CSD). which involves scaling and root surface debridement over a period of weeks in a quadrant-wise or sextant-wise manner.

- Full Mouth Disinfection (FMDis), which comprises scaling and root surface debridement in the shortest period of time and the use of chlorhexidine on the tongue, tonsils and subgingivally with the aim of minimising the bacterial load in the various bacterial reservoirs in and around the oral cavity.
- Full Mouth Debridement (FMDeb), which is essentially the same as the FMDis approach, but without the adjunctive use of chlorhexidine.

Which treatment regime is best for patients with chronic periodontitis? Thankfully, all of the modalities achieve much the same outcomes. FMDis and FMDeb do offer slightly better statistical results, but whether they are clinically significant is debatable. Therefore, all three treatment modalities may be recommended for chronic periodontitis (without preference), provided adequate preventive measures are delivered. A different conclusion is conferred to patients with aggressive periodontitis where FMDeb with the adjunctive use of systemic antimicrobials is commonly advocated as the preferred non-surgical treatment modality.

Outcomes of treatment?

Treatment outcome is dependent upon many factors, including the extent of the initial pocketing, location, plaque control and treatment quality. Local anatomy is highly significant, for example, furcation involvement is also particularly important in terms of treatment outcomes. Data from Claffey et al (1990) indicates that the outcome of periodontal treatment in teeth that are furcally involved is considerably worse than where teeth have no furcation involved. Many patients will require several cycles of

CSD/FMDis/FMDeb to reduce periodontal

where the expectation is that one cycle of treatment is enough; often, it won't be. The requirement for multiple treatment cycles needs to be communicated to patients so that they have realistic expectations relating to the time involved and the financial elements required to achieve a successful periodontal outcome. Of course, the requirements of a periodontal maintenance regime also need to be communicated fully.

To conclude, there are a significant number

the consensus that this state has the potential to deteriorate to chronic periodontitis, the management of gingivitis patients should be a focus of attention. Positive improvements in periodontal health can be provided using a variety of treatment approaches, but consistently affecting the dental plaque biofilm is key. Unfortunately, the pathogenesis of periodontal disease is complex, and while there are tools to assist clinicians to predict those individuals who are likely to have periodontitis, there is currently

a consequence, patients with periodontitis are a significant challenge and outcomes are often like a box of chocolates.

Reader enquiry: 120

Professor Anthony Roberts is professor of Periodontology at University College Cork in Ireland. Since qualifying as a dentist in 1996, Prof Roberts has worked in general practice as well as hospital and university settings.

READER ENGLISH NO. 41

Diamond...



a dentist's best friend



Capsules at Half Price!

Get a Kemdent Disinfectant Starter Pack FREE when you spend £175.00 plus VAT on Diamond Glass Ionomer products during June 2014

To see more offers on Kemdent Glass Ionomer Cements PLUS two new Kemdent Videos on how to use Kemdent's Diamond Rapid Set Capsules and Diamond Carve visit our website www.kemdent.co.uk



FOR MORE INFORMATION CALL: +44 (0)1793 770090 QUOTE: GICProbe06

For information on Kemdent products visit www.kemdent.co.uk. Offer ends 30th June 2014. Quote Reference: GICProbe06

Prices exclude VAT. Minimum order for free delivery £100 excluding VAT (UK Mainland only) Additional charges may apply for delivery outside of Mainland UK

(+44) 01793 770 256 f: (+44) 01793 772 256













Offers valid From 7th Nay - 31st August 2014

New Advanced Defence Gum Treatment cuts gingival bleeding by 50.9% in just 4 weeks1

New Advanced Defence Gum Treatment cuts gingival bleeding by 50.9% in just 4 weeks1

The Probe | June 14 dentalrepublic.co.uk June 14 | The Probe