

Title	Evolutionary stasis of a heritable morphological trait in a wild fish population despite apparent directional selection
Authors	O'Sullivan, Ronan J.;Aykanat, Tutku;Johnston, Susan E.;Kane, Adam;Poole, Russell;Rogan, Ger;Prodöhl, Paulo A.;Primmer, Craig R.;McGinnity, Philip;Reed, Thomas E.
Publication date	2019-06-11
Original Citation	O'Sullivan, R. J., Aykanat, T., Johnston, S. E., Kane, A., Poole, R., Rogan, G., Prodöhl, P. A., Primmer, C. R., McGinnity, P. and Reed, T. E. (2019) 'Evolutionary stasis of a heritable morphological trait in a wild fish population despite apparent directional selection', Ecology and Evolution, 9(12), pp. 7096-7111. doi: 10.1002/ ece3.5274
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://onlinelibrary.wiley.com/doi/full/10.1002/ece3.5274 - 10.1002/ece3.5274
Rights	© 2019 The Authors. Ecology and Evolution published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited https://creativecommons.org/licenses/ by/4.0/
Download date	2025-06-01 01:31:21
Item downloaded from	https://hdl.handle.net/10468/11527



University College Cork, Ireland Coláiste na hOllscoile Corcaigh



Figure S3: Pedigree of Atlantic salmon, *Salmo salar*, from the Burrishoole catchment, Ireland. Red lines represent maternities and blue lines represent paternities. Hatchery maternal/paternal links are represented with lighter shading. Hatchery and wild parents are represented with triangles and circles, respectively. Years on the y-axis represent the spawning cohort to which an individual belonged. The unrooted individuals at the top of the pedigree represent dams and sires whose offspring spawned in one of the study cohorts.