Title	Infection by the castrating parasitic nematode Sphaerularia bombi changes gene expression in Bombus terrestris bumblebee queens
Authors	Colgan, T. J.;Carolan, J. C.;Sumner, S.;Blaxter, M. L.;Brown, M. J. F.
Publication date	2019-09-30
Original Citation	Colgan, T. J., Carolan, J. C., Sumner, S., Blaxter, M. L. and Brown, M. J. F. (2020) 'Infection by the castrating parasitic nematode Sphaerularia bombi changes gene expression in Bombus terrestris bumblebee queens', Insect Molecular Biology, 29(2), pp. 170-182. doi: 10.1111/imb.12618
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1111/imb.12618
Rights	© 2019, The Royal Entomological Society. Published by John Wiley & Sons, Inc. This is the peer reviewed version of the following item: Colgan, T. J., Carolan, J. C., Sumner, S., Blaxter, M. L. and Brown, M. J. F. (2020) 'Infection by the castrating parasitic nematode Sphaerularia bombi changes gene expression in Bombus terrestris bumblebee queens', Insect Molecular Biology, 29(2), pp. 170-182, doi: 10.1111/imb.12618, which has been published in final form at https://doi.org/10.1111/imb.12618. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Download date	2026-08-09 08:13:08
Item downloaded from	https://hdl.handle.net/10468/15137

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Supplemental Information for:

Infection by the castrating parasitic nematode *Sphaerularia bombi* changes gene expression in *Bombus terrestris* bumblebee queens

Thomas J. Colgan, James C. Carolan, Seirian Sumner, Mark L. Blaxter and Mark J.F. Brown

Supplemental Figure S1:



Supplemental Fig. 1. Principal component analysis for normalised gene-level estimated counts for transcriptomic profiles of diapause (blue) and post-diapause queens (grey). Each point represents an individual queen and infection status is indicated by shape. The first two principal components are displayed with principal component 1 (PC1) explaining 76% of the variance in the dataset and separates both time-points into two distinct clusters.