CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Boolean rings are definitely commutative!
Authors	MacHale, Desmond
Publication date	2015
Original Citation	MacHale, D. (2015) 'Boolean Rings are Definitely Commutative!', Bulletin of the Irish Mathematical Society, 76, pp. 77-78.
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://banach.ucd.ie/bull76/index.php
Rights	© 2015 Irish Mathematical Society
Download date	2026-11-01 12:31:31
Item downloaded from	https://hdl.handle.net/10468/9603



UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Irish Math. Soc. Bulletin
Number 76, Winter 2015, 77–78
ISSN 0791-5578

Boolean Rings are Definitely Commutative!

DESMOND MACHALE

Abstract. A ring $\{R, +, .\}$ is called Boolean if $r^2 = r$ for all $r \in R$. We present four proofs that a Boolean ring is commutative.

A ring $\{R, +, .\}$ is called Boolean if $r^2 = r$ for all $r \in R$. In this bicentenary year of Boole's birth we present four proofs that a Boolean ring is commutative. Our first proof is the standard one found in many textbooks.

Proof 1. For all $r \in R$ we have $r = r^2 = (-r)^2 = -r$, so $r + r = 0$. Next, for all x and y in R, $x+y = (x+y)^2 = x^2+xy+yx+y^2$, so by cancellation in the group $\{R, +\}$, we have $xy+yx = 0 = xy+xy$, by the above. Again by cancellation we have $xy = yx$, as required. □

Proof 2. As in Proof 1, $xy + yx = 0$, for all x and y in R. Since for all $r \in R$, $0.r = 0 = r.0$ we have $(xy + yx)x = x(xy + yx)$ or $xyx + y.x^2 = x^2.y + xyx$. Cancelling xyx and remembering that $x^2 = x$, we get $xy = yx$, as required. □

Proof 3. Since for all r, $r^2 = r$ it follows that if $r^2 = 0$ then $r = 0$. Now for all x and y in R we have $(xy - xyx)^2 = xyxy + xyxxyx - xyxyx - xyxxy = xyxy + xyxyx - xyxyx - xyxy = 0$. So $xy - xyx = 0$ and $xy = xyx$. Then $(yx - xyx)^2 = yxyx + xyxxyx - yxxyx - xyxyx = yxyx + xyxyx - yxyx - xyxyx = 0$. So $yx - xyx = 0$ and $yx = xyx$. Thus $xy = yx$ as required. □

Proof 4. For $a, b \in R$ if $ab = 0$, then $ba = (ba)^2 = b(ab)a = 0$. Now, $0 = xy - xy = xy - x^2y = x(y - xy)$, so $0 = (y - xy)x = yx - xyx$. Also, $0 = yx - yx = yx - yx^2 = (y - yx)x$, so $0 = x(y - yx) = xy - xyx$. Thus $xy = yx$ for all x and y in R. □

We note it is immediate in all four proofs that $xy = yx = xyx = yxy$, for all x and y.

2010 *Mathematics Subject Classification.* 19E50.
Key words and phrases. Boolean Rings.
Received on 8-6-2015.

©2015 Irish Mathematical Society

Desmond MacHale received his Ph.D. from the University of Keele and is Emeritus Professor of Mathematics at University College Cork where he taught for nearly forty years. His research interests include commutativity in groups and rings, automorphisms of groups, Euclidean geometry, number theory, and mathematical humour.

School of Mathematical Sciences, University College Cork.
E-mail address: d.machale@ucc.ie

Copyright of Bulletin of the Irish Mathematical Society is the property of IMS Bulletin and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.