

Title	Maternity ultrasound in the Republic of Ireland 2016: a review
Authors	Hayes-Ryan, Deirdre;McNamara, Karen;Russell, Nóirín;Kenny, Louise C.;O'Donoghue, Keelin
Publication date	2017-08
Original Citation	Hayes-Ryan, D., McNamara, K.,Russell, N., Kenny, L. and O'Donoghue, K. (2017) 'Maternity ultrasound in the Republic of Ireland 2016: a review', Irish Medical Journal, 110(7), 598 (9pp). doi:10147/622508
Type of publication	Article (peer-reviewed)
Link to publisher's version	10147/622508
Rights	© 2017, Irish Medical Journal. All rights reserved.
Download date	2025-08-20 08:05:12
ltem downloaded from	https://hdl.handle.net/10468/5889



University College Cork, Ireland Coláiste na hOllscoile Corcaigh



# Maternity Ultrasound in the Republic of Ireland 2016; A Review

D Hayes-Ryan<sup>\*1,3</sup>, K McNamara<sup>\*1,2,3</sup>, N Russell<sup>2,3</sup>, L Kenny<sup>1,3</sup>, K O'Donoghue<sup>1,2,3</sup>

\*Joint lead authors

<sup>1</sup>The Irish Centre for Fetal and Neonatal Translational Research (INFANT), University College Cork <sup>2</sup>Pregnancy Loss Research Group, Department of Obstetrics and Gynaecology, University College Cork

<sup>3</sup>Cork University Maternity Hospital, Wilton, Cork.

## Abstract

Antenatal ultrasound, comprising of a dating ultrasound in the late first trimester followed by a fetal anomaly scan, is a recognised and necessary component of good antenatal care. We conducted a telephone survey of all 19 obstetric units to ascertain the status of maternity ultrasound provision in Ireland. Fetal anomaly ultrasound is offered universally to all women in 7/19 (37%) units, selectively to some women in 7/19 (37%) units and not offered at all in the remaining 5/19 (26%) units. Overall  $\approx$  41,700 (64%) women receive a fetal anomaly ultrasound nationally. Universal first trimester ultrasound, performed in a dedicated ultrasound department by a suitably qualified sonographer, is offered to 47% of women nationally. This study highlights the lack of development in Irish maternity ultrasound services over the last decade. Substantial investment by health care policy makers is urgently needed.

### Introduction

Obstetric ultrasound is a recognised and necessary component of good antenatal care 1,2. Internationally, the widely accepted <u>minimal</u> schedule for antenatal ultrasound comprises of two examinations; a dating ultrasound in the late first trimester, followed by a fetal anomaly scan, usually performed between 20-22 weeks' gestation<sup>3-5</sup>. Routine first trimester ultrasonography is usually performed between the gestational age of  $10^{+0}$  and  $13^{+6}$  weeks. This is the most appropriate time point at which to accurately date a pregnancy<sup>3</sup>, and may have a role to play in reducing the incidence of unnecessary induction of labour for suspected prolonged pregnancy<sup>6, 7</sup>. In the event of a multiple pregnancy diagnosis this is also the ideal time to accurately assign chorionicity, which is vital to ensure appropriate management<sup>6, 8</sup>. Some major fetal anomalies are also detectable in the late first trimester<sup>6</sup> and when diagnosed can facilitate parental choice with respect to continuing with, or terminating the pregnancy, in jurisdictions where this is legally permissible.

Sonographers performing first trimester ultrasound need to have appropriate training in the use of diagnostic ultrasonography and they also need to maintain clinical competency through ongoing medical educational activities<sup>6</sup>. Second trimester fetal anomaly ultrasound is routinely recommended by all obstetric policy-makers<sup>2-4, 9</sup>. If a fetal anomaly is detected, planned delivery of the infant at the right time and in the correct place can be facilitated, thus enabling rapid access to appropriate neonatal intensive care which may ultimately reduce neonatal morbidity and mortality<sup>10, 11</sup>. Diagnosis can also provide adequate time to psychologically prepare parents for the challenges of the pregnancy, intrapartum, and postnatal period. A timely diagnosis of a lethal fetal anomaly, which is devastating for parents, can allow them time to adjust and make memories with their infant.<sup>12, 13</sup> In certain jurisdictions, the detection of a lethal fetal anomaly can also provide parents with the option to continue with the pregnancy or to terminate<sup>3</sup>.

The fetal anomaly scan is best performed between 20 <sup>+0</sup> to 22 <sup>+6</sup> weeks' gestation as this time point allows the best visualisation of fetal anatomical structures<sup>3, 4</sup>. This ultrasound requires a high skill level and needs to be performed by an individual with "specialised training for the practice of diagnostic ultrasonography in pregnant women"<sup>4</sup>. Sonographers performing fetal anomaly ultrasound need to maintain competence by "Continuing Professional Development" and by performing a minimum of 100 ultrasounds per year<sup>14</sup>. Ultrasound machines used should be of high quality, undergo regular maintenance and should have real time, adjustable acoustic power and freeze frame capability. They should also have the capacity to store and print images<sup>4</sup>. Fetal anomaly ultrasound should also follow a strict format and minimum requirements have been set by the major obstetric ultrasound professional bodies<sup>2,4,14</sup>.

Since 2000 the Royal College of Obstetricians and Gynaecologists (RCOG) recommended that all pregnant women should be offered these two ultrasounds for the accurate dating of a pregnancy as well as the diagnosis of fetal anomaly<sup>15</sup>. This two stage ultrasound approach is also endorsed by the National Institute for Health and Clinical Excellence (NICE) in 2008<sup>3</sup>. In January 2016, the Department of Health in the Republic of Ireland published the National Maternity Strategy which also recommended that all women have "equal access to standardised ultrasound services, to accurately date the pregnancy (and) to assess the fetus for ultrasound diagnosable anomalies as part of a planned Prenatal Fetal Diagnostic

Service<sup>"16</sup>. Previous research reported by Lalor et al in 2007<sup>17</sup> and Walsh et al in 2013<sup>18</sup> revealed a large variation in the number and type of ultrasounds provided by the maternity units around Ireland. Recent high-profile media reports, highlighting a lack of diagnosis of lethal fetal anomalies over the last number of years, suggest that availability to fetal anomaly ultrasound remains inconsistent. For these reasons it is appropriate and timely to once again determine the scale of the provision of maternity ultrasound services in the Republic of Ireland.

#### Methods

A telephone survey of all Irish obstetric units was undertaken in February 2017 by the two lead authors. Respondents were asked to provide verbal answers to standard questions relating to the provision of maternity ultrasound services in their unit for the year 2016. The domains focused on were; first trimester ultrasound, fetal anomaly ultrasound and early pregnancy assessment (EPA) ultrasound. In all cases a senior clinician, either a consultant obstetrician or a lead sonographer was contacted in order to ensure that respondents had a thorough knowledge of the services provided in their unit. Results were analysed using IBM-SPSS Version 22. Percentages are given to the nearest whole percentage point. For the purposes of this study, and based on international guidelines<sup>4, 6</sup> these were the definitions used; a routine ultrasound referred to an ultrasound offered universally to all women attending that particular unit at a specific gestational age. A fetal anomaly ultrasound referred to a detailed structural assessment of the fetus, performed by an appropriately trained sonographer<sup>14</sup> between 20-22 weeks' gestation that adhered to a written checklist proforma based upon a recognised internationally accepted guideline such as ISUOG<sup>4</sup> or NHS FASP <sup>9</sup>. This is also sometimes referred to as a standard fetal anomaly ultrasound. A first trimester ultrasound referred to an ultrasound performed by a suitably trained sonographer, before 14 weeks' gestation where the main purpose was to accurately date the pregnancy<sup>6</sup>.

#### Results

In total 100% (n=19) of Irish maternity units were successfully contacted and completed the survey. Approximate annual delivery rates for 2016 were provided to us from each respondent for their respective units and ranged from 1062 to 9186 births with the total number of combined births  $\approx$  65,500. All 19 units have a dedicated Early Pregnancy Assessment Ultrasound service available. A dedicated maternity ultrasound department is present in 16/19 (84%) of the units. Fetal anomaly ultrasound is offered universally to all women in 7/19 (37%) units, selectively to some women in 7/19 (37%) units and not offered at all in the remaining 5/19 (26%) units (Figure 1). In units offering selective anomaly scanning the variation in percentage of women receiving a scan ranges from 10-42%. Figure 2 illustrates the geographic variation that exists nationally in provision of fetal anomaly ultrasound scans. Overall  $\approx$  41,700 (64%) women receive a fetal anomaly ultrasound nationally.

Of the 14 units providing some level of anomaly ultrasound, all units perform the scans between 20-22 weeks' gestation. The time allocated per ultrasound is 20-30 minutes in 7/14 (50%) units and greater than 30 minutes in 6/14 (43%) units. In one unit the respondent was not sure as to how much time was allocated for each anomaly ultrasound. In 6/14 (43%) units verbal information is provided to women undergoing an anomaly ultrasound while both written and verbal information is provided to women in the remaining 8/14 (57%) units. Fetal anomaly ultrasound is performed by a midwife sonographer in 9/14 (62%) units, a radiographer in 1/14 (7%) units, a Fetal Medicine Specialist in 2/14 (14%) units and a combination

of these in the remaining 2/14 (14%) units.



ultrasound scans from the total pregnant population (65,500).

There is some variation around the limited provision of anatomy ultrasounds in the units that do not provide a standard fetal anomaly ultrasound. In one of the units, a nearby larger unit provides 100 ultrasounds per annum, which represents  $\approx$  7% of their booking population. The clinical indication for these ultrasounds is based on maternal age. Another unit provides a limited anatomical survey to all women between 18-22 weeks' gestation. Another unit offers 50% of women, based on maternal age and medical/obstetric history, a combined ultrasound at 18-24 weeks' gestation which dates the pregnancy and identifies some anatomical components but does not adhere to any formal guideline checklist.



Figure 2: Fetal Anomaly Ultrasound Provision 2016: n refers to the number of women who deliver in units in Dublin or outside of Dublin. The % is reflective of the proportion of women who receive or do not receive fetal anomaly ultrasound scans in the pregnant population in these units. First trimester pregnancy scanning is offered universally to all women in 10/19 units (53%), offered but does not meet international criteria for first trimester ultrasound (see above for definition) in 5/19 units (26%) and is not offered in the remaining 4/19 (21%) units (Figure 3). In the units providing some level of first trimester ultrasound, it is performed in the Ultrasound Department in 12/15 (80%) units while in 3/15 (20%) units it is performed in the antenatal clinic. There is wide variation nationally as to who performs the first trimester ultrasound (Figure 4). While not provided routinely in the remaining 4/19 (21%) units, women are able to obtain a first trimester ultrasound on the basis of a specific clinical indication such as a previous miscarriage, ectopic pregnancy or are able to attend the early pregnancy assessment unit if they experienced any complications in the index pregnancy.



units. The % is reflective of the proportion of women who receive or do not receive first trimester ultrasound scans from the total pregnant population (65,500).

In 9/19 (47%) units, there are ultrasound machines in the antenatal clinic (ANC) for clinicians to use. In 8/19 (42%) units ultrasound in the ANC is limited. In 2/19 (11%) units, there is no availability of ultrasound in antenatal clinic. Only 1/19 (5%) unit offers a routine third trimester growth ultrasound to all patients. We also asked respondents if there had been any staff expansion in the ultrasound department since the publication of the National Maternity Strategy in 2016. In 2/19 (11%) units some staff expansion has occurred in the last year. However, in 2/19 (11%) units the ultrasound staffing levels have declined over the same time period.



Figure 4: Flowchart detailing the variation in provision of First Trimester Ultrasound in 2016 in Ireland

#### Discussion

This study highlights the current variations that exist in relation to maternity ultrasound services in Ireland, and in particular with regard to the provision of fetal anomaly ultrasound. In 2012 Walsh<sup>18</sup> identified 7/20 (35%) maternity units routinely offered fetal anomaly ultrasounds to all women, with a further 9/20 (45%) offering selective fetal anomaly ultrasound. While the number of units offering routine fetal anomaly ultrasonography in 2016 has not changed, the number offering selective fetal anomaly ultrasound has declined from nine to seven. There are a number of potential reasons why our numbers differ from the previous study. Firstly one maternity unit closed in 2014. Secondly, for the purposes of this study we excluded units based on 1) the outsourcing of anomaly scans to a larger affiliated unit and 2) those that only provided a limited anatomical survey. This may account for the apparent decrease in the number offering selective fetal anomaly ultrasound scans. From discussion with these units, we are certain that there has been no overall expansion in the provision of fetal anomaly ultrasound, nor any increase in the qualified staff who provide these scans. Approximately one third of pregnant women in Ireland are still not offered routine fetal anomaly ultrasound, which is a routine part of antenatal care in other countries<sup>3</sup>. We also identified that routine first trimester ultrasound performed in a dedicated ultrasound department by a suitably qualified sonographer is offered to only 47% of women nationally. A further 26% of women are offered a first trimester ultrasound in either antenatal clinic and/or by a doctor who may have no formal training in ultrasound. The number of women not being offered routine first trimester ultrasonography stands at 26% and this is also unacceptably high. Walsh<sup>18</sup> noted that first trimester ultrasound was offered routinely in 65% (13/20) of maternity units but did not specify where these were performed or who performed them; this may explain the discrepancy between their study and the present work.

Previous studies have advocated the need for uniform national and local guidelines on the provision of routine maternity ultrasound services in Ireland<sup>17,18</sup>. While we welcome the publication of the National Maternity Strategy and the other Irish national clinical guidelines recently published that advocate for different types of antenatal ultrasound in specific 8,16,19-21

settings , we strongly recommend the development and implementation of an Irish guideline on obstetric ultrasound. A national guideline should set out the standard components of the anomaly ultrasound scan, and the optimum gestation for provision, as well as specify the training of sonographers and set requirements for ultrasound machines and image storage. An algorithm describing appropriate referral pathways to fetal medicine expertise in the event of an abnormal finding needs to also be described. International guidelines already exist in this area which can be consulted to expedite development of Irish guidelines. Corcoran<sup>22</sup> reported that the introduction of a standardised local practice guideline and appropriately trained sonographers significantly increased the rates of antenatal detection of fetal congenital cardiac disease in Ireland.

It is disappointing to conclude that our study findings reflect those of the previous two national ultrasound surveys (2007, 2012)<sup>17, 18</sup> and confirm that there has been no expansion in Irish maternity ultrasound services in the last decade. If we are serious about equity of care and access to specialist services for all Irish women, regardless of geographical location or financial means, there must now be substantial investment by healthcare policymakers. Without recognition of the deficit that currently exists in both the numbers of suitably qualified sonographers as well as in the provision of facilities and equipment, we will be unable to meet one of the fundamental aims of the 2016 National Maternity Strategy – equal access to standardised ultrasound services to every pregnant woman in Ireland.

Correspondence:

Dr Karen McNamara, Pregnancy Loss Research Group, Department of Obstetrics and Gynaecology, University College Cork, Ireland.

Email: Karen.mcnamara@ucc.ie

#### Conflict of Interest

#### The authors have no conflict of interest to declare

#### References

1. Ahman A, Persson M, Edvardsson K, Lalos A, Graner S, Small R, Mogren I. Two sides of the same coin--an interview study of Swedish obstetricians' experiences using ultrasound in pregnancy management. BMC pregnancy and childbirth. 2015;15:304.

2. Practice Bulletin No. 175: Ultrasound in Pregnancy. Obstet Gynecol. 2016;128(6):e241-e56.

3. National Institute for Health and Care Excellence. Antenatal care for uncomplicated pregnancies, Clinical Guideline CG62, NICE 2008.

4. Salomon LJ, Alfirevic Z, Berghella V, Bilardo C, Hernandez-Andrade E, Johnsen SL, Kalache K, Leung KY, Malinger G, Munoz H, Prefumo F, Toi A, Lee W. Practice guidelines for performance of the routine mid-trimester fetal ultrasound scan. Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology. 2011;37(1):116-26.

5. The Royal Australian and New Zealand College of Obstetricians and Gynaecologists. Mid-trimester fetal morphology ultrasound screening. Royal Australian and New Zealand College of Obstetricians and Gynaecologists. Clinical Guideline

#### Maternity Ultrasound in the Republic of Ireland 2016; A Review - Irish Medical Journal

(C-Obs 49) https://www.ranzcog.edu.au/RANZCOG\_SITE/media/RANZCOG-

MEDIA/Women%27s%20Health/Statement%20and%20guidelines/Clinical-Obstetrics/Fetal-Morphology-Ultrasound-(C-Obs-57)-New-Mar-14.pdf?ext=.pdf

6. Salomon LJ, Alfirevic Z, Bilardo CM, Chalouhi GE, Ghi T, Kagan KO, Lau TK, Papageorghiou AT, Raine-Fenning NJ, Stirnemann J, Suresh S, Tabor A, Timor-Tristsch IE, Toi A, Yeo G. ISUOG practice guidelines: performance of first-trimester fetal ultrasound scan. Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology. 2013;41(1):102-13.

7. Committee opinion no 611: method for estimating due date. Obstet Gynecol. 2014;124(4):863-6.

8. Institute of Obstetricians and Gynaecologists, Royal College of Physicians of Ireland, and Directorate of Strategy and Clinical Care Health Service xecutive. Management of Multiple Pregnancy. 2012.

9. National Health Service Screening Programme. Fetal Anomaly Screening Programme Handbook, June 2015.

10. van Velzen CL, Haak MC, Reijnders G, Rijlaarsdam ME, Bax CJ, Pajkrt E, Hruda J, Galindo-Garre F, Bilardo CM, deGroot CJ, Blom NA, Clur SA. Prenatal detection of transposition of the great arteries reduces mortality and morbidity. Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology. 2015;45(3):320-5.

11. Verheijen PM, Lisowski LA, Stoutenbeek P, Hitchcock JF, Brenner JI, Copel JA, Kleinman CS, Meijboom EJ, Bennik GB. Prenatal diagnosis of congenital heart disease affects preoperative acidosis in the newborn patient. The Journal of thoracic and cardiovascular surgery. 2001;121(4):798-803.

12. McNamara K, O'Donoghue K, O'Connell O, Greene RA. Antenatal and Intrapartum Care of Pregnancy Complicated by Lethal Fetal Anomaly. The Obstetrician and Gynaecologist 2013;15:189-194. 2013.

13. Nuzum D, Meaney S, O'Donoghue K. The Spiritual and Theological Challenges of Stillbirth for Bereaved Parents. Journal of religion and health. 2017.

14. Wax J, Minkoff H, Johnson A, Coleman B, Levine D, Helfgott A, O'Keefe D, Henningsen C, Benson C. Consensus report on the detailed fetal anatomic ultrasound examination: indications, components, and qualifications. Journal of ultrasound in medicine : official journal of the American Institute of Ultrasound in Medicine. 2014;33(2):189-95.
15. RCOG (1997). Report of the RCOG Working Party on Ultrasound Screening for Fetal Abnormalities. RCOG Press, London.

16. Department of Health. National Maternity Strategy – Creating a Better Future Together 2016-2026. 2016.

17. Lalor J, Devane D, McParland P. Ultrasound screening for fetal abnormality in Ireland: a national survey. Ir J Med Sci. 2007;176(3):175-9.

Walsh CA, McAuliffe F, Kinsella V, McParland P. Routine obstetric ultrasound services. Irish medical journal.
 2013;106(10):311-3.

19. Institute of Obstetricians and Gynaecologists, Royal college of Physicians of Ireland, Directorate of clinical strategy and programmes HSE. Delivery after Previous Caesarean Section, 2011.

20. Institute of Obstetricians and Gynaecologists, Royal college of Physicians of Ireland, Directorate of clinical strategy and programmes HSE. Ultrasound Diagnosis of Early Pregnancy Loss, 2010.

21. Institute of Obstetricians and Gynaecologists, Royal college of Physicians of Ireland, Directorate of clinical strategy

#### Maternity Ultrasound in the Republic of Ireland 2016; A Review - Irish Medical Journal

and programmes HSE. Diagnosis and Management of Ectopic Pregnancy, 2014.

22. Corcoran S, Briggs K, H OC, Mullers S, Monteith C, Donnelly J, Dicker P, Franklin O, Malone FD, Breathnach FM. Prenatal detection of major congenital heart disease - optimising resources to improve outcomes. European journal of obstetrics, gynecology, and reproductive biology. 2016;203:260-3.

(P598)

Theme by Tyler Moore



This Month Archive Subscribe For Authors About Us My CPD Hosted Content Home