

Title	Retention of microbiota diversity by lactose-free milk in a mouse model of elderly gut microbiota
Authors	Ntemiri, Alexandra;Rivière, Céline;Stanton, Catherine;Ross, R. Paul;O'Connor, Eibhlís M.;O'Toole, Paul W.
Publication date	2019-01-21
Original Citation	Ntemiri, A., Rivière, C., Stanton, C., Ross, R. P., O'Connor, E. M. and O'Toole, P. W. (2019) 'Retention of Microbiota Diversity by Lactose-Free Milk in a Mouse Model of Elderly Gut Microbiota', Journal of Agricultural and Food Chemistry, 67(7), pp. 2098-2112. https://doi.org/10.1021/acs.jafc.8b06414
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://pubs.acs.org/doi/10.1021/acs.jafc.8b06414 - 10.1021/acs.jafc.8b06414
Rights	© 2019 American Chemical Society. This document is the Accepted Manuscript version of a Published Work that appeared in final form in Journal of Agricultural and Food Chemistry, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see https://pubs.acs.org/doi/10.1021/acs.jafc.8b06414
Download date	2026-08-26 05:41:13
Item downloaded from	https://hdl.handle.net/10468/7761

Retention of microbiota diversity by lactose-free milk in a mouse model of elderly gut microbiota

Alexandra Ntemiri, Celine Ribiere, Catherine Stanton,
R. Paul Ross, Eibhlís O'Connor, and Paul W. O'Toole

J. Agric. Food Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jafc.8b06414 • Publication Date (Web): 21 Jan 2019

Downloaded from <http://pubs.acs.org> on January 28, 2019

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Retention of microbiota diversity by lactose-free milk in a mouse model of elderly gut microbiota

Alexandra Ntemiri ^{†,‡}, Celine Ribiere ^{†,‡}, Catherine Stanton ^{‡,§}, R. Paul Ross ^{†,‡}, Eibhlís M. O'Connor^{‡, #, [⊥]}, Paul W. O'Toole ^{†,‡}.

[†] School of Microbiology, Food Science Building, University College Cork, Western Road, T12 Y337 Cork, Ireland

[‡]APC Microbiome Ireland, University College Cork, Western Road, T12 Y337 Cork, Ireland

[§] Teagasc, Moorepark Food Research Centre, Fermoy, Co Cork P61 C996, Ireland.

[#]Department of Biological Sciences, School of Natural Sciences, University of Limerick, V94 T9PX, Limerick, Ireland.

[⊥]Health Research Institute, University of Limerick, V94 T9PX, Limerick, Ireland.

*Corresponding author: Tel: +353-21-490-3000; Fax: +353 21 490 3997; E-mail: pwotoole@ucc.ie

Abstract: Prebiotics may improve ageing-related dysbiosis. Milk is a source of nutrients including oligosaccharides whose prebiotic potential remains largely unexplored. We used a murine model to explore the effect of milk products on high diversity and lower diversity faecal microbiota from healthy and frail elderly subjects, respectively. Mice were treated with antibiotics and subsequently “humanised” with human faecal microbiota. The mice received lactose-free or whole milk, glycomacropeptide, or soy protein (control) supplemented diets for one month. The faecal microbiota was analysed by 16S rRNA gene amplicon sequencing. Lactose-free milk diet was as efficient as the control diet in retaining faecal microbiota diversity in mice. Both milk diets had a significant effect on the relative abundance of health-relevant taxa (e.g. Ruminococcaceae, Lachnospiraceae). The glycomacropeptide prebiotic activity previously observed *in vitro* was not replicated *in vivo*. However, these data indicate the novel prebiotic potential of bovine milk for human nutrition.

Key words: ageing, faecal microbiota, “humanised” mice, milk, lactose free milk, glycomacropeptide, soy protein, prebiotic potential

15 **Introduction**

16 During the last decade in particular, the bacterial component of the human gut microbiota has
17 been intensively studied providing a more comprehensive understanding of the compositional
18 and functional profile of the gut microbiota in health and disease ¹. Habitual diet, and to a
19 lesser extent host genetics, are two factors that shape the gut microbiota ²⁻⁴. The term
20 dysbiosis refers to shifts in the composition, function and phylogenetic diversity of the gut
21 microbiota of non-healthy individuals in comparison to healthy controls ¹. Metabolic disease,
22 functional gastrointestinal (GIT) disorders and cognitive disorders have been linked to
23 dysbiosis ¹.

24 The low-grade chronic inflammation that characterises older age reflects the decline of
25 immune system fitness to respond to stressors, a phenomenon called inflammaging ⁵. Altered
26 intestinal permeability and dysbiotic gut microbiota are potential inflammaging drivers ⁵⁻⁷.
27 Gut microbiota dysbiosis can be further fuelled by an inflamed GIT environment ⁸. For
28 prevention and treatment of dysbiosis and associated conditions, therapeutic interventions
29 based on probiotics, prebiotics or faecal microbiota transplants (FMT) are being developed ¹.

30 Bovine milk combines many nutrients including high quality protein, calcium, potassium and
31 magnesium, vitamins such as vitamin D, fatty acids such as conjugated linoleic acid, complex
32 sialylated oligosaccharides and glycoproteins ^{9, 10}. There is accumulating data pointing
33 towards certain health benefits of milk and dairy consumption ^{9, 11, 12}. Importantly, milk
34 consumption may contribute to muscle and bone density maintenance in older consumers ¹³,
35 ¹⁴. Despite the potential health benefits, studies have shown that elderly people often fail to
36 meet current recommendations for daily dairy consumption partly due to misconceptions
37 about lactose mal-absorption and dairy fat content ^{15, 16}.

The effect of milk ingestion on the gut microbiota has not been extensively studied. A few recent studies have focused on the potential prebiotic effect of milk oligosaccharides on the gut microbiota. Using animal models “humanised” with infant gut microbiota, Charbonneau *et al.*¹⁷ showed that sialylated bovine milk oligosaccharides (BMO) promoted weight gain associated with *Bacteroides fragilis* and *Escherichia coli* responsiveness to BMOs under malnutrition conditions. Karav *et al.*¹⁸ showed that BMO released from glycoproteins could mimic human milk oligosaccharide (HMO) selectivity for *Bifidobacterium* strains in the infant gut microbiota. Boudry *et al.*¹⁹ observed that a BMO-supplemented diet enhanced gut barrier function, increased caecal and colonic microbiota diversity and *Lactobacillus* relative abundance in a murine model of diet-induced obesity.

Glycomacropeptide (GMP) is a by-product of cheese making and it is released in whey protein from the kappa-casein fraction of milk by the activity of chymosin²⁰. Glycomacropeptide has five mucin-type glycans, high content of sialic acid, high content of branched chain amino acids (BCAA) and essential amino acids (EAA) and is deficient in of aromatic amino acids²⁰. Glycomacropeptide consumption is associated with a number of health benefits including anti-inflammatory and anti-GIT pathogen activity, and it is currently being used as an alternative source of protein for phenylketonuria (PKU) nutritional management^{21, 22}.

There is inconclusive data on the prebiotic potential of GMP. Selective growth promotion of *Bifidobacterium* and *Lactobacillus* by GMP has been reported based on tests in pure cultures but the selectivity failed to be confirmed when complex faecal microbiota communities were tested *in vitro*²³⁻²⁵. Some *in vivo* studies showed that GMP administration increased the colonic abundance of *Lactobacillus* and *Bifidobacterium*^{26, 27}. Sawin *et al.*²⁸ reported that a GMP-supplemented diet significantly decreased the caecal and colonic relative abundance of

Desulfovibrio and Proteobacteria in wild type and PKU mice. Interestingly, GMP selectivity for promoting growth of Bacteroidetes taxa has also been reported ^{27, 28}.

In this study, conventional mice were treated with a cocktail of antibiotics to deplete the gut microbiota before receiving faecal microbiota from either a healthy or a frail elderly donor. The effect of a diet supplemented with either lactose-free milk, whole milk, GMP or soy protein (control) on the transplanted faecal microbiota of elderly humans was then investigated in these “humanised” mice.

Materials and Methods

Animals and experimental design

The Health Products Regulatory Authority (HPRA) of Ireland under the European Union Regulations authorised this project with authorisation number AE19130/P033. Eighty-six 6 week-old female and male C57BL/6 mice were purchased from Envigo RMS Ltd, Oxon, UK and caged in groups of six of the same sex. The mice were weighed during the first week and every subsequent week of the trial.

The mice were acclimatised to the animal facility for one week before the commencement of the six weeks antibiotics (Abx) treatment for the depletion of the indigenous murine gut microbiota. Ampicillin (1g/L; Sigma Aldrich, St Louis, USA), Metronidazole (1g/L; Molekula, Darlington, UK), Vancomycin (500 mg/L; Molekula, Darlington, UK), Imipenem (250 mg/L; Molekula, Darlington, UK) and Ciprofloxacin-HCl (200 mg/L; Santa Cruz Biotechnology, Dallas, USA) were diluted in tap water and filter-sterilised before being given to animals as described before ²⁹. Animals had *ad libidum* access to Abx-water. Limited administration of HydroGel (ClearH2O.com) during the Abx treatment was used for the prevention of weight loss. Animals that lost more than 35% of their body weight were euthanised. One-day wash-out with filtered tap-water followed the Abx treatment.

Subsequently, the animals were “humanised” for three days with faecal microbiota from either a healthy elderly donor (EM425) or a frail elderly donor (EM297). The mice were introduced to the experimental diets on the wash-out day; the mice were ca. 14 weeks-old at the beginning of the dietary intervention. The duration of the dietary intervention was one month. Faecal samples were taken on the first day of acclimatisation (baseline), throughout the Abx treatment (time points T1 and T2), one week after humanisation (T3), in the mid-part of the dietary intervention (T4) and at the end of the trial (T5) (**Figure S1**).

Six male and 6 female mice were allocated to each diet (**Table S1**). The animals were fed maintenance diet (ssniff, Spezialdiäten GmbH, German) during the acclimatisation period and during the Abx treatment. The experimental diets (ssniff, Spezialdiäten GmbH, Germany) were supplemented with one of the following powders: **i. lac-free diet:** 20% lactose-free whole milk powder (Valio Ltd., Helsinki, Finland); **ii. wmilk diet:** 20% whole milk powder (Valio Ltd., Helsinki, Finland); **iii. GMP diet:** 20% CGMP-10 (Arla Food Ingredients, Denmark); **iv. control diet:** 20% soy protein isolate (provided by ssniff, Spezialdiäten GmbH, Germany). In order to achieve the isoenergetic and isonitrogenous profile of the test diets, all diets were additionally supplemented with soy protein isolate (**Table 1**) which was the basic protein ingredient for the maintenance diet. The complete compositional data content of the experimental diets is provided in **Appendix A**. All diets were sterilized by irradiation. The lactose-free milk powder (Valio Ltd., Helsinki, Finland) was produced from whole milk powder (Valio Ltd., Helsinki, Finland) by partial removal of lactose using filtration and by enzymatic hydrolysis of the remaining lactose to glucose and galactose.

Faecal slurry preparation and mouse “humanisation”

One faecal sample from a healthy 81 yr old (EM425; community “COM” type microbiota) and one from a frail 82 yr old subject (EM297; longstay “LS” type microbiota) were collected under the approval of the local Clinical Research Ethics Committee. The samples were processed under anaerobic conditions for the preparation of 10% w/v faecal slurries in PBS with 20% glycerol. The faecal slurries were kept in aliquots at -80°C and thawed in the anaerobic cabinet before gavage to mice. The animals received faecal slurry over three days by oral gavage from a total of 300 µl.

Faecal sample collection and genomic DNA extraction

Faecal samples were collected during the trial at various time points (T0 to T5) as indicated in **Figure S1**. One to two faecal pellets were collected at each time point and were immediately frozen in liquid nitrogen before being transferred to -80°C. Total DNA was extracted from murine faecal pellets and caecum content using the QIamp Fast DNA Stool (Qiagen, Manchester, UK) kit. Murine pellets were weighed before performing the extraction using the QIamp Fast DNA Stool (Qiagen, Manchester, UK) extraction kit protocol. The samples were placed in sterile tubes containing 0.1 mm, 0.5 mm and 1.0 mm zirconia/glass beads (Thistle Scientific, UK). Eight hundred ml of InhibitEX buffer was added to the samples and then homogenised under one pulse of 1 min and a final of 30 sec using a Mini-Beadbeater (Biospec Products, USA). In the intervals of the homogenisation steps the samples were placed on ice for 1 min. The samples were then placed in a 95°C heat-block for 5 min. The subsequent steps of the DNA extraction were carried out as described in the Qiagen protocol using 15 µl of proteinase K with 200 µl of AL buffer, 200 µl of lysate and 200 µl of ethanol at the relevant steps.

Library preparation for 16S rRNA gene amplicon sequencing

The V3/V4 variable region of the 16S rRNA gene was targeted for Illumina MiSeq System sequencing (San Diego, California, USA). The region was amplified using the universal 16S ribosomal RNA gene primer pair S-D-Bact-0341-b-S-17 (5'-CCTACGGGNGGCWGCAG-3') / S-D-Bact-0785-a-A-21 (5'-GACTACHVGGGTATC TAATC C-3') 5' ³⁰. The Phusion High-Fidelity PCR Master Mix (ThermoFisher Scientific, Waltham, MA, USA) was used for the amplification. The PCR products were purified and the subsequent library preparation steps were performed according to the Illumina MiSeq system protocol. Dual-index barcodes were attached to the amplicon (Nextera XT V.2 Index Kits sets A and D, Illumina) and then purified using the Agencourt AMPure XP-PCR Purification system (Beckman Coulter, Inc., California, USA). The indexed amplicons were quantified using the Qubit dsDNA HS Assay Kit (Thermo Fischer Scientific, MA, U.S.A.). The pooled library containing the same concentration of each amplicon was sequenced (2 x 250 bp) on a MiSeq Illumina platform in the Teagasc Food Research Centre sequencing facility (Fermoy, Ireland).

Microbiota composition analysis

FLASH programme ³¹ was used to join the paired-end reads that were subsequently quality filtered in Qiime (v.1.9.1) using the split_libraries_fastq.py script ³². Forward primer was removed using cutadapt ³³ and reverse primer with QIIME's script truncate_reverse_primer.py. The USEARCH sequence analysis tool was used for further quality filtering and *de novo* operational taxonomic unit (OTU) clustering. The sequences were filtered by length, sorted by size and single unique sequences were removed. The remaining sequences were clustered into OTUs based on 97% identity. Subsequently, UCHIME with the GOLD reference database was used for chimera removal. The initial quality filtered sequences were mapped against the OTUs (97 % identity). OTU representative sequences were taxonomically classified (classify.seqs) from phylum down to genus level with a 80% confidence threshold by using the mothur suite of tools (v1.36.1) and

the RDP database (trainset 14) [34]. SPINGO was used in order to classify the OTUs at species level ³⁵. Alpha and beta diversity analysis was performed in QIIME on a rarefied OTU table at 1300 reads in order to include the majority of the samples collected throughout the study. The sequences were aligned using the PyNast tool ³⁶ in Qiime in order to generate alpha (α) diversity indices i.e., Shannon, Simpson, PD whole tree, Chao1 and Observed Species, and beta (β) diversity indices i.e. Weighted UniFrac and Unweighted UniFrac.

Statistical analysis

The statistical software R (3.2.5; R Core Team) was used for statistical analysis and visualisation of data. The reads per sample were converted into relative abundances and ggplot2 (v. 2.2.1) was used for the visualisation of the bacterial composition of the microbiota. Principal coordinates analysis (PCoA) was performed with Weighted and Unweighted UniFrac distances with the made4 package (v. 1.50.1). Analysis of similarities (ANOSIM) was used (vegan 2.4-3) for the detection of significant differences in the observed clustering between the groups. A Kruskal-Wallis test followed by a Dunn's post hoc test (dunn.test 1.3.4) with Benjamini-Hochberg p value adjustment for multiple testing was applied in order to detect significant differences in the faecal microbiota diversity and taxa relative abundances due to diet at T4 and T5. Values were considered significant when Kruskal-Wallis $p < 0.5$, Kruskal-Wallis $p_{adj} < 0.1$ and Dunn's post hoc test $p_{adj} < 0.1$. Significance indications: "●" $p_{adj} < 0.1$; "*" $p_{adj} < 0.05$; "***" $p_{adj} < 0.01$; "****" $p_{adj} < 0.001$; "*****" $p_{adj} < 0.0001$. Graph Pad Prism (v 5.03) was used to visualise alpha diversity scores and animal body weight and to perform a Kruskal-Wallis test on body weight values at the end of the trial.

Results and Discussion

Effects of treatment on animal body weight

Due to initial aversion of the animals to the Abx water and subsequent abstinence from chow during the first 2 weeks of the Abx treatment, some animal body weight loss was observed (**Figure S2**) and 8 animals were euthanised. The animals began to recover their body weight after the 2nd week of Abx treatment (**Figure S2**). At the end of the trial (T5), consumption of the milk supplemented diets resulted in increased body weight compared to baseline but not significantly different to the other diets effect (**Figure S2**). Upon dissection all animals had normal spleen weight (data not shown). The number of animals in each dietary group at T5 for which the gut microbiota was analysed is shown in **Table S1**.

Characterisation of human donor microbiota and its engraftment in mice

General features of the compositional and phylogenetic diversity differences between the faecal microbiota of frail long-term care unit-residing elderly subjects and healthy community-living were described in the ElderMet study ³⁷. The ElderMet subjects EM297 (LS) and EM425 (COM) were re-enrolled and their faecal microbiota was re-analysed in order to confirm suitability for the mouse trial. Compositional differences at taxon level between COM and LS type faecal microbiota have been described before ²⁵. Compared to the COM type microbiota, the LS microbiota had lower Firmicutes relative abundance, and it was enriched in Proteobacteria, Synergistetes and Bacteroidetes (**Figure 1**), but as expected for a subject of this type, the alpha diversity was lower (**Table S2**). Thus, the inocula were considered suitable for proceeding with mouse humanisation.

One week after humanisation (T3), the murine faecal microbiota had diverged from baseline and did not revert back to the original murine phylogenetic profile by the completion of the trial (T5), as demonstrated by analysis of the Weighted and Unweighted UniFrac distances at time-point 3 (T3; **Figure S3**) and T5 (**Figure 2**). Importantly, the COM or LS type phylogenetic separation in β -diversity was retained at T5 as shown by PCoA of the

Unweighted UniFrac distances (**Figure 2B**) in spite of the effect of diet that drove the abundance-based clustering (**Figure 2A**). The faecal microbiota from mice fed the wmlk and the lac-free diets clustered closer together compared to the mice fed the control or GMP diets irrespective of humanisation microbiota type (**Figure 2A**). A detailed description of the differential effect of the diet supplements on the faecal microbiota follows further below. Importantly, the murine host may have played a role in re-shaping the relative abundances of taxa in the xenomicrobiota (i.e. the human microbiota, LS or COM type). It has been reported that the gut of germ-free or antibiotic-treated mice may be refractory to *Clostridium* clusters XIVa and IV taxa colonisation but more susceptible to Bacteroidetes colonisation³⁸. The Bacteroidetes and especially Porphyromonadaceae enrichment across most dietary groups at T5 and the compositional profile of the Lachnospiraceae at T5 comprised of mostly two taxa groups i.e. unclassified *Clostridium* cluster XIVa and unclassified Lachnospiraceae (**Figure S5**), can be viewed as the result of selective pressure applied by the murine host to reshape the relative abundances of the shared with the human xenomicrobiota taxa to best fit murine host physiology and immunity^{39, 40}. *Barnesiella intestinihominis* and *Parabacteroides goldsteinii* were the most abundant Porphyromonadaceae taxa in the microbiota of COM type humanised mouse whereas *Parabacteroides merdae* was the most abundant Porphyromonadaceae in LS type humanised mouse (**Figure S5**).

The lactose-free milk supplemented diet sustained higher microbiota diversity

In mice colonised with either human microbiota type, the lac-free diet sustained significantly higher faecal microbiota diversity compared to the wmlk or GMP diets, performing as efficiently as the soy protein supplemented diet (control) (**Figure 3A and B**). For example, for COM type humanised mouse microbiota at T5, feeding with lac-free diet resulted in ca. 4.5 Shannon diversity index which was significantly higher compared to the other three diets, whereas the effect of the lac-free diet feeding on indices PD (ca. 8) and Observed Species (ca.

90) was similar to that of the control diet (**Figure 3A**). A similar trend was observed for LS type humanised mouse microbiota; upon the lac-free diet the Shannon diversity index was ca. 4.1, the PD was ca. 5.5 and the Observed Species index was 58 (**Figure 3B**). Some animal and human studies indicate the prebiotic potential of soy and its products⁴¹. Decreased gut microbiota diversity is a biomarker of ageing and frailty, and dietary supplements contributing to sustaining diversity may slow down ageing-related health loss^{37, 42}. The efficiency of the lac-free milk diet in retaining microbiota diversity can be attributed partly to the BMOs available for microbiota fermentation. Bovine milk oligosaccharides have attracted interest as potential prebiotics due to their structural complexity and similarity to HMOs^{43, 44}. The effect of the diets on the diversity of the faecal microbiota of female and male mice analysed separately was similar to that presented here for the aggregated microbiota across gender (data not shown). The alpha diversity indices Chao 1 and Simpson were also analysed and indicated similar significant results as those presented in **Figure 3** (data not shown).

Differential effect of dietary supplementation on microbiota composition

Diet had a significant effect on the composition of the murine faecal microbiota as shown in **Figure 2**. In order to profile the detailed effect of the diets on the faecal microbiota of mice at T5, PCoAs of the UniFrac distances for the faecal microbiota per humanisation microbiota type were generated, i.e. COM type humanisation (**Figure 4 A and B**) and LS type humanisation (**Figure 4 C and D**). The ANOSIM test result confirmed the significant effect of the diets on the β -diversity (**Figures 4**). Within the diet-based clustering, the mouse gender had a significant impact on further sub-clustering of the microbiota in most of the dietary groups (gender is denoted by shape in **Figures 4**). The ANOSIM test result showed that the effect of gender was significant ($p < 0.05$) for most of the dietary groups and that the variation observed was due to the gender effect ($R \approx 1$) (**Table 2**). Gender did not have a significant

effect on the GMP and lac-free dietary groups for LS colonised mice (Weighted and Unweighted, respectively) (**Table 2; Figure 4 C and D**).

Prebiotic potential of the lactose-free milk

The murine faecal microbiota (either colonisation type) was enriched in Ruminococcaceae upon receiving the lac-free diet (17% and 4.6% average relative abundance, COM and LS type colonisation, respectively) (**Figure 5, S4**). The responsiveness of the Ruminococcaceae in the female mice was comparable to control diet ($p>0.1$) (**Figure 6A and 7A**), whereas in the male mice the Ruminococcaceae responsiveness to the lac-free diet was either comparable to wmlk diet ($p>0.1$) (COM type colonisation: **Figure 6B**) or significantly higher compared to all diets ($p<0.09$) (LS type colonisation; **Figure 7B**). The Ruminococcaceae enrichment associated with the lac-free diet could indicate that such a dietary supplementation might contribute to colonic health and healthy ageing as suggested by previous studies for these taxa^{45, 46}. In the LS humanised mice the Ruminococcaceae reached high relative abundance after the lac-free diet but not as high as in COM type humanised mice, potentially due to low starting abundance of this family in the LS microbiota (**Figures 5 and 7**). In COM type colonised female mice and LS type colonised mice *Flavonifractor plautii* was a dominant Ruminococcaceae taxon whereas *Anaerotruncus colihominis* and *Oscillibacter* taxa were differentially abundant in female and male COM type colonised mice (**Figures 8 and S5**). Others have reported probiotic-associated *Anaerotruncus* enrichment of dysbiotic murine microbiota after Abx treatment⁴⁷ whereas *Oscillibacter* and *Flavonifractor* taxa have been associated with resistant starch degradation and identified in the gut microbiota of lean individuals^{48, 49}. However, caution is needed before extrapolating these findings to human dietary interventions because some studies

report enrichment of *Oscillibacter* in cancer associated microbiota and *Anaerotruncus* enrichment in the microbiota of frail elderly^{50, 51}.

Effect of the whole milk supplemented diet on the gut microbiota

The whole milk supplemented diet was less efficient than the lac-free diet in sustaining microbiota diversity based on the alpha diversity indices at T5 (**Figure 3**). For COM type humanised mouse microbiota, the wmilk diet resulted in ca. 4.1, 6.7 and 64 Shannon, PD and Observed Species indices, respectively, that were significantly lower compared to the respective indices upon the lac-free diet supplementation (**Figure 3A**). Similar results were observed for LS type colonised mouse microbiota (**Figure 3B**). However, the comparative increase in the relative abundance of certain taxa after the wmilk diet warrants further investigation. The Ruminococcaceae family were responsive to the wmilk diet only in the male mice (**Figures 5, 6 and 7**). *Ruminococcus bromii* (6% average relative abundance; detected only in the COM type colonised male mice) and other unclassified taxa were the dominant Ruminococcaceae representative taxa (**Figures 8C and S5C**). *Ruminococcus bromii* is a dominant gut microbiota taxon highly responsive to resistant starch, essential in nutrient degradation and as such it can be a prebiotic target leading to substrate release in the colon and increase in cross-feeding mediated by short chain fatty acids (SCFA) production^{52, 53}.

The significant effect of the wmilk diet on the relative abundance of the Erysipelotrichaceae (Firmicutes) is evident in **Figure S4**. In animals with COM type colonisation, the Erysipelotrichaceae were the most abundant Firmicutes family in the male mouse microbiota (23% average relative abundance) with *Allobaculum* as the most abundant representative taxon (22% average relative abundance) (not detected in the female mice) (**Figure 8C**), whereas *Turicibacter* was dominant in female mice (1.6% average relative abundance) (not

301 detected in male mice) (**Figure 8B**). In animals with LS type colonisation, low in abundance
302 Erysipelotrichaceae taxa were significantly enriched in the male mouse microbiota upon
303 wmlk diet feeding (0.6% average relative abundance) (**Figure 7B**). *Allobaculum stercorialis*
304 was the dominant Erysipelotrichaceae taxon in the female mouse microbiota (32% average
305 relative abundance) similarly to what was observed in the microbiota of COM type
306 humanised male mice that were fed the wmlk diet (**Figures 8B, S5B**). The
307 Erysipelotrichaceae enrichment of the faecal microbiota of the wmlk diet fed mice and the
308 responsiveness of *Allobaculum* taxa to the whole milk supplementation may also be a
309 positive effect on the microbiota. In murine trials, high-fat diet-associated dysbiosis and
310 impaired gut barrier function were restored after BMO supplementation correlating to
311 *Allobaculum* (and Ruminococcaceae) enrichment of the microbiota^{19; 54}. Inulin
312 supplementation had similar effect on *Allobaculum* taxa associated with improved vascular
313 function⁵⁵. However, experimental extrapolation to human diet is needed due to *Allobaculum*
314 association with thrombosis risk through trimethylamine-N-oxide (TMAO) metabolism and
315 to other erysipelotrichial associations with the obesity phenotype in murine models^{54, 55}.

316 Interestingly, the wmlk diet, similarly to the lac-free diet, sustained the highest
317 Lachnospiraceae relative abundance in the murine microbiota compared to the GMP and
318 control diets. Analytically, COM type humanisation: ca. 12% and 15%, upon wmlk and lac-
319 free diet, respectively; LS type humanisation: ca. 15.5% and 29% upon wmlk and lac-free
320 diet, respectively (**Figures 5 and S4**). Taxa belonging to unclassified Lachnospiraceae and
321 unclassified *Clostridium* cluster XIVa were the representative Lachnospiraceae taxa (**Figure**
322 **8**).

323 Here the Acidaminococcaceae taxon *Phascolarctobacterium faecium*, an asaccharolytic
324 bacterium, was responsive to the wmlk diet (5.6% average relative abundance in COM type

colonised mouse) and the GMP diet (5% and 6% average relative abundance in COM and LS type colonised mouse, respectively), and to a lesser extent to the control diet (1.8% and 1.1% average relative abundance in COM and LS type colonised mouse, respectively) (**Figures 8 and S5**). We have previously reported Acidaminococcaceae responsiveness to GMP supplementation in an *in vitro* colon system ²⁵. The Acidaminococcaceae family relative abundance was significantly increased upon wmlk diet mainly when compared to the control and lac-free diet in COM type humanised mice (**Figures 6**). In LS colonised mice, the relative abundance of the Acidaminococcaceae increased significantly upon GMP diet in the female mice (**Figure 7A**); the Acidaminococcaceae family was also abundant in the male mouse microbiota (**Figure 5B and C; Figure S4B**). In previous mouse model studies, the taxon *Phascolarctobacterium* was linked to soy diet-associated metabolic markers improvement ⁵⁸, whereas Lecomte *et al.* reported that the taxon increased after high-fat diet ⁵⁹. Tran *et al.* ⁶⁰ reported the positive association of the taxon to high fibre diet in both murine and human intervention studies.

Desulfovibrionaceae responsiveness to the dairy diets

All three milk-associated diets presented in this study had some effect on the relative abundance of the Desulfovibrionaceae family of the Proteobacteria. In animals that received COM type colonisation, the GMP diet had the strongest effect on the Desulfovibrionaceae taxon *Bilophila wadsworthia* that reached 11.5% average relative abundance upon the GMP diet (**Figure S5**). The control and wmlk diet had the weakest effect on the Desulfovibrionaceae/*Bilophila* taxa responsiveness in the female (4% average relative abundance) and male mice (1.8% average relative abundance), respectively (**Figures 6 and 8**). In LS type colonised mice, the taxon *Bilophila* was not detected and *Desulfovibrio* was the Desulfovibrionaceae representative, responsive to the lac-free diet, reaching an average 8% relative abundance (**Figures 5, 8**). Contrary to what we report in the current study, Sawin

350 *et al.* ²⁸ reported that over a month of 20% GMP supplemented diet resulted in reduced
351 Desulfovibrionaceae/*Desulfovibrio* relative abundance and amelioration of inflammatory
352 markers in wild type (WT) and phenylketonuria (PKU) C57Bl/6 mice and compared to casein
353 and amino acids supplemented diets, respectively.

354 *Desulfovibrio* spp. is a taxon negatively correlated to health in elderly people ⁵⁰. Previous
355 studies have reported that intake of milk-derived high-fat diet resulted in *Bilophila*
356 *wadsworthia* associated dysbiosis and inflammation in mouse models ^{61, 62}. Recently,
357 Natividad *et al* ⁶³ using a mouse model observed that milk fat promoted *B. wadsworthia*,
358 inflammation and bile acid metabolism deregulation. Importantly, colonic health was
359 promoted by reduced *Bilophila* relative abundances in human gut microbiota ⁶⁴. Importantly,
360 although some mouse model studies show that milk fat may be associated with negative
361 health outcomes associated with increased *Bilophila* abundance, other studies support non-
362 deleterious effects of dairy fat on health ^{73, 74, 75}. The balance of the potentially positive and
363 negative effects of milk diets on the faecal microbiota needs to be measured and considered
364 in human studies.

365 **Bacteroidetes responsiveness to the dairy diets**

366 Phylum Bacteroidetes encompasses many fibrolytic and saccharolytic taxa ⁶⁸. The
367 Bacteroidetes were more responsive to the dairy diets than to control. In COM type colonised
368 mice, Bacteroidetes had $\geq 55\%$ average relative abundance upon the dairy diets and 40% upon
369 control diet, whereas for LS type colonised mice, the Bacteroidetes had $\geq 42\%$ average
370 relative abundance upon the dairy diets and 32% upon control diet (**Figure 5**). In the female
371 mice with COM type humanisation, the families Porphyromonadaceae, Rikenellaceae and
372 other unclassified Bacteroidales were highly responsive to the GMP diet (**Figure 6A**),
373 whereas for either gender the Bacteroidaceae (mainly *Bacteroides thetaiotaomicron* and
374 *Bacteroides uniformis*) were significantly increased upon the wmlk diet (**Figure 6, S5B**).

The Porphyromonadaceae responsiveness upon the GMP diet was high also in the male mice with COM type humanisation, whereas the Bacteroidaceae were least responsive to the GMP diet compared to the other diets (**Figure 6B**). In the LS type colonised mice, the families Bacteroidaceae, Porphyromonadaceae and Rikenellaceae were responsive to the dairy diets, whereas the Rikenellaceae were also highly responsive to the control diet (the latter response was also observed in the male animals with COM type humanisation) (**Figures S4B, 6B**). Increase in *Bacteroides* population due to GMP-supplementation has been reported in mouse model studies ^{27, 28}. Milk and subsequently GMP contain highly sialylated carbohydrate components ¹⁰. Interestingly, different Bacteroidaceae taxa have been associated with sialidase activity with or without sialic acid utilisation ability ^{69, 70}. In the current study, compared to the other diets, feeding with the GMP diet resulted in Enterobacteriaceae enrichment in the faecal microbiota of COM type colonised mice (4% average relative abundance) (**Figures 5 and S4A**); *Escherichia/Shigella* taxa were the dominant Enterobacteriaceae representatives (**Figure 8**). The sialic acid content of the GMP may have contributed to the reported comparative increase in Enterobacteriaceae relative abundance ⁶⁹. Charbonneau *et al.* ⁷¹ reported growth improvement in a gnotobiotic mouse and piglet models through cross-feeding between *Bacteroides* taxa that could degrade sialic acid-containing carbohydrates from sialylated BMO and secondary utilisers of sialic acid-containing compounds like *E.coli*. Importantly, a similar response upon GMP diet was not observed in LS humanised mice; the Enterobacteriaceae were more responsive to the lac-free diet (1.7% average relative abundance) (**Figures 5 and 8**). These findings in combination to existing literature may highlight the need for caution when designing the dose for human trials using milk because the sialylated carbohydrate component may induce the growth of pathobionts of the *E.coli/Shigella* group ⁷¹.

Enrichment of the microbiota in Verrucomicrobiaceae in some dietary groups

400 Enrichment of the Verrucomicrobiaceae family (*Akkermansia*) ($\geq 38\%$ average relative
401 abundance) was observed at T5 upon the control diet in the faecal microbiota of COM type
402 humanised mouse and in LS type humanised mouse microbiota upon the GMP and control
403 diets (**Figures 5 and 8**). The values of the Shannon alpha diversity index (that measure
404 evenness ⁷²) at T5 may have been biased due to the high relative abundance of *Akkermansia*
405 (i.e. decreased alpha diversity values in the dietary groups where the *Akkermansia* bloom was
406 observed) and for this reason the alpha diversity was analysed at an earlier time-point, i.e. in
407 the middle of the dietary intervention (T4) (**Figure S6**) in order to confirm the effect of the
408 lac-free diet on sustaining the microbiota diversity. The Verrucomicrobiaceae bloom was
409 detected at T4 only for mice humanised with COM type microbiota but not in LS type
410 humanised mice. For either colonisation type at T4, both milk supplemented diets (i.e. the
411 lac-free and the wmlk diets) sustained significantly higher alpha diversity indices (Shannon,
412 PD and Observed Species) compared to both GMP and control diet (**Figure S7**). The
413 *Akkermansia* bloom could be attributed to the interruption of the Abx treatment due to
414 HydroGel treatment in some dietary groups, to coprophagy and to cross-contamination
415 between cages. Interestingly, Dubourg *et al.* ⁷³ reported increased *A. muciniphila* colonisation
416 in the gut microbiota of patients after broad-spectrum Abx treatment.

417 **Translatability to human nutrition**

418 Milk is a widely accessible dietary product, it combines an array of valuable nutrients and
419 many recent human cohort studies have demonstrated the health benefits of dairy
420 consumption including milk ⁷⁴⁻⁷⁶. Importantly, BMOs may be an efficient prebiotic substrate
421 similar to HMOs ^{43, 44}. We have investigated the gut microbiota modulatory potential of
422 (lactose-free and whole) milk in a murine model. A lactose-free milk supplemented diet was
423 found a better prebiotic candidate compared to whole milk supplemented diet as it retained
424 higher microbiota diversity. However, the whole milk diet affected the abundance of

significant health-relevant microbiota taxa. Importantly, in LS type colonised mice, the w milk diet sustained a higher Firmicutes relative abundance compared to the Bacteroidetes relative abundance (**Figure 5**) which may indicate the potential of milk to improve the Firmicutes dominance over the Bacteroidetes in the microbiota of frail elderly ⁵⁰. When translating the findings of the current study to humans, the fact that mice may not fully tolerate lactose ⁷⁷, and that the lactose fermentation in w milk diet fed mice may have prevented BMOs utilisation by the gut microbiota, must be taken into consideration. Importantly, we did not observe any adverse outcomes in mice consuming the w milk diet such as diarrhoea or abstinence from eating; furthermore the stool consistency of mice fed the w milk diet (by macroscopic observation before DNA extraction) was similar to that of mice receiving the other diets. Whether or not the two milk types, i.e., whole milk and lactose-free milk, have a differential effect on the gut microbiota of lactase persistent subjects remains to be clarified in a clinical trial. Interestingly, dairy companies employ different preparation techniques for the manufacture of lactose-free milk. Whole milk can either be filtered in order for lactose to be retained and the remaining lactose is hydrolysed enzymatically (e.g. Valio Ltd. product used here) or whole milk is enzymatically treated without prior filtering. The two methods may result in lactose-free milk with differing residual content of simple sugars that, upon consumption, may result in differential effect on the gut microbiota. Future *in vivo* studies can clarify the potential differential effect on the gut microbiota of the two preparations. Importantly, in this study, similarly to previous observations, we observed gender dependent differential responses to diet partly attributed to hormonal effects ^{78, 79}. Addressing gender-dependant responsiveness of the faecal microbiota to diet in murine models could increase translatability of the findings to human nutrition.

448 **Acknowledgements:** We thank Arla Foods Ingredients, Denmark and Valio Ltd Finland for
449 providing Lacprodan CGMP-10 and milk powders, respectively.

450 **Funding Sources:** This work was financially supported by the Government of Ireland
451 National Development Plan by way of a Department of Agriculture, Food and the Marine
452 (DAFM) under a Food Institutional Research Measure (FIRM) award (11/F/053) for the
453 ELDERFOOD project.

454 **Supporting Information Description:** Table S1 Treatment groups indicating the diet,
455 gender and the colonisation with human faecal microbiota type, Table S2 Alpha diversity
456 indices of the faecal microbiotas of the human donors and the murine baseline faecal
457 microbiota, Figure S1 Timeline of the mouse trial, Figure S2 Animal body weight during the
458 11 weeks of the trial, Figure S3 Principal coordinates analysis (PCoA) of the UniFrac
459 distances one week after “humanisation” (T3), Figure S4 Differentially abundant taxa at
460 family level at the end of the trial (T5) for aggregated female and male mice faecal
461 microbiota, Figure S5 Faecal microbiota composition at species level at the end of the trial
462 (T5), Figure S6 Composition of the murine faecal microbiota at phylum level at the mid part
463 of the dietary intervention (T4), Figure S7 Alpha diversity indices at the mid part of the
464 dietary intervention (T4). **Accession numbers:** Sequence Read Archive (SRA) (NCBI) under
465 accession no. PRJNA504771.

466 **Conflict of interest:** The authors declare no conflict of interest.

References

1. Marchesi, J. R.; Adams, D. H.; Fava, F.; Hermes, G. D. A.; Hirschfield, G. M.; Hold, G.; Quraishi, M. N.; Kinross, J.; Smidt, H.; Tuohy, K. M.; Thomas, L. V.; Zoetendal, E. G.; Hart, A. The gut microbiota and host health: a new clinical frontier. *Gut* **2016**, *65*, 330–9.
2. Sonnenburg, J. L.; Bäckhed, F. Diet-microbiota interactions as moderators of human metabolism. *Nature* **2016**.
3. Kurilshikov, A.; Wijmenga, C.; Fu, J.; Zhernakova, A. Host genetics and gut microbiome: Challenges and perspectives. *Trends Immunol.* **2017**, *38*, 633–647.
4. Rothschild, D.; Weissbrod, O.; Barkan, E.; Korem, T.; Zeevi, D.; Costea, P. I.; Godneva, A.; Kalka, I. N.; Bar, N.; Zmora, N.; Pevsner-Fischer, M.; Israeli, D.; Kosower, N.; Malka, G.; Wolf, B. C.; Avnit-Sagi, T.; Lotan-Pompan, M.; Weinberger, A.; Halpern, Z.; Carmi, S.; Elinav, E.; Segal, E. Environmental factors dominate over host genetics in shaping human gut microbiota composition. *Nature* **2018**, *555*, 210–215.
5. Franceschi, C.; Campisi, J. Chronic inflammation (Inflammaging) and its potential contribution to age-associated diseases. *Journals Gerontol. - Ser. A Biol. Sci. Med. Sci.* **2014**, *69*, S4–S9.
6. Li, H.; Qi, Y.; Jasper, H. Preventing age-related decline of gut compartmentalization limits microbiota dysbiosis and extends lifespan. *Cell Host Microbe* **2016**, *19*, 240–253.
7. Thevaranjan, N.; Puchta, A.; Schulz, C.; Naidoo, A.; Szamosi, J. C.; Verschoor, C. P.; Loukov, D.; Schenck, L. P.; Jury, J.; Foley, K. P.; Schertzer, J. D.; Larché, M. J.; Davidson, D. J.; Verdú, E. F.; Surette, M. G.; Bowdish, D. M. E. Age-associated microbial dysbiosis promotes intestinal permeability, systemic inflammation, and macrophage dysfunction. *Cell Host Microbe* **2017**, *21*, 455–466.e4.

- 493 8. Zeng, M. Y.; Inohara, N.; Nuñez, G. Mechanisms of inflammation-driven bacterial
494 dysbiosis in the gut. *Mucosal Immunol.* **2016**, *10*, 1–9.
- 495 9. Rozenberg, S.; Body, J. J.; Bruyère, O.; Bergmann, P.; Brandi, M. L.; Cooper, C.;
496 Devogelaer, J. P.; Gielen, E.; Goemaere, S.; Kaufman, J. M.; Rizzoli, R.; Reginster, J.
497 Y. Effects of dairy products consumption on health: Benefits and beliefs—A
498 commentary from the Belgian Bone Club and the European Society for Clinical and
499 Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases.
500 *Calcif. Tissue Int.* **2016**, *98*, 1–17.
- 501 10. O’Riordan, N.; Kane, M.; Joshi, L.; Hickey, R. M. Structural and functional
502 characteristics of bovine milk protein glycosylation. *Glycobiology* **2014**, *24*, 220–236.
- 503 11. Song, M.; Garrett, W. S.; Chan, A. T. Nutrients, foods, and colorectal cancer
504 prevention. *Gastroenterology* **2015**, *148*, 1244–1260.
- 505 12. Larsson, S. C.; Crippa, A.; Orsini, N.; Wolk, A.; Michaëlsson, K. Milk consumption
506 and mortality from all causes, cardiovascular disease, and cancer: A systematic
507 review and meta-analysis. *Nutrients* **2015**, *7*.
- 508 13. Sahni, S.; Mangano, K. M.; Tucker, K. L.; Douglas, P. K.; Casey, V. A.; Hannan, M.
509 T. Protective association of milk intake on the risk of hip fracture: Results from the
510 Framingham Original Cohort. *J Bone Min. Res.* **2015**, *29*, 37–54.
- 511 14. Lana, A.; Rodriguez-Artalejo, F.; Lopez-Garcia, E. Dairy consumption and risk of
512 frailty in older adults: A prospective cohort study. *J. Am. Geriatr. Soc.* **2015**, *63*,
513 1852–1860.
- 514 15. Power, S. E.; Jeffery, I. B.; Ross, R. P.; Stanton, C.; O’Toole, P. W.; O’Connor, E.
515 M.; Fitzgerald, G. F. Food and nutrient intake of Irish community-dwelling elderly
516 subjects: who is at nutritional risk? *J Nutr Heal. Aging* **2014**, *18*, 561–572.

16. Laird, E.; Casey, M. C.; Ward, M.; Hoey, L.; Hughes, C. F.; Mccarroll, K.; Cunningham, C.; Strain, J. J.; McNulty, H.; Molloy, A. M. Dairy intakes in older Irish adults and effects on vitamin micronutrient status: Data from the TUDA study. *J. Nutr. Heal. Aging* **2016**, 1–8.
17. Charbonneau, M. R.; O'Donnell, D.; Blanton, L. V.; Totten, S. M.; Davis, J. C. C.; Barratt, M. J.; Cheng, J.; Guruge, J.; Talcott, M.; Bain, J. R.; Muehlbauer, M. J.; Ilkayeva, O.; Wu, C.; Struckmeyer, T.; Barile, D.; Mangani, C.; Jorgensen, J.; Fan, Y. M.; Maleta, K.; Dewey, K. G.; Ashorn, P.; Newgard, C. B.; Lebrilla, C.; Mills, D. A.; Gordon, J. I. Sialylated milk oligosaccharides promote microbiota-dependent growth in models of infant undernutrition. *Cell* **2016**, 164, 859–871.
18. Karav, S.; Le Parc, A.; Leite Nobrega de Moura Bell, J. M.; Frese, S. A.; Kirmiz, N.; Block, D. E.; Barile, D.; Mills, D. A. Oligosaccharides released from milk glycoproteins are selective growth substrates for infant-associated bifidobacteria. *Appl. Environ. Microbiol.* **2016**, 82, 3622–3630.
19. Boudry, G.; Hamilton, M. K.; Chichlowski, M.; Wickramasinghe, S.; Barile, D.; Kalanetra, K. M.; Mills, D. A.; Raybould, H. E. Bovine milk oligosaccharides decrease gut permeability and improve inflammation and microbial dysbiosis in diet-induced obese mice. *J. Dairy Sci.* **2017**, 100, 2471–2481.
20. Neelima, Sharma, R., Rajput, Y.S., and Mann, B. Chemical and functional properties of glycomacropeptide (GMP) and its role in the detection of cheese whey adulteration in milk: a review. *Dairy Sci Technol* **2013**, 93, 21–43.
21. van Calcar, S. C.; Ney, D. M. Food products made with glycomacropeptide, a low-phenylalanine whey protein, provide a new alternative to amino acid-based medical foods for nutrition management of phenylketonuria. *J. Acad. Nutr. Diet.* **2012**, 112.

22. Hvas, C. L.; Dige, A.; Bendix, M.; Wernlund, P. G.; Christensen, L. a; Dahlerup, J. F.; Agnholt, J. Casein glycomacropeptide for active distal ulcerative colitis: a randomised pilot study. *Eur. J. Clin. Invest.* **2016**, *46*, 555–563.
23. Hernandez-Hernandez, O.; Sanz, M. L.; Kolida, S.; Rastall, R. a; Moreno, F. J. In vitro fermentation by human gut bacteria of proteolytically digested caseinomacropeptide nonenzymatically glycosylated with prebiotic carbohydrates. *J. Agric. Food Chem.* **2011**, *59*, 11949–55.
24. Robitaille, G. Growth-promoting effects of caseinomacropeptide from cow and goat milk on probiotics. *J. Dairy Res.* **2013**, *80*, 58–63.
25. Ntemiri, A.; Chonchúir, F. N.; O’Callaghan, T. F.; Stanton, C.; Ross, R. P.; O’Toole, P. W. Glycomacropeptide sustains microbiota diversity and promotes specific taxa in an artificial colon model of elderly gut microbiota. *J. Agric. Food Chem.* **2017**, *65*.
26. Gustavo Hermes, R.; Molist, F.; Francisco Pérez, J.; Gómez de Segura, A.; Ywazaki, M.; Davin, R.; Nofrías, M.; Korhonen, T. K.; Virkola, R.; Martín-Orúe, S. M. Casein glycomacropeptide in the diet may reduce *Escherichia coli* attachment to the intestinal mucosa and increase the intestinal lactobacilli of early weaned piglets after an enterotoxigenic *E. coli* K88 challenge. *Br. J. Nutr.* **2013**, *109*, 1001–12.
27. Jiménez, M.; Cervantes-García, D.; Muñoz, Y. H.; García, A.; Haro, L. M.; Salinas, E. Novel mechanisms underlying the therapeutic effect of glycomacropeptide on allergy: Change in gut microbiota, upregulation of TGF- β , and inhibition of mast cells. *Int. Arch. Allergy Immunol.* **2017**, *171*, 217–226.
28. Sawin, E. a; De Wolfe, T. J.; Aktas, B.; Stroup, B. M.; Murali, S. G.; Steele, J. L.; Ney, D. M. Glycomacropeptide is a prebiotic that reduces *Desulfovibrio* bacteria, increases cecal short-chain fatty acids, and is anti-inflammatory in mice. *Am. J. Physiol. Liver Physiol.* **2015**, *309*, G590–G601.

29. Bereswill, S.; Fischer, A.; Plickert, R.; Haag, L.-M.; Otto, B.; Kühl, A. a.; Dashti, J. I.; Zautner, A. E.; Muñoz, M.; Loddenkemper, C.; Groß, U.; Göbel, U. B.; Heimesaat, M. M. Novel murine infection models provide deep insights into the “Ménage à Trois” of *Campylobacter jejuni*, microbiota and host innate immunity. *PLoS One* **2011**, *6*.
30. Klindworth, A.; Pruesse, E.; Schweer, T.; Peplies, J.; Quast, C.; Horn, M.; Glöckner, F. O. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Res.* **2012**, *41*, 1–11.
31. Magoc, T.; Salzberg, S. L. FLASH: Fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* **2011**, *27*, 2957–2963.
32. Caporaso JG, Kuczynski J, Stombaugh J, et al (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7:335–336.
33. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **2011**, *17*, 10.
34. Schloss, P. D.; Westcott, S. L.; Ryabin, T.; Hall, J. R.; Hartmann, M.; Hollister, E. B.; Lesniewski, R. A.; Oakley, B. B.; Parks, D. H.; Robinson, C. J.; Sahl, J. W.; Stres, B.; Thallinger, G. G.; Van Horn, D. J.; Weber, C. F. Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* **2009**, *75*, 7537–7541.
35. Allard, G.; Ryan, F. J.; Jeffery, I. B.; Claesson, M. J. SPINGO: A rapid species-classifier for microbial amplicon sequences. *BMC Bioinformatics* **2015**, *16*, 1–8.
36. Caporaso, J. G.; Bittinger, K.; Bushman, F. D.; Desantis, T. Z.; Andersen, G. L.; Knight, R. PyNAST: A flexible tool for aligning sequences to a template alignment. *Bioinformatics* **2010**, *26*, 266–267.

37. Claesson, M. J.; Jeffery, I. B.; Conde, S.; Power, S. E.; O'Connor, E. M.; Cusack, S.; Harris, H. M. B.; Coakley, M.; Lakshminarayanan, B.; O'Sullivan, O.; Fitzgerald, G. F.; Deane, J.; O'Connor, M.; Harnedy, N.; O'Connor, K.; O'Mahony, D.; van Sinderen, D.; Wallace, M.; Brennan, L.; Stanton, C.; Marchesi, J. R.; Fitzgerald, A. P.; Shanahan, F.; Hill, C.; Ross, R. P.; O'Toole, P. W. Gut microbiota composition correlates with diet and health in the elderly. *Nature* **2012**, *488*, 178–84.
38. Lundberg, R.; Toft, M. F.; August, B.; Hansen, A. K.; Hansen, C. H. F. Antibiotic-treated versus germ-free rodents for microbiota transplantation studies. *Gut Microbes* **2016**, *7*, 68–74.
39. Nguyen, T. L. A.; Vieira-Silva, S.; Liston, A.; Raes, J. How informative is the mouse for human gut microbiota research? *Dis. Model. Mech.* **2015**, *8*, 1–16.
40. Messer, J. S.; Liechty, E. R.; Vogel, O. A.; Chang, E. B. Evolutionary and ecological forces that shape the bacterial communities of the human gut. *Mucosal Immunol.* **2017**, *10*, 567–579.
41. Huang, H.; Krishnan, H. B.; Pham, Q.; Yu, L. L.; Wang, T. T. Y. Soy and gut microbiota: Interaction and implication for human health. *J. Agric. Food Chem.* **2016**, *64*, 8695–8709.
42. Jackson, M.; Jeffery, I. B.; Beaumont, M.; Bell, J. T.; Clark, A. G.; Ley, R. E.; O'Toole, P. W.; Spector, T. D.; Steves, C. J. Signatures of early frailty in the gut microbiota. *Genome Med.* **2016**, *8*, 8.
43. Zivkovic, A. M.; Barile, D. Bovine milk as a source of functional oligosaccharides for improving human health. *Adv. Nutr.* **2011**, *2*, 284–9.
44. Smilowitz, J. T.; Lebrilla, C. B.; Mills, D. A.; German, J. B.; Freeman, S. L. Breast milk oligosaccharides: Structure-function relationships in the neonate. *Annu. Rev. Nutr.* **2014**, *34*, 143–169.

45. Kong, F.; Hua, Y.; Zeng, B.; Ning, R.; Li, Y.; Zhao, J. Gut microbiota signatures of longevity. *Curr. Biol.* **2016**, *26*, R832–R833.
46. Mottawea, W.; Chiang, C.-K.; Mühlbauer, M.; Starr, A. E.; Butcher, J.; Abujamel, T.; Deeke, S. A.; Brandel, A.; Zhou, H.; Shokralla, S.; Hajibabaei, M.; Singleton, R.; Benchimol, E. I.; Jobin, C.; Mack, D. R.; Figeys, D.; Stintzi, A. Altered intestinal microbiota-host mitochondria crosstalk in new onset Crohn's disease. *Nat. Commun.* **2016**, *7*, 13419.
47. Grazul, H.; Kanda, L. L.; Gondek, D. Impact of probiotic supplements on microbiome diversity following antibiotic treatment of mice. *Gut Microbes* **2016**, *7*, 101–114.
48. Walker, A. W.; Ince, J.; Duncan, S. H.; Webster, L. M.; Holtrop, G.; Ze, X.; Brown, D.; Stares, M. D.; Scott, P.; Bergerat, A.; Louis, P.; McIntosh, F.; Johnstone, A. M.; Lobley, G. E.; Parkhill, J.; Flint, H. J. Dominant and diet-responsive groups of bacteria within the human colonic microbiota. *ISME J.* **2011**, *5*, 220–30.
49. Kasai, C.; Sugimoto, K.; Moritani, I.; Tanaka, J.; Oya, Y.; Inoue, H.; Tameda, M.; Shiraki, K.; Ito, M.; Takei, Y.; Takase, K. Comparison of the gut microbiota composition between obese and non-obese individuals in a Japanese population, as analyzed by terminal restriction fragment length polymorphism and next-generation sequencing. *BMC Gastroenterol.* **2015**, *15*, 1–10.
50. Jeffery, I. B.; Lynch, D. B.; O'Toole, P. W. Composition and temporal stability of the gut microbiota in older persons. *ISME J.* **2016**, *10*, 170–182.
51. Flemer, B.; Lynch, D. B.; Brown, J. M. R.; Jeffery, I. B.; Ryan, F. J.; Claesson, M. J.; O'Riordain, M.; Shanahan, F.; O'Toole, P. W. Tumour-associated and non-tumour-associated microbiota in colorectal cancer. *Gut* **2017**, *66*, 633–643.

52. Ze, X.; Duncan, S. H.; Louis, P.; Flint, H. J. *Ruminococcus bromii* is a keystone species for the degradation of resistant starch in the human colon. *ISME J.* **2012**, *6*, 1535–1543.
53. Scott, K. P.; Antoine, J.-M.; Midtvedt, T.; van Hemert, S. Manipulating the gut microbiota to maintain health and treat disease. *Microb. Ecol. Health Dis.* **2015**, *26*, 25877.
54. Hamilton, M. K.; Ronveaux, C. C.; Rust, B. M.; Newman, J. W.; Hawley, M.; Barile, D.; Mills, D. A.; Raybould, H. E. Prebiotic milk oligosaccharides prevent development of obese phenotype, impairment of gut permeability, and microbial dysbiosis in high fat-fed mice. *Am. J. Physiol. - Gastrointest. Liver Physiol.* **2017**, *312*, G474–G487.
55. Catry, E.; Bindels, L. B.; Tailleux, A.; Lestavel, S.; Neyrinck, A. M.; Goossens, J.-F.; Lobysheva, I.; Plovier, H.; Essaghir, A.; Demoulin, J.-B.; Bouzin, C.; Pachikian, B. D.; Cani, P. D.; Staels, B.; Dessy, C.; Delzenne, N. M. Targeting the gut microbiota with inulin-type fructans: preclinical demonstration of a novel approach in the management of endothelial dysfunction. *Gut* **2018**, *67*, 271–283.
56. Woting, A.; Pfeiffer, N.; Loh, G.; Klaus, S.; Blaut, M. *Clostridium ramosum* promotes high-fat diet-induced obesity in gnotobiotic mouse models. *MBio* **2014**, *5*, e01530-14-e01530-14.
57. Zhu, W.; Gregory, J. C.; Org, E.; Buffa, J. A.; Gupta, N.; Wang, Z.; Li, L.; Fu, X.; Wu, Y.; Mehrabian, M.; Sartor, R. B.; McIntyre, T. M.; Silverstein, R. L.; Tang, W. H. W.; DiDonato, J. A.; Brown, J. M.; Lusi, A. J.; Hazen, S. L. Gut microbial metabolite TMAO enhances platelet hyperreactivity and thrombosis risk. *Cell* **2016**, *165*, 111–124.

58. Cross, T.-W. L.; Zidon, T. M.; Welly, R. J.; Park, Y.-M.; Britton, S. L.; Koch, L. G.; Rottinghaus, G. E.; de Godoy, M. R. C.; Padilla, J.; Swanson, K. S.; Vieira-Potter, V. J. Soy improves cardiometabolic health and cecal microbiota in female low-fit rats. *Sci. Rep.* **2017**, *7*, 9261.
59. Lecomte, V.; Kaakoush, N. O.; Maloney, C. A.; Raipuria, M.; Huinao, K. D.; Mitchell, H. M.; Morris, M. J. Changes in gut microbiota in rats fed a high fat diet Correlate with obesity-associated metabolic parameters. *PLoS One* **2015**, *10*, e0126931.
60. Tam T.T. Tran, Fabien J. Cousin, Denise B. Lynch, Ravi Menon, Jennifer Brulc, Jillian R-M. Brown, Eileen O'Herlihy, Ludovica F. Butto, Katie Power, Ian B. Jeffery, Eibhlís M. O'Connor, Paul W. O'Toole (**Submitted**). Dietary modulation of the low diversity microbiota of frail older people: responsive species and innate immune effects.
61. Devkota, S.; Wang, Y.; Musch, M. W.; Leone, V.; Fehlner-Peach, H.; Nadimpalli, A.; Antonopoulos, D. A.; Jabri, B.; Chang, E. B. Dietary-fat-induced taurocholic acid promotes pathobiont expansion and colitis in *Il10^{-/-}* mice. *Nature* **2012**, *487*, 104–8.
62. Caesar, R.; Tremaroli, V.; Kovatcheva-Datchary, P.; Cani, P. D.; Bäckhed, F. Crosstalk between gut microbiota and dietary lipids aggravates WAT inflammation through TLR signaling. *Cell Metab.* **2015**, *22*, 658–668.
63. Natividad, J. M.; Lamas, B.; Pham, H. P.; Michel, M.-L.; Rainteau, D.; Bridonneau, C.; da Costa, G.; van Hylckama Vlieg, J.; Sovran, B.; Chamignon, C.; Planchais, J.; Richard, M. L.; Langella, P.; Veiga, P.; Sokol, H. *Bilophila wadsworthia* aggravates high fat diet induced metabolic dysfunctions in mice. *Nat. Commun.* **2018**, *9*, 2802.

64. Vandeputte, D.; Falony, G.; Vieira-Silva, S.; Wang, J.; Sailer, M.; Theis, S.; Verbeke, K.; Raes, J. Prebiotic inulin-type fructans induce specific changes in the human gut microbiota. *Gut* **2017**, gutjnl-2016-313271.
65. Bhinder, G.; Allaire, J. M.; Garcia, C.; Lau, J. T.; Chan, J. M.; Ryz, N. R.; Bosman, E. S.; Graef, F. A.; Crowley, S. M.; Celiberto, L. S.; Berkman, J. C.; Dyer, R. A.; Jacobson, K.; Surette, M. G.; Innis, S. M.; Vallance, B. A. Milk Fat Globule Membrane supplementation in formula modulates the neonatal gut microbiome and normalizes intestinal development. *Sci. Rep.* **2017**, 7, 1–15.
66. Lee, H.; Zavaleta, N.; Chen, S.; Lönnerdal, B.; Slupsky, C. Effect of bovine milk fat globule membranes as a complementary food on the serum metabolome and immune markers of 6-11-month-old Peruvian infants. *npj Sci. Food* **2018**, 2, 6.
67. Lovegrove, J. A.; Hobbs, D. A. Plenary Lecture 2: Milk and dairy produce and CVD: new perspectives on dairy and cardiovascular health. *Proc. Nutr. Soc.* **2016**, 44, 1–12.
68. Flint, H. J.; Scott, K. P.; Duncan, S. H.; Louis, P.; Forano, E. Microbial degradation of complex carbohydrates in the gut. *Gut Microbes* **2012**, 3, 289–306.
69. Huang, Y.-L.; Chassard, C.; Hausmann, M.; von Itzstein, M.; Hennot, T. Sialic acid catabolism drives intestinal inflammation and microbial dysbiosis in mice. *Nat. Commun.* **2015**, 6, 8141.
70. Juge, N.; Tailford, L.; Owen, C. D. Sialidases from gut bacteria: a mini-review. *Biochem. Soc. Trans.* **2016**, 44, 166–175.
71. Charbonneau, M. R.; O'Donnell, D.; Blanton, L. V.; Totten, S. M.; Davis, J. C. C.; Barratt, M. J.; Cheng, J.; Guruge, J.; Talcott, M.; Bain, J. R.; Muehlbauer, M. J.; Ilkayeva, O.; Wu, C.; Struckmeyer, T.; Barile, D.; Mangani, C.; Jorgensen, J.; Fan, Y. M.; Maleta, K.; Dewey, K. G.; Ashorn, P.; Newgard, C. B.; Lebrilla, C.; Mills, D. A.;

- Gordon, J. I. Sialylated Milk Oligosaccharides Promote Microbiota-Dependent Growth in Models of Infant Undernutrition. *Cell* **2016**, *164*, 859–871.
72. Knight, C. A. L. and R. Species Divergence and the Measurement of Microbial Diversity. *FEMS Microbiol Rev.* **2009**, *32*, 557–578.
73. Dubourg, G.; Lagier, J.-C.; Armougom, F.; Robert, C.; Audoly, G.; Papazian, L.; Raoult, D. High-level colonisation of the human gut by Verrucomicrobia following broad-spectrum antibiotic treatment. *Int. J. Antimicrob. Agents* **2013**, *41*, 149–155.
74. Haug, A.; Høstmark, A. T.; Harstad, O. M. Bovine milk in human nutrition – a review. *Lipids Health Dis.* **2007**, *6*, 25.
75. Visioli, F.; Strata, A. Milk, dairy products, and their functional effects in humans: a narrative review of recent evidence. *Adv. Nutr.* **2014**, *5*, 131–143.
76. Guo, J.; Astrup, A.; Lovegrove, J. A.; Gijsbers, L.; Givens, D. I.; Soedamah-Muthu, S. S. Milk and dairy consumption and risk of cardiovascular diseases and all-cause mortality: dose–response meta-analysis of prospective cohort studies. *Eur. J. Epidemiol.* **2017**, *32*, 269–287.
77. de Heijning, B.; Kegler, D.; Schipper, L.; Voogd, E.; Oosting, A.; Beek, E. Acute and chronic effects of dietary lactose in adult rats are not explained by residual intestinal lactase Activity. *Nutrients* **2015**, *7*, 5542–5555.
78. Org, E.; Mehrabian, M.; Parks, B. W.; Shipkova, P.; Liu, X.; Drake, T. A.; Lusi, A. J. Sex differences and hormonal effects on gut microbiota composition in mice. *Gut Microbes* **2016**, *7*, 313–322.
79. Wang, J.; Wang, J.; Pang, X.; Zhao, L.; Tian, L.; Wang, X. Sex differences in colonization of gut microbiota from a man with short-term vegetarian and inulin-supplemented diet in germ-free mice. *Sci. Rep.* **2016**, *6*, 36137.

734 **FIGURE CAPTIONS**

735 Figure 1 Composition of the human and murine baseline faecal microbiota at family level.
 736 Bacterial families that were present at >1% relative abundance are presented. Murine:
 737 aggregated faecal microbiota across all mice at baseline; COM: community type faecal
 738 microbiota; LS: longstay type faecal microbiota.

739 Figure 2 Principal coordinates analysis (PCoA) of the UniFrac distances at the end of the trial
 740 (T5). A. Weighted UniFrac distances; B. Unweighted UniFrac distances. Blue: murine
 741 baseline microbiota; green colour coding: community (COM) type faecal microbiota; red
 742 colour coding: longstay (LS) type faecal microbiota. The inocula of human faecal microbiota
 743 (donors) are denoted with “▲” and the relevant red and green colour..

744 Figure 3 Alpha diversity indices at the end of the trial (T5). A: community (COM) type
 745 colonisation; B: longstay (LS) type colonisation. LAC_FREE: lactose free milk
 746 supplemented diet; WMILK: whole milk supplemented diet; GMP: glycomacropeptide
 747 supplemented diet; CONTROL: soy protein diet. Asterisks next to alpha diversity indices
 748 denote the result of the Kruskal-Wallis comparison; markings above boxplots denote the
 749 Dunn’s post hoc result. Significant values: “●” $p_{adj}<0.1$; “*” $p_{adj}<0.05$; “**” $p_{adj}<0.01$;
 750 “***” $p_{adj}<0.001$; “****” $p_{adj}<0.0001$.

751 Figure 4 Principal coordinates analysis (PCoA) of the faecal microbiota of mice at the end of
 752 the trial (T5). Community type (COM) humanisation: A: Weighted UniFrac distances; B:
 753 Unweighted UniFrac distances. Longstay type (LS) humanisation: C: Weighted UniFrac
 754 distances; D: Unweighted UniFrac distances. LAC_FREE: lactose free milk supplemented
 755 diet; WMILK: whole milk supplemented diet; GMP: glycomacropeptide supplemented diet;
 756 CONTROL: soy protein diet.”●”: female mice microbiota; “▲”: male mice microbiota. The
 757 ANOSIM results of the diet effect are noted.

758 Figure 5 Faecal microbiota composition at family level at the end of the trial (T5). COM:
759 community type colonisation; LS: longstay type colonisation. A. microbiota composition of
760 female and male mouse aggregated; B. female mouse microbiota composition; C. male
761 mouse microbiota composition. LAC_FREE: lactose free milk supplemented diet; WMILK:
762 whole milk supplemented diet; GMP: glycomacropeptide supplemented diet; CONTROL:
763 soy protein diet. Bacterial families that were present at >1% relative abundance are presented.

764 Figure 6 Differentially abundant taxa at family level at the end of the trial (T5) for
765 community (COM) type humanisation. A: female mouse microbiota; B: male mouse
766 microbiota. Asterisks next to taxa denote the result of Kruskal-Wallis comparison; markings
767 above boxplots denote the Dunn's post hoc result. Significant values: "●" $p_{adj}<0.1$; "*" $p_{adj}<0.05$;
768 "***" $p_{adj}<0.01$; "****" $p_{adj}<0.001$; "*****" $p_{adj}<0.0001$.

769 Figure 7 Figure Differentially abundant taxa at family level at the end of the trial (T5) for
770 longstay (LS) type humanisation. A: female mouse microbiota; B: male mouse microbiota.
771 Asterisks next to taxa denote the result of Kruskal-Wallis comparison; markings above
772 boxplots denote the Dunn's post hoc result. Significant values: "●" $p_{adj}<0.1$; "*" $p_{adj}<0.05$;
773 "***" $p_{adj}<0.01$; "****" $p_{adj}<0.001$; "*****" $p_{adj}<0.0001$.

774 Figure 8 Faecal microbiota composition at genus level at the end of the trial (T5). COM:
775 community type colonisation; LS: longstay type colonisation. A. microbiota composition of
776 female and male mouse aggregated; B. female mouse microbiota composition; C. male
777 mouse microbiota composition. LAC_FREE: lactose free milk supplemented diet; WMILK:
778 whole milk supplemented diet; GMP: glycomacropeptide supplemented diet; CONTROL:
779 soy protein diet. Bacterial genera that were present at >1% relative abundance are presented.

780 Figure S 1 Timeline of the mouse trial. Time points of faecal sample collection: baseline (T0)
781 during acclimatisation; T1 and T2 during the antibiotic treatment; T3, T4 and T5 during the
782 dietary intervention.

783 Figure S 2 Animal body weight during the 11 weeks of the trial. The body weight from the
784 first week of Abx treatment (W0) till the completion of the trial (W11) is shown. Body
785 weight change between baseline and end of trial (T5) is shown at the bottom. A and B: COM:
786 community type humanisation; LS: longstay type humanisation, respectively.

787 Figure S 3 Principal coordinates analysis (PCoA) of the UniFrac distances one week after
788 “humanisation” (T3). A: Weighted UniFrac distances; B: Unweighted UniFrac distances.
789 Blue: murine baseline microbiota; green colour coding: community (COM) type faecal
790 microbiota; red colour coding: longstay (LS) type faecal microbiota. Colour shading denotes
791 the test diets. The inocula of human faecal microbiota are denoted with “▲” and the relevant
792 red and green colour.

793 Figure S 4 Differentially abundant taxa at family level at the end of the trial (T5) for
794 aggregated female and male mouse faecal microbiota. A: community (COM) type
795 colonisation; B: longstay (LS) type colonisation. Asterisks next to taxa denote the result of
796 Kruskal-Wallis comparison; markings above boxplots denote the Dunn’s post hoc result.
797 Significant values: “●” $p_{adj}<0.1$; “*” $p_{adj}<0.05$; “**” $p_{adj}<0.01$; “***” $p_{adj}<0.001$; “*****”
798 $p_{adj}<0.0001$. COM type humanisation: Differences in the Erysipelotrichaceae relative
799 abundance shown in Figures 6 and S4A are due to filtering for taxa presence in the compared
800 groups for the statistical analysis. LS type humanisation: Due to filtering of the taxa tables for
801 the statistical analysis, *Allobaculum* that was not detected in the LS type colonised male mice
802 and fed wmilk diet (Figure 8C), was not represented in Figure 7.

803 Figure S 5 Faecal microbiota composition at species level at the end of the trial (T5). COM:
804 community type colonisation; LS: longstay type colonisation. A. microbiota composition of
805 female and male mouse aggregated; B. female mouse microbiota composition; C. male
806 mouse microbiota composition LAC_FREE: lactose free milk supplemented diet; WMILK:
807 whole milk supplemented diet; GMP: glycomacropeptide supplemented diet; CONTROL:
808 soy protein diet. Bacterial species present at >1% relative abundance are shown. Colour
809 coding: purple: Bacteroidetes; green: Firmicutes; red: Proteobacteria; blue: Verrucomicrobia.

810 Figure S 6 Composition of the murine faecal microbiota at family level at the mid part of the
811 dietary intervention (T4). COM: community type microbiota; LS: longstay type microbiota.
812 LAC_FREE: lactose free milk diet; WMILK: whole milk diet, GMP: glycomacropeptide diet;
813 CONTROL: soy protein diet. Only phyla present at >1% relative abundance are presented.

814 Figure S 7 Alpha diversity indices at the mid part of the dietary intervention (T4). A:
815 community (COM) type colonisation; B: longstay (LS) type colonisation. LAC_FREE:
816 lactose free milk supplemented diet; WMILK: whole milk supplemented diet; GMP:
817 glycomacropeptide supplemented diet; CONTROL: soy protein diet. Asterisks next to alpha
818 diversity indices denote the result of the Kruskal-Wallis comparison; markings above
819 boxplots denote the Dunn's post hoc result: Significant values: "●" $p_{adj}<0.1$; "*" $p_{adj}<0.05$;
820 "***" $p_{adj}<0.01$; "****" $p_{adj}<0.001$; "*****" $p_{adj}<0.0001$.

TABLES

Table 1 Composition of the customised experimental diets (ssniff Spezialdiäten GmbH).

Ingredient %	Control	20% whole milk powder	20% lactose-free milk powder	20% CGMP-10 powder
Soy protein isolate	20.0	14.5	12.75	1.8
Whole milk powder	-	20.0	-	-
Lactose-free milk powder	-	-	20.0	-
Glycomacropeptide CGMP-10	-	-	-	20.0
Maltodextrin	33.1	33.2	32.85	32.0
Sucrose	10.5	2.5	4.3	10.3
Cellulose powder	6.0	4.9	5.0	5.2
Soybean oil	7.5	2.3	2.3	7.5
Contents %				
Crude protein	18.1	18.1	18.1	18.1
Crude fat	7.6	7.6	7.6	7.6
Crude ash	4.6	5.4	5.6	5.5
Starch	31.8	31.9	31.6	30.8
Sugar (total)	12.3	12.3	12.3	12.3
Lactose	-	8.0	-	-
Energy (MJ/kg)	16.0	16.0	16.0	16.0
kcal % Protein	19	19	19	19
kcal % Fat	18	18	18	18
kcal % Carbohydrates	63	63	63	63

Table 2 ANOSIM test results for the effect of gender on the observed PCoA clustering of the faecal microbiota at the end of the trial (T5).

	COM				LS			
	Weighted		Unweighted		Weighted		Unweighted	
			ANOSIM					
	p value	R	p value	R	p value	R	p value	R
Control	0.005	0.98	0.006	1.0	0.004	0.75	0.003	1.0
GMP	0.014	0.98	0.01	0.94	0.41	0.0	0.005	1.0
Lactose-free milk	0.009	1.0	0.008	1.0	0.04	0.37	0.19	0.10
Whole milk diet	0.008	1.0	0.005	1.0	0.028	1.0	0.035	1.0

FIGURE GRAPHICS

FIGURE 1





FIGURE 2

FIGURE 3





FIGURE 4

FIGURE 5




FIGURE 6




FIGURE 7





FIGURE 8



Supporting Information

Table S 1 Treatment groups indicating the diet, gender and the colonisation with human faecal microbiota type. The number of animals in each dietary group at T5 for which the gut microbiota was analysed is shown. The experimental arms of groups A, B, C and D were performed simultaneously followed by the experimental arms of groups E, F, G and H. The animals that received HydroGel treatment during Abx period, is indicated by #.

	Community microbiota (COM)	type human	Longstay type human microbiota (LS)	
	female	male	female	male
Lactose free milk	A (n=5)	A (n=5)	B (n=4)	B (n=5)
Whole milk	C (n=6)	C (n=4)	D (n=4)	D (n=4)
GMP	E (n=5)#	E (n=4)#	G (n=6)	G (n=6)
Control	H (n=6)###	H (n=6)	F (n=6)##	F (n=5)###
“#”, “##”, “###” three, six and nine days, respectively of HydroGel intake				

Table S 2 Alpha diversity indices of the faecal microbiotas of the human donors and the murine baseline faecal microbiota. EM297: longstay type faecal microbiota (LS); EM425: community type faecal microbiota (COM); murine: aggregated faecal microbiota across all mice at baseline.

Alpha Diversity Index	MURINE	EM297 LS	EM425 COM
Shannon	6.0	4.0	6.0
Simpson	1.0	1.0	1.0
PD Whole tree	15.0	9.0	17.0
Observed species	207.0	73.0	169.0
Chao 1	346.0	104.0	264.0

FIGURE S1




FIGURE S2




FIGURE S3




FIGURE S4




FIGURE S5

FIGURE S6





FIGURE S7



GRAPHIC FOR TABLE OF CONTENTS

