

Title	A window beneath the skin: how computed tomography assessment of body composition can assist in the identification of hidden wasting conditions in oncology that profoundly impact outcomes?
Authors	Daly, Louise E.;Prado, Carla M.;Ryan, Aoife M.
Publication date	2018-05-10
Original Citation	Daly, L. E., Prado, C. M. and Ryan, A. M. (2018) 'A window beneath the skin: how computed tomography assessment of body composition can assist in the identification of hidden wasting conditions in oncology that profoundly impact outcomes?', Proceedings of the Nutrition Society, 77(2), pp. 135-151. doi:10.1017/S0029665118000046
Type of publication	Conference item
Link to publisher's version	10.1017/S0029665118000046
Rights	© 2018, the Authors. Published by Cambridge University Press on behalf of the Nutrition Society. All rights reserved.
Download date	2026-09-25 03:47:23
Item downloaded from	https://hdl.handle.net/10468/6149



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh



Conference on ‘What governs what we eat?’ Postgraduate Symposium

A window beneath the skin: how computed tomography assessment of body composition can assist in the identification of hidden wasting conditions in oncology that profoundly impact outcomes

L. E. Daly^{1,2†}, C. M. Prado³ and A. M. Ryan^{1,4*}

¹*School of Food and Nutritional Sciences, College of Science, Engineering and Food Science, University College Cork, Ireland*

²*APC Microbiome Institute, University College Cork, Cork, Ireland*

³*Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, Alberta, Canada*

⁴*Cork Cancer Research Centre, University College Cork, Cork, Ireland*

Advancements in image-based technologies and body composition research over the past decade has led to increased understanding of the importance of muscle abnormalities, such as low muscle mass (sarcopenia), and more recently low muscle attenuation (MA), as important prognostic indicators of unfavourable outcomes in patients with cancer. Muscle abnormalities can be highly prevalent in patients with cancer (ranging between 10 and 90 %), depending on the cohort under investigation and diagnostic criteria used. Importantly, both low muscle mass and low MA have been associated with poorer tolerance to chemotherapy, increased risk of post-operative infectious and non-infectious complications, increased length of hospital stay and poorer survival in patients with cancer. Studies have shown that systemic antineoplastic treatment can exacerbate losses in muscle mass and MA, with reported loss of skeletal muscle between 3 and 5 % per 100 d, which are increased exponentially with progressive disease and proximity to death. At present, no effective medical intervention to improve muscle mass and MA exists. Most research to date has focused on treating muscle depletion as part of the cachexia syndrome using nutritional, exercise and pharmacological interventions; however, these single-agent therapies have not provided promising results. Rehabilitation care to modify body composition, either increasing muscle mass and/or MA should be conducted, and its respective impact on oncology outcomes explored. Although the optimal timing and treatment strategy for preventing or delaying the development of muscle abnormalities are yet to be determined, multimodal interventions initiated early in the disease trajectory appear to hold the most promise.

Sarcopenia: Muscle attenuation: Cancer: Cachexia: Body composition: Survival

Over the past decade, there has been a growing interest in the measurement of body composition in patients with cancer. This has largely been in response to advancements in image-based technologies, including

gold standard computed tomography (CT) that allow the precise quantification of both muscle and adipose tissue. This research has led to increased understanding of the importance of abnormal body composition

Abbreviations: CT, computed tomography; DLT, dose-limiting toxicity; GI, gastrointestinal; HR, hazard ratio; HU, Hounsfield unit; MA, muscle attenuation; SMA, skeletal muscle area; SMI, skeletal muscle index.

*Corresponding author: Dr Aoife Ryan, email a.ryan@ucc.ie

†Presented by Louise Daly at The Nutrition Society Summer Meeting, Queens University Belfast, Belfast, Ireland, on 21–23 June 2017

phenotypes, such as low muscle mass (sarcopenia), and more recently low muscle attenuation (MA) as important prognostic indicators of unfavourable outcomes in patients with cancer^(1–4).

Cancer is a disease associated with ageing. As a result, the aetiology of muscle loss in patients with cancer can be two-fold. Firstly, resulting from the age-related decline in muscle mass (sarcopenia), and secondly due to the metabolic changes induced by malignancy and cancer cachexia. Muscle loss related to sarcopenia begins relatively early in life, as muscle mass begins to decline from the age of 40 at a rate of 6 % per decade⁽⁵⁾ and accelerates to a rate of 25–40 % per decade above age 70 years⁽⁶⁾. The precise definition of sarcopenia remains controversial and a topic of much debate today; however, a generally accepted criterion in geriatric populations is a level of muscle mass greater than two standard deviations below that of a healthy young reference population⁽⁷⁾. Cancer cachexia is a multifactorial syndrome that is characterised by the loss of muscle with or without the loss of fat mass leading to progressive functional impairment⁽⁸⁾. It is driven by a variable combination of reduced food intake and abnormal metabolism⁽⁸⁾. Systemic inflammation is commonly described as part of the pathogenesis of cancer cachexia, and may suppress appetite, increase body's metabolic needs and energy expenditure, and accelerate muscle protein catabolism in patients^(8–10). Cancer cachexia represents a spectrum of conditions and can range in severity and clinical presentation from pre-cachexia, identified by early clinical and metabolic signs (e.g. anorexia and impaired glucose tolerance) to refractory cachexia, where extensive muscle and fat depletion is evident, and patients are often immunocompromised⁽⁸⁾. Recognition of these stages of cachexia is important as these stages have different implications in the anabolic therapy response.

Reduced skeletal MA is a relatively newly characterised and distinctive abnormality in patients with cancer⁽¹¹⁾. It represents a 'qualitative' measure of skeletal muscle, as low radiation attenuation is reflective of intramuscular adipose tissue infiltration, and therefore, poor 'quality' skeletal muscle⁽¹¹⁾. It has been speculated that low MA precedes the development of sarcopenia, as the increase in lipid content occurs before a decline in muscle mass^(12,13). Importantly, in recent years, low MA is emerging as an important prognostic indicator in patients with cancer, and in some cases, a better prognostic indicator compared with muscle mass alone^(12–18).

Low muscle mass and low MA can occur at any given BMI (kg/m²)^(1,19), and now with 40–60 % of cancer patients presenting with overweight and obesity⁽²⁰⁾, identification of these conditions is becoming increasingly difficult. Patients may therefore not appear malnourished as muscle depletion is often hidden behind mantles of adipose tissue and can often go undiagnosed and untreated. Body composition assessment is therefore crucial within this patient group.

This review will focus on the diagnostic criteria, prevalence and clinical consequences associated with CT-defined muscle abnormalities (low muscle mass and low MA) in relation to chemotherapy tolerance, post-operative

outcomes and survival in oncology patients. We will examine the changes in body composition that occur in patients undergoing active anti-cancer treatment, and finally, we will briefly examine the evidence behind current treatments aimed at restoring muscle mass and MA, with a particular emphasis on nutritional interventions, physical activity, pharmacological agents and multimodal treatments.

Measurement of skeletal muscle in oncology

The three most commonly used methods to measure body composition in patients with cancer are bioelectrical impedance analysis, dual-energy x-ray absorptiometry and CT. Each technique has its advantages and disadvantages, and methods differ in terms of cost, reliability, validity, availability and training required⁽²¹⁾. CT images are considered a gold standard method for body composition assessment, and allow the precise quantification of tissue area, volume and attenuation. Unlike bioelectrical impedance analysis or dual-energy x-ray absorptiometry, CT can measure body composition at a tissue-organ level, particularly total and regional adipose and skeletal muscle tissues. The use of CT for body composition assessment in non-cancer populations is limited by the high radiation dose, high cost and lack of availability. However, in oncology, CT scans are obtained routinely during diagnostic and surveillance purposes, and therefore represent a unique and exploitable opportunity to assess body composition within this patient group.

Measuring muscle mass by CT is usually done by measuring total skeletal muscle area (SMA) at the third lumbar vertebra (L3). Using a commercially available image analysis software (e.g. Sliceomatic (TomoVision), OsiriX (Pixmeo), Image J (National Institutes of Health), among others), muscle and adipose tissues can be evaluated based on Hounsfield unit (HU) thresholds. Muscles in this area include the psoas, paraspinal muscles (erector spinae, quadratus lumborum) and the abdominal wall muscles (transversus abdominus, external and internal oblique, rectus abdominus). Measurements are commonly taken at the third lumbar vertebra, as the SMA obtained at this level is a good correlate for whole body muscle in healthy individuals (r 0.92)⁽²²⁾. From this skeletal muscle index (SMI; total SMA (cm²)/height (m²)) can be calculated and patients are often compared on this basis. Mean MA is typically derived by averaging HU of the SMA at the third lumbar vertebra. Although SMI and MA are continuous variables and could be modelled as such to predict survival/outcome, many clinicians find the interpretation of continuous prognostic covariates difficult, and prefer categorical or binary covariates based on a threshold/cut point to stratify patients into distinct risk groups when making treatment decisions.

Defining low muscle mass (sarcopenia) in patients with cancer

Many oncological studies define sarcopenia based solely on low muscle mass, and studies often lack information on muscle strength or physical function. Therefore, the criteria to define sarcopenia in this group differ from

those proposed in geriatric populations^(23,24). In the oncology setting, consensus-based cut points to define low muscle mass or sarcopenia are lacking, and a variety have been devised. Studies^(25–28) have defined sarcopenia based on cut points developed by Baumgartner *et al.*⁽⁷⁾ in elderly individuals (2-sd below a healthy reference population), by converting the original dual-energy x-ray absorptiometry cut points ($<7.26 \text{ kg/m}^2$ for men and $<5.45 \text{ kg/m}^2$ for women) to corresponding CT cut points using published regression equations⁽²⁹⁾. These cut points are $\text{SMI} <55.4 \text{ cm}^2/\text{m}^2$ for men and $<38.9 \text{ cm}^2/\text{m}^2$ for women⁽²⁹⁾, and are those used in the cancer cachexia consensus definition⁽⁸⁾. To date, no healthy population reference values for muscle mass obtained from CT exist.

Studies have employed a statistical technique known as optimal stratification, which is used to identify threshold values associated with elevated risk of poor outcome (e.g. mortality) to define sarcopenia. Using this technique, Prado *et al.*⁽¹⁹⁾ identified cut points for SMI that best predicted survival in a cohort of 250 obese ($\text{BMI} >30 \text{ kg/m}^2$) cancer patients. These cut points are $<52.4 \text{ cm}^2/\text{m}^2$ for men and $<38.5 \text{ cm}^2/\text{m}^2$ for women⁽¹⁹⁾, and have been widely applied in the literature. In 2013, Martin *et al.*⁽¹⁾ identified both sex and BMI-specific cut points for SMI that best predicted survival (men: $<43 \text{ cm}^2/\text{m}^2$ if $\text{BMI} \leq 24.9 \text{ kg/m}^2$ and $<53 \text{ cm}^2/\text{m}^2$ if $\text{BMI} \geq 25 \text{ kg/m}^2$; women: $<41 \text{ cm}^2/\text{m}^2$)⁽¹⁾ in a large cohort of 1473 patients with lung and gastrointestinal (GI) cancer, which are more applicable to non-obese cohorts. More recently, in 2017, Caan *et al.*⁽³⁰⁾ identified sex, BMI and cancer-specific cut points for survival in a very large cohort of early-stage colorectal cancer patients ($n = 3262$; men: $<52.3 \text{ cm}^2/\text{m}^2$ if $\text{BMI} <30 \text{ kg/m}^2$ and $<54.3 \text{ cm}^2/\text{m}^2$ if $\text{BMI} \geq 30 \text{ kg/m}^2$; women: $<38.6 \text{ cm}^2/\text{m}^2$ if $\text{BMI} <30 \text{ kg/m}^2$ and $<46.6 \text{ cm}^2/\text{m}^2$ if $\text{BMI} \geq 30 \text{ kg/m}^2$)⁽³⁰⁾.

In addition to sex and BMI, ethnicity should be considered when applying cut points to a specific cohort. Fujiwara *et al.*⁽³¹⁾ defined CT-derived cut points for a large ($n = 1257$) homogenous cohort of Japanese patients with mixed stage (I–IV) hepatocellular carcinoma that best predicted survival ($<36.2 \text{ cm}^2/\text{m}^2$ for men and $<29.0 \text{ cm}^2/\text{m}^2$ for women), providing a reference population for these individuals. Notably, these cut points are lower^(31,32) compared with those derived from large Caucasian populations^(1,3,34).

Using optimal stratification methodology, several other studies have reported cut points for SMI associated with mortality in a number of cancer cohorts (see Table 1), these range $36\text{--}55.8 \text{ cm}^2/\text{m}^2$ for men and $29\text{--}46.6 \text{ cm}^2/\text{m}^2$ for women. Several factors influence patient's muscularity (ethnicity, age, sex, physical activity and magnitude of adiposity)⁽³⁵⁾; hence, published cut points may not be applicable to all cancer populations. For example, cut points published by Prado *et al.*⁽¹⁹⁾ were devised in a cohort of obese patients with cancer, and may have limited relevance when applied to populations with varying prevalence of obesity, but have subsequently been used in studies comprised of predominantly non-obese individuals^(36–40).

Other studies have defined sarcopenia based on more data-orientated approaches, dichotomizing SMI based on predetermined percentiles, such as quartiles⁽⁴¹⁾, tertiles^(17,42)

or based on the median^(43,44). Studies focused solely on one muscle group, the psoas muscle, generating a psoas muscle index, often considered patients within the lowest quartile of psoas muscle index to be sarcopenic^(45–47). However, no studies have related psoas muscle area to whole body measures, and it has only been weakly correlated with total lumbar muscle area⁽⁴⁸⁾. Therefore, it has been argued that the use of the psoas muscle as a sentinel muscle for the diagnosis of sarcopenia is flawed⁽⁴⁹⁾. Other investigations have included additional measures of physical function (hand grip strength and/or gait speed) in conjunction with skeletal muscle from CT scan analysis when defining sarcopenia in patients with cancer^(50,51). The rationale for including the measures of both muscle mass and function is that muscle strength does not depend solely on muscle mass, and the relationship between muscle strength and mass is not linear⁽⁶⁾. Hence, several expert groups have proposed the use of both muscle mass and function to define sarcopenia in older adults^(23,24,52).

Prevalence of low muscle mass (sarcopenia) in patients with cancer

As discussed previously, comparison among studies reporting the prevalence of sarcopenia is often difficult because of inconsistent methodology used to evaluate body composition (dual-energy x-ray absorptiometry, bioelectrical impedance analysis, CT), and diagnostic criteria/cut points used to define sarcopenia. Table 2 summarises the prevalence of sarcopenia among patients with cancer using the same methodology (CT defined SMI) but using varying published cut points. The prevalence of sarcopenia is highly variable across primary cancer sites, and has been shown to range from 11 to 90%^(31,53). Of note is the heterogeneity in the prevalence of sarcopenia among cohorts of the same primary cancer site and stage, e.g. 19–71% in advanced colorectal cancer^(54,55), 33–90% in metastatic kidney cancer^(53–56) and 21–89% in advanced pancreatic cancer^(57,58). This large variability could be attributed to the varying cut points used to define sarcopenia; limitations related to sample size; or patient characteristics such as age, sex, ethnicity, BMI and concurrent co-morbidities, thus limiting the ability to draw conclusions as to the true prevalence according to the cancer site. Sarcopenia appears to be the most prevalent in patients with any stage of pancreatic or lung cancer, while in other cancer types (e.g. gastric and breast), it appears to be more frequent in advanced stage disease compared with earlier loco regional disease. Sarcopenia can be present at any given BMI, and the prevalence of sarcopenic obesity (sarcopenia and obesity ($\text{BMI} \geq 30 \text{ kg/m}^2$)) has been shown to vary between 1 and 29% in studies including individuals from all BMI categories, and between 15 and 36% in studies including obese individuals only⁽⁵⁹⁾.

Low muscle attenuation in patients with cancer

Similar to the measures of SMI, no widely agreed upon cut points are available for defining low MA. Martin *et al.*⁽¹⁾ provided the first set of cut points for low MA

Table 1. Cut points for skeletal muscle index (SMI) at the third lumbar vertebra (L3) associated with mortality in patients with cancer

Author	Country	Tumour site/stage	n	Males SMI (cm ² /m ²)	Females SMI (cm ² /m ²)
Prado <i>et al.</i> ⁽¹⁹⁾	Canada	Respiratory and GI/stages I–IV*	250	<52.4	<38.5
Van Vledder <i>et al.</i> ⁽⁵⁴⁾	The Netherlands	Colorectal cancer with liver metastasis	196	<43.75	<41.1
Martin <i>et al.</i> ⁽¹⁾	Canada	Lung and GI/stages I–IV	1473	<43.0 (BMI <25 kg/m ²) <53.0 (BMI ≥25 kg/m ²)	<41.0
Lanic <i>et al.</i> ⁽³³⁾	France	Diffuse large B-cell lymphoma	82	<55.8	<38.9
Iritani <i>et al.</i> ⁽³²⁾	Japan	Hepatocellular carcinoma/stages I–IV	217	≤36.0	≤29.0
Fujiwara <i>et al.</i> ⁽³¹⁾	Japan	Hepatocellular carcinoma/stages I–IV	1257	<36.2	<29.6
Kimura <i>et al.</i> ⁽¹⁶¹⁾	Japan	Lung cancer/stage III–IV	134	<41.0	<38.0
Choi <i>et al.</i> ⁽⁵⁷⁾	Korea	Pancreatic cancer/advanced	484	<42.2	<33.9
Coelen <i>et al.</i> ⁽³⁴⁾	The Netherlands	Perihilar cholangiocarcinoma/stages I–IV	100	<46.8	<39.1
Cann <i>et al.</i> ⁽³⁰⁾	USA	Colorectal cancer/stages I–III	3262	<52.3 (BMI <30 kg/m ²) <54.3 (BMI ≥30 kg/m ²)	<38.6 (BMI <30 kg/m ²) <46.6 (BMI ≥30 kg/m ²)

GI, gastrointestinal.

* Obese individuals only (BMI >30 kg/m²).

(using optimal stratification) that related to poor survival in a large cohort of lung and GI cancer patients (n 1473; <41 HU for BMI <25 kg/m² and <33 for BMI ≥25 kg/m²)⁽¹⁾. Since then, cancer-specific cut points for low MA that relate optimally to survival have been reported for lung^(18,60), ovarian^(61,62), perianapillary⁽¹⁶⁾, pancreatic⁽⁶³⁾, gastro-oesophageal⁽²⁸⁾ and large B-cell lymphoma⁽¹⁵⁾. These range from <28.0 to 44.1 HU in men^(16,18,60) and <23.8 to 40.5 HU in women^(16,18,60–62). Other studies have defined low MA based on the sample sex-specific median^(44,64), tertile⁽¹⁷⁾ or quartile⁽⁶⁵⁾.

The prevalence of low MA among patients with cancer varies greatly, and is dependent on the cohort under investigation and cut point used, but has been shown to range from 10 to 86%^(18,66). Using the cut points devised by Martin *et al.*⁽¹⁾, which have been applied most widely in the literature, the prevalence of low MA has been reported to be between 46 and 53% in two large cohorts of patients with cancer (mixed tumour sites and stages (I–IV))^(1,67). In the setting of advanced disease, the prevalence of low MA has been reported to be 33% in melanoma⁽⁴⁾, 55% in pancreatic⁽⁶⁸⁾, 59% in gastric⁽¹³⁾, 60% in breast⁽¹⁴⁾ and 86% in prostate cancer⁽⁶⁶⁾, while in patients with operable colorectal cancer, low MA is present in 58–78% of patients^(69,70).

Impact of muscle abnormalities on clinical outcomes

Notwithstanding the controversies in determining and defining low muscle mass and low MA in oncology, it has been well established and reported over the past decade that these muscle abnormalities are unequivocally associated with negative clinical outcomes in patients with cancer.

Muscle abnormalities and tolerance to chemotherapy

Chemotherapy can often be associated with severe toxicity (grades III–IV) that can result in dose delays, dose reductions and treatment termination, referred to as dose-limiting toxicities (DLT). DLT may lead to hospitalisations and can be life threatening. Exploratory

studies provided the initial evidence of an association between low muscularity and increased incidence of severe toxicity/DLT to chemotherapy, and subsequent work has confirmed these observations in multiple cancer sites and treatments (Fig. 1). In advanced disease, low muscularity has been associated with poorer tolerance to chemotherapy in patients with breast^(71,72), renal^(56,73,74), liver⁽⁷⁵⁾, lung⁽⁷⁶⁾, colorectal^(55,77), thyroid⁽⁷⁸⁾ and melanoma skin cancer^(4,79). Sarcopenia, in patients with peritoneal metastasis from colorectal cancer, was associated with significantly more chemotherapy toxicities (57 v. 26%, $P=0.004$) and particularly neutropenia (36 v. 17%, $P=0.04$) in a cohort of 97 patients receiving hyperthermic intraperitoneal chemotherapy.

Even in early-stage disease (stages I–III), sarcopenia is associated with poorer tolerance to chemotherapy^(2,80–84). Interestingly, in patients with early-stage breast cancer (n 151) and receiving anthracycline- and taxane-based chemotherapy, with every five-unit decrease in SMI, the risk of any grade III–IV toxicity increased by 27% (relative risk 1.27 (95% CI 1.09, 1.49), $P=0.002$)⁽⁸²⁾. In a large study of patients with non-metastatic colon cancer (n 533) treated with adjuvant FOLOX treatment, patients with low muscle mass (lowest sex-specific tertile) were twice as likely to experience dose reductions (OR 2.28, $P=0.01$), dose delays (OR 2.24, $P=0.002$) and early discontinuation of treatment (OR 2.34, $P=0.03$)⁽²⁾. Although the majority of studies have consistently demonstrated an association between low muscle mass and increased incidence of severe toxicity/DLT (twenty-six of thirty-two studies; see Supplementary material for summary of each study), fewer small studies found contradictory findings^(85–88).

Emerging data suggest low MA to be also associated with poorer tolerance to antineoplastic agents^(4,66,82). In metastatic melanoma patients (n 84), we have previously shown that patients with low MA more frequently experienced high-grade toxicities (75 v. 31%, $P=0.001$) and immune-related toxicities to ipilimumab (54 v. 23%, $P=0.017$) compared with those without low MA. More importantly, these patients were more susceptible to experience a DLT (37.5 v. 10.4%, $P=0.011$)⁽⁴⁾. In patients with metastatic prostate cancer treated with

Table 2. Prevalence of sarcopenia according to cancer site, country and definition used

Author	Country	Stage/n	% Sarcopenic*	Sarcopenia definition†
Colorectal cancer				
Van Vledder <i>et al.</i> ⁽⁵⁴⁾	The Netherlands	Advanced/196	19	4
Lieffers <i>et al.</i> ⁽¹¹⁰⁾	Canada	Stages II–IV/234	39	1
Thoresen <i>et al.</i> ⁽¹⁶²⁾	Norway	Advanced/50	20	1
Barret <i>et al.</i> ⁽⁵⁵⁾	France	Metastatic/51	71	2
Thoresen <i>et al.</i> ⁽¹⁶³⁾	Norway and Canada	Advanced/77	39	1
Miyamoto <i>et al.</i> ⁽⁴¹⁾	Japan	Stages I–III/220	25	6
Reisinger <i>et al.</i> ⁽¹⁶⁴⁾	The Netherlands	Stages I–IV/310	48	1
Buskermolen <i>et al.</i> ⁽⁸⁸⁾	The Netherlands	Metastatic/67	57	3
Chemama <i>et al.</i> ⁽⁷⁷⁾	France	Advanced/97	40	3
Malietzis <i>et al.</i> ⁽¹⁶⁵⁾	UK	Resectable/763	65	3
Cann <i>et al.</i> ⁽³⁰⁾	USA	Stages I–III/3262	42	12
McSorley <i>et al.</i> ⁽⁶⁹⁾	UK	Resectable/322	47	3
Oesophageal/oesophagogastric cancer				
Awad <i>et al.</i> ⁽¹¹⁸⁾	UK	Stages I–III/47	57	1
Yip <i>et al.</i> ⁽⁸⁵⁾	UK	Stages I–III/35	26	1
Reisinger <i>et al.</i> ⁽¹¹⁹⁾	The Netherlands	Stages I–IV/123	56	1
Tamandl <i>et al.</i> ⁽²⁸⁾	Austria	Non-metastatic/200	65	2
Tan <i>et al.</i> ⁽⁸⁴⁾	UK	Stages I–III/89	49	1
Anandavadivelan <i>et al.</i> ⁽⁸³⁾	Sweden	Resectable/72	43	1
Grotenhuis <i>et al.</i> ⁽¹⁶⁶⁾	The Netherlands	Resectable/120	45	1
Nishigori <i>et al.</i> ⁽¹⁶⁷⁾	Japan	Resectable/199	75	1
Elliott <i>et al.</i> ⁽⁴⁰⁾	Ireland	Resectable/252	16	1
Gastric cancer				
Tegels <i>et al.</i> ⁽¹⁶⁸⁾	The Netherlands	Stages I–IV/152	58	3
Hayashi <i>et al.</i> ⁽¹³⁾	Japan	Advanced/53	70	3
Nishigori <i>et al.</i> ⁽¹⁶⁹⁾	Japan	Resectable/157	62	1
Palmela <i>et al.</i> ⁽⁸¹⁾	Portugal	Stages II–III/48	23	3
Kudou <i>et al.</i> ⁽¹⁷⁰⁾	Japan	Stages I–III/148	28	3
Lung cancer				
Baracos <i>et al.</i> ⁽²⁷⁾	Canada	NSC/stages III–IV/441	47	2
Stene <i>et al.</i> ⁽³⁷⁾	Norway	NSC/stages IIIb–IV/35	71	1
Arrieta <i>et al.</i> ⁽¹⁷¹⁾	Mexico	NSC/metastatic/84	69	1
Kim <i>et al.</i> ⁽¹⁰¹⁾	Korea	SC/stages I–III/149	79	2
Kidney cancer				
Antoun <i>et al.</i> ⁽⁷³⁾	France	Metastatic/55	55	2
Antoun <i>et al.</i> ⁽³⁸⁾	Canada	Advanced/80	53	1
Huillard <i>et al.</i> ⁽⁷⁴⁾	France	Metastatic/61	53	2
Sharma <i>et al.</i> ⁽⁹⁶⁾	USA	Metastatic/93	29	3
Auclin <i>et al.</i> ⁽⁵³⁾	France	Metastatic/124	90	2
Cushen <i>et al.</i> ⁽⁵⁶⁾	Ireland	Metastatic/55	33	1
Pancreatic cancer				
Tan <i>et al.</i> ⁽⁹⁵⁾	Canada	Advanced/111	56	1
Dalal <i>et al.</i> ⁽³⁶⁾	USA	Locally advanced/41	63	1
Di Sebastiano <i>et al.</i> ⁽¹⁷²⁾	Canada	Stages IIb–IV/50	48	2
Cooper <i>et al.</i> ⁽¹²⁵⁾	USA	Resectable/89	52	2
Rollins <i>et al.</i> ⁽⁶⁸⁾	UK	Advanced/228	61	3
Wesseltoft-Rao <i>et al.</i> ⁽⁵⁸⁾	Norway	Advanced/45	89	2
Choi <i>et al.</i> ⁽⁵⁷⁾	Korea	Advanced/484	21	7
Liver cancer				
Mir <i>et al.</i> ⁽⁷⁵⁾	France	Advanced/40	28	2
Harimoto <i>et al.</i> ⁽¹⁷³⁾	Japan	Stages I–IV/186	40	4
Fujiwara <i>et al.</i> ⁽³¹⁾	Tokyo	Mixed stages/1257	11	8
Iritani <i>et al.</i> ⁽³²⁾	Japan	Stages I–IV/217	11	9
Voron <i>et al.</i> ⁽⁹⁹⁾	France	Non-metastatic/109	54	1 + 2
Mixed cancer cohorts				
Prado <i>et al.</i> ⁽¹⁹⁾	Canada	Lung and GI/ Stages I–IV/250 obese	15	1
Martin <i>et al.</i> ⁽¹⁾	Canada	Lung and GI/ Stages I–IV/1473	41	3
Veasey-Rodrigues <i>et al.</i> ⁽³⁹⁾	USA	Mixed tumour sites/ Advanced/306	47	1

Table 2. (Cont.)

Author	Country	Stage/n	% Sarcopenic*	Sarcopenia definition†
Daly <i>et al.</i> ⁽¹²⁷⁾	Ireland	Foregut/stage I–IV/225	40	3
Ni Bhuachalla <i>et al.</i> ⁽⁶⁷⁾	Ireland	Mixed tumour sites/stages I–IV/725	41	3
Breast cancer				
Prado <i>et al.</i> ⁽⁷¹⁾	Canada	Metastatic/55	26	1
Del Fabbro <i>et al.</i> ⁽¹⁷⁴⁾	USA	Loco regional/129	14	1
Shachar <i>et al.</i> ⁽⁷²⁾	USA	Metastatic/40	58	3
Rier <i>et al.</i> ⁽¹⁴⁾	The Netherlands	Metastatic/166	67	3
Diffuse large B-cell lymphoma				
Camus <i>et al.</i> ⁽¹⁷⁵⁾	France	NR/80/>70 years	55	5
Lanic <i>et al.</i> ⁽³³⁾	France	NR/82/>70 years	55	5
Other cancers				
Moryoussef <i>et al.</i> ⁽¹⁷⁶⁾	France	GIST/advanced or high-risk resected/31	39	3
Fukushima <i>et al.</i> ⁽¹⁰³⁾	Japan	Urothelial/advanced/88	60	3
Psutka <i>et al.</i> ⁽¹⁰²⁾	USA	Urothelial/ Stages I–III/205	69	2
Cushen <i>et al.</i> ⁽⁶⁶⁾	Ireland	Prostate cancer/ Metastatic/63	47	3
Rutten <i>et al.</i> ⁽⁴³⁾	The Netherlands	Ovarian cancer/ Stages IIb–IV/123	50	10
Coelen <i>et al.</i> ⁽³⁴⁾	The Netherlands	Cholangiocarcinoma/stages I–IV/100	42	11
Daly <i>et al.</i> ⁽⁴⁾	Ireland	Melanoma/metastatic/89	24	3

NSC, non-small cell; SC, small cell; GI, gastrointestinal; NR, not recorded; GIST, gastrointestinal stromal tumour.

* Prevalence of sarcopenia in both sexes combined.

† Sarcopenia definitions stratified as follows:

1: Skeletal muscle index (SMI) <52.4 cm²/m² for males; <38.5 cm²/m² for females⁽¹⁹⁾.

2: SMI <55.4 cm²/m² for males; <38.9 cm²/m² for females (Baumgartner sarcopenia cut points for elderly healthy subjects⁽⁷⁾ converted to CT cut points using regression equations⁽²⁹⁾.

3: SMI <43 cm²/m² BMI ≤24.9 kg/m² and <53 cm²/m² for BMI >25 kg/m² for males; <41 cm²/m² for females⁽¹⁾.

4: SMI <43.75 cm²/m² for males; <41.1 cm²/m² for females⁽⁵⁴⁾.

5: SMI <55.8 cm²/m² for males; <38.9 cm²/m² for females⁽³³⁾.

6: SMI <49.5 cm²/m² and for women <42.1 cm²/m² (lowest quartile)⁽⁴¹⁾.

7: SMI <42.2 cm²/m² for males; <33.9 cm²/m² for females⁽⁵⁷⁾.

8: SMI <36.2 cm²/m² for males; <29.0 cm²/m² for females⁽³¹⁾.

9: SMI <36.0 cm²/m² for males; <29.0 cm²/m² for females⁽³²⁾.

10: SMI <41.5 cm²/m² (median) for females⁽⁴³⁾.

11: SMI <46.8 cm²/m² for males; <39.1 cm²/m² for females⁽³⁴⁾.

12: SMI <52.3 cm²/m² for men and <38.6 cm²/m² for men with a BMI <30 kg/m² and <54.3 cm²/m² for men and <46.6 cm²/m² for women with a BMI ≥30 kg/m²⁽³⁰⁾.

docetaxel (*n* 63), a combination of both low muscle mass and low MA was associated with an increased risk of DLT (59 v. 29 %, *P* = 0.021)⁽⁶⁶⁾. Similarly, in early-stage disease, Shachar *et al.*⁽⁸²⁾ demonstrated that in breast cancer (*n* 151), the risk of hospitalisations due to chemotherapy toxicity increased 19 % with every five-unit decrease in MA (relative risk 1.19 (95 % CI 1.00, 1.43), *P* = 0.05).

Increased toxicity in patients with sarcopenia may be attributed to alterations in the distribution, metabolism and clearance of systemic chemotherapy drugs⁽⁸⁰⁾. The practice of administration of cytotoxic chemotherapy based on body surface area, and targeted therapy as a flat dose, ignores several sources of inter-individual variation. Using body surface area as the only method to individualise chemotherapy drug dose is insufficient to avoid severe toxicity, but the continued use mainly relies on the lack of other more precise methods for dose individualisation⁽⁸⁹⁾. It is recently acknowledged that variability in body composition (lean mass, fat mass and total body water) of cancer patients may be a source of disparities in the metabolism of cytotoxic agents resulting in increased toxicity^(73,80,90). The rationale is that body weight comprises two major compartments, lean mass

and fat mass, which may be the two major sites of distribution of hydrophilic and lipophilic drugs, respectively^(80,91). Therefore, changes in body composition may lead to changes in the volume of distribution and adversely impact the effectiveness and tolerance of cancer therapies.

Administration of hydrophilic chemotherapy drugs that are mainly distributed to the lean mass compartment in sarcopenic patients would result in a disproportionately small volume of the drug distribution in relation to their body weight or body surface area^(71,80). This has been hypothesised to lead to considerable variation in milligram of chemotherapeutic agent per kg lean mass, and a higher dose per kg lean mass is associated with more frequent severe toxic side effects^(76,80,92,93). In patients with advanced non-small-cell lung cancer (*n* 424), when the dose of non-platinum drugs was expressed as mg/kg lean mass, a 3-fold range was observed, and the dose of non-platinum chemotherapeutic agent per kg lean mass was a significant predictor of haematologic toxicity. Every 1 % increase in drug dose per kg lean mass above the mean was associated with a 3 % increased risk of grade III–IV haematologic toxicity, while patients with




Fig. 1. Sarcopenic patients experience more dose-limiting toxicities (DLT) to fluorouracil (5-FU) in colon cancer⁽⁸⁰⁾; capecitabine in breast cancer⁽⁷¹⁾; sorafenib in renal cell carcinoma⁽⁷³⁾; 5-fluorouracil, epirubicin, cyclophosphamide (FEC) in breast cancer⁽⁹⁰⁾; sorafenib in hepatocellular carcinoma⁽⁷⁵⁾; sunitinib in renal cell carcinoma⁽⁷⁴⁾; vandetanib in medullary thyroid cancer⁽⁷⁸⁾; fluoropyrimidine in colorectal cancer⁽⁶⁵⁾; phase I drugs in mixed cancer types⁽¹⁷⁷⁾; imatinib in gastrointestinal stromal tumours (grade I-II toxicity)⁽¹⁷⁶⁾; epirubicin, cisplatin, capecitabine (ECX) and cisplatin, 5-fluorouracil (CF) in oesophagogastric cancer⁽⁸⁴⁾; taxane-based chemotherapy (paclitaxel, docetaxel, nab-paclitaxel) in breast cancer⁽⁷²⁾; hyperthermic intraperitoneal chemotherapy (oxaliplatin and irinotecan) in colorectal cancer⁽⁷⁷⁾; neoadjuvant chemotherapy (NACT) (mixed types) in gastric cancer⁽⁸¹⁾; ipilimumab in metastatic melanoma⁽⁴⁾.

doses >20 % above the mean were at almost double the risk of experiencing a grade III–IV haematologic toxicity⁽⁹³⁾. Pharmacokinetic data have supported this hypothesis, with sarcopenic patients experiencing higher plasma concentrations of antineoplastic drugs, and experiencing more toxicity^(75,78). The same hypothesis holds true for toxicity to lipophilic drugs. In a study of ovarian cancer patients receiving doxorubicin and trabectedin, the risk of DLT decreased with increased fat mass:lean mass ratio. As lipophilic drugs are mainly distributed in the adipose tissue, leaner individuals would present with reduced volume of distribution for the drug and consequently increase their risk for DLT⁽⁹¹⁾.

In addition to the pharmacokinetic hypothesis to explain the increased toxicity in sarcopenic patients, other mechanisms have been suggested. Low levels of lean mass and altered concentration of plasma proteins (e.g. albumin) may affect the distribution of highly protein-bound drugs, and may explain the increased toxicity of the vandetanib⁽⁷⁸⁾, sorafenib⁽⁷⁵⁾ and epirubicin⁽⁹⁰⁾. Systemic inflammation is known to inhibit hepatic enzymes and may contribute to higher drug exposure, and subsequently excess toxicity in patients with sarcopenia^(75,78). In addition, sarcopenic patients are generally more susceptible to acute medical events and perhaps chemotherapy toxicity may be an additional generalised intolerance⁽⁹⁴⁾. Whether altered chemotherapy dosing based on lean mass in patients receiving treatment is

effective in preventing toxicity is currently being investigated (ClinicalTrials.gov identifier: NCT01624051).

Sarcopenia and survival

The impact of sarcopenia on survival in oncology was first identified in a cohort of 250 obese lung and GI cancer patients by Prado *et al.* in 2008⁽¹⁹⁾. Within this study, sarcopenic obese patients had a lower median overall survival compared with their non-sarcopenic counterparts (11.3 v. 21.6 months, $P < 0.001$), and sarcopenic obesity independently predicted survival when adjusted for known prognostic covariates (hazard ratio (HR) 4.2 (95 % CI 2.4, 7.2), $P < 0.0001$)⁽¹⁹⁾. Since then, sarcopenia has repeatedly been shown to be independently prognostic of reduced survival in a number of cancer sites and stages including pancreatic⁽⁹⁵⁾, kidney^(96,97), liver^(98–100), lung⁽¹⁰¹⁾, oesophageal⁽²⁸⁾, colorectal^(41,54,70), urothelial^(102,103) and lung and GI cancers⁽¹⁾. Its important prognostic ability was reported in a study of 196 patients following resection for colorectal cancer liver metastases, whereby sarcopenic patients had a median survival of 24 months, which was in stark contrast to 60 months in those without sarcopenia ($P < 0.001$)⁽⁵⁴⁾. In patients with hepatocellular carcinoma (n 116), similar results were obtained with a median survival of 16 months in patients with sarcopenia compared with 28 months in those without sarcopenia ($P = 0.003$)⁽⁹⁸⁾. More recently, a large cohort of stage I–III colorectal cancer patients

(*n* 3262) showed that sarcopenic patients had a 27 % higher risk of overall mortality than their non-sarcopenic counterparts⁽³⁰⁾.

The impact of sarcopenia on survival has been extensively reviewed previously^(104–106) and of note is a recent systematic review and meta-analysis⁽¹⁰⁴⁾, which examined the prognostic value of sarcopenia in 7843 patients with solid tumours from a total of thirty-eight studies. Sarcopenia was found to be associated with poor overall survival (HR 1.44 (95 % CI 1.32, 1.56), $P < 0.001$), the effect of which remained in various tumour types and across disease stages. Additionally, sarcopenia was associated with cancer-specific survival (HR 1.93 (95 % CI 1.38, 2.70), $P < 0.001$), as well as disease-free survival (HR 1.16 (95 % CI 1.00, 1.30), $P = 0.014$). However, it is noteworthy to mention the limitations of this study. The authors included several studies in the review that contained duplicated data, and this is prohibited in a meta-analysis as it may lead to overestimation of an effect⁽¹⁰⁷⁾. In addition, the studies were largely heterogeneous in terms of the tumours considered, stage of disease, treatments and cut points used to define sarcopenia, which ranged from 29 to 42.1 cm²/m² in women and 36 to 55.4 cm²/m² in men⁽¹⁰⁴⁾.

In spite of the abundance of evidence supporting the role of sarcopenia as an important and independent prognostic factor in cancer, this finding has not always been consistently reported and several studies have reported no impact on survival^(16–18,37,68,88). It is possible that the impact of sarcopenia may vary depending on cancer diagnosis, treatment and overall prognosis. For example, in colorectal cancer patients, sarcopenia was shown to be predictive of survival in patients undergoing curative resection⁽⁴¹⁾, but not in patients with unresectable disease receiving chemotherapy⁽¹⁰⁸⁾. Alternatively, this may be a consequence of the choice of cut points to define sarcopenia in many of these studies. As discussed earlier, cut points obtained in different studies are highly dependent on the methodology and the population from which they were devised, and when used in dissimilar populations, they may represent a suboptimal approach in identifying the true prevalence of low muscle mass and its relationship to survival within these cohorts.

Muscle quality v. quantity

Low MA, often referred to as myosteatosis, is emerging as an important prognostic indicator in patients with cancer and in some cases more superior to predicting survival compared with muscle mass alone^(12–18). The first report by Sabel *et al.*⁽¹⁰⁹⁾ in melanoma patients showed poorer disease-free survival ($P = 0.04$) and distant disease-free survival ($P = 0.0002$) in patients with the lowest tertile of psoas MA. Antoun *et al.*⁽⁶⁴⁾ corroborated these findings in a cohort of metastatic renal cell carcinoma patients, and reported that the median survival of patients with a MA below the median (<38 HU in men; <36 HU in women) was half that of those with MA above the median (29 v. 14 months, $P = 0.001$). In a large cohort of 1473 patients with lung and GI cancer, Martin *et al.*⁽¹⁾ reported that patients with a MA below

the identified cut point (<41 HU for BMI <25 kg/m² and <33 HU for BMI ≥25 kg/m²) were at a 36 % increased risk of mortality (HR 1.36 (95 % CI 1.19, 1.55), $P < 0.001$). Recently, low MA has been shown to be prognostic of reduced survival in a variety of different tumours, including lung^(18,60), kidney⁽⁶⁴⁾, breast⁽¹⁴⁾, gastric⁽¹³⁾, pancreatic^(17,65), periampullary⁽¹⁶⁾, cholangiocarcinoma⁽⁶⁸⁾, lymphoma^(12,15) and melanoma⁽¹⁰⁹⁾. In a cohort of 734 patients with advanced lung cancer, Sjöblom *et al.*⁽¹⁸⁾ reported that even minor changes in MA are associated with mortality risk. An incremental increase of 1 HU was associated with a 2 % reduction in the risk of death (HR 0.98 (95 % CI 0.97, 0.99), $P < 0.001$). Accordingly, the mortality risk reduction of a 5 HU and 10 HU increase in MA were 8 % and 16 %, respectively⁽¹⁸⁾.

Sarcopenia, low muscle attenuation and post-operative outcomes

Sarcopenia has also been identified as a predictor of post-operative infections, complications, readmissions, longer length of hospital stay and higher health care costs^(3,100,110–114). Elliott *et al.*⁽⁴⁰⁾ demonstrated that sarcopenia, in a cohort of 252 patients with oesophageal cancer undergoing resection, was independently associated with increased risk of major postoperative complications (grade ≥IIIb; OR 5.30 (95 % CI 1.94, 14.45), $P = 0.001$) and pulmonary complications (OR 2.17 (95 % CI 1.12, 4.23), $P = 0.023$). Similarly, in patients with colorectal cancer undergoing cytoreductive surgery for peritoneal carcinomatosis (*n* 206), sarcopenia was associated with severe post-operative complications, and sarcopenic patients underwent significantly more reoperations (26 v. 12 %, $P = 0.012$)⁽³⁾. Consequently, length of hospital stay has shown to be significantly longer in patients with sarcopenia^(40,46,96,110,111,115). Following colorectal cancer surgery (*n* 234), the length of hospital stay for sarcopenic patients was 15.9 d compared with 12.3 d in patients without sarcopenia ($P = 0.038$)⁽¹¹⁰⁾. Ultimately, adverse post-operative outcomes in patients with sarcopenia lead to increased health care costs. In a Western-Europe healthcare system, sarcopenia was independently associated with increased hospital costs of €4061 per patient ($P = 0.015$) in a study of 452 patients who underwent cancer surgery of the alimentary tract⁽¹¹⁵⁾.

Akin to sarcopenia, low MA is emerging as a predictor of poor post-operative outcomes^(16,44,109,116). In patients with colon cancer undergoing open resection (*n* 91), low MA was an independent risk factor for one or more complications ($P < 0.001$)⁽⁴⁴⁾. Van Rijssen *et al.*⁽¹⁶⁾ reported similar findings, whereby patients with low MA experienced significantly more major post-operative complications (58 v. 37 %, $P = 0.005$) following resection for periampullary cancer (*n* 166) compared with those without low MA. Examining MA as a continuous variable, Sabel *et al.*⁽¹⁰⁹⁾ reported that a 10 HU decrease in psoas MA was associated with an 8.1 % increase in complication rate in patients with stage III melanoma (*n* 101). In line with this, low MA was associated with prolonged length of hospital stay following colorectal cancer

surgery in a large cohort of 805 patients (7 v. 6 d ($P = 0.034$))⁽⁷⁰⁾, and to a greater extent following pancreatic cancer resection, whereby patients with low MA (post-neoadjuvant chemoradiotherapy) had a mean length of stay of 42 d compared with 23 d in those without low MA ($P = 0.001$)⁽⁶⁵⁾.

Muscle loss during chemotherapy

The precision associated with CT analysis of body composition (1–1.5 %) has allowed recent investigations to focus on the nature and magnitude of changes in body composition during the disease trajectory in patients with cancer. Studies have consistently shown that losses in muscle mass and MA are exacerbated by antineoplastic treatment^(81,88,117–119). In oesophagogastric cancer patients, neoadjuvant chemotherapy treatment which is typically delivered over a 6–8-week period can decrease SMA by 9.59 cm² ($P < 0.0001$), which is the equivalent to a loss of 2.9 kg of lean mass⁽¹¹⁸⁾. These results have been corroborated in a study of 252 oesophageal cancer patients, whereby the prevalence of sarcopenia almost doubled (15.9–30.8 %) during a course of neoadjuvant chemotherapy⁽⁴⁰⁾.

In patients with cancer, muscle is lost at a very high rate of 3–5 % per 100 d during systemic chemotherapy^(4,36,43,95), and losses are exponentially increased with progressive disease and proximity to death^(117,120). In a recent study of metastatic colorectal cancer patients (n 63), on average patients lost muscle at a rate of 6.1 % during 3 months of chemotherapy⁽⁸⁸⁾; it is noteworthy that the rate of decline is 24-fold more rapid than that observed in healthy ageing adults who tend to lose muscle at a rate of 1–1.4 % per year^(6,121).

Alterations in muscle mass may be a consequence of some cancer-directed therapies. Sorafenib, a multi-kinase inhibitor, has been shown to provoke muscle wasting in metastatic renal cell carcinoma patients, through the downstream suppression of PI3K, AKTm and mTOR, key mediators in activating muscle protein synthesis by amino acids and other stimuli⁽³⁸⁾. Abiraterone⁽¹²²⁾ and MK-0646 (anti-insulin-like growth factor 1 receptor)⁽¹²³⁾ have also been reported to stimulate muscle loss through maximal androgen suppression and inhibition of anti-insulin-like growth factor 1 receptor, respectively. In contrast, two chemotherapy agents (vandetanib and selumetinib) have resulted in significant muscle gain in patients with advanced cancer^(78,124).

Loss of muscle during systemic anti-cancer treatment is associated with increased mortality in patients with pancreatic^(36,125), lung^(37,126), colorectal^(88,108), ovarian⁽⁴³⁾, melanoma⁽⁴⁾ and foregut cancer⁽¹²⁷⁾ (see Table 3 for a summary of available studies reporting that muscle loss during treatment is prognostic of reduced survival). Loss of muscle >2 %/100 d was independently associated with reduced survival in ovarian cancer patients receiving neoadjuvant chemotherapy (n 123; HR 1.77 (95 % CI 1.02, 3.09), $P = 0.043$)⁽⁴³⁾, and we have previously reported that in patients with advanced cancers of the foregut, those with a skeletal muscle loss >6 %/100 d are at more than double the risk of mortality (HR 2.66 (95 % CI 1.42,

4.97), $P = 0.002$)⁽¹²⁷⁾. Importantly, skeletal muscle depletion during chemotherapy has also been associated with reductions in physical function in elderly patients with advanced non-small-cell lung cancer⁽¹²⁸⁾.

Strategies to improve muscle mass and attenuation

Currently, there is no consensus treatment for attenuating or reversing the muscle wasting caused by sarcopenia and/or cachexia in patients with cancer. Research to date has focused on strategies to treat sarcopenia in the context of cancer cachexia using nutritional, exercise and pharmacological interventions. However, these single-agent therapies have not provided promising results, and perhaps multifaceted treatment strategies may be more effective in treating cachexia. Ideally, interventions for cachexia should be initiated in the early stages of weight loss (i.e. pre-cachexia); however, patients are often referred for cachexia interventions late in their disease trajectory (i.e. refractory cachexia). At this stage, patients are in a catabolic state and respond poorly to anti-cancer treatment⁽⁸⁾. Experts have speculated that the failure of treatments to show benefit at clinical evaluation may not have been because of the drugs ability to ameliorate or treat cancer cachexia, but because they were introduced to patients too late⁽¹²⁹⁾.

Rehabilitation care to modify body composition, either increasing muscle mass and/or MA should be conducted, and its respective impact on oncology outcomes explored. Previous investigations have shown that intramuscular adipose tissue was responsive to exercise in non-cancer individuals⁽¹³⁰⁾, and that cancer patients have anabolic potential⁽¹¹⁷⁾. Current evidence in evaluating the efficacy of prehabilitation on postoperative outcomes among cancer patients is still limited^(131,132). Future clinical trials are warranted to test whether the window of time between cancer diagnosis and initiation of treatment is an opportunity to commence interventions to increase muscle mass and MA.

Nutritional interventions

A meta-analysis of oral nutritional interventions in malnourished patients with cancer identified thirteen randomised controlled trials and included 1414 patients. The analysis, conducted by Baldwin *et al.*⁽¹³³⁾, concluded that oral nutritional interventions had no significant effect on body weight, energy intake or mortality compared with standard care. Given the wide range of pathophysiological alterations that occur in cancer, complex and individualised targeted strategies incorporating modulations of the metabolic components of cachexia (e.g. inflammation) may be required to allow nutrition interventions to be more effective⁽¹³⁴⁾. To date, n -3 fatty acids, such as EPA have received a lot of attention for the treatment of cancer cachexia due to their potent anti-inflammatory properties. In cancer, plasma n -3 fatty acid levels appear to be depleted in patients with sarcopenia and may contribute to accelerate muscle mass⁽¹³⁵⁾. The results of several small clinical trials suggested that

Table 3. Summary of studies reporting muscle loss during treatment as prognostic of reduced survival

Author	Cancer type/ stage/n	Treatment	Time between scans	Main findings
Dalal <i>et al.</i> ⁽³⁶⁾	Pancreatic/ advanced/41	Bevacizumab, capecitabine	Median 104 d (IQR 97–112)	Median loss of skeletal muscle was 3.8 % per 100 d ($P = 0.003$). Loss of skeletal muscle >3.8 % was associated with poor OS (10.1 v. 16.3 months, $P = 0.02$) on univariate analysis
Stene <i>et al.</i> ⁽³⁷⁾	NSC lung/ advanced/35	Carboplatin, vinorelbine, gemcitabine	Median 88 $\pm$ 22 d	Mean reduction in SMA was 4.2 cm ² (95 % CI -7.3 , -1.9 , $P < 0.002$) from baseline to follow-up. A trend that loss of muscle >2 % was associated with reduced survival (5.8 v. 10.7 months, $P = 0.073$)
Cooper <i>et al.</i> ⁽¹²⁵⁾	Pancreatic/ resectable/89	Gemcitabine, cisplatin	Median 4.2 months (min 2.2 months, max 11.5 months)	Mean decrease in SMI was 1.2 cm ² /m ² ($P < 0.01$). Degree of muscle loss correlated with disease-free survival (HR 0.89 (95 % CI 0.80, 1.00), $P = 0.04$), while visceral adipose loss was associated with overall survival (HR 0.97 (95 % CI 0.95, 0.99), $P = 0.001$) and progression-free survival (HR 0.98 (95 % CI 0.96, 0.99), $P = 0.01$) on univariate analysis
Miyamoto <i>et al.</i> ⁽¹⁰⁸⁾	Colorectal/ unresectable/148	Oxaliplatin, irinotecan base chemotherapy	Median 2.1 months (IQR 1–3 months)	Median change in skeletal muscle was 4.2 %. Skeletal muscle loss >5 % (highest quartile) independently predicted reduced OS (HR 2.1 (95 % CI 1.19, 3.62); $P = 0.010$)
Blauwhoff Buskermolen <i>et al.</i> ⁽⁸⁸⁾	Colorectal/ metastatic/63	Capecitabine, irinotecan, bevacizumab	Median 78 d (IQR 67–92)	On average patients lost 6.1 % of SMA in 3 months. Muscle loss >9 % (highest tertile) was significantly associated with lower survival on multivariate analysis (HR 4.47 (95 % CI 2.21, 9.05), $P < 0.001$)
Rutten <i>et al.</i> ⁽⁴³⁾	Ovarian/stages II–IV/123	Standard neoadjuvant chemotherapy	84 d (SE $\pm$ 1.77)	Mean change in SMA was -5.2 ± 9.8 %/100 d. Change in SM >2 % was significantly associated with reduced survival on multivariate analysis (HR 1.77 (95 % CI 1.02, 3.09), $P = 0.043$)
Nattenmuller <i>et al.</i> ⁽¹²⁶⁾	Lung/stages Ib–IV/200	Mixed type standard chemotherapy	Mean 4.3 months	Loss of SMA and SMI from baseline to follow-up scans was 4.1 cm ² ($P < 0.001$) and 1.4 cm ² /m ² ($P < 0.001$), respectively. Decrease in SMI was associated with poor survival (HR 1.063; $P < 0.001$)
Daly <i>et al.</i> ⁽⁴⁾	Melanoma/stage IV/59	Ipilimumab	Mean 146 $\pm$ 40 d	Mean loss of SMA was 3.3 ± 5.8 %/100 d. A loss of SMA >7.5 % was a significant predictor of overall survival in multivariable (HR 2.1 (95 % CI 1.1, 4.6), $P = 0.046$)
Daly <i>et al.</i> ⁽¹²⁷⁾	Foregut/stages I–IV/163	Mixed type standard chemotherapy	Median 118 d (IQR 92 to 158)	Mean loss of SMA was 3.9 % (95 % CI -4.9 , -2.8)/100 d. In patients receiving palliative chemotherapy ($n = 89$), a loss of SMA >6 %/100 d independently predicted survival (HR 2.66 (95 % CI 1.42, 4.97), $P = 0.002$)

IQR, interquartile range; NSC, non-small cell; SM, skeletal muscle; SMA, skeletal muscle area; HR, hazard ratio; SMI, skeletal muscle index; SE, standard error; OS, overall survival.

using EPA supplements or oral nutritional supplements containing EPA in patients with advanced cancer improved appetite, energy intake, body weight, lean mass and/or physical activity^(136–139). The n -3 fatty acid supplementation has also been shown to be effective in reducing intramuscular adipose tissue while maintaining muscle mass (compared with standard care) in a small cohort of patients with non-small-cell lung cancer⁽¹³⁸⁾, supporting what is observed in preclinical experimental animal models⁽¹⁴⁰⁾. However, the clinical evidence supporting n -3 fatty acids remain inconclusive. Three systematic reviews published in 2007, 2009 and 2012 concluded that there was insufficient evidence to support a recommendation for n -3 fatty acids to treat muscle depletion and cancer cachexia^(141–144). Due to the inconsistencies in the reported effects of n -3 fatty acids^(145,146), the ESPEN oncology nutrition guidelines issue only a

weak recommendation for their use to stabilise or improve appetite, food intake, lean mass and body weight⁽¹⁴⁴⁾.

Achieving adequate protein intake is an essential component to achieving muscle anabolism; however, no optimal amino acid and protein requirements have been established to prevent or treat sarcopenia in patients with cancer. Current recommendations suggest intakes of 1–1.5 g/kg/d in patients with cancer⁽¹⁴⁴⁾; however, many patients fail to reach these requirements⁽¹⁴⁷⁾. Moreover, recent research has suggested that protein requirements should be based on the measures of lean mass as opposed to body weight^(148,149).

Physical activity

A recent meta-analysis⁽¹⁵⁰⁾ has shown resistance exercise to be effective in increasing muscle strength and mass in

cancer patients undergoing neoadjuvant and adjuvant therapy, with mean increases in lower limb strength of 26.2 kg ($P=0.00001$) and lean mass by 0.8 kg ($P<0.00001$)⁽¹⁵⁰⁾. In a study of early-stage breast cancer patients receiving adjuvant chemotherapy, resistance exercise resulted in the reversal of sarcopenia and dynapenia in 43 and 59 % of patients, respectively⁽¹⁵¹⁾. To date, most studies have been conducted in early-stage breast and prostate cancer and evidence of the effect of physical activity on muscle mass and strength in patients with advanced disease and cancer cachexia is lacking^(152,153). A recent Cochrane systemic review⁽¹⁵³⁾, which aimed to examine the impact of exercise on lean mass in patients with cancer cachexia, showed disappointing results. The authors screened more than 3000 individual articles, but found no trials that met the inclusion criteria for analysis⁽¹⁵³⁾. Initial findings indicate that exercise is safe and well tolerated in patients with advanced disease⁽¹⁵⁴⁾ and improves physical performance and several domains of quality of life⁽¹⁵⁵⁾, but further work is needed. Evidence of interventions aimed at improving MA in patients with cancer are scarce; however, evidence in older adults suggests that exercise may represent a counter measure to improve MA, as studies have shown that muscle fatty infiltration is amenable to change after 12 weeks of thrice-weekly resistance training⁽¹³⁰⁾.

Recently, a few studies have demonstrated that exercise interventions are capable of improving cancer outcomes. Cho *et al.*⁽¹⁵⁶⁾ reported that preoperative exercise significantly improved operative risk factors and decreased the frequency of serious postoperative complications in gastric cancer patients. In breast cancer, a supervised moderate- to high-intensity exercise programme during adjuvant chemotherapy was associated with a beneficial effect on chemotherapy completion rates with a significantly smaller proportion of patients requiring dose adjustments compared with usual care (12 v. 34 %) ⁽¹⁵⁷⁾. The impact of exercise prehabilitation on cancer outcome warrants further investigation.

Pharmacological agents

To date, no pharmacotherapies have been approved to specifically treat the cancer cachexia syndrome. Non-steroidal anti-inflammatory drugs have been suggested as a potential treatment for cancer cachexia, with the aim of reducing systemic inflammation. In a recent systematic review of thirteen studies, eleven showed either improvement or stabilisation in weight or lean mass in patients with cancer⁽¹⁵⁸⁾; however, the evidence is still insufficient to recommend non-steroidal anti-inflammatory drugs to treat cancer cachexia outside of clinical trials. Promising results have been obtained for anamorelin (a ghrelin receptor agonist); phase III trials (ROMANA 1 and ROMANA 2) have shown that anamorelin significantly increases lean mass and improves anorexia/cachexia symptoms compared with placebo over a 12-week intervention period. However, muscle function (measured by hand grip strength) failed to improve during the intervention⁽¹⁵⁹⁾. Food and drug administration regulators generally require improvements in lean mass and

functional outcomes as co-primary endpoints for the approval of new medications to treat cancer cachexia. As a result, anamorelin has not received the food and drug administration's approval to date. Phase III trials have recently been completed for enobosarm, a selective androgen receptor modulator (NCT01355484), and MABp1, an anti IL-1 α monoclonal antibody (NCT01767857), the results of which are yet to be published.

Multimodal interventions

Results so far suggest that single-agent therapies may be insufficient to counteract cancer cachexia and that early multimodal interventions may be necessary to combat its multifactorial and complex pathogenesis. A multimodal approach including the use of individualised nutritional interventions to promote energy balance and ensure optimal protein intake, decreasing inflammation and hyper metabolic stress with the aid of anti-inflammatory agents (non-steroidal anti-inflammatory drugs and EPA) and increasing physical activity to stabilise/increase muscle mass, strength and physical performance has been recommended⁽¹³⁴⁾. One such intervention currently under investigation is the MENAC trial (Multimodal Exercise/Nutrition/Anti-inflammatory treatment for Cachexia), whereby phase II feasibility studies have yielded encouraging results and suggest multimodal interventions are feasible and safe in patients with cancer⁽¹⁶⁰⁾. The phase III trial (NCT02330926) is currently being conducted across a number of international sites.

Conclusion

In summary, the evidence reviewed here shows that muscle abnormalities are highly prevalent in adult patients with cancer, across cancer sites and stages, and overweight and obesity do not preclude their presence. However, the heterogeneity in the diagnostic criteria limits the ability to accurately compare the reported prevalence rates between different cohorts and study findings overall. Efforts are required to standardise muscle measurements from CT images, as well as the diagnostic criteria for sarcopenia and low MA in patients with cancer. Nonetheless, the clinical importance of muscle abnormalities in these patients is evident, given their associations with negative clinical outcomes, such as poorer tolerance to chemotherapy, increased risk of post-operative complications and infections, increased length of hospital stays and poor prognosis.

In an age of increasingly personalised medicine, CT scans, obtained as part of routine patient care, provide valuable individualised information on muscle mass and MA with well-established prognostic implications. Incorporating routine assessment of muscle abnormalities from CT images into clinical practice has the potential to play a major role in stratifying patients at risk of poorer clinical outcomes, who may benefit from targeted interventions. At present, no effective medical intervention to improve muscle mass and MA exists; however,

in recent years, substantial progress has been made, with the results of several clinical trials awaited in the near future. The optimal timing and treatment strategy for preventing or delaying the development of muscle abnormalities are yet to be determined, but multimodal interventions appear to hold the most promise.

Supplementary material

The supplementary material for this article can be found at <https://doi.org/10.1017/S0029665118000046>

Acknowledgements

The authors would like to thank the Irish section of the Nutrition Society for inviting the present review paper as part of the postgraduate review competition. The authors would also like to acknowledge the support of the Health Research Board Clinical Research Facility, Cork.

Financial Support

This publication has emanated from research conducted with the financial support of Science Foundation Ireland (SFI) under grant number SFI/12/RC/2273.

Conflict of interest

None.

Authorship

L. E. D. reviewed the literature and wrote the manuscript. A. M. R. and C. M. P. participated in writing and revising the manuscript. All authors approved the final manuscript.

References

- Martin L, Birdsell L, Macdonald N *et al.* (2013) Cancer cachexia in the age of obesity: skeletal muscle depletion is a powerful prognostic factor, independent of body mass index. *J Clin Oncol* **31**, 1539–1547.
- Cespedes Feliciano EM, Lee VS, Prado CM *et al.* (2017) Muscle mass at the time of diagnosis of nonmetastatic colon cancer and early discontinuation of chemotherapy, delays, and dose reductions on adjuvant FOLFOX: The C-SCANS study. *Cancer* **123**, 4868–4877.
- van Vugt JL, Braam HJ, van Oudheusden TR *et al.* (2015) Skeletal muscle depletion is associated with severe post-operative complications in patients undergoing cytoreductive surgery with hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis of colorectal cancer. *Ann Surg Oncol* **22**, 3625–3631.
- Daly LE, Power DG, O'Reilly Á *et al.* (2017) The impact of body composition parameters on ipilimumab toxicity and survival in patients with metastatic melanoma. *Br J Cancer* **116**, 310–317.
- Janssen I & Ross R (2005) Linking age-related changes in skeletal muscle mass and composition with metabolism and disease. *J Nutr Health Aging* **9**, 408–419.
- Goodpaster BH, Park SW, Harris TB *et al.* (2006) The loss of skeletal muscle strength, mass, and quality in older adults: the health, aging and body composition study. *J Gerontol A Biol Sci Med Sci* **61**, 1059–1064.
- Baumgartner RN, Koehler KM, Gallagher D *et al.* (1998) Epidemiology of sarcopenia among the elderly in New Mexico. *Am J Epidemiol* **147**, 755–763.
- Fearon K, Strasser F, Anker SD *et al.* (2011) Definition and classification of cancer cachexia: an international consensus. *Lancet Oncol* **12**, 489–495.
- Argilés JM, Busquets S, Stemmler B *et al.* (2014) Cancer cachexia: understanding the molecular basis. *Nat Rev Cancer* **14**, 754–762.
- Donohoe CL, Ryan AM & Reynolds JV (2011) Cancer cachexia: mechanisms and clinical implications. *Gastroenterol Res Pract* **2011**, 601434.
- Aubrey J, Esfandiari N, Baracos VE *et al.* (2014) Measurement of skeletal muscle radiation attenuation and basis of its biological variation. *Acta Physiol (Oxf)* **210**, 489–497.
- Chu MP, Lieffers J, Ghosh S *et al.* (2015) Skeletal muscle radio-density is an independent predictor of response and outcomes in follicular lymphoma treated with chemoimmunotherapy. *PLoS ONE* **10**, e0127589.
- Hayashi N, Ando Y, Gyawali B *et al.* (2016) Low skeletal muscle density is associated with poor survival in patients who receive chemotherapy for metastatic gastric cancer. *Oncol Rep* **35**, 1727–1731.
- Rier HN, Jager A, Sleijfer S *et al.* (2017) Low muscle attenuation is a prognostic factor for survival in metastatic breast cancer patients treated with first line palliative chemotherapy. *Breast* **31**, 9–15.
- Chu MP, Lieffers J, Ghosh S *et al.* (2017) Skeletal muscle density is an independent predictor of diffuse large B-cell lymphoma outcomes treated with rituximab-based chemoimmunotherapy. *J Cachexia Sarcopenia Muscle* **8**, 298–304.
- Van Rijssen LB, van Huijgevoort NC, Coelen RJ *et al.* (2017) Skeletal muscle quality is associated with worse survival after pancreatoduodenectomy for periampullary, nonpancreatic cancer. *Ann Surg Oncol* **24**, 272–280.
- van Dijk DP, Bakens MJ, Coolsen MM *et al.* (2017) Low skeletal muscle radiation attenuation and visceral adiposity are associated with overall survival and surgical site infections in patients with pancreatic cancer. *J Cachexia Sarcopenia Muscle* **8**, 317–326.
- Sjöblom B, Grønberg BH, Wentzel-Larsen T *et al.* (2016) Skeletal muscle radiodensity is prognostic for survival in patients with advanced non-small cell lung cancer. *Clin Nutr* **35**, 1386–1393.
- Prado CM, Lieffers JR, McCargar LJ *et al.* (2008) Prevalence and clinical implications of sarcopenic obesity in patients with solid tumours of the respiratory and gastrointestinal tracts: a population-based study. *Lancet Oncol* **9**, 629–635.
- Ryan AM, Power DG, Daly L *et al.* (2016) Cancer-associated malnutrition, cachexia and sarcopenia: the skeleton in the hospital closet 40 years later. *Proc Nutr Soc* **75**, 199–211.
- Prado CM & Heymsfield SB (2014) Lean tissue imaging: a new era for nutritional assessment and intervention. *J Parenter Enteral Nutr* **38**, 940–953.

22. Shen W, Punyanitya M, Wang Z *et al.* (2004) Total body skeletal muscle and adipose tissue volumes: estimation from a single abdominal cross-sectional image. *J Appl Physiol* (1985) **97**, 2333–2338.
23. Cruz-Jentoft AJ, Baeyens JP, Bauer JM *et al.* (2010) Sarcopenia: European consensus on definition and diagnosis: report of the European working group on sarcopenia in older people. *Age Ageing* **39**, 412–423.
24. Fielding RA, Vellas B, Evans WJ *et al.* (2011) Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc* **12**, 249–256.
25. Prado CM, Lieffers JR, Bowthorpe L *et al.* (2013) Sarcopenia and physical function in overweight patients with advanced cancer. *Can J Diet Pract Res* **74**, 69–74.
26. Prado CM, Baracos VE, Reiman T *et al.* (2010) Overweight and obese patients with solid tumors may have sarcopenia, poor prognosis and early features of cachexia. *International Journal of Body Composition Research* **8**, 7–15.
27. Baracos VE, Reiman T, Mourtzakis M *et al.* (2010) Body composition in patients with non-small cell lung cancer: a contemporary view of cancer cachexia with the use of computed tomography image analysis. *Am J Clin Nutr* **91**, 1133S–1137S.
28. Tamandl D, Paireder M, Asari R *et al.* (2016) Markers of sarcopenia quantified by computed tomography predict adverse long-term outcome in patients with resected oesophageal or gastro-oesophageal junction cancer. *Eur Radiol* **26**, 1359–1367.
29. Mourtzakis M, Prado CM, Lieffers JR *et al.* (2008) A practical and precise approach to quantification of body composition in cancer patients using computed tomography images acquired during routine care. *Appl Physiol Nutr Metab* **33**, 997–1006.
30. Caan BJ, Meyerhardt JA, Kroenke CH *et al.* (2017) Explaining the obesity paradox: the association between body composition and colorectal cancer survival (C-SCANS study). *Cancer Epidemiol Biomarkers Prev* **26**, 1008–1015.
31. Fujiwara N, Nakagawa H, Kudo Y *et al.* (2015) Sarcopenia, intramuscular fat deposition, and visceral adiposity independently predict the outcomes of hepatocellular carcinoma. *J Hepatol* **63**, 131–140.
32. Iritani S, Imai K, Takai K *et al.* (2015) Skeletal muscle depletion is an independent prognostic factor for hepatocellular carcinoma. *J Gastroenterol* **50**, 323–332.
33. Lanic H, Kraut-Tauzia J, Modzelewski R *et al.* (2014) Sarcopenia is an independent prognostic factor in elderly patients with diffuse large B-cell lymphoma treated with immunochemotherapy. *Leuk Lymphoma* **55**, 817–823.
34. Coelen RJ, Wiggers JK, Nio CY *et al.* (2015) Preoperative computed tomography assessment of skeletal muscle mass is valuable in predicting outcomes following hepatectomy for perihilar cholangiocarcinoma. *HPB (Oxford)* **17**, 520–528.
35. Heymsfield SB, Gonzalez MC, Lu J *et al.* (2015) Skeletal muscle mass and quality: evolution of modern measurement concepts in the context of sarcopenia. *Proc Nutr Soc* **74**, 355–366.
36. Dalal S, Hui D, Bidaut L *et al.* (2012) Relationships among body mass index, longitudinal body composition alterations, and survival in patients with locally advanced pancreatic cancer receiving chemoradiation: a pilot study. *J Pain Symptom Manage* **44**, 181–191.
37. Stene GB, Helbostad JL, Amundsen T *et al.* (2014) Changes in skeletal muscle mass during palliative chemotherapy in patients with advanced lung cancer. *Acta Oncol* **54**, 340–348.
38. Antoun S, Birdsell L, Sawyer MB *et al.* (2010) Association of skeletal muscle wasting with treatment with sorafenib in patients with advanced renal cell carcinoma: results from a placebo-controlled study. *J Clin Oncol* **28**, 1054–1060.
39. Veasey Rodrigues H, Baracos VE, Wheler JJ *et al.* (2013) Body composition and survival in the early clinical trials setting. *Eur J Cancer* **49**, 3068–3075.
40. Elliott JA, Doyle SL, Murphy CF *et al.* (2017) Sarcopenia: prevalence, and impact on operative and oncologic outcomes in the multimodal management of locally advanced esophageal cancer. *Ann Surg* **266**, 822–830.
41. Miyamoto Y, Baba Y, Sakamoto Y *et al.* (2015) Sarcopenia is a negative prognostic factor after curative resection of colorectal cancer. *Ann Surg Oncol* **22**, 2663–2668.
42. Harada K, Ida S, Baba Y *et al.* (2016) Prognostic and clinical impact of sarcopenia in esophageal squamous cell carcinoma. *Dis Esophagus* **29**, 627–633.
43. Rutten IJ, van Dijk DP, Kruitwagen RF *et al.* (2016) Loss of skeletal muscle during neoadjuvant chemotherapy is related to decreased survival in ovarian cancer patients. *J Cachexia Sarcopenia Muscle* **7**, 458–466.
44. Boer BC, de Graaff F, Brusse-Keizer M *et al.* (2016) Skeletal muscle mass and quality as risk factors for postoperative outcome after open colon resection for cancer. *Int J Colorectal Dis* **31**, 1117–1124.
45. Amini N, Spolverato G, Gupta R *et al.* (2015) Impact total psoas volume on short- and long-term outcomes in patients undergoing curative resection for pancreatic adenocarcinoma: a new tool to assess sarcopenia. *J Gastrointest Surg* **19**, 1593–1602.
46. Joglekar S, Asghar A, Mott SL *et al.* (2015) Sarcopenia is an independent predictor of complications following pancreatectomy for adenocarcinoma. *J Surg Oncol* **111**, 771–775.
47. Peng P, Hyder O, Firoozmand A *et al.* (2012) Impact of sarcopenia on outcomes following resection of pancreatic adenocarcinoma. *J Gastrointest Surg* **16**, 1478–1486.
48. Rutten IJG, Ubachs J, Kruitwagen RFPM *et al.* (2017) Psoas muscle area is not representative of total skeletal muscle area in the assessment of sarcopenia in ovarian cancer. *J Cachexia Sarcopenia Muscle* **8**, 630–638.
49. Baracos VE (2017) Psoas as a sentinel muscle for sarcopenia: a flawed premise. *J Cachexia Sarcopenia Muscle* **8**, 527–528.
50. Huang DD, Zhou CJ, Wang SL *et al.* (2017) Impact of different sarcopenia stages on the postoperative outcomes after radical gastrectomy for gastric cancer. *Surgery* **161**, 680–693.
51. Fukuda Y, Yamamoto K, Hirao M *et al.* (2016) Sarcopenia is associated with severe postoperative complications in elderly gastric cancer patients undergoing gastrectomy. *Gastric Cancer* **19**, 986–993.
52. Chen LK, Liu LK, Woo J *et al.* (2014) Sarcopenia in Asia: consensus report of the Asian working group for sarcopenia. *J Am Med Dir Assoc* **15**, 95–101.
53. Auclin E, Bourillon C, De Maio E *et al.* (2017) Prediction of everolimus toxicity and prognostic value of skeletal muscle index in patients with metastatic renal cell carcinoma. *Clin Genitourin Cancer* **15**, 350–355.
54. van Vledder MG, Levolver S, Ayez N *et al.* (2012) Body composition and outcome in patients undergoing resection of colorectal liver metastases. *Br J Surg* **99**, 550–557.
55. Barret M, Antoun S, Dalban C *et al.* (2014) Sarcopenia is linked to treatment toxicity in patients with metastatic colorectal cancer. *Nutr Cancer* **66**, 583–589.

56. Cushen SJ, Power DG, Teo MY *et al.* (2017) Body composition by computed tomography as a predictor of toxicity in patients with renal cell carcinoma treated with sunitinib. *Am J Clin Oncol* **40**, 47–52.
57. Choi Y, Oh DY, Kim TY *et al.* (2015) Skeletal muscle depletion predicts the prognosis of patients with advanced pancreatic cancer undergoing palliative chemotherapy, independent of body mass index. *PLoS ONE* **10**, e0139749.
58. Wesselt-Rao N, Hjermstad MJ, Ikeda T *et al.* (2015) Comparing two classifications of cancer cachexia and their association with survival in patients with unresected pancreatic cancer. *Nutr Cancer* **67**, 472–480.
59. Carneiro IP, Mazurak VC & Prado CM (2016) Clinical implications of sarcopenic obesity in cancer. *Curr Oncol Rep* **18**, 62.
60. Bowden JCS, Williams LJ, Simms A *et al.* (2017) Prediction of 90 day and overall survival after chemoradiotherapy for lung cancer: role of performance status and body composition. *Clin Oncol (R Coll Radiol)* **29**, 576–584.
61. Kumar A, Moynagh MR, Multinu F *et al.* (2016) Muscle composition measured by CT scan is a measurable predictor of overall survival in advanced ovarian cancer. *Gynecol Oncol* **142**, 311–316.
62. Aust S, Knogler T, Pils D *et al.* (2015) Skeletal muscle depletion and markers for cancer cachexia are strong prognostic factors in epithelial ovarian cancer. *PLoS ONE* **10**, e0140403.
63. Okumura S, Kaido T, Hamaguchi Y *et al.* (2017) Visceral adiposity and sarcopenic visceral obesity are associated with poor prognosis after resection of pancreatic cancer. *Ann Surg Oncol* **24**, 3732–3740.
64. Antoun S, Lanoy E, Iacovelli R *et al.* (2013) Skeletal muscle density predicts prognosis in patients with metastatic renal cell carcinoma treated with targeted therapies. *Cancer* **119**, 3377–3384.
65. Akahori T, Sho M, Kinoshita S *et al.* (2015) Prognostic significance of muscle attenuation in pancreatic cancer patients treated with neoadjuvant chemoradiotherapy. *World J Surg* **39**, 2975–2982.
66. Cushen SJ, Power DG, Murphy KP *et al.* (2016) Impact of body composition parameters on clinical outcomes in patients with metastatic castrate-resistant prostate cancer treated with docetaxel. *Clin Nutr* **13**, e39–e45.
67. Ní Bhuachalla É, Daly LE, Power DG *et al.* (2017) Computed tomography diagnosed cachexia and sarcopenia in 725 oncology patients: is nutritional screening capturing hidden malnutrition? *J Cachexia Sarcopenia Muscle* [Epublication ahead of print version].
68. Rollins KE, Tewari N, Ackner A *et al.* (2015) The impact of sarcopenia and myosteatosis on outcomes of unresectable pancreatic cancer or distal cholangiocarcinoma. *Clin Nutr* **35**, 1103–1109.
69. McSorley ST, Black DH, Horgan PG *et al.* (2017) The relationship between tumour stage, systemic inflammation, body composition and survival in patients with colorectal cancer. *Clin Nutr* [Epublication ahead of print version].
70. Malietzis G, Currie AC, Athanasiou T *et al.* (2016) Influence of body composition profile on outcomes following colorectal cancer surgery. *Br J Surg* **103**, 572–580.
71. Prado CM, Baracos VE, McCargar LJ *et al.* (2009) Sarcopenia as a determinant of chemotherapy toxicity and time to tumor progression in metastatic breast cancer patients receiving capecitabine treatment. *Clin Cancer Res* **15**, 2920–2926.
72. Shachar SS, Deal AM, Weinberg M *et al.* (2017) Skeletal muscle measures as predictors of toxicity, hospitalization, and survival in patients with metastatic breast cancer receiving taxane-based chemotherapy. *Clin Cancer Res* **23**, 658–665.
73. Antoun S, Baracos VE, Birdsell L *et al.* (2010) Low body mass index and sarcopenia associated with dose-limiting toxicity of sorafenib in patients with renal cell carcinoma. *Ann Oncol* **21**, 1594–1598.
74. Huillard O, Mir O, Peyromaure M *et al.* (2013) Sarcopenia and body mass index predict sunitinib-induced early dose-limiting toxicities in renal cancer patients. *Br J Cancer* **108**, 1034–1041.
75. Mir O, Coriat R, Blanchet B *et al.* (2012) Sarcopenia predicts early dose-limiting toxicities and pharmacokinetics of sorafenib in patients with hepatocellular carcinoma. *PLoS ONE* **7**, e37563.
76. Sjöblom B, Grønberg BH, Benth J *et al.* (2015) Low muscle mass is associated with chemotherapy-induced haematological toxicity in advanced non-small cell lung cancer. *Lung Cancer* **90**, 85–91.
77. Chemama S, Bayar MA, Lanoy E *et al.* (2016) Sarcopenia is associated with chemotherapy toxicity in patients undergoing cytoreductive surgery with hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis from colorectal cancer. *Ann Surg Oncol* **23**, 3891–3898.
78. Massicotte MH, Borget I, Broutin S *et al.* (2013) Body composition variation and impact of low skeletal muscle mass in patients with advanced medullary thyroid carcinoma treated with vandetanib: results from a placebo-controlled study. *J Clin Endocrinol Metab* **98**, 2401–2408.
79. Valentine H, François G, Nora K *et al.* (2017) Sarcopenic overweight is associated with early acute limiting toxicity of anti-PD1 checkpoint inhibitors in melanoma patients. *Invest New Drugs* **35**, 436–441.
80. Prado CM, Baracos VE, McCargar LJ *et al.* (2007) Body composition as an independent determinant of 5-fluorouracil-based chemotherapy toxicity. *Clin Cancer Res* **13**, 3264–3268.
81. Palmela C, Velho S, Agostinho L *et al.* (2017) Body composition as a prognostic factor of neoadjuvant chemotherapy toxicity and outcome in patients with locally advanced gastric cancer. *J Gastric Cancer* **17**, 74–87.
82. Shachar SS, Deal AM, Weinberg M *et al.* (2017) Body composition as a predictor of toxicity in patients receiving anthracycline and taxane-based chemotherapy for early-stage breast cancer. *Clin Cancer Res* **23**, 3537–3543.
83. Anandavadevelan P, Brismar TB, Nilsson M *et al.* (2016) Sarcopenic obesity: a probable risk factor for dose limiting toxicity during neo-adjuvant chemotherapy in oesophageal cancer patients. *Clin Nutr* **35**, 724–730.
84. Tan BH, Brammer K, Randhawa N *et al.* (2015) Sarcopenia is associated with toxicity in patients undergoing neo-adjuvant chemotherapy for oesophago-gastric cancer. *Eur J Surg Oncol* **41**, 333–338.
85. Yip C, Goh V, Davies A *et al.* (2014) Assessment of sarcopenia and changes in body composition after neoadjuvant chemotherapy and associations with clinical outcomes in oesophageal cancer. *Eur Radiol* **24**, 998–1005.
86. Parsons HA, Tsimberidou AM, Pontikos M *et al.* (2012) Evaluation of the clinical relevance of body composition parameters in patients with cancer metastatic to the liver treated with hepatic arterial infusion chemotherapy. *Nutr Cancer* **64**, 206–217.
87. Veasey-Rodrigues H, Parsons HA, Janku F *et al.* (2013) A pilot study of temsirolimus and body composition. *J Cachexia Sarcopenia Muscle* **4**, 259–265.
88. Blauwhoff-Buskermolen S, Versteeg KS, de van der Schueren MA *et al.* (2016) Loss of muscle mass during

- chemotherapy is predictive for poor survival of patients with metastatic colorectal cancer. *J Clin Oncol* **34**, 1339–1344.
89. Felici A, Verweij J & Sparreboom A (2002) Dosing strategies for anticancer drugs: the good, the bad and body-surface area. *Eur J Cancer* **38**, 1677–1684.
 90. Prado CM, Lima IS, Baracos VE *et al.* (2011) An exploratory study of body composition as a determinant of epirubicin pharmacokinetics and toxicity. *Cancer Chemother Pharmacol* **67**, 93–101.
 91. Prado CM, Baracos VE, Xiao J *et al.* (2014) The association between body composition and toxicities from the combination of doxil and trabectedin in patients with advanced relapsed ovarian cancer. *Appl Physiol Nutr Metab* **39**, 693–698.
 92. Ali R, Baracos VE, Sawyer MB *et al.* (2016) Lean body mass as an independent determinant of dose-limiting toxicity and neuropathy in patients with colon cancer treated with FOLFOX regimens. *Cancer Med* **5**, 607–616.
 93. Sjøblom B, Benth J, Grønberg BH *et al.* (2017) Drug dose per kilogram lean body mass predicts hematologic toxicity from carboplatin-doublet chemotherapy in advanced non-small-cell lung cancer. *Clin Lung Cancer* **18**, e129–e136.
 94. Antoun S, Borget I & Lanoy E (2013) Impact of sarcopenia on the prognosis and treatment toxicities in patients diagnosed with cancer. *Curr Opin Support Palliat Care* **7**, 383–389.
 95. Tan BH, Birdsell LA, Martin L *et al.* (2009) Sarcopenia in an overweight or obese patient is an adverse prognostic factor in pancreatic cancer. *Clin Cancer Res* **15**, 6973–6979.
 96. Sharma P, Zargar-Shoshtari K, Caracciolo JT *et al.* (2015) Sarcopenia as a predictor of overall survival after cytoreductive nephrectomy for metastatic renal cell carcinoma. *Urol Oncol* **33**, 339.e317–323.
 97. Fukushima H, Nakanishi Y, Kataoka M *et al.* (2016) Prognostic significance of sarcopenia in patients with metastatic renal cell carcinoma. *J Urol* **195**, 26–32.
 98. Meza-Junco J, Montano-Loza AJ, Baracos VE *et al.* (2013) Sarcopenia as a prognostic index of nutritional status in concurrent cirrhosis and hepatocellular carcinoma. *J Clin Gastroenterol* **47**, 861–870.
 99. Voron T, Tselikas L, Pietrasz D *et al.* (2015) Sarcopenia impacts on short- and long-term results of hepatectomy for hepatocellular carcinoma. *Ann Surg* **261**, 1173–1183.
 100. Levolger S, van Vledder MG, Muslem R *et al.* (2015) Sarcopenia impairs survival in patients with potentially curable hepatocellular carcinoma. *J Surg Oncol* **112**, 208–213.
 101. Kim EY, Kim YS, Park I *et al.* (2015) Prognostic significance of CT-determined sarcopenia in patients with small-cell lung cancer. *J Thorac Oncol* **10**, 1795–1799.
 102. Psutka SP, Carrasco A, Schmit GD *et al.* (2014) Sarcopenia in patients with bladder cancer undergoing radical cystectomy: impact on cancer-specific and all-cause mortality. *Cancer* **120**, 2910–2918.
 103. Fukushima H, Yokoyama M, Nakanishi Y *et al.* (2015) Sarcopenia as a prognostic biomarker of advanced urothelial carcinoma. *PLoS ONE* **10**, e0115895.
 104. Shachar SS, Williams GR, Muss HB *et al.* (2016) Prognostic value of sarcopenia in adults with solid tumours: a meta-analysis and systematic review. *Eur J Cancer* **57**, 58–67.
 105. Kazemi-Bajestani SM, Mazurak VC & Baracos V (2016) Computed tomography-defined muscle and fat wasting are associated with cancer clinical outcomes. *Semin Cell Dev Biol* **54**, 2–10.
 106. Gibson DJ, Burden ST, Strauss BJ *et al.* (2015) The role of computed tomography in evaluating body composition and the influence of reduced muscle mass on clinical outcome in abdominal malignancy: a systematic review. *Eur J Clin Nutr* **69**, 1079–1086.
 107. Tramèr MR, Reynolds DJ, Moore RA *et al.* (1997) Impact of covert duplicate publication on meta-analysis: a case study. *BMJ* **315**, 635–640.
 108. Miyamoto Y, Baba Y, Sakamoto Y *et al.* (2015) Negative impact of skeletal muscle loss after systemic chemotherapy in patients with unresectable colorectal cancer. *PLoS ONE* **10**, e0129742.
 109. Sabel MS, Lee J, Cai S *et al.* (2011) Sarcopenia as a prognostic factor among patients with stage III melanoma. *Ann Surg Oncol* **18**, 3579–3585.
 110. Liefers JR, Bathe OF, Fassbender K *et al.* (2012) Sarcopenia is associated with postoperative infection and delayed recovery from colorectal cancer resection surgery. *Br J Cancer* **107**, 931–936.
 111. Peng PD, van Vledder MG, Tsai S *et al.* (2011) Sarcopenia negatively impacts short-term outcomes in patients undergoing hepatic resection for colorectal liver metastasis. *HPB (Oxford)* **13**, 439–446.
 112. Lou N, Chi CH, Chen XD *et al.* (2017) Sarcopenia in overweight and obese patients is a predictive factor for postoperative complication in gastric cancer: a prospective study. *Eur J Surg Oncol* **43**, 188–195.
 113. Ida S, Watanabe M, Yoshida N *et al.* (2015) Sarcopenia is a predictor of postoperative respiratory complications in patients with esophageal cancer. *Ann Surg Oncol* **22**, 4432–4437.
 114. Sharma P, Zargar-Shoshtari K, Caracciolo JT *et al.* (2015) Sarcopenia as a predictor of complications in penile cancer patients undergoing inguinal lymph node dissection. *World J Urol* **33**, 1585–1592.
 115. van Vugt JLA, Buettner S, Levolger S *et al.* (2017) Low skeletal muscle mass is associated with increased hospital expenditure in patients undergoing cancer surgery of the alimentary tract. *PLoS ONE* **12**, e0186547.
 116. Margadant CC, Bruns ER, Sloothaak DA *et al.* (2016) Lower muscle density is associated with major postoperative complications in older patients after surgery for colorectal cancer. *Eur J Surg Oncol* **42**, 1654–1659.
 117. Prado CM, Sawyer MB, Ghosh S *et al.* (2013) Central tenet of cancer cachexia therapy: do patients with advanced cancer have exploitable anabolic potential? *Am J Clin Nutr* **98**, 1012–1019.
 118. Awad S, Tan BH, Cui H *et al.* (2012) Marked changes in body composition following neoadjuvant chemotherapy for oesophagogastric cancer. *Clin Nutr* **31**, 74–77.
 119. Reisinger KW, Bosmans JW, Uittenbogaart M *et al.* (2015) Loss of skeletal muscle mass during neoadjuvant chemoradiotherapy predicts postoperative mortality in esophageal cancer surgery. *Ann Surg Oncol* **22**, 4445–4452.
 120. Liefers JR, Mourtzakis M, Hall KD *et al.* (2009) A viscerally driven cachexia syndrome in patients with advanced colorectal cancer: contributions of organ and tumor mass to whole-body energy demands. *Am J Clin Nutr* **89**, 1173–1179.
 121. Frontera WR, Hughes VA, Fielding RA *et al.* (2000) Aging of skeletal muscle: a 12-yr longitudinal study. *J Appl Physiol* (1985) **88**, 1321–1326.
 122. Pezaro C, Mukherji D, Tunariu N *et al.* (2013) Sarcopenia and change in body composition following maximal androgen suppression with abiraterone in men with castration-resistant prostate cancer. *Br J Cancer* **109**, 325–331.

123. Fogelman DR, Holmes H, Mohammed K *et al.* (2014) Does IGFR1 inhibition result in increased muscle mass loss in patients undergoing treatment for pancreatic cancer? *J Cachexia Sarcopenia Muscle* **5**, 307–313.
124. Prado CM, Bekaii-Saab T, Doyle LA *et al.* (2012) Skeletal muscle anabolism is a side effect of therapy with the MEK inhibitor: selumetinib in patients with cholangiocarcinoma. *Br J Cancer* **106**, 1583–1586.
125. Cooper AB, Slack R, Fogelman D *et al.* (2015) Characterization of anthropometric changes that occur during neoadjuvant therapy for potentially resectable pancreatic cancer. *Ann Surg Oncol* **22**, 2416–2423.
126. Nattenmüller J, Wochner R, Muley T *et al.* (2017) Prognostic impact of CT-quantified muscle and fat distribution before and after first-line-chemotherapy in lung cancer patients. *PLoS ONE* **12**, e0169136.
127. Daly LE, Ní Bhuachalla É, Power DG *et al.* (2017) Loss of skeletal muscle during systemic chemotherapy is prognostic of poor survival in patients with foregut cancer. *J Cachexia Sarcopenia Muscle* [Epublication ahead of print version].
128. Naito T, Okayama T, Aoyama T *et al.* (2017) Skeletal muscle depletion during chemotherapy has a large impact on physical function in elderly Japanese patients with advanced non-small-cell lung cancer. *BMC Cancer* **17**, 571.
129. Martin L & Sawyer MB (2015) Cancer cachexia: emerging pre-clinical evidence and the pathway forward to clinical trials. *J Natl Cancer Inst* **107**, Issue 12, djv322.
130. Marcus RL, Addison O, Kidde JP *et al.* (2010) Skeletal muscle fat infiltration: impact of age, inactivity, and exercise. *J Nutr Health Aging* **14**, 362–366.
131. Bruns ERJ, Argillander TE, Van Den Heuvel B *et al.* (2017) Oral nutrition as a form of pre-operative enhancement in patients undergoing surgery for colorectal cancer: a systematic review. *Surg Infect (Larchmt)* **19**, 1–10.
132. Santa Mina D, Clarke H, Ritvo P *et al.* (2014) Effect of total-body prehabilitation on postoperative outcomes: a systematic review and meta-analysis. *Physiotherapy* **100**, 196–207.
133. Baldwin C, Spiro A, Ahern R *et al.* (2012) Oral nutritional interventions in malnourished patients with cancer: a systematic review and meta-analysis. *J Natl Cancer Inst* **104**, 371–385.
134. Arends J, Baracos V, Bertz H *et al.* (2017) ESPEN expert group recommendations for action against cancer-related malnutrition. *Clin Nutr* **36**, 1187–1196.
135. Murphy RA, Mourtzakis M, Chu QS *et al.* (2010) Skeletal muscle depletion is associated with reduced plasma (n-3) fatty acids in non-small cell lung cancer patients. *J Nutr* **140**, 1602–1606.
136. Silva JeA, Trindade EB, Fabre ME *et al.* (2012) Fish oil supplement alters markers of inflammatory and nutritional status in colorectal cancer patients. *Nutr Cancer* **64**, 267–273.
137. van der Meij BS, Langius JA, Spreeuwenberg MD *et al.* (2012) Oral nutritional supplements containing n-3 polyunsaturated fatty acids affect quality of life and functional status in lung cancer patients during multimodality treatment: an RCT. *Eur J Clin Nutr* **66**, 399–404.
138. Murphy RA, Mourtzakis M, Chu QS *et al.* (2011) Nutritional intervention with fish oil provides a benefit over standard of care for weight and skeletal muscle mass in patients with nonsmall cell lung cancer receiving chemotherapy. *Cancer* **117**, 1775–1782.
139. Ryan AM, Reynolds JV, Healy L *et al.* (2009) Enteral nutrition enriched with eicosapentaenoic acid (EPA) preserves lean body mass following esophageal cancer surgery: results of a double-blinded randomized controlled trial. *Ann Surg* **249**, 355–363.
140. Almasud AA, Giles KH, Miklavcic JJ *et al.* (2017) Fish oil mitigates myosteatosis and improves chemotherapy efficacy in a preclinical model of colon cancer. *PLoS ONE* **12**, e0183576.
141. Ries A, Trottenberg P, Elsner F *et al.* (2012) A systematic review on the role of fish oil for the treatment of cachexia in advanced cancer: an EPCRC cachexia guidelines project. *Palliat Med* **26**, 294–304.
142. Dewey A, Baughan C, Dean T *et al.* (2007) Eicosapentaenoic acid (EPA, an omega-3 fatty acid from fish oils) for the treatment of cancer cachexia. *Cochrane Database Syst Rev*, CD004597.
143. Mazzotta P & Jeney CM (2009) Anorexia-cachexia syndrome: a systematic review of the role of dietary polyunsaturated Fatty acids in the management of symptoms, survival, and quality of life. *J Pain Symptom Manage* **37**, 1069–1077.
144. Arends J, Bachmann P, Baracos V *et al.* (2017) ESPEN guidelines on nutrition in cancer patients. *Clin Nutr* **36**, 11–48.
145. Fearon KC, Barber MD, Moses AG *et al.* (2006) Double-blind, placebo-controlled, randomized study of eicosapentaenoic acid diester in patients with cancer cachexia. *J Clin Oncol* **24**, 3401–3407.
146. Jatoti A, Rowland K, Loprinzi CL *et al.* (2004) An eicosapentaenoic acid supplement versus megestrol acetate versus both for patients with cancer-associated wasting: a North Central Cancer Treatment Group and National Cancer Institute of Canada collaborative effort. *J Clin Oncol* **22**, 2469–2476.
147. Prado CM, Liefers JR, Bergsten G *et al.* (2012) Dietary patterns of patients with advanced lung or colorectal cancer. *Can J Diet Pract Res* **73**, e298–e303.
148. Geisler C, Prado CM & Müller MJ (2016) Inadequacy of body weight-based recommendations for individual protein intake-lessons from body composition analysis. *Nutrients* **9**, 23.
149. Prado CM, Cushen SJ, Orsso CE *et al.* (2016) Sarcopenia and cachexia in the era of obesity: clinical and nutritional impact. *Proc Nutr Soc* **75**, 188–198.
150. Padilha CS, Marinello PC, Galvão DA *et al.* (2017) Evaluation of resistance training to improve muscular strength and body composition in cancer patients undergoing neoadjuvant and adjuvant therapy: a meta-analysis. *J Cancer Surviv* **11**, 339–349.
151. Adams SC, Segal RJ, McKenzie DC *et al.* (2016) Impact of resistance and aerobic exercise on sarcopenia and dynapenia in breast cancer patients receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *Breast Cancer Res Treat* **158**, 497–507.
152. Stene GB, Helbostad JL, Balstad TR *et al.* (2013) Effect of physical exercise on muscle mass and strength in cancer patients during treatment – a systematic review. *Crit Rev Oncol Hematol* **88**, 573–593.
153. Grande AJ, Silva V & Maddocks M (2015) Exercise for cancer cachexia in adults: executive summary of a Cochrane Collaboration systematic review. *J Cachexia Sarcopenia Muscle* **6**, 208–211.
154. Heywood R, McCarthy AL & Skinner TL (2017) Safety and feasibility of exercise interventions in patients with advanced cancer: a systematic review. *Support Care Cancer* [Epublication ahead of print version].
155. Salakari MR, Surakka T, Nurminen R *et al.* (2015) Effects of rehabilitation among patients with advanced cancer: a systematic review. *Acta Oncol* **54**, 618–628.

156. Cho H, Yoshikawa T, Oba MS *et al.* (2014) Matched pair analysis to examine the effects of a planned preoperative exercise program in early gastric cancer patients with metabolic syndrome to reduce operative risk: the Adjuvant Exercise for General Elective Surgery (AEGES) study group. *Ann Surg Oncol* **21**, 2044–2050.
157. van Waart H, Stuiver MM, van Harten WH *et al.* (2015) Effect of low-intensity physical activity and moderate- to high-intensity physical exercise during adjuvant chemotherapy on physical fitness, fatigue, and chemotherapy completion rates: results of the paces randomized clinical trial. *J Clin Oncol* **33**, 1918–1927.
158. Solheim TS, Fearon KC, Blum D *et al.* (2013) Non-steroidal anti-inflammatory treatment in cancer cachexia: a systematic literature review. *Acta Oncol* **52**, 6–17.
159. Temel JS, Abernethy AP, Currow DC *et al.* (2016) Anamorelin in patients with non-small-cell lung cancer and cachexia (ROMANA 1 and ROMANA 2): results from two randomised, double-blind, phase 3 trials. *Lancet Oncol* **17**, 519–531.
160. Solheim TS, Laird BJA, Balstad TR *et al.* (2017) A randomized phase II feasibility trial of a multimodal intervention for the management of cachexia in lung and pancreatic cancer. *J Cachexia Sarcopenia Muscle* **8**, 778–788.
161. Kimura M, Naito T, Kenmotsu H *et al.* (2015) Prognostic impact of cancer cachexia in patients with advanced non-small cell lung cancer. *Support Care Cancer* **23**, 1699–1708.
162. Thoresen L, Frykholm G, Lydersen S *et al.* (2012) The association of nutritional assessment criteria with health-related quality of life in patients with advanced colorectal carcinoma. *Eur J Cancer Care (Engl)* **21**, 505–516.
163. Thoresen L, Frykholm G, Lydersen S *et al.* (2013) Nutritional status, cachexia and survival in patients with advanced colorectal carcinoma. Different assessment criteria for nutritional status provide unequal results. *Clin Nutr* **32**, 65–72.
164. Reisinger KW, van Vugt JL, Tegels JJ *et al.* (2015) Functional compromise reflected by sarcopenia, frailty, and nutritional depletion predicts adverse postoperative outcome after colorectal cancer surgery. *Ann Surg* **261**, 345–352.
165. Malietzis G, Johns N, Al-Hassi HO *et al.* (2016) Low muscularity and myosteatosis is related to the host systemic inflammatory response in patients undergoing surgery for colorectal cancer. *Ann Surg* **263**, 320–325.
166. Grotenhuis B, Shapiro J, van Adrichem S *et al.* (2016) Sarcopenia/muscle mass is not a prognostic factor for short- and long-term outcome after esophagectomy for cancer. *World J Surg* **40**, 2698–2704.
167. Nishigori T, Okabe H, Tanaka E *et al.* (2016) Sarcopenia as a predictor of pulmonary complications after esophagectomy for thoracic esophageal cancer. *J Surg Oncol* **113**, 678–684.
168. Tegels JJ, van Vugt JL, Reisinger KW *et al.* (2015) Sarcopenia is highly prevalent in patients undergoing surgery for gastric cancer but not associated with worse outcomes. *J Surg Oncol* **112**, 403–407.
169. Nishigori T, Tsunoda S, Okabe H *et al.* (2016) Impact of sarcopenic obesity on surgical site infection after laparoscopic total gastrectomy. *Ann Surg Oncol* **23**, 524–531.
170. Kudou K, Saeki H, Nakashima Y *et al.* (2017) Prognostic significance of sarcopenia in patients with esophagogastric junction cancer or upper gastric cancer. *Ann Surg Oncol* **24**, 1804–1810.
171. Arrieta O, De la Torre-Vallejo M, López-Macias D *et al.* (2015) Nutritional status, body surface, and low lean body mass/body mass index are related to dose reduction and severe gastrointestinal toxicity induced by afatinib in patients with non-small cell lung cancer. *Oncologist* **20**, 967–974.
172. Di Sebastiano KM, Yang L, Zbuk K *et al.* (2013) Accelerated muscle and adipose tissue loss may predict survival in pancreatic cancer patients: the relationship with diabetes and anaemia. *Br J Nutr* **109**, 302–312.
173. Harimoto N, Shirabe K, Yamashita YI *et al.* (2013) Sarcopenia as a predictor of prognosis in patients following hepatectomy for hepatocellular carcinoma. *Br J Surg* **100**, 1523–1530.
174. Del Fabbro E, Parsons H, Warneke CL *et al.* (2012) The relationship between body composition and response to neoadjuvant chemotherapy in women with operable breast cancer. *Oncologist* **17**, 1240–1245.
175. Camus V, Lanic H, Kraut J *et al.* (2014) Prognostic impact of fat tissue loss and cachexia assessed by computed tomography scan in elderly patients with diffuse large B-cell lymphoma treated with immunochemotherapy. *Eur J Haematol* **93**, 9–18.
176. Moryoussef F, Dhooge M, Volet J *et al.* (2015) Reversible sarcopenia in patients with gastrointestinal stromal tumor treated with imatinib. *J Cachexia Sarcopenia Muscle* **6**, 343–350.
177. Cousin S, Hollebecque A, Koscielny S *et al.* (2014) Low skeletal muscle is associated with toxicity in patients included in phase I trials. *Invest New Drugs* **32**, 382–387.