


Title	Combination of electroporation delivered metabolic modulators with low-dose chemotherapy in osteosarcoma
Authors	Gill, Kheshwant S.;Fernandes, Philana;Bird, Brian;Soden, Declan M.;Forde, Patrick F.
Publication date	2018
Original Citation	Gill, K. S., Fernandes, P., Bird, B., Soden, D. M. and Forde, P. F. (2018) 'Combination of electroporation delivered metabolic modulators with low-dose chemotherapy in osteosarcoma', Oncotarget, 9(59), pp. 31473-31489. doi: 10.18632/oncotarget.25843
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.oncotarget.com/index.php?journal=oncotarget&page=article&op=view&path[]=25843&path[]=80841 - 10.18632/oncotarget.25843
Rights	© 2018, the Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. https://creativecommons.org/licenses/by/3.0/ - https://creativecommons.org/licenses/by/3.0/
Download date	2024-05-04 03:54:51
Item downloaded from	https://hdl.handle.net/10468/6947

Combination of electroporation delivered metabolic modulators with low-dose chemotherapy in osteosarcoma

SUPPLEMENTARY MATERIALS



Supplementary Figure 1: Sensitivity of OS cells to low-dose chemotherapy increases when treated with metabolic modulators using EP first. Sensitivity of (A) K7M2 and (B) Saos2 cell lines treated with metabolic modulators actively (with EP) or passively (without EP), combined with low-dose Cisplatin was evaluated by stained colonies formed in Cisplatin-treated media over a period of time. Picture of each well shown is a representative image of at least nine similar wells (three independent experiments).