

Title	Tracking cocrystallization of active pharmaceutical ingredients with Bbenzoic acid coformer using Broadband Acoustic resonance Dissolution Spectroscopy (BARDS)
Authors	Kent, Christopher;Rao Khandavilli, Udaya B. Rao;Alfarsi, Anas;Hanna-Brown, Melissa;McSweeney, Seán;Krüse, Jacob;Lawrence, Simon E.;Fitzpatrick, Dara
Publication date	2018-09-18
Original Citation	Kent, C., Rao Khandavilli, U. B., Alfarsi, A., Hanna-Brown, M., McSweeney, S., Krüse, J., Lawrence, S. and Fitzpatrick, D. (2018) 'Tracking Cocrystallization of Active Pharmaceutical Ingredients with Benzoic Acid Coformer Using Broadband Acoustic Resonance Dissolution Spectroscopy (BARDS)', Crystal Growth & Design, 18(11), pp. 6528-6537. doi: 10.1021/acs.cgd.8b00668
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://pubs.acs.org/doi/10.1021/acs.cgd.8b00668 - 10.1021/ acs.cgd.8b00668
Rights	© 2018 American Chemical Society. This document is the Accepted Manuscript version of a Published Work that appeared in final form in Crystal Growth & Design, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see https://pubs.acs.org/doi/pdf/10.1021/acs.cgd.8b00668
Download date	2024-04-19 09:08:00
Item downloaded from	https://hdl.handle.net/10468/9648



## Tracking Cocrystallization of Active Pharmaceutical Ingredients with Benzoic Acid Coformer using Broadband Acoustic Resonance Dissolution Spectroscopy (BARDS)

Christopher Kent‡, Uday B. Rao Khandavilli† Anas Alfarsi‡, Melissa Hanna-Brown#,

Seán McSweeney¥, Jacob Krüse, Simon Lawrence‡†, and Dara Fitzpatrick‡\*

- ‡ School of Chemistry, Analytical and Biological Chemistry Research Facility, University College Cork, Cork, Ireland.
- † Synthesis and Solid State Pharmaceutical Cluster, University College Cork, Ireland.
- # Pfizer, Sandwich, Kent, United Kingdom.
- ¥ BARDS Acoustic Science Labs, Bio-Innovation Centre, University College Cork.

## IR spectra for Table 1:




Figure 1: IR Analysis for a 1:1 mixture of INA: BA analysed after 1 minute of milling in the Retsch MM400 mixer mill




Figure 2: IR Analysis for a 1:1 mixture of INA: BA analysed after 2 minutes of milling in the Retsch MM400 mixer mill.




Figure 3: IR Analysis for a 1:1 mixture of INA: BA analysed after 5 minutes of milling in the Retsch MM400 mixer mill.




Figure 4: IR Analysis for a 1:1 mixture of INA: BA analysed after 10 minutes of milling in the Retsch MM400 mixer mill.




Figure 5: IR Analysis for a 1:1 mixture of INA: BA analysed after 15 minutes of milling in the Retsch MM400 mixer mill.




Figure 6: IR Analysis for a 1:1 mixture of INA: BA analysed after 20 minutes of milling in the Retsch MM400 mixer mill.




Figure 7: IR Analysis for a 1:1 mixture of INA: BA analysed after 40 minutes of milling in the Retsch MM400 mixer mill.

## Analysis of 1:1 mixture of Theophylline and Benzoic Acid:




Figure 8: Infrared analysis after 5 minutes of LAG using a Retsch MM400 of a 1:1 TP:BA mixture.




Figure 9: Infrared analysis after 10 minutes of LAG using a Retsch MM400 of a 1:1 TP:BA mixture.




Figure 10: Infrared analysis after 15 minutes of LAG using a Retsch MM400 of a 1:1 TP:BA mixture.




Figure 11: Infrared analysis after 20 minutes of LAG using a Retsch MM400 of a 1:1 TP:BA mixture.