

Title	Complete genome sequence of Escherichia coli phage APC_JM3.2 isolated from a chicken cecum
Authors	Mullally, Jayne;Stockdale, Stephen R.;Smith, Muireann K.;Draper, Lorraine A.;Ross, R. Paul;Hill, Colin
Publication date	2018-01-04
Original Citation	MULLALLY, J., STOCKDALE, S.R., SMITH, M.K., DRAPER, L.A., ROSS, P.R. and HILL, C., 2018. Complete Genome Sequence of Escherichia coli Phage APC_JM3.2 Isolated from a Chicken Cecum. Genome Announcements, 6(1), pp.1-2. DOI: 10.1128/genomeA.01292-17
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1128/genomeA.01292-17
Rights	© 2018 Mullally et al. This work is licensed under the Creative Commons Attribution License ( <a href="https://creativecommons.org/licenses/by/3.0/">https://creativecommons.org/licenses/by/3.0/</a> ) (the "License"). Notwithstanding the ProQuest Terms and Conditions, you may use this content in accordance with the terms of the License. - <a href="https://creativecommons.org/licenses/by/3.0/">https://creativecommons.org/licenses/by/3.0/</a>
Download date	2024-05-02 19:43:51
Item downloaded from	<a href="https://hdl.handle.net/10468/5531">https://hdl.handle.net/10468/5531</a>



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh



# Complete Genome Sequence of *Escherichia coli* Phage APC\_JM3.2 Isolated from a Chicken Cecum

Jayne Mullally,<sup>a</sup> Stephen R. Stockdale,<sup>a,b,c</sup> Muireann K. Smith,<sup>a,b</sup> Lorraine A. Draper,<sup>a,b</sup> Paul R. Ross,<sup>a,b,c</sup> Colin Hill<sup>a,b,c</sup>

<sup>a</sup>School of Microbiology, University College Cork, Co. Cork, Ireland

<sup>b</sup>APC Microbiome Institute, Biological Sciences Institute, University College Cork, Co. Cork, Ireland

<sup>c</sup>Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland

**ABSTRACT** Avian pathogenic *Escherichia coli* (APEC) bacteria are a significant challenge to the poultry industry. Bacteriophages (phages) have the potential to control APEC strains, increasing animal welfare and economic productivity. Here, we report the isolation of an *E. coli*-infecting phage, APC\_JM3.2, isolated from the cecum of a broiler chicken in Ireland.

In this study, the bacteriophage APC\_JM3.2 was isolated from the cecum of a broiler chicken in Ireland. Phage APC\_JM3.2 was detected using the bacterium *Escherichia coli* strain JM3.2 as its host, which was also isolated from the same chicken cecum sample. Virulence factors associated with *E. coli* JM3.2, and its classification as an APEC strain, were not investigated in this study.

The morphology of phage APC\_JM3.2 was determined by transmission electron microscopy (UCD Conway Institute of Biomolecular and Biomedical Research, Dublin). The short noncontractile tail of phage APC\_JM3.2 indicates that it belongs to the *Podoviridae* family. Biological characterization of phage APC\_JM3.2 revealed that it has a latent period of between 15 and 20 min and a burst size of approximately 45 phages per infected cell.

The APC\_JM3.2 phage genome was sequenced by using Illumina MiSeq technology (Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork). Reads were trimmed and filtered by using Trimmomatic (1) to remove adaptor sequences and reads less than 70 bp when a sliding window of 4 bp and minimum Phred score of 30 was applied. Subsequent read quality was assessed using FastQC (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>), and assembly was performed using SPAdes Genome Assembler v3.9.0 (2). An average genome coverage of 1,254× for phage APC\_JM3.2 was achieved. Putative phage coding sequences were identified using Prodigal v1.20 (3), and functions were predicted by querying translated nucleotide sequences against the BLAST nr and UniProt TrEMBL databases. No tRNA- or transfer-messenger RNA (tmRNA)-encoding sequences could be detected using ARAGORN v1.2.36 (4).

Phage APC\_JM3.2 has a 39,761-bp double-stranded DNA (dsDNA) genome with a GC content of 47.3%. APC\_JM3.2 is predicted to encode 57 open reading frames (ORFs), which translate into proteins ranging in size from 37 to 911 amino acids. Thirty-five of the APC\_JM3.2 ORFs are predicted on the forward strand, while the remaining 22 ORFs have a reverse orientation. Forty-three ORFs of APC\_JM3.2 were assigned putative functions by BLASTp, while 11 of the remaining 14 ORFs not assigned a BLASTp function were assigned a putative function by the UniProt database.

The identification of an integrase on the genome of phage APC\_JM3.2 indicates that it can lysogenize target bacteria, limiting its usefulness in the control of APEC strains. However, phage APC\_JM3.2 is dissimilar to other known *E. coli* phages, with 96% identity across only 47% of its genome to its closest detected BLASTn search result,

**Received** 17 October 2017 **Accepted** 10 November 2017 **Published** 4 January 2018

**Citation** Mullally J, Stockdale SR, Smith MK, Draper LA, Ross PR, Hill C. 2018. Complete genome sequence of *Escherichia coli* phage APC\_JM3.2 isolated from a chicken cecum. *Genome Announc* 6:e01292-17. <https://doi.org/10.1128/genomeA.01292-17>.

**Copyright** © 2018 Mullally et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Colin Hill, [c.hill@ucc.ie](mailto:c.hill@ucc.ie).

J.M. and S.R.S. contributed equally to this work.

*Enterobacteria* phage IME10. Therefore, the screening of chicken ceca should yield more phages that are genetically distinct and have therapeutic potential against bacterial infections caused by avian pathogenic *E. coli*.

**Accession number(s).** The complete genome sequence of *Escherichia coli* phage APC\_JM3.2 has been deposited in GenBank under the accession number [MG197996](#).

## ACKNOWLEDGMENTS

This publication was supported by Science Foundation Ireland under grant SFI/12/RC/2273.

The funders of this study had no role in the design, data collection, interpretation, or decision to publish this work. All authors have seen and approved the manuscript and declare no conflict of interest.

## REFERENCES

1. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
2. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
3. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
4. Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res* 32:11–16. <https://doi.org/10.1093/nar/gkh152>.