CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Genome sequence of Staphylococcus saprophyticus DPC5671, a strain Isolated from cheddar cheese
Authors	Bertuzzi, Andrea S.;Guinane, Caitriona M.;Crispie, Fiona;Kilcawley, Kieran N.;McSweeney, Paul L. H.;Rea, Mary C.
Publication date	2017-04-20
Original Citation	Bertuzzi, A. S., Guinane, C. M., Crispie, F., Kilcawley, K. N., McSweeney, P. L. H. and Rea, M. C. (2017) 'Genome Sequence of Staphylococcus saprophyticus DPC5671, a Strain Isolated from Cheddar Cheese', Genome Announcements, 5(16), e00193-17. doi:10.1128/genomeA.00193-17
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1128/genomeA.00193-17
Rights	© 2017 Bertuzzi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license. - http://creativecommons.org/licenses/by/4.0/
Download date	2026-11-04 22:27:37
Item downloaded from	https://hdl.handle.net/10468/3916

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

AMERICAN SOCIETY FOR MICROBIOLOGY | genomeAnnouncements™

PROKARYOTES

Genome Sequence of *Staphylococcus saprophyticus* DPC5671, a Strain Isolated from Cheddar Cheese

Andrea S. Bertuzzi,[a,b] Caitriona M. Guinane,[c] Fiona Crispie,[a,d] Kieran N. Kilcawley,[a] Paul L. H. McSweeney,[b] Mary C. Rea[a,d]

Teagasc Food Research Centre, Moorepark, Fermoy, Cork, Ireland[a]; School of Food and Nutritional Science, University College Cork, Cork, Ireland[b]; Department of Biological Sciences, Cork Institute of Technology, Cork, Ireland[c]; APC Microbiome Institute, University College Cork, Cork, Ireland[d]

ABSTRACT The draft genome sequence of *Staphylococcus saprophyticus* DPC5671, isolated from cheddar cheese, was determined. *S. saprophyticus* is a common Gram-positive bacterium detected on the surface of smear-ripened cheese and other fermented foods.

Received 25 February 2017 **Accepted** 7 March 2017 **Published** 20 April 2017

Citation Bertuzzi AS, Guinane CM, Crispie F, Kilcawley KN, McSweeney PLH, Rea MC. 2017. Genome sequence of *Staphylococcus saprophyticus* DPC5671, a strain isolated from cheddar cheese. Genome Announc 5: e00193-17. https://doi.org/10.1128/genomeA.00193-17.

Copyright © 2017 Bertuzzi et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Mary C. Rea, mary.rea@teagasc.ie.

Staphylococci are Gram-positive catalase-positive bacteria with halotolerance that enables their growth in salted fermented food. Strains belonging to the *Staphylococcus* genus are commonly identified in soft cheese varieties made from cow, ewe, or goat milk, and together with *Brevibacterium*, *Corynebacterium*, and *Microbacterium*, *Staphylococcus* is considered the most important genus making up the microbiota of the cheese surface (1). *S. saprophyticus* is a species frequently detected on the surface of smear-ripened cheese and other fermented foods (2–4). Here, we present the draft genome sequence of *S. saprophyticus* DPC5671, which will allow a full safety assessment and further analysis on its role in cheese ripening.

The draft genome of *S. saprophyticus* DPC5671 was sequenced using paired-end 454 pyrosequencing to a coverage of 23×. Sequencing took place at the Teagasc 454 sequencing facility on a genome sequencer FLX platform (Roche Diagnostics, West Sussex, United Kingdom), according to the manufacturer's protocols. This was followed by initial assembly into 24 contigs using the Newbler program (Roche Life Science). The software Prodigal (5) was used to predict open reading frames within the draft genome, and the RAST annotation server (6) was used to determine complementary gene calling and automated annotation. The draft genome was manually analyzed using the ARTEMIS genome browser (7), and comparative analysis with the genome of *S. saprophyticus* ATCC 15305 (8) was performed using the Artemis Comparison Tool (ACT) (9). The PHAST (PHAge Search Tool) Web server (10) was used to determine the presence of putative phage within the genome. The Comprehensive Antibiotic Resistance Database (CARD) software (11) was used to determine the presence of genes potentially involved in antibiotic resistance, and the presence of known staphylococcal virulence factors was analyzed using the BlastP Web server (12).

The draft genome of *S. saprophyticus* DPC5671 includes 2,676,318 bp, with an average G+C content of 33.1%. It consists of a single circular chromosome and does not appear to harbor any plasmids. Within the draft genome, there are 2,647 coding regions predicted, in addition to four rRNA and 59 tRNA genes. The genome sequence also includes two putative novel phages of ~43.6 kb and ~42.9 kb within the chromosome. Overall, the genome of *S. saprophyticus* DPC5671 shows high similarity to *S. saprophyticus* ATCC 15305 in genome size, G+C content, and gene synteny (8).

S. saprophyticus DPC5671 is coagulase negative, nonhemolytic, and does not appear to produce any toxins associated with *Staphylococcus aureus*. Genomic analysis re-

Downloaded from http://genomea.asm.org/ on May 4, 2017 by IRIS

vealed no obvious transferable antibiotic resistance loci by the methods used in this study. Previous studies showed an involvement of *S. saprophyticus* in urinary tract infections, showing a specific adhesin, *uafA*, to be associated with adherence to the eukaryotic cell in the urinary tract (8). The *S. saprophyticus* DPC5671 genome was found to have a predicted coding sequence (CDS) for adhesion, with similarities examined with the BlastP Web server to *uafA* in *S. saprophyticus* ATCC 15305 with 39% of query cover and 97% of protein identity.

The availability of the genome sequence of DPC5671 will allow its role in flavor development in cheese ripening to be studied.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number MUXI00000000. The version described in this paper is version MUXI01000000.

ACKNOWLEDGMENT

Andrea Bertuzzi is currently in receipt of a Teagasc Walsh Fellowship.

REFERENCES

1. Desmasures N, Bora N, Ward A. 2015. Smear ripened cheeses, p 1–18. *In* Bora N, Dodd C, Desmasures N (ed), Diversity, dynamics and functional role of actinomycetes on European smear ripened cheeses. Springer, Switzerland. https://doi.org/10.1007/978-3-319-10464-5_1.
2. Leroy S, Giammarinaro P, Chacornac JP, Lebert I, Talon R. 2010. Biodiversity of indigenous staphylococci of naturally fermented dry sausages and manufacturing environments of small-scale processing units. Food Microbiol 27:294–301. https://doi.org/10.1016/j.fm.2009.11.005.
3. Larpin-Laborde S, Imran M, Bonaïti C, Bora N, Gelsomino R, Goerges S, Irlinger F, Goodfellow M, Ward AC, Vancanneyt M, Swings J, Scherer S, Guéguen M, Desmasures N. 2011. Surface microbial consortia from Livarot, a French smear-ripened cheese. Can J Microbiol 57:651–660. https://doi.org/10.1139/w11-050.
4. Cogan TM, Goerges S, Gelsomino R, Larpin S, Hohenegger M, Bora N, Jamet E, Rea MC, Mounier J, Vancanneyt M, Guéguen M, Desmasures N, Swings J, Goodfellow M, Ward AC, Sebastiani H, Irlinger F, Chamba JF, Beduhn R, Scherer S. 2014. Biodiversity of the surface microbial consortia from Limburger, Reblochon, Livarot, Tilsit, and Gubbeen cheeses, p 219–250. *In* Donnelly C (ed), Cheese and microbes. American Society of Microbiology, Washington, DC.
5. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC Bioinformatics 11:119. https://doi.org/10.1186/1471-2105-11-119.
6. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. BMC Genomics 9:75. https://doi.org/10.1186/1471-2164-9-75.
7. Carver T, Harris SR, Berriman M, Parkhill J, McQuillan JA. 2012. Artemis: an integrated platform for visualization and analysis of high-throughput sequence-based experimental data. Bioinformatics 28:464–469. https://doi.org/10.1093/bioinformatics/btr703.
8. Kuroda M, Yamashita A, Hirakawa H, Kumano M, Morikawa K, Higashide M, Maruyama A, Inose Y, Matoba K, Toh H, Kuhara S, Hattori M, Ohta T. 2005. Whole genome sequence of *Staphylococcus saprophyticus* reveals the pathogenesis of uncomplicated urinary tract infection. Proc Natl Acad Sci U S A 102:13272–13277. https://doi.org/10.1073/pnas.0502950102.
9. Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis Comparison Tool. Bioinformatics. 21:3422–3423. https://doi.org/10.1093/bioinformatics/bti553.
10. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. Nucleic Acids Res 39:W347–W352. https://doi.org/10.1093/nar/gkr485.
11. McArthur AG, Waglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, Bhullar K, Canova MJ, De Pascale G, Ejim L, Kalan L, King AM, Koteva K, Morar M, Mulvey MR, O'Brien JS, Pawlowski AC, Piddock LJ, Spanogiannopoulos P, Sutherland AD, Tang I, Taylor PL, Thaker M, Wang W, Yan M, Yu T, Wright GD. 2013. The Comprehensive Antibiotic Resistance Database. Antimicrob Agents Chemother 57:3348–3357. https://doi.org/10.1128/AAC.00419-13.
12. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res 25:3389–3402. https://doi.org/10.1093/nar/25.17.3389.

Downloaded from http://genomea.asm.org/ on May 4, 2017 by IRIS