

|                             |                                                                                                                                                                                                                                                                                                                                                                                  |
|-----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Title                       | Influence of dietary cardoon meal on growth performance and selected meat quality parameters of lambs, and the antioxidant potential of cardoon extract in ovine muscle homogenates                                                                                                                                                                                              |
| Authors                     | Salami, Saheed A.;Valenti, Bernardo;O'Grady, Michael N.;Kerry, Joseph P.;Mattioli, Simona;Licitra, Giuseppe;Luciano, Giuseppe;Priolo, Alessandro                                                                                                                                                                                                                                 |
| Publication date            | 2019-03-25                                                                                                                                                                                                                                                                                                                                                                       |
| Original Citation           | Salami, S. A., Valenti, B., O'Grady, M. N., Kerry, J. P., Mattioli, S., Licitra, G., Luciano, G. and Priolo, A. (2019) 'Influence of dietary cardoon meal on growth performance and selected meat quality parameters of lambs, and the antioxidant potential of cardoon extract in ovine muscle homogenates', Meat Science, 153, pp. 126-134. doi: 10.1016/j.meatsci.2019.03.017 |
| Type of publication         | Article (peer-reviewed)                                                                                                                                                                                                                                                                                                                                                          |
| Link to publisher's version | <a href="http://www.sciencedirect.com/science/article/pii/S030917401830860X">http://www.sciencedirect.com/science/article/pii/S030917401830860X</a> - 10.1016/j.meatsci.2019.03.017                                                                                                                                                                                              |
| Rights                      | © 2019 Elsevier Ltd. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license - <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>                                                                                                                                        |
| Download date               | 2026-08-29 17:23:36                                                                                                                                                                                                                                                                                                                                                              |
| Item downloaded from        | <a href="https://hdl.handle.net/10468/7816">https://hdl.handle.net/10468/7816</a>                                                                                                                                                                                                                                                                                                |



**UCC**

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

**Influence of dietary cardoon meal on growth performance and selected meat quality parameters of lambs, and the antioxidant potential of cardoon extract in ovine muscle homogenates**

Saheed A. Salami<sup>a,b\*</sup>, Bernardo Valenti<sup>a</sup>, Michael N. O'Grady<sup>b</sup>, Joseph P. Kerry<sup>b</sup>, Simona Mattioli<sup>c</sup>, Giuseppe Licitra<sup>a</sup>, Giuseppe Luciano<sup>a,c</sup> and Alessandro Priolo<sup>a</sup>

<sup>a</sup>*Department Di3A, Animal Production Science, University of Catania, Via Valdisavoia 5, 95123 Catania, Italy*

<sup>b</sup>*Food Packaging Group, School of Food and Nutritional Sciences, College of Science, Engineering and Food Science, University College Cork, Cork, Ireland*

<sup>c</sup>*Department of Agricultural, Environmental and Food Science (DSA3), University of Perugia, Borgo XX Giugno 74, 06121 Perugia, Italy*

**\*Corresponding author:** *E-mail address:* s.salami@umail.ucc.ie; *Telephone:* +353899466807

21 **Abstract**

22 Fatty acids and oxidative stability were determined in meat from lambs fed a diet containing  
23 15% dehydrated alfalfa (CON,  $n=8$ ) or cardoon meal (CMD,  $n=7$ ). Furthermore, the antioxidant  
24 activity of a phenolic-rich cardoon meal extract (1.32 GAE mg/ml) was examined in muscle  
25 homogenates (0, 0.5, 1, 5% v/w) subjected to iron/ascorbate-induced oxidation. Feeding CMD  
26 did not affect lamb performances and carcass traits but reduced ( $P < 0.05$ ) the vaccenic and  
27 rumenic acids and increased stearic acid in muscle. Lipid oxidation was higher in raw meat from  
28 the CMD-fed lambs after 7 days of storage ( $P < 0.05$ ). Feeding CMD did not affect the colour  
29 stability of raw meat and the oxidative stability of cooked meat and of muscle homogenates  
30 incubated with pro-oxidant catalysts. Adding 5% cardoon extract in muscle homogenates  
31 increased (+114.3%;  $P = 0.03$ ) the total phenolic content and reduced (-77.6%;  $P < 0.01$ ) lipid  
32 oxidation, demonstrating the antioxidant potential of compounds present in cardoon meal.

34 **Keywords:** *Lamb meat, Growth performance, Fatty acids, Lipid oxidation, Cardoon,*  
35 *Polyphenols*

## 1. Introduction

Meat from ruminant animals (cattle, sheep, goat and buffalo) can be a viable dietary source of bioactive fatty acids including conjugated linoleic acids (CLA), monounsaturated fatty acids (MUFA) and *n*-3 long-chain polyunsaturated fatty acids (PUFA; Bessa, Alves, & Santos-Silva, 2015). However, ruminant meat also contains high levels of saturated fatty acids (SFA) and *trans*-fatty acids (TFA) linked to increased risk of cardio-metabolic diseases in humans (McAfee et al., 2010). The fatty acid profile of ruminant meat is significantly influenced by extensive conversion of dietary PUFA to SFA during biohydrogenation of lipids that occur in the fore-stomach (rumen) (Shingfield, Bonnet, & Scollan, 2013). However, feeding strategies such as supplementation of PUFA-rich vegetable oil could increase the intramuscular deposition of PUFA and CLA and reduce SFA levels in ruminant meat (Bessa et al., 2015).

Lipid oxidation is the main chemical process associated with oxidative deterioration of meat, resulting in undesirable rancid off-flavour and colour deterioration that limit shelf-life and negatively impact consumer acceptability of meat (Morrissey, Sheehy, Galvin, Kerry, & Buckley, 1998). The balance between antioxidant and pro-oxidant components in muscle tissues mainly influences lipid stability. Animal diets could influence the biochemical components of muscle tissues through enrichment with antioxidant compounds, such as  $\alpha$ -tocopherol (vitamin E), that enhance the oxidative stability of meat (Luciano et al., 2013; Salami et al., 2016).

Phenolic compounds are secondary metabolites that are ubiquitous in several plant species and may be found in considerable amounts in agro-industrial by-products (AIBP; Balasundram, Sundram, & Samman, 2006). The AIBP can be a valuable and economical resource for animal feeding or for the extraction of bioactive phenolic compounds used in functional food

applications. Feeding phenolic-rich AIBP improves the fatty acid composition of ruminant meat through inhibition of ruminal biohydrogenation (Lanza et al., 2015) and may enhance the oxidative stability of meat through controversial antioxidant mechanisms (Vasta & Luciano, 2011). Phenolics could exhibit *in vivo* antioxidant activities through direct deposition of these compounds in tissues or through possible indirect antioxidant mechanisms including the sparing effect of polyphenols on other antioxidants, such as ascorbic acid and tocopherols (Valenti et al., 2018). Furthermore, phenolic-rich extracts obtained from AIBP have been directly incorporated into meat products during processing to inhibit oxidative deterioration and extend shelf-life (Balzan et al., 2017; Kanatt, Chander, Radhakrishna, & Sharma, 2005; Rodríguez-Carpena, Morcuende, & Estévez, 2011).

Cardoon meal is a by-product retained after the extraction of oil from the seeds of cultivated cardoon (*Cynara cardunculus* var. *altilis*), a perennial herb native to the Mediterranean region and widespread in parts of Europe, Americas and Oceania. Cardoon meal has been proposed as an alternative feed resource because of its potential as a valuable source of fibre, protein, amino acids and bioactive compounds such as polyphenols and unsaturated fatty acids (Genovese et al., 2015). Cardoon meal may also contain residual amount of other antioxidant compounds such as  $\alpha$ -tocopherol found in considerable concentration in cardoon seed oil (Maccarone et al., 1999). Moreover, extracts obtained from different morphological parts (leaf, seed, stem and flower) of cardoon have demonstrated *in vitro* antioxidant and antimicrobial activities attributed to the phenolic constituents present (Falleh et al., 2008; Pandino, Lombardo, Mauromicale, & Williamson, 2011; Ramos et al., 2014). To our knowledge, no information has been published on the effect of dietary cardoon meal or cardoon extract on meat quality. Therefore, the first objective of this study was to determine the effect of dietary cardoon meal on lamb growth

performance, carcass characteristics, and meat fatty acid composition and oxidative stability. In addition, the antioxidant potential of a phenolic-rich cardoon meal extract on lipid oxidation in an ovine muscle model system was investigated.

## **2. Materials and Methods**

### ***2.1. Animals, diets and experimental design***

The experiment was conducted indoors in the experimental farm of the University of Catania. The animals were handled by trained personnel according to the European Union legislation for the protection of animals used for scientific purposes (2010/63/ EU Directive) and the study was approved by the University of Catania (FIR-2014-PI/LB/Di3A). Fifteen male Sarda x Comisana lambs (age 75 d and average weight  $19.58 \pm 2.01$  kg) were randomly assigned to two experimental groups. Each animal was reared in an individual pen and adapted to the experimental diets for a period of 9 d by progressive substitution of the weaning feed with the experimental feeds until a total replacement of the weaning diet was achieved. The control group (CON,  $n = 8$ ), was raised on a commercial concentrate-based diet containing the following ingredients (*as-fed* basis): barley (48.0%), dehydrated alfalfa (15.0%), wheat bran (23.0%), soybean meal (10.0%), molasses (2.0%) and vitamin premix (2.0%). The cardoon meal group (CMD,  $n = 7$ ), received the same diet as the CON lambs except that the 15% dehydrated alfalfa was completely replaced by cardoon meal. The chemical composition of the experimental diets is outlined in Table 1. The CON and CMD diets were supplied in form of pellets and lambs had *ad libitum* access to feeds and water for 75 days pre-slaughter. Experimental feed samples were collected at two-week intervals during the feeding trial and stored in vacuum packs at  $-30$  °C prior to chemical analysis. Diets were supplied daily and the amount of refusal was measured before morning feeding to calculate dry matter intake (DMI). The body weight (BW) of the

lambs was measured at the start of the experiment and recorded weekly (at 09:00 h before providing fresh feed) to calculate average daily gain (ADG).

## ***2.2. Slaughter, and carcass sampling and measurement***

The lambs were slaughtered (stunned by captive bolt before exsanguination) in a commercial abattoir, where they had free access to the experimental diets and water until approximately 3 h before slaughter. Carcass weight was recorded following removal of the visceral organs and dressing percentage was calculated as the percentage of carcass weight to final BW. Visual appraisal of hot hanging carcasses was performed by a certified meat grader to determine the conformation and fat cover scores according to the European Union's EUROP carcass classification system (Commission Regulation (EC) No 823/98, 1998) . Carcass conformation and fatness scores were based on a five-point scale and each score class was further classified into high, medium and low to obtain a 15-point score for a more precise description of carcass traits. Carcasses were stored at 4°C for 24 h *post-mortem* and ultimate pH (pH<sub>24</sub>) was measured on the *longissimus thoracis* muscle using a pH-meter (Orion 9106). Colour descriptors lightness (L\*), redness (a\*), yellowness (b\*), Chroma (C\*), and Hue angle (H\*) were recorded on the *longissimus thoracis* muscle, using Minolta CM-2022 spectrophotometer (d/8° geometry; Minolta Co., Ltd. Osaka, Japan) on specular components excluded (SCE) mode, illuminant A and a 10° standard observer. After 24 h of storage at 4 °C, the entire *longissimus thoracis et lumborum* (LTL) muscle from the right half-carcass was removed, packed under vacuum and stored at -80 °C for analyses of intramuscular fatty acids and tocopherols. The entire LTL from the left half-carcass was vacuum-packed and stored at 4 °C for 3 days, pending oxidative stability measurements.

## 2.3. Feed analysis

### 2.3.1. Analysis of chemical composition, fatty acids and vitamin E

Dry matter (DM), crude protein, ether extract and ash content were determined in the experimental diets following the method of AOAC (1995). The neutral detergent fibre, acid detergent fibre and acid detergent lignin were analysed according to Van Soest, Robertson, and Lewis (1991). The fatty acid composition of the experimental diets was determined by a one-step extraction–transesterification procedure using chloroform (Sukhija & Palmquist, 1988) and 2% (v/v) sulfuric acid in methanol (Shingfield et al., 2003). Gas chromatographic (GC) analysis of fatty acid methyl esters (FAME) was performed as described later (section 2.3) for fatty acid profile of lamb meat. Individual fatty acids of experimental diets were expressed as mg/g of DM.

Vitamin E ( $\alpha$ -,  $\gamma$ - and  $\delta$ -tocopherols) was analysed in the experimental diets as described by Cherif et al. (2018). Briefly, freeze-dried samples were homogenized and saponified with ethanolic KOH stabilized with BHT. Tocopherols were extracted using hexane/ethyl acetate (9/1, v/v), dried under N<sub>2</sub> and dissolved in acetonitrile. Vitamin E was analysed by HPLC (Perkin Elmer series 200), equipped with an autosampler (model AS 950-10, Tokyo, Japan) and a Synergy Hydro-RP column (4  $\mu$ m, 4.6  $\times$  100 mm; Phenomenex, Bologna, Italy). Tocopherols were eluted at a flow rate of 2 ml/min and identified using a fluorescence detector (model Jasco, FP-1525) set at an excitation and emission  $\lambda$  of 295 nm and 328 nm, respectively. Quantification was based on external calibration curves of pure standard compounds (Sigma-Aldrich, Bornem, Belgium) solubilised in ethanol.

### 2.3.2. Analysis of total phenol content and in vitro antioxidant activity

Phenolic compounds were extracted from cardoon meal, dehydrated alfalfa and experimental diets using aqueous methanol (50:50, v/v) and acetone (70:30, v/v) solvents (Jiménez-Escrig, Jiménez-Jiménez, Pulido, & Saura-Calixto, 2001). Polyphenol-rich extracts were analysed for total phenol content (TPC) using the Folin-Ciocalteu method (Singleton, Orthofer, & Lamuela-Raventós, 1999) with minor modifications. Briefly, extracts (0.5 ml) were mixed with Folin-Ciocalteu reagent (2.5 ml, 20% in distilled water) and sodium carbonate (2 ml, 7.5% in distilled water) was added after 5 min. The mixture was stored in the dark for 2 h at room temperature and absorbance measurements were recorded at 750 nm using a UV-vis spectrophotometer (Cary 300 Bio, UV-vis spectrophotometer, Varian Instruments, CA, USA) against a blank containing all reagents and distilled water. A calibration curve using standard solutions of aqueous gallic acid (20 – 100 µg/ml) was plotted and results were expressed as g of gallic acid equivalents (GAE)/kg of DM feed.

*In vitro* antioxidant activity of polyphenol-rich extracts was measured using 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radical scavenging (Yen & Wu, 1999) and ferric reducing antioxidant power (FRAP) total antioxidant assays (Benzie & Strain, 1999), with minor modifications. For the DPPH assay, extract (0.6 ml) and distilled water (2.4 ml) were mixed with 0.2 mM DPPH in methanol (3 ml) and stored in the dark for 1 h at room temperature. Absorbance measurements were recorded at 517 nm using a UV-vis spectrophotometer (Cary 300 Bio) against a methanol blank. An assay blank containing distilled water (3 ml) and 0.2 mM DPPH in methanol (3 ml) was used for calculation purposes. A calibration curve using standard solutions of methanolic Trolox (10 – 50 µg/ml) was plotted and results were expressed as g of Trolox equivalents (TE)/kg of DM feed.

For the FRAP assay, extract (0.45 ml) was mixed with 8.55 ml FRAP reagent (a mixture of 30 mM acetate buffer (pH 3.6), 10 mM 2,4,6-Tris (2-pyridyl)-s-triazine in 40 mM HCl and 20 mM FeCl<sub>3</sub>.6H<sub>2</sub>O in distilled water in the ratio 10:1:1, respectively incubated at 37 °C for 10 min prior to use). The mixture was stored in the dark for 30 min at room temperature and absorbance measurements were recorded at 593 nm using a UV-vis spectrophotometer (Cary 300 Bio) against a blank containing all reagents. A calibration curve using standard solutions of methanolic Trolox (0.033 – 0.1 mg/ml) was plotted and results were expressed as g of Trolox equivalents (TE)/kg of DM feed.

#### ***2.4. Analysis of fatty acids and vitamin E in lamb meat***

Intramuscular fat was extracted from 10 g of minced LTL with a mixture of methanol and chloroform (2:1, v/v) according to the method of Folch, Lees, and Sloane Stanley (1957). Lipids (30 mg) were converted to FAME by base-catalysed transesterification (Christie, 1982) using 0.5 mL of sodium methoxide in methanol 0.5 N and 1 mL of hexane containing 1mg /mL nonadecanoic acid (C19:0) as an internal standard. Gas chromatographic analysis was conducted as described by Valenti et al. (2018) using a GC 8000 Top ThermoQuest (Milan, Italy) gas-chromatograph equipped with a flame ionization detector and a high polar column (WCOT-fused silica CP-Select CB for FAME Varian, Middelburg, the Netherlands; 100m×0.25mm i.d.; film thickness 0.25 µm). Helium was the carrier gas at a constant flow of 1 ml/min. Total FAME profile in a 1 µL sample volume (2 µL for feed samples) at a split ratio of 1:80 was determined using the following conditions: the oven temperature was programmed at 40 °C and held for 4 min, then increased to 120 °C at 10 °C/min, held for 1 min, then increased up to 180 °C at 5 °C/min, held for 18 min, then increased up to 200 °C at 2 °C/min, held for 15 min, and then increased up to 230 °C at 2 °C/min, held for 19 min. The injector and detector temperatures were

set at 270 °C and 300 °C, respectively. The identification of individual FAME was based on the retention time comparison with commercially available standard mixture of FAME (Nu-Chek Prep Inc., Elysian, MN, USA; Larodan Fine Chemicals, Malmo, Sweden) and comparison with chromatograms published by Kramer, Hernandez, Hernandez, Kraft, and Dugan (2008) and Alves and Bessa (2007). Fatty acids were expressed as g/100 g of total fatty acids. The dietary risk for cardiovascular diseases was assessed by calculating the atherogenic index (the relationship between FA with pro-atherogenic and anti-atherogenic properties) and thrombogenic index (the relationship between FA with pro-thrombogenic and anti-thrombogenic properties). The atherogenic index (AI) and thrombogenic index (TI) were calculated as outlined by Ulbricht and Southgate (1991):

$$AI = \frac{C12:0 + (4 \times C14:0) + C16:0}{n - 6 \text{ PUFA} + n - 3 \text{ PUFA} + \text{MUFA}}$$

$$TI = \frac{C14:0 + C16:0 + C18:0}{(0.5 \times \text{MUFA}) + (0.5 \times n - 6 \text{ PUFA}) + (3 \times n - 3 \text{ PUFA}) + \left(\frac{n - 3 \text{ PUFA}}{n - 6 \text{ PUFA}}\right)}$$

Vitamin E (α-, γ- and δ-tocopherols) and retinol were extracted from lamb muscle according to the method of Schüep and Rettenmaier (1994). Chromatographic analysis of vitamin E was performed as described in section 2.2.1 for feeds. Retinol was analyzed in the same chromatographic run and identified using the UV-VIS detector set at λ 325 nm (Cherif et al., 2018). Identification and quantification were achieved using external calibration curves of standard compounds (Sigma-Aldrich) solubilised in ethanol.

## 2.5. Measurement of lamb meat oxidative stability

Oxidative stability parameters were measured in raw and cooked lamb meat, as well as in LTL homogenates incubated with pro-oxidant catalysts, as described by Valenti et al. (2018). Briefly, for analyses on raw and cooked meat, 6 slices (2 cm thickness) were cut from the LTL which was previously stored under vacuum at 4 °C as described in section 2.1. Three slices were placed in polystyrene trays, covered with PVC film and stored at 4 °C. Each slice was used for measuring lipid oxidation and colour stability at one of three-time points: day 0 (after 2 hours of blooming), and days 4 and 7. The remaining 3 slices were packed under vacuum and cooked for 30 min at 70°C in a water bath. After cooling in a cold-water bath, one slice was used immediately for measurement of lipid oxidation, while the other 2 slices were stored at 4°C in the same conditions described for the raw meat samples, and lipid oxidation was measured after 2 and 4 days. For both raw and cooked meat, lipid oxidation was measured as thiobarbituric acid reactive substances (TBARS) values according to the procedure of Siu and Draper (1978) and results were expressed as µg malonaldehyde (MDA)/g of meat. Colour was measured in raw meat using a Minolta CM-2022 spectrophotometer (d/8° geometry; Minolta Co., Ltd. Osaka, Japan) set in the specular components excluded (SCE) mode, illuminant A and a 10° standard observer. The colour descriptors lightness (L\*), redness (a\*), yellowness (b\*), Chroma (C\*), and Hue angle (H\*) were recorded, as well as the reflectance spectra from 400 to 700 nm. The ratio  $(K/S)_{572} \div (K/S)_{525}$  was calculated to monitor the accumulation of metmyoglobin (MetMb) on the meat surface over time, with values of the ratio decreasing with increasing proportion of MetMb. The ratio (K/S) between the absorption (K) and the scattering (S) coefficients at the selected wavelengths was calculated as:

$$(K/S)_\lambda = (1 - R_\lambda)^2 / 2R_\lambda$$

The resistance of meat to lipid and myoglobin oxidation was also assessed in LTL homogenates incubated in the presence of  $\text{Fe}^{3+}$  and ascorbate (Fe/Asc) as catalysts of oxidative reactions. Briefly, the minced LTL (7.5 g) was homogenized with 37.5 g of MES buffer (pH 5.6). Homogenates were equilibrated to 37 °C and two aliquots (3 ml and 4 ml) were collected for measuring the initial extent of lipid and myoglobin oxidation (0 minutes). Ferric chloride hexahydrate and L-sodium ascorbate were added at equimolar concentration to reach the final concentration of 45  $\mu\text{M}$ . The homogenates were incubated under continuous stirring in a temperature controlled IKA KS-4000 orbital shaker (IKA-Werke GmbH & Co. KG, Staufen, Germany) set at 37°C and 190 rpm. After 30 and 60 minutes of incubation, two aliquots (3 ml and 4 ml) were collected for lipid and myoglobin oxidation analyses. Lipid oxidation was measured in the 3 ml aliquots using the method of Siu and Draper (1978). The 4 ml aliquots were centrifuged at  $6800 \times g$  at 4 °C, filtered through Whatman 541 filter paper and directly scanned in a UV/VIS spectrophotometer (UV-1601, Shimadzu Co., Milan, Italy). The absorbances at 503, 525, 557, 582, and 730 nm were used to calculate the proportion of metmyoglobin (MetMb %; Tang, Faustman, & Hoagland, 2004).

## ***2.6. Determination of antioxidant potential of cardoon extract in a muscle model system***

### ***2.6.1. Preparation of cardoon extract***

Phenolic compounds were extracted in triplicate by suspending finely-ground cardoon meal (0.5 g) in 100% methanol (20 ml) and incubated at room temperature in an orbital shaker (Max Q 6000 Shaker Thermo Fisher Scientific, Ireland) at 200 rpm for 4 h. After 4 h, the mixture was filtered through Whatman No. 1 filter paper. Pooled solvent extracts were concentrated by placing in a 50 ml round-bottomed flask and solvent (methanol) was removed by rotary evaporation (Labo-Rota C-311, Resona Technics, Switzerland) for 1 h at 55 °C. The dried

extract was re-suspended in 20 ml methanol and analysed for TPC using the Folin-Ciocalteu reagent as described for experimental feeds (section 2.2.1.). Result was expressed as mg GAE/ml extract.

### **2.6.2. Preparation of muscle homogenates**

Fresh lamb LTL ( $n = 3$ ) was obtained from a meat retail outlet (Cork, Ireland) and stored at 4 °C prior to analysis. Muscle homogenates (25%) were prepared in triplicate following a minor modification of the method described by O'Grady, Monahan, and Brunton (2001). Briefly, LTL (15 g) was chopped and homogenised in 0.12 M KCL 5 mM histidine (45 ml), pH 5.5, using an Ultra-turrax T25 homogeniser (Janke & Kunkel GmbH, IKA® Labortechnik, Staufen, Germany) at 24,000 rpm for 5 min. The muscle tissue and buffer were surrounded by crushed ice to control the temperature during homogenisation. Cardoon extract (stock concentration of 1.32 GAE mg/ml) was added to the homogenates (39.2 g) at 0, 0.5, 1 and 5% v/w of the final volume (40 ml). Lipid oxidation in muscle homogenate samples was initiated by the addition of equimolar FeCl<sub>3</sub>/sodium ascorbate (45 µM) pro-oxidants. The TPC and lipid oxidation in muscle homogenate were measured after 4 h of storage at 4 °C.

### **2.6.3. Measurement of TPC and lipid oxidation in muscle homogenate**

The total phenolic content in muscle homogenates was measured in triplicate by mixing 10% TCA (0.5 ml), 0.05 M phosphate buffer (3 ml) and muscle homogenate (2 g). The mixture was centrifuged at 7800 ×g for 10 min at 4 °C using an Avanti® J-E Centrifuge (Beckman Coulter Inc., Palo Alto, CA, USA). The supernatant was filtered through Whatman No. 1 paper and the filtrate was analysed for TPC using the Folin-Ciocalteu reagent as described for experimental feeds (section 2.2.2). Results were expressed as g GAE/g muscle. Lipid oxidation was measured

284 in triplicate following the TBARS assay described by Siu and Draper (1978) and results were  
285 expressed as  $\mu\text{g MDA/g meat}$ .

## 286 **2.7. Statistical analysis**

287 The average data of DMI, BW and feed efficiency for individual lambs were subjected to one-  
288 way ANOVA test to analyse the effect of dietary treatment on growth performance parameters.  
289 The effect of dietary treatment on carcass traits, intramuscular fatty acids and fat-soluble  
290 vitamins, were also analysed with one-way ANOVA test, using individual lambs as the  
291 experimental units. Data on the oxidative stability parameters (raw, cooked and muscle  
292 homogenates) were analysed with a full-repeated measures ANOVA. Effects of diet represented  
293 the ‘between-subjects’ factor and the effect of storage time/incubation was measured using the  
294 ‘within-subjects’ factor and the interaction between diet and storage time/incubation was tested.  
295 Individual lambs were considered as the experimental units in all the statistical analyses  
296 performed for the effect of dietary cardoon meal on growth performance and meat quality.  
297 Analyses relating to the effect of cardoon extract addition on TPC and lipid oxidation (TBARS)  
298 in LTL homogenates was performed in triplicate and mean sample values ( $n = 3$ ) for each of the  
299 four treatment groups (CON, CE0.5, CE1.0 and CE5.0) were subjected to a one-way ANOVA.  
300 Pearson’s correlation analysis was performed to assess the relationship between the TPC and  
301 TBARS in LTL homogenates. Significance was declared when  $P \leq 0.05$ , while a tendency for  
302 effects were considered when  $0.05 < P \leq 0.10$ . All statistical analyses were performed using the  
303 SPSS software (IBM Statistics version 22).

### 3. Results and Discussion

#### 3.1. Antioxidant activity of experimental feedstuffs and diets

The TPC of cardoon meal was 4-fold higher ( $P < 0.01$ ; 60.4 vs 15.5 g GAE/kg DM) than dehydrated alfalfa, which may explain the 2.5-fold increase in TPC when dehydrated alfalfa was replaced by cardoon meal in CMD, compared to CON (Table 1). The phenolic concentration of cardoon meal was 5-fold higher than previously reported for cardoon press cake (Genovese et al., 2015), a similar by-product obtained from the mechanical extraction of oil from cardoon seeds. Variation in the TPC of cardoon extracts may be related to factors such as plant geographical origin, varieties and maturity stage, agricultural methodologies, and phenol extraction procedures (Ramos et al., 2014). As expected, assessment of the *in vitro* antioxidant activities showed that cardoon meal exhibited a higher ( $P < 0.01$ ) DPPH free radical scavenging activity and FRAP total antioxidant activity compared to dehydrated alfalfa (Figure 1a). Similarly, replacement of dehydrated alfalfa with cardoon meal in the experimental diets resulted in higher ( $P < 0.01$ ) antioxidant activities in CMD compared to CON (Figure 1b).

Flavonoids and hydroxycinnamic acids are the main classes of phenolic compounds, which contribute to the antioxidant effect of extracts obtained from cardoon leaf, seed, stem and flower (Falleh et al., 2008; Pandino et al., 2011; Pinelli et al., 2007; Ramos et al., 2014). It has been shown that the TPC of cardoon extracts strongly correlated with DPPH antiradical activity (Falleh et al., 2008; Ramos et al., 2014) and FRAP total antioxidant capacity (Pandino et al., 2011). In addition,  $\alpha$ -tocopherol is another potent antioxidant which can be found in high residual levels in cardoon meal as cardoon oil has been shown to contain considerable amounts of vitamin E (Maccarone et al., 1999). This is consistent with the 4-fold increase in  $\alpha$ -tocopherol content found in CMD compared to CON (Table 1).

### 3.2. Growth performance and carcass traits

Lambs fed CMD exhibited lower ( $P < 0.05$ ) DMI compared to CON-fed lambs. Lower DMI in lambs fed CMD may be related to low feed palatability due to the high content of dietary phenolic compounds. Cajarville, González, Repetto, Rodríguez, and Martínez (1999) reported a similar decrease in the voluntary intake of sheep fed *ad libitum* green forage of cardoon possibly due to the presence of high concentration of phenolic compounds in cardoon leaves (Kukić et al., 2008). Phenolic compounds, such as tannins, may confer unpleasant taste or bind to salivary proteins forming a polyphenol-protein complex that induce astringency sensations and trigger low feed intake in animals (Makkar, 2003). However, dietary treatment did not affect ( $P > 0.05$ ) growth performance parameters in terms of final BW, ADG and feed efficiency (Table 2). Differences in DMI did not reflect on growth performance parameters possibly due to the low number of experimental units or that reduced DMI in CMD-fed lambs did not compromise nutrient digestibility and utilization. Accordingly, Cajarville, González, Repetto, Alvir, and Rodríguez (2000) reported that up to 25% cardoon seeds could be included in ruminant diets without negative effect on ruminal fermentation pattern and nutrient digestibility. Furthermore, there was no effect of dietary treatment on carcass characteristics such as carcass weight, dressing percentage, carcass conformation, fatness scores, muscle ultimate pH and colour characteristics (Table 2). Notably, values of ultimate pH were slightly higher than the normal pH values (5.4 – 5.8) for post-mortem muscle (Faustman & Cassens, 1990) particularly for lambs fed CON due to the fact that one carcass displayed an ultimate pH of 6.3. Nonetheless, the values of colour variables are within the satisfactory range for average consumer acceptance of fresh meat, particularly in terms of lightness ( $L^*$ ) and redness ( $a^*$ ) (Khlijji, Van de Ven, Lamb, Lanza, & Hopkins, 2010).

### 3.3. Effect of dietary cardoon meal on lamb meat fatty acid composition and fat-soluble vitamin levels

Intramuscular fat, fatty acid profile and fat-soluble vitamin contents in LTL muscle from lambs fed CON and CMD diets are presented in Table 3. In general, dietary treatment did not influence ( $P > 0.05$ ) intramuscular fat content or the total composition of SFA, MUFA, PUFA, odd- and branched-chain fatty acids (OBCFA). Dietary treatment did not influence ( $P > 0.05$ ) the concentration of individual SFA (C12:0, C14:0 and C16:0) that results in hypercholesterolemic effects in humans (Mensink, 2005). This could in part explain why meat from lambs fed CON and CMD had similar ( $P > 0.05$ ) nutritional indices (atherogenic index and thrombogenic index) for risk of cardiovascular diseases. However, lambs fed CMD displayed higher concentration ( $P < 0.05$ ) of intramuscular C18:0 (stearic acid), and lower proportions ( $P < 0.05$ ) of total *trans*-18:1, *trans*-10 C18:1, *trans*-11 C18:1 (vaccenic acid) and *cis*-9 *trans*-11 C18:2 CLA (rumenic acid) compared to CON-fed lambs. In addition, the proportion of C18:3 *n*-3 ( $\alpha$ -linolenic acid) tended ( $P = 0.070$ ) to be greater in the muscle of lambs fed CON compared to CMD-fed lambs. Consequently, the ratio of *n*-6:*n*-3 PUFA ( $P = 0.052$ ) tended to be lower in CON-fed lambs compared to CMD-fed lambs. Regarding fat-soluble vitamins in LTL, the concentration of retinol ( $P = 0.048$ ) was greater in muscle from lambs fed CMD relative to CON but diet did not affect ( $P > 0.05$ ) vitamin E ( $\alpha$ -,  $\gamma$ - and  $\delta$ -tocopherols) concentration (Table 3).

Indeed, changes in intramuscular fatty acids might be due to possible differences in ruminal biohydrogenation. The present study showed that dietary treatment significantly affected the intramuscular concentration of individual fatty acids (stearic, *trans*-10 C18:1, vaccenic and rumenic acids) with potential nutritional implications. It has been suggested that dietary consumption of stearic acid does not increase plasma low-density lipoprotein and cholesterol

levels in humans in contrast to the negative effect of other SFA that are risk factors for cardiovascular diseases (Mensink, 2005). Thus, the greater amount of stearic acid in lambs fed CMD may not have detrimental effects on human health. Muscle from lambs fed CON exhibited a higher proportion of vaccenic and rumenic acids that are known for their potential health benefits in humans (Bessa et al., 2015). Both vaccenic and rumenic acids are intermediate FA synthesized during ruminal biohydrogenation but rumenic acid can be further synthesized in muscle tissues through endogenous desaturation of vaccenic acid by the enzyme  $\Delta$ -9-desaturase (Corl et al., 2001). Thus, higher concentration of intramuscular rumenic acid in CON-fed lambs may be due to greater ruminal outflow and absorption of vaccenic acid that serves as the main precursor for the endogenous synthesis of rumenic acid in muscle tissues. Rumenic acid is the major naturally occurring CLA isomer found in ruminant meat and milk (Bessa et al., 2015) and dietary consumption of this FA potentially prevent human diseases including cancer, cardiovascular diseases, obesity, bone density loss, and diabetes (McGuire & McGuire, 2000).

From a nutritional perspective, the accumulation *t*-10 C18:1 in ruminant edible products is undesirable due to the potential cytotoxic effect of this fatty acid (Vahmani et al., 2016). Thus, a lower concentration of *t*-10 C18:1 in meat of lambs fed CMD may be viewed as a positive outcome compared to CON. Concentrate feeding systems are known to promote the alternative ruminal biohydrogenation pathway that increase the synthesis of intermediates such as *t*-10 18:1, at the expense of *t*-11 18:1, referred to as the *t*-10 shift (Bessa et al. 2015). Bessa et al. (2015) postulated that the ratio of *t*-10/*t*-11 18:1 >1 in ruminant meat or milk is an indicator of occurrence of *trans*-10 shift during ruminal biohydrogenation. Considering that concentrate diets were fed in this study, it is not surprising that *t*-10/*t*-11 18:1 was > 1 in meat from lambs fed CON and CMD, suggesting that both diets induced *t*-10 shift. However, feeding CMD tended (*P*

396 = 0.084) to reduce the occurrence of the *t*-10 shift, suggesting that the presence of phenolics in  
397 CMD might have modified the ruminal biohydrogenation pathway.

398 A number of factors may account for the lack of positive effect of CMD to inhibit ruminal  
399 biohydrogenation and increase PUFA and CLA contents in meat. It is possible that  
400 hydroxycinnamic acids, the major cardoon phenolic compounds, were metabolised by the  
401 consortium of microbes residing in the fore-stomach (rumen) of lambs as shown with human  
402 faecal microbiota (Gonthier et al., 2006). Furthermore, effects of dietary phenolics to increase  
403 PUFA and CLA contents in ruminant meat have been largely reported for diets containing a high  
404 concentration of tannins (Vasta, Nudda, Cannas, Lanza, & Priolo, 2008). Though the tannin  
405 content of diets was not measured in the present study, it has been shown that cardoon extracts  
406 contain a very low concentration of tannins (Falleh et al., 2008). Thus, results from the present  
407 study emphasized the variation in the effect of diets containing a high concentration of bioactive  
408 compounds, which may be dependent on the type and/or concentration of phenolic compounds  
409 present. In contrast, alfalfa fed in various processed forms (fresh, hay, dehydrated, silage) has  
410 been documented to enrich lamb muscle and offal with lower SFA and higher content of PUFA  
411 and rumenic acid (Cerci, Ciftci, Bahsi, & Kilinc, 2011; Ciftci et al., 2010; Realini, Bianchi,  
412 Bentancur, & Garibotto, 2017). The positive effects of dietary alfalfa on the fatty acid profile of  
413 ruminant meat may be due to the presence of saponins and flavonoids which inhibit ruminal  
414 biohydrogenation (Petersen & Jensen, 2014).

#### 415 ***3.4. Effect of dietary cardoon meal on oxidative stability of lamb meat***

416 The results of oxidative stability of raw meat, cooked meat and muscle homogenates are  
417 presented in Table 4. As expected, lipid oxidation (TBARS values) significantly increased ( $P <$   
418 0.001) in raw meat as a function of storage time. Although in raw meat an overall effect of the

dietary treatment was not found on lipid oxidation, and interactive effect of the dietary treatment and time of storage was found ( $P < 0.05$ ), with higher TBARS values measured in raw meat slices from the CMD-fed lambs compared to the CON treatment after 7 days of storage (Figure 2). This result is not easy to explain, considering the lack of effect of the dietary treatment on the concentration of PUFA and  $\alpha$ -tocopherol in muscle. It can be speculated that a higher content of CLA (cis-9 trans-11 18:2) in the meat from CON-fed lambs could be responsible for the reduced lipid oxidation as CLA may exert a protective effect on muscle oxidation (Joo, Lee, Ha, & Park, 2002). Hur et al. (2004) also reported that the CLA present in meat does not participate in oxidation processes and reduces the formation of fatty acid free radicals, which results in reduced lipid oxidation. Noteworthy, as shown in Table 4, neither the dietary, nor the diet  $\times$  time interaction affected the colour stability of raw lamb meat measured using instrumental colour descriptors (lightness  $L^*$ , redness  $a^*$ , yellowness  $b^*$ , saturation  $C^*$  and hue angle  $H^*$ ) and an index for metmyoglobin accumulation ( $(K/S)_{572} \div (K/S)_{525}$ ). The measured parameters mostly related to meat browning were only affected by the storage period, with redness, saturation and the  $(K/S)_{572} \div (K/S)_{525}$  decreasing over time ( $P < 0.05$ ), while hue increased ( $P < 0.05$ ). Additionally, as shown in Table 4, the use of stronger pro-oxidant conditions (i.e. cooking and incubation of muscle homogenates with  $FeCl_3$ /sodium ascorbate pro-oxidants) greatly promoted lipid and myoglobin oxidation which markedly increased over time ( $P < 0.001$ ). However, the different susceptibility to lipid oxidation observed in raw meat was not evident in cooked meat and in muscle homogenates, which suggests that such oxidative challenges might have overcome inherent differences between treatments in the susceptibility to oxidation.

In general, feeding CMD did not improve the oxidative stability of meat despite the greater content of phenolic compounds and the consequently higher antioxidant capacity of the CMD

diet compared to CON (Table 1 and Figure 1b). Indeed, the *in vivo* antioxidant potential of dietary phenolics is still controversial. There are evidences that phenolics may exhibit antioxidant effect through direct absorption and deposition into muscle tissues or through indirect antioxidant mechanisms including the sparing effect of polyphenols on other antioxidants, such as ascorbic acid and tocopherols (Valenti et al., 2018). As previously highlighted in this study, possible microbial metabolism of cardoon phenolics in the gut of lambs may also account for the lack of functional *in vivo* effect in lambs fed CMD.

### **3.5. Effect of cardoon extract on lipid oxidation in a muscle-based system**

Antioxidant effect of the phenolic-rich extract obtained from cardoon meal was tested in LTL homogenates subjected to FeCl<sub>3</sub>/sodium ascorbate-induced oxidation by incubating with iron/ascorbate pro-oxidants for 4 h. Addition of 5% cardoon extract significantly increased (+114.3%;  $P < 0.05$ ) the total phenolic content (TPC) in LTL homogenates compared to the control (Figure 3a). Similarly, addition of 5% extract inhibited (-77.6%;  $P < 0.001$ ) lipid oxidation (TBARS values) relative to the control (Figure 3b). Pearson's correlation analysis showed that TPC negatively correlated ( $r = -0.724$ ;  $P = 0.008$ ) with TBARS values, suggesting that cardoon phenolics contributed to the inhibition of lipid oxidation in LTL homogenates. The present study suggests that the antioxidant effect of cardoon extract in an ovine muscle system is dose-dependent as only the addition of 5% cardoon extract significantly inhibited lipid oxidation relative to the control. Similarly, Falleh et al. (2008) demonstrated that cardoon extracts exhibited *in vitro* antioxidant activities (DPPH radical and superoxide anion scavenging activities) in a concentration-dependent manner.

Several plant extracts have been shown to exhibit antioxidant efficacy with potential application as natural alternatives to potentially-toxic synthetic antioxidants in meat products (Jiang &

Xiong, 2016). Indeed, the present study is the first to demonstrate the antioxidant potential of cardoon extract in muscle model systems. Moreover, the addition of a cardoon extract can enhance the healthiness of meat products as cardoon phenolics may exert functional therapeutic properties such as antimicrobial, antimutagenic, hepatoprotective, choleretic and anti-cholestatic actions in humans (Adzet, Camarasa, & Laguna, 1987; Falleh et al., 2008). Therefore, further research is required to examine the use of cardoon extract as a natural additive for developing functional meat products with extended shelf-life and health-promoting properties.

#### **4. Conclusions**

The present study demonstrated that cardoon meal is a rich source of phenolic compounds with potent antioxidant activity. The inclusion of 15% cardoon meal replacing dehydrated alfalfa in a concentrate diet had no adverse effect on lamb growth performance but depressed feed intake. Dietary cardoon meal did not influence the intramuscular composition of SFA, MUFA, PUFA and nutritional indices (atherogenic index and thrombogenic index) but presented a lower concentration of potentially health-promoting fatty acids (vaccenic and rumenic acids) in lamb meat. Moreover, dietary inclusion of cardoon meal did not improve the oxidative stability of raw and cooked meat stored aerobically at 4 °C for up to 7 and 4 days, respectively. In addition, a phenolic-rich extract obtained from cardoon meal exhibited potent antioxidant activity against lipid oxidation in an ovine muscle model system. Further research is required to evaluate the antioxidant effect of cardoon extract on the shelf-life and quality parameters of meat products.

#### **Acknowledgements**

This work was conducted as part of a PhD thesis project supported by the Agricultural Transformation by Innovation (AGTRAIN) Erasmus Mundus Joint Doctorate Program, funded

by the EACEA (Education, Audiovisual and Culture Executive Agency) of the European Commission.

## References

Adzet, T., Camarasa, J., & Laguna, J. C. (1987). Hepatoprotective activity of polyphenolic compounds from *Cynara scolymus* against CCl<sub>4</sub> toxicity in isolated rat hepatocytes. *Journal of Natural Products*, 50, 612-617.

Alves, S.P., & Bessa, R.J. (2007). Identification of cis-12, cis-15 octadecadienoic acid and other minor polyenoic fatty acids in ruminant fat. *European Journal of Lipid Science and Technology*, 109, 879-883.

AOAC, Association of Official Analytical Chemists (1995). *Official methods of analysis*, AOAC (16th ed.), Arlington, VA, USA.

Balasundram, N., Sundram, K., & Samman, S. (2006). Phenolic compounds in plants and agricultural by-products: Antioxidant activity, occurrence, and potential uses. *Food Chemistry*, 99, 191-203.

Balzan, S., Taticchi, A., Cardazzo, B., Urbani, S., Servili, M., Di Lecce, G., . . . Fasolato, L. (2017). Effect of phenols extracted from a by-product of the oil mill on the shelf-life of raw and cooked fresh pork sausages in the absence of chemical additives. *LWT-Food Science and Technology*, 85, 89-95.

Benzie, I. F., & Strain, J. (1999). Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods in Enzymology*, 299, 15-27.

- Bessa, R. J., Alves, S. P., & Santos-Silva, J. (2015). Constraints and potentials for the nutritional modulation of the fatty acid composition of ruminant meat. *European Journal of Lipid Science and Technology*, 117, 1325-1344.
- Cajarville, C., González, J., Repetto, J. L., Rodríguez, C. A., & Martínez, A. (1999). Nutritive value of green forage and crop by-products of *Cynara cardunculus*. *Annales de Zootechnie*, 48, 353-365.
- Cajarville, C., González, J., Repetto, J.L., Alvir, M.R., & Rodríguez, C.A. (2000). Nutritional evaluation of cardoon (*Cynara cardunculus*) seed for ruminants. *Animal Feed Science and Technology*, 87, 203-213.
- Cerci, I. H., Ciftci, M., Bahsi, M., & Kilinc, U. (2011). Cholesterol and fatty acid composition of lamb serum and offal as affected by alfalfa and concentrate. *Veterinarski Arhiv*, 81, 575-584.
- Cherif, M., Valenti, B., Abidi, S., Luciano, G., Mattioli, S., Pauselli, M., . . . Salem, H. B. (2018). Supplementation of *Nigella sativa* seeds to Barbarine lambs raised on low-or high-concentrate diets: Effects on meat fatty acid composition and oxidative stability. *Meat Science*, 139, 134-141.
- Christie, W. W. (1982). A simple procedure for rapid transmethylation of glycerolipids and cholesteryl esters. *Journal of Lipid Research*, 23, 1072-1075.
- Ciftci, M., Cerci, I., Kilinc, U., Yilmaz, O., Gurdogan, F., Seven, P., . . . Ozcelik, M. (2010). Effects of alfalfa (fresh, silage, hay) on the fatty acid and conjugated linoleic acid amounts in lamb muscles and fats. *Revue de Médecine Vétérinaire*, 161, 432-437.

- Commission Regulation (EC) (1998). *Commission Regulation (EC) 823/98 of 20 April 1998 amending Regulation (EEC) 461/93 laying down detailed rules for the Community scale for the classification of carcasses of ovine animals.*
- Corl, B. A., Baumgard, L. H., Dwyer, D. A., Griinari, J. M., Phillips, B. S., & Bauman, D. E. (2001). The role of  $\Delta$  9-desaturase in the production of cis-9, trans-11 CLA. *The Journal of Nutritional Biochemistry*, 12, 622-630.
- Falleh, H., Ksouri, R., Chaieb, K., Karray-Bouraoui, N., Trabelsi, N., Boulaaba, M., & Abdelly, C. (2008). Phenolic composition of *Cynara cardunculus* L. organs, and their biological activities. *Comptes Rendus Biologies*, 331, 372-379.
- Faustman, C., & Cassens, R. (1990). The biochemical basis for discoloration in fresh meat: a review. *Journal of Muscle Foods*, 1, 217-243.
- Folch, J., Lees, M., & Sloane Stanley, G. (1957). A simple method for the isolation and purification of total lipids from animal tissues. *The Journal of Biological Chemistry*, 226, 497-509.
- Genovese, C., Platania, C., Venticinque, M., Calderaro, P., Argento, S., Scandurra, S., & Raccuia, S. (2015). Evaluation of cardoon seeds presscake for animal feeding. *Acta Horticulturae*, 1147, 323-328.
- Gonthier, M.-P., Remesy, C., Scalbert, A., Cheynier, V., Souquet, J.-M., Poutanen, K., & Aura, A.-M. (2006). Microbial metabolism of caffeic acid and its esters chlorogenic and caftaric acids by human faecal microbiota in vitro. *Biomedicine and Pharmacotherapy*, 60, 536-540.
- Hur, S., Ye, B., Lee, J., Ha, Y., Park, G., & Joo, S. (2004). Effects of conjugated linoleic acid on color and lipid oxidation of beef patties during cold storage. *Meat Science*, 66, 771-775.

- 553 Jiang, J., & Xiong, Y. L. (2016). Natural antioxidants as food and feed additives to promote  
554 health benefits and quality of meat products: A review. *Meat Science*, 120, 107-117.
- 555 Jiménez-Escrig, A., Jiménez-Jiménez, I., Pulido, R., & Saura-Calixto, F. (2001). Antioxidant  
556 activity of fresh and processed edible seaweeds. *Journal of the Science of Food and  
557 Agriculture*, 81, 530-534.
- 558 Joo, S.T., Lee, J.I., Ha, Y.L. & Park, G.B. (2002). Effects of dietary conjugated linoleic acid on  
559 fatty acid composition, lipid oxidation, color, and water-holding capacity of pork loin.  
560 *Journal of Animal Science*, 80, 108-112.
- 561 Kanatt, S. R., Chander, R., Radhakrishna, P., & Sharma, A. (2005). Potato peel extract a natural  
562 antioxidant for retarding lipid peroxidation in radiation processed lamb meat. *Journal of  
563 Agricultural and Food Chemistry*, 53, 1499-1504.
- 564 Khlijji, S., Van de Ven, R., Lamb, T., Lanza, M., & Hopkins, D. (2010). Relationship between  
565 consumer ranking of lamb colour and objective measures of colour. *Meat Science*, 85,  
566 224-229.
- 567 Kramer, J.K., Hernandez, M., Cruz-Hernandez, C., Kraft, J. & Dugan, M.E. (2008). Combining  
568 results of two GC separations partly achieves determination of all cis and trans 16: 1, 18:  
569 1, 18: 2 and 18: 3 except CLA isomers of milk fat as demonstrated using Ag-ion SPE  
570 fractionation. *Lipids*, 43, 259-273.
- 571 Kukić, J., Popović, V., Petrović, S., Mucaji, P., Ćirić, A., Stojković, D., & Soković, M. (2008).  
572 Antioxidant and antimicrobial activity of *Cynara cardunculus* extracts. *Food Chemistry*,  
573 107, 861-868.

- Lanza, M., Scerra, M., Bognanno, M., Buccioni, A., Cilione, C., Biondi, L., . . . Luciano, G. (2015). Fatty acid metabolism in lambs fed citrus pulp. *Journal of Animal Science*, 93, 3179-3188.
- Luciano, G., Pauselli, M., Servili, M., Mourvaki, E., Serra, A., Monahan, F., . . . Mele, M. (2013). Dietary olive cake reduces the oxidation of lipids, including cholesterol, in lamb meat enriched in polyunsaturated fatty acids. *Meat Science*, 93, 703-714.
- Maccarone, E., Fallico, B., Fanella, F., Mauromicale, G., Raccuia, S., & Foti, S. (1999). Possible alternative utilization of *Cynara* spp.: II. Chemical characterization of their grain oil. *Industrial Crops and Products*, 10, 229-237.
- Makkar, H. (2003). Effects and fate of tannins in ruminant animals, adaptation to tannins, and strategies to overcome detrimental effects of feeding tannin-rich feeds. *Small Ruminant Research*, 49, 241-256.
- McAfee, A. J., McSorley, E. M., Cuskelly, G. J., Moss, B. W., Wallace, J. M., Bonham, M. P., & Fearon, A. M. (2010). Red meat consumption: An overview of the risks and benefits. *Meat Science*, 84, 1-13.
- McGuire, M., & McGuire, M. (2000). Conjugated linoleic acid (CLA): A ruminant fatty acid with beneficial effects on human health. *Journal of Animal Science*, 77, 1-8.
- Mensink, R. P. (2005). Effects of stearic acid on plasma lipid and lipoproteins in humans. *Lipids*, 40, 1201-1205.
- Morrissey, P., Sheehy, P., Galvin, K., Kerry, J., & Buckley, D. (1998). Lipid stability in meat and meat products. *Meat Science*, 49, S73-86.
- O'Grady, M., Monahan, F., & Brunton, N. (2001). Oxymyoglobin oxidation and lipid oxidation in bovine muscle—mechanistic studies. *Journal of Food Science*, 66, 386-392.

- Pandino, G., Lombardo, S., Mauromicale, G., & Williamson, G. (2011). Phenolic acids and flavonoids in leaf and floral stem of cultivated and wild *Cynara cardunculus* L. genotypes. *Food Chemistry*, 126, 417-422.
- Petersen, M. B., & Jensen, S. K. (2014). Biohydrogenation of fatty acids is dependent on plant species and feeding regimen of dairy cows. *Journal of Agricultural and Food Chemistry*, 62, 3570-3576.
- Pinelli, P., Agostini, F., Comino, C., Lanteri, S., Portis, E., & Romani, A. (2007). Simultaneous quantification of caffeoyl esters and flavonoids in wild and cultivated cardoon leaves. *Food Chemistry*, 105, 1695-1701.
- Ramos, P. A., Santos, S. A., Guerra, Â. R., Guerreiro, O., Freire, C. S., Rocha, S. M., . . . Silvestre, A. J. (2014). Phenolic composition and antioxidant activity of different morphological parts of *Cynara cardunculus* L. var. *altilis* (DC). *Industrial Crops and Products*, 61, 460-471.
- Realini, C., Bianchi, G., Bentancur, O., & Garibotto, G. (2017). Effect of supplementation with linseed or a blend of aromatic spices and time on feed on fatty acid composition, meat quality and consumer liking of meat from lambs fed dehydrated alfalfa or corn. *Meat Science*, 127, 21-29.
- Rodríguez-Carpena, J., Morcuende, D., & Estévez, M. (2011). Avocado by-products as inhibitors of color deterioration and lipid and protein oxidation in raw porcine patties subjected to chilled storage. *Meat Science*, 89, 166-173.
- Salami, S., Guinguina, A., Agboola, J., Omede, A., Agbonlahor, E., & Tayyab, U. (2016). In vivo and postmortem effects of feed antioxidants in livestock: a review of the implications on authorization of antioxidant feed additives. *Animal*, 10, 1375-1390.

- Schüep, W., & Rettenmaier, R. (1994). Analysis of vitamin E homologs in plasma and tissue: High-performance liquid chromatography *Methods in Enzymology*, 234, 294-302.
- Shingfield, K., Ahvenjärvi, S., Toivonen, V., Ärölä, A., Nurmela, K., Huhtanen, P., & Griinari, J. M. (2003). Effect of dietary fish oil on biohydrogenation of fatty acids and milk fatty acid content in cows. *Animal Science*, 77, 165-179.
- Shingfield, K., Bonnet, M., & Scollan, N. (2013). Recent developments in altering the fatty acid composition of ruminant-derived foods. *Animal*, 7, 132-162.
- Singleton, V. L., Orthofer, R., & Lamuela-Raventós, R. M. (1999). [14] Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent *Methods in Enzymology*, 299, 152-178.
- Siu, G., & Draper, H. (1978). A survey of the malonaldehyde content of retail meats and fish. *Journal of Food Science*, 43, 1147-1149.
- Sukhija, P. S., & Palmquist, D. (1988). Rapid method for determination of total fatty acid content and composition of feedstuffs and feces. *Journal of Agricultural and Food Chemistry*, 36, 1202-1206.
- Tang, J., Faustman, C., & Hoagland, T. (2004). Krzywicki revisited: Equations for spectrophotometric determination of myoglobin redox forms in aqueous meat extracts. *Journal of Food Science*, 69, C717-720.
- Ulbricht, T., & Southgate, D. (1991). Coronary heart disease: seven dietary factors. *The Lancet*, 338, 985-992.
- Vahmani, P., Meadus, W. J., da Silva, M. L., Mitchell, A. D., Mapiye, C., Duff, P., . . . Dugan, M. E. (2016). A trans10-18: 1 enriched fraction from beef fed a barley grain-based diet

- induces lipogenic gene expression and reduces viability of HepG2 cells. *Biochemistry and Biophysics Reports*, 7, 84-90.
- Valenti, B., Natalello, A., Vasta, V., Campidonico, L., Roscini, V., Mattioli, S., . . . Luciano, G. (2018). Effect of different dietary tannin extracts on lamb growth performances and meat oxidative stability: comparison between mimosa, chestnut and tara. *Animal*, 1-9.
- Van Soest, P. V., Robertson, J., & Lewis, B. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74, 3583-3597.
- Vasta, V., & Luciano, G. (2011). The effects of dietary consumption of plants secondary compounds on small ruminants' products quality. *Small Ruminant Research*, 101, 150-159.
- Vasta, V., Nudda, A., Cannas, A., Lanza, M., & Priolo, A. (2008). Alternative feed resources and their effects on the quality of meat and milk from small ruminants. *Animal Feed Science and Technology*, 147, 223-246.
- Yen, G. C., & Wu, J. Y. (1999). Antioxidant and radical scavenging properties of extracts from *Ganoderma tsugae*. *Food Chemistry*, 65, 375-379.

**Table 1.** Chemical composition of experimental diets

| Parameter                           | Diets <sup>1</sup> |        |
|-------------------------------------|--------------------|--------|
|                                     | CON                | CMD    |
| Dry matter (DM), % <i>as-fed</i>    | 89.65              | 89.63  |
| Crude protein, % DM                 | 15.67              | 16.45  |
| Ether extract, % DM                 | 2.68               | 3.84   |
| Ash, % DM                           | 7.01               | 6.31   |
| NDF, % DM                           | 30.36              | 27.32  |
| ADF, % DM                           | 15.97              | 12.39  |
| ADL, % DM                           | 3.62               | 4.15   |
| Total phenolic content <sup>2</sup> | 5.21               | 13.08  |
| <i>Fatty acids (mg/g DM)</i>        |                    |        |
| C14:0                               | 0.034              | 0.034  |
| C16:0                               | 4.357              | 5.062  |
| <i>cis</i> -9 C16:1                 | 0.035              | 0.034  |
| C18:0                               | 0.454              | 0.705  |
| C18:1 <i>n</i> -9                   | 3.855              | 5.362  |
| <i>cis</i> -11 C18:1                | 0.209              | 0.206  |
| C18:2 <i>n</i> -6                   | 12.190             | 16.852 |
| C18:3 <i>n</i> -3                   | 1.255              | 1.065  |
| C20:0                               | 0.088              | 0.086  |
| <i>Vitamins (µg/g DM)</i>           |                    |        |
| α-Tocopherol                        | 5.267              | 20.877 |
| γ-Tocopherol                        | 0.155              | 0.119  |
| δ-Tocopherol                        | 0.012              | 0.005  |

<sup>1</sup>Diets were: CON (control diet), CMD (cardoon meal diet)

<sup>2</sup>Expressed as grams gallic acid equivalents/kg DM

NDF: neutral detergent fibre; ADF: acid detergent fibre; ADL: acid detergent lignin

**Table 2.** Effect of dietary treatment on growth performance, carcass characteristics and colour of *longissimus thoracis et lumborum* muscle of lambs.

|                                 | Dietary <sup>1</sup> |       |       |         |
|---------------------------------|----------------------|-------|-------|---------|
| Parameter                       | CON                  | CMD   | SEM   | P-value |
| <i>Growth performance</i>       |                      |       |       |         |
| Dry matter intake, g/d          | 1078.3               | 932.2 | 31.15 | 0.013   |
| Initial body weight, kg         | 20.1                 | 20.1  | 0.54  | 0.974   |
| Final body weight, kg           | 35.5                 | 33.9  | 0.74  | 0.303   |
| Average daily gain, g/d         | 204.5                | 184.0 | 8.04  | 0.215   |
| Feed efficiency <sup>2</sup>    | 190.8                | 198.7 | 8.24  | 0.649   |
| <i>Carcass traits</i>           |                      |       |       |         |
| Carcass weight, kg              | 17.1                 | 16.1  | 0.41  | 0.264   |
| Dressing percent, %             | 48.1                 | 47.5  | 0.29  | 0.333   |
| Conformation score <sup>3</sup> | 2.3                  | 1.9   | 0.13  | 0.102   |
| Fatness score <sup>4</sup>      | 2.5                  | 2.4   | 0.04  | 0.738   |
| Ultimate pH                     | 5.91                 | 5.83  | 0.04  | 0.310   |
| Lightness, L <sup>*</sup>       | 37.95                | 38.72 | 0.58  | 0.528   |
| Redness, a <sup>*</sup>         | 11.90                | 12.53 | 0.43  | 0.481   |
| Yellowness, b <sup>*</sup>      | 6.93                 | 7.32  | 0.34  | 0.588   |
| Saturation, C <sup>*</sup>      | 13.78                | 14.51 | 0.53  | 0.512   |
| Hue angle, H <sup>*</sup>       | 29.96                | 30.23 | 0.56  | 0.817   |

<sup>1</sup>Diets were: CON (control diet), CMD (cardoon meal diet)

<sup>2</sup>Calculated as: g BW gain/kg DMI

<sup>3</sup>Conformation score: E = 5, excellent shape and muscularity; U = 4; R = 3; O = 2; P = 1, poor shape and muscularity.

<sup>4</sup>Fatness score: 1 = low, 5 = very high

**Table 3.** Effect of dietary treatment on the intramuscular fat content, fatty acid composition and fat soluble vitamins in the *longissimus thoracis et lumborum* muscle of lambs

| Parameter                                         | Dietary <sup>1</sup> |       | SEM   | P-value |
|---------------------------------------------------|----------------------|-------|-------|---------|
|                                                   | CON                  | CMD   |       |         |
| IMF (g/100g of muscle)                            | 2.07                 | 2.06  | 0.117 | 0.946   |
| <i>Fatty acids (g/100 g of total fatty acids)</i> |                      |       |       |         |
| C12:0                                             | 0.13                 | 0.13  | 0.007 | 0.920   |
| C14:0                                             | 2.90                 | 2.81  | 0.121 | 0.727   |
| <i>cis</i> -9 C14:1                               | 0.11                 | 0.10  | 0.007 | 0.367   |
| C15:0                                             | 0.36                 | 0.36  | 0.011 | 0.848   |
| <i>iso</i> C15:0                                  | 0.07                 | 0.06  | 0.002 | 0.146   |
| <i>anteiso</i> C15:0                              | 0.11                 | 0.11  | 0.005 | 0.907   |
| C16:0                                             | 23.60                | 22.81 | 0.245 | 0.110   |
| <i>cis</i> -9 C16:1                               | 1.77                 | 1.57  | 0.075 | 0.174   |
| C17:0                                             | 1.15                 | 1.22  | 0.052 | 0.544   |
| <i>iso</i> C17:0                                  | 0.37                 | 0.33  | 0.010 | 0.065   |
| <i>anteiso</i> C17:0                              | 0.49                 | 0.45  | 0.011 | 0.060   |
| C18:0                                             | 12.13                | 13.80 | 0.267 | <0.001  |
| <i>cis</i> -9 C18:1                               | 38.78                | 39.56 | 0.447 | 0.403   |
| <i>cis</i> -11 C18:1                              | 1.58                 | 1.58  | 0.047 | 0.994   |
| <i>trans</i> -6 + 7 + 8 18:1                      | 0.28                 | 0.24  | 0.011 | 0.027   |
| <i>trans</i> -9 C18:1                             | 0.26                 | 0.22  | 0.011 | 0.066   |
| <i>trans</i> -10 C18:1                            | 1.58                 | 0.72  | 0.149 | 0.001   |
| <i>trans</i> -11 C18:1                            | 0.72                 | 0.48  | 0.055 | 0.006   |
| C18:2 <i>n</i> -6                                 | 6.85                 | 6.89  | 0.323 | 0.944   |
| <i>cis</i> -9 <i>trans</i> -11 C18:2*             | 0.43                 | 0.30  | 0.024 | 0.002   |
| <i>cis</i> -11 <i>trans</i> -13 C18:2             | 0.02                 | 0.01  | 0.002 | 0.064   |
| C18:3 <i>n</i> -3                                 | 0.53                 | 0.47  | 0.015 | 0.070   |
| C20:0                                             | 0.09                 | 0.10  | 0.004 | 0.183   |
| C20:3 <i>n</i> -6                                 | 0.15                 | 0.15  | 0.013 | 0.992   |
| C20:4 <i>n</i> -6                                 | 1.40                 | 1.39  | 0.136 | 0.963   |
| C20:5 <i>n</i> -3                                 | 0.08                 | 0.07  | 0.010 | 0.589   |
| C22:4 <i>n</i> -6                                 | 0.13                 | 0.15  | 0.014 | 0.624   |
| C22:5 <i>n</i> -6                                 | 0.04                 | 0.04  | 0.005 | 0.541   |
| C22:5 <i>n</i> -3                                 | 0.22                 | 0.21  | 0.024 | 0.878   |
| C22:6 <i>n</i> -3                                 | 0.05                 | 0.06  | 0.006 | 0.577   |
| <i>Summary</i>                                    |                      |       |       |         |
| Σ SFA <sup>2</sup>                                | 41.80                | 42.59 | 0.255 | 0.128   |
| Σ MUFA <sup>3</sup>                               | 47.21                | 46.35 | 0.444 | 0.357   |
| Σ PUFA <sup>4</sup>                               | 10.71                | 10.67 | 0.524 | 0.975   |
| Σ OBCFA <sup>5</sup>                              | 2.80                 | 2.80  | 0.060 | 0.978   |
| <i>trans</i> -10/ <i>trans</i> -11 18:1           | 2.27                 | 1.49  | 0.226 | 0.084   |
| Total <i>trans</i> 18:1                           | 3.041                | 1.821 | 0.201 | <0.001  |
| Atherogenic index <sup>6</sup>                    | 0.62                 | 0.61  | 0.014 | 0.716   |
| Thrombogenic index <sup>7</sup>                   | 1.24                 | 1.29  | 0.019 | 0.196   |

*Fat soluble vitamins, ng/g muscle*

|                      |        |        |        |       |
|----------------------|--------|--------|--------|-------|
| Retinol              | 229.05 | 311.21 | 21.131 | 0.048 |
| $\alpha$ -Tocopherol | 276.94 | 365.51 | 34.200 | 0.207 |
| $\gamma$ -Tocopherol | 1.99   | 1.48   | 0.191  | 0.191 |
| $\delta$ -Tocopherol | 19.90  | 26.54  | 1.970  | 0.093 |

<sup>1</sup>Diets were: CON (control diet), CMD (cardoon meal diet)

<sup>2</sup> $\Sigma$  SFA = C12:0 + C14:0 + C15:0 + C16:0 + C17:0 + C18:0 + C20:0

<sup>3</sup> $\Sigma$  MUFA = *c*-9 C14:1 + *c*-9 C16:1 + *c*-9 C18:1 + *c*-11 C18:1 + *t*-9 C18:1 + *t*-10 C18:1 + *t*-11 C18:1

<sup>4</sup> $\Sigma$  PUFA = C18:2 *n*-6 + *c*-9, *t*-11 C18:2 + *c*-11, *t*-13 C18:2 + C18:3 *n*-3 + C20:3 *n*-6 + C20:4 *n*-6 + C20:5 *n*-3 + C22:4 *n*-6 + C22:5 *n*-6 + C22:5 *n*-3 + C22:6 *n*-3

<sup>5</sup> $\Sigma$  OBCFA = *iso* C15:0 + *anteiso* C15:0 + *iso* C17:0 + *anteiso* C17:0

<sup>6</sup>Atherogenic index = (C12:0 + [4  $\times$  C14:0] + C16:0)/(*n*-3 PUFA + *n*-6 PUFA + MUFA).

<sup>7</sup>Thrombogenic index = (C14:0 + C16:0 + C18:0)/([0.5  $\times$  MUFA] + [0.5  $\times$  *n*-6 PUFA] + [3  $\times$  *n*-3 PUFA] + [*n*-3/*n*-6 PUFA]).

IMF: intramuscular fat; SFA: saturated fatty acids; MUFA: monounsaturated fatty acids; PUFA: polyunsaturated fatty acids; OBCFA: Odd-and branched-chain fatty acids

\**c*-9, *t*-11 C18:2 co-eluted with *t*-8,*c*-10 C18:2 and *t*-7,*c*-9 C18:2.

**Table 4.** Effect of the dietary treatment and time of storage or incubation on the oxidative stability parameters of meat

|                                             | Diet (D) <sup>1</sup> |       | Storage/incubation time (T) <sup>2</sup> |                    |                    | SEM   | P values <sup>3</sup> |         |       |
|---------------------------------------------|-----------------------|-------|------------------------------------------|--------------------|--------------------|-------|-----------------------|---------|-------|
|                                             | CON                   | CMD   | 1                                        | 2                  | 3                  |       | D                     | T       | D × T |
| <i>Raw meat sclices</i>                     |                       |       |                                          |                    |                    |       |                       |         |       |
| TBARS, µg/g                                 | 0.92                  | 1.13  | 0.18 <sup>c</sup>                        | 0.83 <sup>b</sup>  | 2.03 <sup>a</sup>  | 0.136 | 0.129                 | < 0.001 | 0.036 |
| Lightness, <i>L</i> *                       | 43.70                 | 42.96 | 43.87                                    | 43.41              | 42.79              | 0.299 | 0.229                 | 0.364   | 0.872 |
| Redness, <i>a</i> *                         | 12.45                 | 13.19 | 15.09 <sup>a</sup>                       | 12.67 <sup>b</sup> | 10.62 <sup>c</sup> | 0.340 | 0.075                 | < 0.001 | 0.635 |
| Yellowness, <i>b</i> *                      | 11.82                 | 12.20 | 12.42                                    | 12.14              | 11.43              | 0.194 | 0.133                 | 0.098   | 0.626 |
| Saturation, <i>C</i> *                      | 17.21                 | 17.99 | 19.55 <sup>a</sup>                       | 17.56 <sup>b</sup> | 15.62 <sup>c</sup> | 0.356 | 0.146                 | < 0.001 | 0.630 |
| Hue Angle, <i>H</i> *                       | 43.78                 | 43.06 | 39.33 <sup>c</sup>                       | 43.81 <sup>b</sup> | 47.19 <sup>a</sup> | 0.550 | 0.183                 | < 0.001 | 0.549 |
| (K/S) <sub>572</sub> ÷ (K/S) <sub>525</sub> | 0.93                  | 0.92  | 0.99 <sup>a</sup>                        | 0.91 <sup>b</sup>  | 0.87 <sup>c</sup>  | 0.009 | 0.392                 | < 0.001 | 0.880 |
| <i>Muscle homogenates</i>                   |                       |       |                                          |                    |                    |       |                       |         |       |
| TBARS, µg/g                                 | 2.43                  | 2.73  | 0.15 <sup>b</sup>                        | 3.63 <sup>a</sup>  | 3.96 <sup>a</sup>  | 0.271 | 0.102                 | < 0.001 | 0.437 |
| Metmyoglobin, %                             | 55.41                 | 55.56 | 12.11 <sup>c</sup>                       | 65.10 <sup>b</sup> | 89.21 <sup>a</sup> | 4.911 | 0.925                 | < 0.001 | 0.747 |
| <i>Cooked meat slices</i>                   |                       |       |                                          |                    |                    |       |                       |         |       |
| TBARS, µg/g                                 | 3.58                  | 3.78  | 1.42 <sup>c</sup>                        | 3.85 <sup>b</sup>  | 5.76 <sup>a</sup>  | 0.281 | 0.337                 | < 0.001 | 0.930 |

<sup>1</sup>Diets were: CON (control diet), CMD (cardoon meal diet)<sup>2</sup>Times 1, 2, 3 correspond to: 0, 4, 7 days (raw meat stored at 4°C under aerobic conditions), 0, 30 and 60 minutes (muscle homogenates incubated with Fe/Asc at 37°C under continuous stirring) and 0, 2, 4 days (cooked meat stored at 4°C under aerobic conditions)<sup>3</sup>*P* values for the effects of the dietary treatment (D), time of storage or incubation (T) and of the D × T interaction.<sup>a, b, c</sup> Within row, different superscript letter indicate differences (*P* < 0.05) between times of storage

## Figure captions:

**Figure 1.** Antioxidant activity of **(a)** dietary test ingredients (feedstuff): dehydrated alfalfa (ALF) and cardoon meal (CM) **(b)** experimental diets: control (CON) and cardoon meal diet (CMD). Values are presented as means with standard error bars. <sup>a,b</sup>For antioxidant activity assay, mean values with different letters are significantly different ( $P < 0.05$ ). DPPH: 2,2-diphenyl-1-picrylhydrazyl assay (free radical scavenging activity); FRAP: Ferric reducing antioxidant power assay (total antioxidant activity); TE: trolox equivalent.


**Figure 2.** Interactive effect of the diet and time of storage ( $D \times T$ ) on the lipid oxidation (TBARS,  $\mu\text{g MDA/g}$  of muscle ) measured in fresh *longissimus thoracis et lumborum* (LTL) muscle slices stored aerobically at 4°C for 7 days. Diets were: CON (control diet), CMD (cardoon meal diet). Values are presented as means with standard error bars. <sup>a, b, c, d</sup> Mean values with different letters are significantly different ( $P < 0.05$ )

**Figure 3.** Effect of cardoon extract (CE) addition on **(a)** total phenolic contents (TPC, GAE/g of muscle) **(b)** lipid oxidation (TBARS,  $\mu\text{g MDA/g}$  of muscle) of ovine *longissimus thoracis et lumborum* (LTL) muscle homogenates after 4 h of incubation with a  $\text{FeCl}_3$ /sodium ascorbate pro-oxidant system.

Treatments are as follows: CON, CE0.5, CE1.0 and CE5.0 represent addition of 0, 0.5%, 1.0%, and 5.0% of cardoon extract to muscle homogenates, respectively. Values are presented as means with standard error bars. <sup>a,b</sup>For each treatment, bars with different letters are significantly different ( $P < 0.05$ ).

GAE: gallic acid equivalents; TBARS: thiobarbituric acid reactive substances.

**a**



**b**

