

Updating of the Irish Food Composition Database for vitamin K1 and vitamin K2

C. Kingston¹, L. Kehoe¹, B.A. McNulty³, A.P. Nugent^{3,4}, K.D. Cashman¹, A. Flynn¹ and J. Walton^{1,2}

¹School of Food and Nutritional Sciences, University College Cork, Republic of Ireland,

²Dept. Biological Sciences, Cork Institute of Technology, Republic of Ireland,

³UCD Institute of Food and Health, University College Dublin, Belfield, Dublin 4, Republic of Ireland and

⁴School of Biological Sciences, Institute for Global Food Security, Queens University Belfast, N. Ireland

There are limited data available on vitamin K intakes in national dietary surveys across Europe partly due to the lack of food composition data for vitamin K. The Irish Food Composition Database (IFCD) (2012) has been developed over the last 20 years informed by foods consumed on the Irish national dietary surveys (www.iuna.net). The IFCD (2012) has been updated for each survey to include brand level data and recipes of composite dishes. Each food and beverage consumed throughout the surveys was assigned a food code (*n* 3443), based on the food descriptor and the nutritional composition of the food. The aim of this study was to update the IFCD (2012) with composition values for vitamin K₁ and vitamin K₂ (menaquinone-4 (MK-4) and menaquinone-5 to 10 (MK-5–10)).

The vitamin K₁ and vitamin K₂ content of foods was determined using published analytical values from The UK Composition of Foods Integrated Dataset⁽¹⁾ (COFID), the United States Department of Agriculture (USDA) National Nutrient Database⁽²⁾, published papers, recipe calculation and manufacturers information. The table below outlines the proportion of vitamin K composition values assigned using each data source.

Source of composition data	Vitamin K1	MK-4%	MK-5-10
Published analytical values	52.5	13.6	14.6
UKCOFID nutrient databank ¹	30.7	—	—
USDA nutrient databank ²	16.7	—	—
Published Papers	5.1	13.6	14.6
Recipes	30.4	27.3	27.1
Recipe calculated from ingredients list	25.2	23.2	23.1
Similar to calculated recipes	3.7	4.0	4.0
Retail Product calculated from ingredients list	1.5	—	—
Manufacturers information	10.5	—	—
Nutritional supplements	9.9	—	—
Infant formula	0.6	—	—
Foods that did not contain any vitamin K (as signed value of 0µg)	4.3	57.2	56.5
Other	2.3	1.8	1.7

Published analytical values for vitamin K₁ provided composition data for 53% of food codes in the IFCD (2012). Each recipe in the IFCD (2012) was individually calculated for vitamin K content providing composition values for 29% of food codes for vitamin K₁. The vitamin K₁ composition values for 2% of food codes were calculated from the ingredients listed on packaging. Foods that did not contain any vitamin K₁ (4%) were assigned a composition value of 0µg/100 g. Manufacturers' information provided vitamin K₁ composition data for 11% of food codes. For MK-4 and MK-5–10, published analytical values provided composition data for 14 and 15% of food codes, respectively. Recipe calculation was used to estimate vitamin K₂ composition values for 27% of food codes. Fifty-seven percent of food codes did not contain vitamin K₂ and were assigned a composition value of 0µg/100 g.

These updates to the IFCD (2012) will allow for estimation of vitamin K₁ and vitamin K₂ intakes and sources in the Irish population and will add to the limited vitamin K composition data currently available in Ireland.

1. Public Health England (2015) PHE. London.

2. USDA National Nutrient Database for Standard References 28 (2015) <https://ndb.nal.usda.gov/ndb/>