

Title	Association between air pollution exposure and risk of hemorrhagic stroke: A systematic review and meta-analysis
Authors	Irfan, Syed Asad;Strobl, Kleiton;Wariach, Khubaib;Zahiri, Mehdi;Kurmi, Om
Publication date	2026-05-25
Original Citation	Irfan, S.A. et al. (2026) "Association between air pollution exposure and risk of hemorrhagic stroke: A systematic review and meta-analysis," Canadian Journal of Cardiology, 42(9), (12pp), https://doi.org/10.1016/j.cjca.2026.05.010 .
Type of publication	Article (peer-reviewed);journal-article
Link to publisher's version	10.1016/j.cjca.2026.05.010
Rights	© 2026, authors. Published by Elsevier Inc. on behalf of the Canadian Cardiovascular Society. This is an open access article under the CC BY license - http://creativecommons.org/licenses/by/4.0/
Download date	2026-10-25 17:42:37
Item downloaded from	https://hdl.handle.net/10468/19500

Systematic Review/Meta-analysis

Association Between Air Pollution Exposure and Risk of Hemorrhagic Stroke: A Systematic Review and Meta-analysis

Syed Asad Irfan, MSc,^a Kleiton Strobl, MSc,^b Khubaib Wariach, BSc,^c Mehdi Zahiri, BSc,^d and Om Kurmi, PhD^{e,f}

^aSchulich School of Medicine and Dentistry, Western University, London, Ontario, Canada

^bSchool of Medicine, University College Cork, Cork, Ireland

^cFaculty of Science, McMaster University, Hamilton, Ontario, Canada

^dDepartment of Health Research Methods, Evidence, and Impact, Faculty of Health Sciences, McMaster University, Hamilton, Ontario, Canada

^eDivision of Respiriology, Department of Medicine, McMaster University, Hamilton, Ontario, Canada

^fLincoln Institute for Rural and Coastal Health, University of Lincoln, Lincoln, United Kingdom

See editorial by Kurasz et al., pages 1943-1944 of this issue.

ABSTRACT

Background: Air pollution has been associated with multisystem health adversities. However, the association between short-term air pollution and hemorrhagic stroke has not been sufficiently evaluated in recent years. Considering that hemorrhagic stroke has a high incidence rate (28.8% of incident strokes) and accounts for almost 7.2 million deaths per year globally, it is important to identify risk factors and use it to guide preventative measures.

Methods: A systematic search was conducted on 4 databases (Embase, OVID Medline, Web of Science, PubMed). Studies selected for review included time series, case-cross-over, and cohort studies, published in English, from all time periods, and investigated the risk of hemorrhagic stroke after exposure to air pollution (particulate matter $\leq 2.5 \mu\text{m}$ in diameter [$\text{PM}_{2.5}$], particulate matter $\leq 10 \mu\text{m}$ in diameter [PM_{10}], nitrogen dioxide [NO_2], sulfur dioxide [SO_2], ozone [O_3], or carbon monoxide [CO]). Data for different risk outcomes (risk ratio, odds ratio) were meta-analyzed using an inverse variance statistical analysis and random effects model.

Results: Among the 2435 studies identified, 31 were included and deemed eligible for review and meta-analysis. The results illustrated a

RÉSUMÉ

Contexte : La pollution atmosphérique est associée à des effets néfastes sur la santé en général. Le lien entre la pollution atmosphérique à court terme et l'accident vasculaire cérébral (AVC) hémorragique n'a toutefois pas été suffisamment étudié ces dernières années. Étant donné la fréquence élevée de l'AVC hémorragique (il représente 28,8 % des nouveaux cas d'AVC) et sachant qu'il est la cause de près de 7,2 millions de décès par an dans le monde, il est important d'en connaître les facteurs de risque et de les utiliser pour orienter les mesures de prévention.

Méthodologie : Nous avons réalisé une recherche systématique dans quatre bases de données (Embase, OVID Medline, Web of Science, PubMed). Les études retenues comprenaient des séries temporelles, des études de cas croisées et des études de cohorte, en anglais et sans égard à la période de temps, portant sur le risque d'AVC hémorragique après exposition à des polluants atmosphériques ($\text{PM}_{2.5}$, PM_{10} , NO_2 , SO_2 , O_3 ou CO). Les données relatives au risque (rapport de risques [RR], rapport de cotes [RC]) ont été soumises à une méta-analyse reposant sur une analyse statistique par variance inverse et un modèle à effets aléatoires.

Keywords: hemorrhagic stroke; air pollution; intracerebral hemorrhage; subarachnoid hemorrhage; stroke risk; cardiovascular health; cerebrovascular health

Corresponding author: Dr Syed Asad Irfan

E-mail: asadirfan14@yahoo.com

LinkedIn: <https://www.linkedin.com/in/syed-asad-irfan/>, <https://www.linkedin.com/in/kleiton-strobl/>

See page 2207 for disclosure information.

Received for publication January 18, 2026. Accepted May 15, 2026.

Air pollution is a mixture of hazardous particles and gaseous contaminants released into the atmosphere from natural and anthropogenic sources.¹ Common sources include power generation facilities, forest fires, and traffic-related and industrial emissions.² These air pollution sources, along with many others, are associated with approximately 4.2 million premature deaths annually.³

Common air pollutants include but are not limited to: particulate matter $\leq 10 \mu\text{m}$ in diameter (PM_{10}), fine particulate matter $\leq 2.5 \mu\text{m}$ in diameter ($\text{PM}_{2.5}$), nitrogen

statistically significant increase in the risk of hemorrhagic stroke per 10 $\mu\text{g}/\text{m}^3$ increase in NO_2 exposure (risk ratio, 1.0142; 95% confidence interval, 1.0000–1.0286; $P = 0.05$), and CO exposure (odds ratio, 1.0008; 95% confidence interval, 1.0000–1.0016; $P = 0.05$).

Conclusions: Among the 6 air pollutants studied, NO_2 and CO exposure was associated with a statistically significant increase in the risk of hemorrhagic stroke. This can help guide effective preventative protocols and public health measures in an effort to reduce the burden of cardiovascular and cerebrovascular disease.

dioxide (NO_2), sulfur dioxide (SO_2), ozone (O_3), and carbon monoxide (CO).⁴ Particulate matter consists of small solid and liquid particles suspended in the air, containing organic and nonorganic compounds.⁵ Common sources include dust from roads, mining and construction sites, and byproducts of fossil fuel combustion.⁶ Gaseous air pollutants include NO_2 , SO_2 , O_3 , and CO, and are commonly emitted from vehicle exhausts, power plants, industrial processes, and wildfires.⁷

The primary mode of entry of air pollution is inhalation, which leads to inflammation in the respiratory tract.⁸ Air pollutants might also translocate across the alveolar-capillary membrane into the vasculature, contributing to systemic inflammatory effects.⁹ Previous literature has shown that air pollutants are associated with elevated levels of proinflammatory proteins, such as C-reactive protein and interleukin-6, as well as decreased levels of regulatory, anti-inflammatory proteins, such as adiponectin.¹⁰ This suggests that ambient air pollutants can impair physiological defense mechanisms and contribute to multisystem health adversities, including respiratory and cardiovascular disease.¹¹

Hemorrhagic stroke—internal bleeding in the brain due to the rupture of a blood vessel—is of particular concern because it has a high incidence rate (28.8% of incident strokes) and accounts for almost 7.2 million deaths annually.^{12,13} There are 2 types of hemorrhagic stroke: intracerebral hemorrhage—a hematoma in the parenchyma which can extend into the ventricles—and subarachnoid hemorrhage—an accumulation of blood in the subarachnoid space.^{14,15} Intracerebral hemorrhage is associated with high morbidity and mortality due to its rapidly progressive nature, inducing neurologic deterioration, and potential sudden loss of consciousness.¹⁴ Considering hemorrhage stroke's high burden of disease, it is important to identify risk factors and use them to guide preventative measures.¹³

The main risk factors for hemorrhagic stroke include: alcohol, smoking, hypertension, substance abuse, and pregnancy.¹² Although air pollution as a risk factor for other cardiovascular diseases—such as myocardial infarction and ischemic stroke—has been extensively studied, the risk of hemorrhagic stroke associated with air pollution has not been sufficiently evaluated.^{16–19} As such, the primary objective of this study was to conduct a comprehensive, up-to-date meta-analysis to assess the effect of air pollution exposure on the risk of hemorrhagic stroke.

Résultats : Parmi les 2 435 études trouvées, 31 ont été incluses et jugées admissibles à l'examen et à la méta-analyse. Les résultats ont mis en évidence une augmentation statistiquement significative du risque d'AVC hémorragique par 10 $\mu\text{g}/\text{m}^3$ d'augmentation de l'exposition au NO_2 (RR = 1,0142; intervalle de confiance [IC] à 95 % : 1,0000 à 1,0286; $p = 0,05$) et au CO (RC = 1,0008; IC à 95 % : 1,0000 à 1,0016; $p = 0,05$).

Conclusions : Parmi les six polluants atmosphériques étudiés, l'exposition au NO_2 et au CO a été associée à une augmentation statistiquement significative du risque d'AVC hémorragique. Ces connaissances peuvent contribuer à orienter la mise en place de protocoles de prévention et de mesures de santé publique efficaces visant à réduire le fardeau des maladies cardiovasculaires et vasculaires cérébrales.

Methods

Search strategy

A systematic literature search was conducted on March 10, 2025 of the Embase, OVID Medline, Web of Science, and PubMed databases using detailed search terms, “hemorrhagic stroke”, “intracerebral hemorrhage”, “subarachnoid hemorrhage”, “air pollution”, “particulate matter”, “carbon monoxide”, “sulfur dioxide”, “nitrogen dioxide”, and “ozone”. The detailed search strategy is outlined in [Supplemental Table S1](#). No limitations on publication year were implemented. Our study protocol was registered on PROSPERO (CRD420251031712) on April 13, 2025, after completion of the literature search; this delayed registration is reported herein for transparency.

Primary outcome measure

The primary outcome measure was hemorrhagic stroke risk, measured according to risk ratio (RR) or odds ratio (OR). Risk was standardized, when possible, to a pollutant concentration increment of 10 $\mu\text{g}/\text{m}^3$ for $\text{PM}_{2.5}$, PM_{10} , NO_2 , SO_2 , O_3 , and CO. Risk was also evaluated on the basis of a 10 parts per billion (ppb) increment in pollutant concentrations for NO_2 , SO_2 , O_3 , and CO. For the primary analysis, risk outcomes from the shortest reported lag were used to grasp the short-term effects of air pollution. Hemorrhagic stroke risk was stratified according to risk scale (RR, OR), air pollutant exposure, and concentration increment (10 $\mu\text{g}/\text{m}^3$, 10 ppb).²⁰

Study selection

The inclusion criteria included: (1) time series, case-crossover, or cohort study designs; (2) studies published in English; (3) exposure to at least 1 of the following air pollutants: $\text{PM}_{2.5}$, PM_{10} , NO_2 , SO_2 , O_3 , or CO; and (4) risk of hemorrhagic stroke as an outcome.

The exclusion criteria included: (1) participants with a previous history of stroke; and (2) studies in which hemorrhagic stroke mortality, rather than incident events, was the outcome. Mortality was excluded because incidence/admission and mortality reflect related but distinct outcomes with different determinants, although this decision might reduce the capture of fatal events that occurred before hospital presentation.

Screening was conducted using the online program, Covidence.²¹ Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were followed during this process.²² Two authors (K.W. and M.Z.) independently screened candidate studies for inclusion and exclusion criteria. Discrepancies were evaluated by 2 different authors (K.S. and S.A.I.). Full-text screening was performed independently by 4 authors (K.S., K.W., M.Z., and S.A.I.), and any conflicts were resolved by discussion with the same 2 authors (K.S. and S.A.I.).

Data extraction

Two authors (K.S. and S.A.I.) independently extracted the following data from eligible studies: study name, design, country, air pollutant(s) analyzed, risk value and 95% confidence intervals, exposure increment of the risk value, lag (time delay between exposure and outcome), sample size, male:female ratio, and risk scale. The most short-term, available lag risk data (eg, lag0 values) were extracted from each study in an effort to minimize the time other confounding variables could have a potential influence on the outcome. The same 2 authors (K.S. and S.A.I.) checked for discrepancies, but none existed.

Risk of bias assessment

Two authors (K.S. and S.A.I.) independently assessed the included studies using the National Heart, Lung, and Blood Institute (NHLBI) Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies.²³ Studies were classified as good if all applicable criteria were met, fair if 1 criterion was rated insufficient, and poor if 2 or more criteria were rated insufficient. However, these overall ratings were interpreted cautiously because the tool might be less sensitive to heterogeneity in exposure assessment methods and covariate adjustment strategies across short-term air pollution studies. Any conflicts were resolved with consensus by the same 2 authors (K.S. and S.A.I.).

Data synthesis

Meta-analysis was conducted using Review Manager version 5.4.1 (The Cochrane Collaboration, London, England). The pooled risk outcome was evaluated only when 2 or more studies used the same risk scale, air pollutant exposure, and pollutant concentration increment. If risk values were reported with interquartile range (IQR) exposure increments, they were converted to 10 µg/m³ using the following formula: $\text{Risk value}_{10 \mu\text{g}/\text{m}^3} = \exp((\ln(\text{Risk value}_{\text{IQR } \mu\text{g}/\text{m}^3 \text{ increment}})) \times 10 \mu\text{g}/\text{m}^3 \text{ increment} / \text{IQR } \mu\text{g}/\text{m}^3 \text{ increment})$. Substantial heterogeneity was considered to be $I^2 > 75\%$. An inverse variance statistical analysis with a random-effects model was used for forest plots. Effect sizes were reported to 4 decimal places to preserve computational precision and ensure reproducibility of results derived from the statistical model. Publication bias was assessed using funnel plot symmetry for all outcomes, except those with fewer than 5 studies, because of poor statistical power.

Quality of evidence assessment

Two authors (K.S. and S.A.I.) independently conducted a quality of evidence assessment using Cochrane's Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approach. Because we synthesized observational studies in this review, certainty was considered to begin at low because of the inherent risk of bias and confounding. Upgrading was considered only when prespecified GRADE criteria were met for large magnitude of effect, dose-response gradient, and residual confounding, whereas downgrading was considered for risk of bias, inconsistency, indirectness, imprecision, and publication bias. Any conflicts were resolved through discussion among the same 2 authors (K.S. and S.A.I.).

Results

Study selection

A summary of study identification and screening is available in the PRISMA chart (Fig. 1). A literature search was conducted in the databases Web of Science, Embase, PubMed, and MEDLINE, which yielded 2435 studies. Among these, 1709 duplicates were removed. In total, 726 studies were screened in the abstract screening phase, and 105 were assessed in the full-text screening. Seventy-four studies were removed for the following reasons: wrong outcomes ($n = 40$), wrong study design ($n = 23$), full-text not available in English ($n = 6$), and lack of clarity regarding previous strokes ($n = 5$). Thirty-one studies were deemed eligible for meta-analysis.

Study characteristics

Table 1 shows the characteristics of the studies included in the meta-analysis.

Study design characteristics. Among the 31 studies included in the systematic review and meta-analysis, 22 (71%) used a case-crossover design, 7 (22.6%) a time-series design, and 2 (6.4%) an observational cohort design. Among these, 15 studies (48.4%) used RR as their effect size for hemorrhagic stroke risk outcome measure, and the other 16 studies (51.6%) used OR.

Participant characteristics. To synthesize qualitative findings, data were used from 585,692 participants. Six studies did not report the male:female participant ratio.²⁴⁻²⁹ Among the 554,542 participants in the other studies, 332,144 (59.9%) were male, and 222,398 (40.1%) were female. With regard to the geographic distribution of data, 22 studies (71%) gathered data from participants in China, 3 studies (9.7%) from South Korea, 3 studies (9.7%) from the United States, 2 studies (6.4%) from Taiwan, 1 study (3.2%) from Poland, 1 study (3.2%) from Japan, and 1 study (3.2%) from Canada.

Methodological risk of bias analysis and quality of evidence assessment

Using the prespecified NHLBI approach, all included studies were classified as "good" quality because they met the applicable criteria (Supplemental Table S2). However, this

should be interpreted cautiously rather than as an absence of bias, because important differences remained across studies in exposure assessment, lag specification, and adjustment for key confounders such as temperature, seasonality, and co-pollutants, which are not comprehensively addressed by the NHLBI tool.

The quality of evidence assessment was conducted using the GRADE approach and is shown in [Supplemental Tables S3-S18](#). Among the 16 meta-analyses conducted, 10 were deemed to be of low quality and 6 were of very low quality. The top reasons for downgrading included inconsistency ($n = 3$) and publication bias ($n = 4$).

Hemorrhagic stroke risk outcomes

Table 2 shows a summary of the risk outcomes for air pollution and hemorrhagic stroke. This review showed that there is an increased risk of hemorrhagic stroke per $10 \mu\text{g}/\text{m}^3$ increase in NO_2 exposure (RR, 1.0142; 95% confidence interval [CI], 1.0000-1.0286; $P = 0.05$) and CO exposure (OR, 1.0008; 95% CI, 1.0000-1.0016; $P = 0.05$). The other air pollutants had a statistically insignificant ($P > 0.05$) increase in risk as follows: $\text{PM}_{2.5}$ (OR, 1.0172; RR, 1.0002), PM_{10} (OR, 1.0076; RR, 1.0008), SO_2 (RR, 1.0060), O_3 (OR, 1.0024; RR, 1.0017), CO (OR, 1.0008; RR, 1.0001). With the exception of SO_2 , the risk outcomes for 10 ppb exposure increments also had a statistically insignificant increase in risk: NO_2 (OR, 1.0634), CO (OR, 1.0017), SO_2 (OR, 0.9928), and O_3 (OR, 1.0131).

$\text{PM}_{2.5}$ - Risk Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Ten studies ($n = 413,578$) reported RR for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Guo et al.,³⁰ Liu et al.,³¹ Liu et al.,³² Lv et al.,³³ Wu et al.,³⁴ and Zhao et al.,³⁵ who reported an increased risk of hemorrhagic stroke, and Ban et al.,³⁶ Jin et al.,³⁷ and Weng et al.,³⁸ who reported a reduced risk. Liu et al.³⁹ reported a null effect. There was an overall increased risk of hemorrhagic stroke (RR, 1.00; 95% CI, 0.9985-1.0018; $I^2 = 20\%$), but it was not statistically significant ($P = 0.85$; [Supplemental Fig. S1](#)). The quality of evidence was rated low on the GRADE scale ([Supplemental Table S3](#)).

PM_{10} - Risk Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Ten studies ($n = 336,687$) reported RR for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Czernych et al.,⁴⁰ Han et al.,⁴¹ Han et al.,⁴² Liu et al.,⁴³ Liu et al.,³¹ Lv et al.,³³ Weng et al.,³⁸ Wu et al.,³⁴ and Zhao et al.,³⁵ who reported an increased risk of hemorrhagic stroke, and Liu et al.,⁴³ who reported a reduced risk. There was an overall increased risk of hemorrhagic stroke (RR, 1.0008; 95% CI, 0.9990-1.0025; $I^2 = 49\%$), but it was not statistically significant ($P = 0.39$; [Supplemental Fig. S2](#)). The quality of evidence was rated very low on the GRADE scale because of publication bias ([Supplemental Table S4](#)).

NO_2 - Risk Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Seven studies ($n = 122,602$) reported RR for a $10 \mu\text{g}/\text{m}^3$ exposure interval: Czernych et al.,⁴⁰ Guo et al.,³⁰ Han et al.,⁴¹ Liu et al.,⁴³ Liu et al.,³² and Lv et al.,³³ who reported an increased risk of hemorrhagic stroke, and Weng et al.³⁸ who reported a decreased risk. There was an overall increased risk of hemorrhagic stroke (RR, 1.0142; 95% CI, 1.0000-1.0286; $I^2 =$

71%), and it was statistically significant ($P = 0.05$; [Fig. 2](#)). The quality of evidence was rated low on the GRADE scale ([Supplemental Table S5](#)).

CO - Risk Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Four studies ($n = 107,500$) reported RR for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Czernych et al.,³⁹ Liu et al.,³² and Lv et al.,³³ who report an increased risk of hemorrhagic stroke, and Liu et al.,⁴³ who reported a reduced risk. There was an overall increased risk of hemorrhagic stroke (RR, 1.0001; 95% CI, 0.9997-1.0005; $I^2 = 62\%$), but it was not statistically significant ($P = 0.53$; [Supplemental Fig. S3](#)). The quality of evidence was rated low on the GRADE scale ([Supplemental Table S6](#)).

SO_2 - Risk Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Six studies ($n = 119,555$) reported RR for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Czernych et al.,⁴⁰ Liu et al.,⁴³ Liu et al.,³² Lv et al.,³³ and Shen et al.,²⁷ who reported an increased risk of hemorrhagic stroke, and Weng et al.,³⁸ who reported a reduced risk. There was an overall increased risk of hemorrhagic stroke (RR, 1.0060; 95% CI, 0.9983-1.0138; $I^2 = 43\%$), but it was not statistically significant ($P = 0.13$; [Supplemental Fig. S4](#)). The quality of evidence was rated low on the GRADE scale ([Supplemental Table S7](#)).

O_3 - Risk Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Five studies ($n = 117,261$) reported RR for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Guo et al.,³⁰ Liu et al.,⁴³ and Weng et al.,³⁸ who reported an increased risk of hemorrhagic stroke, and Liu et al.³² and Lv et al.,³³ who reported a reduced risk. There was an overall increased risk of hemorrhagic stroke (RR, 1.0017; 95% CI, 0.9975-1.0060; $I^2 = 31\%$), but it was not statistically significant ($P = 0.43$; [Supplemental Fig. S5](#)). The quality of evidence was rated low on the GRADE scale ([Supplemental Table S8](#)).

$\text{PM}_{2.5}$ - Odds Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Ten studies ($n = 99,780$) reported ORs for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Chiu et al.,²⁴ Huang et al.,⁴⁴ Huang et al.,²⁵ and Sun et al.,⁴⁵ who reported an increased risk of hemorrhagic stroke, and Fisher et al.,⁴⁶ Li et al.,⁴⁷ Villeneuve et al.,⁴⁸ Wang et al.,⁴⁹ Wang et al.,⁵⁰ and Zheng et al.,⁵¹ who reported a reduced risk. Overall, there was a statistically insignificant increase in the risk of hemorrhagic stroke (OR, 1.0172; 95% CI, 0.9918-1.0431; $P = 0.19$; $I^2 = 90\%$; [Supplemental Fig. S6](#)). The quality of evidence was rated very low on the GRADE scale because of inconsistency ([Supplemental Table S9](#)).

PM_{10} - Odds Ratio ($10 \mu\text{g}/\text{m}^3$ exposure interval). Twelve studies ($n = 98,750$) reported ORs for a $10 \mu\text{g}/\text{m}^3$ exposure interval, with Huang et al.,⁴⁴ Sun et al.,⁴⁵ Tsai et al.,²⁸ Villeneuve et al.,⁴⁸ and Zhong et al.,²⁹ who reported an increased risk of hemorrhagic stroke, and Fisher et al.,⁴⁶ Kim et al.,²⁶ Li et al.,⁴⁷ Wang et al.,⁴⁹ Wang et al.,⁵⁰ Xiang et al.,⁵² and Zheng et al.,⁵¹ who reported a reduced risk. Overall, there was a statistically insignificant increase in the risk of hemorrhagic stroke (OR, 1.0076; 95% CI, 0.9957-1.0196; $P = 0.21$; $I^2 = 82\%$;




Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram.

Supplemental Fig. S7). The quality of evidence was rated very low on the GRADE scale because of inconsistency (Supplemental Table S10).

NO₂ - Odds Ratio (10 µg/m³ exposure interval). Five studies (n = 44,297) reported ORs for a 10 µg/m³ exposure interval, with Zhong et al.,²⁹ who reported an increased risk of hemorrhagic stroke, and Li et al.,⁴⁷ Wang et al.,⁴⁹ Wang et al.,⁵⁰ and Xiang et al.,⁵² who reported a reduced risk. Overall, there was a reduced risk of hemorrhagic stroke (OR, 0.9853; 95% CI, 0.9685-1.0024; *I*² = 4%), but this was not statistically significant (*P* = 0.09; Supplemental Fig. S8). The quality of evidence was rated low on the GRADE scale (Supplemental Table S11).

CO - Odds Ratio (10 µg/m³ exposure interval). Three studies (n = 41,651) reported ORs for a 10 µg/m³ exposure interval, with Li et al.,⁴⁷ who reported an increased risk of hemorrhagic stroke, and Wang et al.⁴⁹ and Wang et al.,⁵⁰ who reported a reduced risk. Overall, there was an increased risk of hemorrhagic stroke (OR, 1.0008; 95% CI, 1.0000-1.0016; *I*² = 0%), and it was statistically significant (*P* = 0.05; Fig. 3). The quality of evidence was rated low on the GRADE scale (Supplemental Table S12).

SO₂ - Odds Ratio (10 µg/m³ exposure interval). Five studies (n = 44,297) reported ORs for a 10 µg/m³ exposure interval, with Wang et al.⁴⁹ and Zhong et al.,²⁹ who reported an increased risk of hemorrhagic stroke, and Li et al.,⁴⁷ Wang

Table 1. Summary of study characteristics

Reference	Study design	Air pollutant	Exposure (increments in $\mu\text{g}/\text{m}^3$)	Lag (time delay)	Risk scale (ie, RR or OR)	Country	Sample size, n	Male sex, n (%)
Ban et al. ³⁶	Case-crossover	PM _{2.5}	10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	19,100	10,254 (53.7%)
Chiu et al. ²⁴	Case-crossover	PM _{2.5}	IQR, 17.46 $\mu\text{g}/\text{m}^3$	Lag0-2 (3-day average)	OR	Taiwan	12,520	NR
Czernych et al. ⁴⁰	Time series	NO ₂	IQR, 14.62 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	Poland	4542	2767 (61%)
		CO	IQR, 164.04 $\mu\text{g}/\text{m}^3$					
		SO ₂	IQR, 3.65 $\mu\text{g}/\text{m}^3$					
		PM ₁₀	IQR, 16.76 $\mu\text{g}/\text{m}^3$					
		PM _{2.5}	IQR, 8.91 $\mu\text{g}/\text{m}^3$					
Fisher et al. ⁴⁶	Case-crossover	PM _{2.5}	IQR, 14.46 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	OR	United States	122	122 (100%)
Guo et al. ³⁰	Time series	PM _{2.5}	10 $\mu\text{g}/\text{m}^3$	Lag0-3 (4-day average)	RR	China	11,415	7071 (61.9%)
		NO ₂	10 $\mu\text{g}/\text{m}^3$					
		O ₃	10 $\mu\text{g}/\text{m}^3$					
Han et al. ⁴¹	Cohort	PM ₁₀	10 $\mu\text{g}/\text{m}^3$	Lag0-month end (monthly average)	RR	South Korea	799	495 (62.0%)
Han et al. ⁴²	Time series	NO ₂	10 $\mu\text{g}/\text{m}^3$	Lag0-month end (monthly average)	RR	South Korea	1477	801 (54.2%)
		PM ₁₀	10 $\mu\text{g}/\text{m}^3$					
		NO ₂	10 ppb					
Huang 2016 ⁴⁴	Case-crossover	O ₃	10 ppb	Lag0 (same day)	OR	China	16,880	10,349 (61.3%)
		PM _{2.5}	IQR, 82.0 $\mu\text{g}/\text{m}^3$					
		PM ₁₀	IQR, 93.5 $\mu\text{g}/\text{m}^3$					
Huang et al. ²⁵	Case-crossover	PM _{2.5}	10 $\mu\text{g}/\text{m}^3$	Lag2 (2-day delay)	OR	China	99	NR
Jin et al. ³⁷	Case-crossover	PM _{2.5}	10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	36,850	22,647 (61.5%)
Kim et al. ²⁶	Cohort	SO ₂	0.01 ppm	Lag0-7 (8-day average)	OR	South Korea	4404	NR
		NO ₂	0.1 ppm					
		O ₃	0.01 ppm					
		CO	1 ppm					
		PM ₁₀	10 $\mu\text{g}/\text{m}^3$					
Li et al. ⁴⁷	Case-crossover	PM _{2.5}	IQR, 27 $\mu\text{g}/\text{m}^3$	Lag0-1 (2-day average)	OR	China	39,732	24,926 (62.7%)
		PM ₁₀	IQR, 37 $\mu\text{g}/\text{m}^3$					
		SO ₂	IQR, 10.8 $\mu\text{g}/\text{m}^3$					
		NO ₂	IQR, 11.2 $\mu\text{g}/\text{m}^3$					
		O ₃	IQR, 38.6 $\mu\text{g}/\text{m}^3$					
		CO	IQR, 0.32 mg/m ³					
		PM _{2.5}	IQR, 47.5 $\mu\text{g}/\text{m}^3$					
Liu et al. ³⁹	Case-crossover	PM ₁₀	IQR, 76.9 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	69,399	47,179 (68.0%)
Liu et al. ⁴³	Case-crossover	PM ₁₀	IQR, 85.7 $\mu\text{g}/\text{m}^3$	Lag0-5 (6-day average)	RR	China	41,746	28,223 (67.6%)
		NO ₂	IQR, 25.4 $\mu\text{g}/\text{m}^3$					
		SO ₂	IQR, 38.3 $\mu\text{g}/\text{m}^3$					
		CO	IQR, 0.81 mg/m ³					
		O ₃	IQR, 67 $\mu\text{g}/\text{m}^3$					
Liu et al. ³¹	Time series	PM _{2.5}	10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	9541	5692 (60.0%)
Liu et al. ³²	Case-crossover	PM ₁₀	10 $\mu\text{g}/\text{m}^3$	Lag0-1 (2-day average)	RR	China	58,090	31,378 (54.0%)
		PM _{2.5}	10 $\mu\text{g}/\text{m}^3$					
		NO ₂	10 $\mu\text{g}/\text{m}^3$					
		O ₃	10 $\mu\text{g}/\text{m}^3$					
		CO	10 $\mu\text{g}/\text{m}^3$					
Lv et al. ³³	Case-crossover	SO ₂	10 $\mu\text{g}/\text{m}^3$	Lag0 (0-2 hour average)	RR	China	3122	1658 (53.1%)
		PM _{2.5}	10 $\mu\text{g}/\text{m}^3$					
		PM ₁₀	10 $\mu\text{g}/\text{m}^3$					
		NO ₂	10 $\mu\text{g}/\text{m}^3$					
		SO ₂	10 $\mu\text{g}/\text{m}^3$					
		O ₃	10 $\mu\text{g}/\text{m}^3$					
		CO	1 mg/m ³					
Montresor-López et al. ⁵⁴	Case-crossover	O ₃	10 ppb	Lag0 (same day)	OR	United States	2582	1153 (44.7%)
Shen et al. ²⁷	Time series	SO ₂	10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	9167	NR

Continued

Table 1. Continued.

Reference	Study design	Air pollutant	Exposure (increments in $\mu\text{g}/\text{m}^3$)	Lag (time delay)	Risk scale (ie, RR or OR)	Country	Sample size, n	Male sex, n (%)
Sun et al. ⁴⁵	Case-crossover	PM _{2.5} PM ₁₀ NO ₂ SO ₂ O ₃	IQR, 8.2 $\mu\text{g}/\text{m}^3$ IQR, 14.3 $\mu\text{g}/\text{m}^3$ IQR, 10.2 ppb IQR, 2.5 ppb IQR, 20.6 ppb	Lag0 (same day)	OR	United States	924	0 (0.0%)
Tsai et al. ²⁸	Case-crossover	PM ₁₀ SO ₂ NO ₂ CO O ₃	IQR, 66.33 $\mu\text{g}/\text{m}^3$ IQR, 6.17 ppb IQR, 17.08 ppb IQR, 0.29 ppm IQR, 21.20 ppb	Lag0-1 (2-day average)	OR	Taiwan	4539	NR
Turin et al. ⁵³	Case-crossover	SO ₂ NO ₂	IQR, 2.2 ppb IQR, 10.4 ppb	Lag0 (same day)	OR	Japan	432	230 (53.1%)
Villeneuve et al. ⁴⁸	Case-crossover	SO ₂ NO ₂ CO O ₃ PM _{2.5} PM ₁₀	IQR, 3.0 ppb IQR, 13.5 ppb IQR, 0.5 ppm IQR, 18.0 ppm IQR, 6.3 $\mu\text{g}/\text{m}^3$ IQR, 16.0 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	OR	Canada	2329	1126 (48.3%)
Wang et al. ⁴⁹	Case-crossover	PM _{2.5} PM ₁₀ SO ₂ NO ₂ CO O ₃	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 1 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 0.1 mg/m ³ 10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	OR	China	1052	693 (65.9%)
Wang et al. ⁵⁰	Case-crossover	PM _{2.5} PM ₁₀ SO ₂ NO ₂ CO O ₃	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 1 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 0.1 mg/m ³ 10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	OR	China	867	572 (66.0%)
Weng et al. ³⁸	Time series	PM _{2.5} PM ₁₀ NO ₂ SO ₂ O ₃	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	2888	1807 (62.5%)
Wu et al. ³⁴	Case-crossover	PM _{2.5} PM ₁₀	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$	Lag0-4 (5-day average)	RR	China	160,264	89,185 (55.6%)
Xiang et al. ⁵²	Case-crossover	PM ₁₀ NO ₂ SO ₂	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	OR	China	2225	1358 (61%)
Zhao et al. ³⁵	Time series	PM _{2.5} PM ₁₀	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	RR	China	42,909	27,351 (63.74%)
Zheng et al. ⁵¹	Case-crossover	PM _{2.5} PM ₁₀ PM _{2.5} PM ₁₀	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ IQR, 22.6 $\mu\text{g}/\text{m}^3$ IQR, 29.8 $\mu\text{g}/\text{m}^3$	Lag0 (same day)	OR	China	25,255	15,107 (59.8%)
Zhong et al. ²⁹	Case-crossover	SO ₂ NO ₂ PM ₁₀	10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$ 10 $\mu\text{g}/\text{m}^3$	Lag2 (2-day delay) Lag1 (1-day delay) Lag2 (2-day delay)	OR	China	421	NR

CO, carbon monoxide; IQR, interquartile range; NO₂, nitrogen dioxide; NR, not reported; O₃, ozone; OR, odds ratio; PM_{2.5}, particulate matter $\leq 2.5 \mu\text{m}$ in diameter; PM₁₀, particulate matter $\leq 10 \mu\text{m}$ in diameter; ppb, parts per billion; ppm, parts per million; RR, risk ratio; SO₂, sulfur dioxide.

et al.,⁵⁰ and Xiang et al.,⁵² who reported a reduced risk. An overall reduction in hemorrhagic stroke risk was observed (OR, 0.9982; 95% CI, 0.9805-1.0163; $I^2 = 9\%$), but it was not statistically significant ($P = 0.85$; Supplemental Fig. S9). The quality of evidence was rated low on the GRADE scale (Supplemental Table S13).

O₃ - Odds Ratio (10 µg/m³ exposure interval). Three studies (n = 41,651) reported ORs for a 10 µg/m³ exposure interval, with Wang et al.,⁵⁰ who reported an increased risk of hemorrhagic stroke, and Li et al.⁴⁷ and Wang et al.,⁴⁹ who reported a reduced risk. There was an overall increased risk of hemorrhagic stroke (OR, 1.0024; 95% CI, 0.9814-1.0239; $I^2 = 61\%$), but it was not statistically significant ($P = 0.82$; Supplemental Fig. S10). The quality of evidence was rated low on the GRADE scale (Supplemental Table S14).

NO₂ - Odds Ratio (10 ppb exposure interval). Five studies (n = 12,628) reported ORs for a 10-ppb exposure interval, with Sun et al.,⁴⁵ Tsai et al.,²⁸ Turin et al.,⁵³ and Villeneuve et al.,⁴⁸ who reported an increased risk of hemorrhagic stroke, and Kim et al.,²⁶ who reported a reduced risk. Overall, an increased risk of hemorrhagic stroke was apparent (OR, 1.0634; 95% CI, 0.9623-1.1751; $I^2 = 86\%$), although not statistically significant ($P = 0.23$; Supplemental Fig. S11). The quality of evidence was rated very low on the GRADE scale because of inconsistency and publication bias (Supplemental Table S15).

CO - Odds Ratio (10 ppb exposure interval). Three studies (n = 11,272) reported ORs for a 10-ppb exposure interval, with Kim et al.,²⁶ Tsai et al.,²⁸ and Villeneuve et al.,⁴⁸ who reported an increased risk of hemorrhagic stroke. Although an overall increased risk of hemorrhagic stroke was evident (OR, 1.0017; 95% CI, 0.9988-1.0046; $I^2 = 83\%$), it was not statistically significant ($P = 0.26$; Supplemental Fig. S12). The quality of evidence was rated very low on

the GRADE scale because of inconsistency (Supplemental Table S16).

SO₂ - Odds Ratio (10 ppb exposure interval). Five studies (n = 12,628) reported ORs for a 10-ppb exposure interval, with Tsai et al.²⁸ and Turin et al.,⁵³ who reported an increased risk of hemorrhagic stroke, and Kim et al.,²⁶ Sun et al.,⁴⁵ and Villeneuve et al.,⁴⁸ who reported a reduced risk. Overall, there was a reduction in the risk of hemorrhagic stroke (OR, 0.9928; 95% CI, 0.8924-1.1046; $I^2 = 0\%$), but this was not statistically significant ($P = 0.89$; Supplemental Fig. S13). The quality of evidence was rated very low on the GRADE scale because of publication bias (Supplemental Table S17).

O₃ - Odds Ratio (10 ppb exposure interval). Five studies (n = 14,778) reported ORs for a 10-ppb exposure interval, with Kim et al.²⁶ and Tsai et al.,²⁸ who reported an increased risk of hemorrhagic stroke, Villeneuve et al.,⁴⁸ who reported a null effect, and Montessor-Lopez et al.⁵⁴ and Sun et al.,⁴⁵ who report a reduced risk. There was an overall increased risk of hemorrhagic stroke (OR, 1.0131; 95% CI, 0.9776-1.0499; $I^2 = 71\%$), but it was not statistically significant ($P = 0.47$; Supplemental Fig. S14). The quality of evidence was rated low on the GRADE scale (Supplemental Table S18).

Sensitivity analyses. To explore potential sources of heterogeneity, 3 sensitivity analyses were performed on lag structure, region, and study design, respectively.

The first sensitivity analysis was performed to account for potential heterogeneity stemming from lag differences. Nineteen of 31 studies (61.3%) reported risk on the basis of lag0 (same day exposure) and were analyzed separately to assess immediate, short-term outcomes. Sensitivity analysis revealed that NO₂ and hemorrhagic stroke showed a stronger, yet statistically insignificant association (RR, 1.0253; 95% CI, 0.9825-1.0700; $P = 0.25$; Supplemental Fig. S15). The RR outcome of CO and hemorrhagic stroke showed a mildly stronger association (RR, 1.0012; 95% CI, 1.0003-1.0021) and was statistically significant ($P = 0.01$; Supplemental Fig. S16), whereas the OR outcome of CO and hemorrhagic stroke was no longer significant (OR, 0.9995; 95% CI, 0.9971-1.0019; $P = 0.70$; Supplemental Fig. S17). There were no other significant differences in this sensitivity analysis. Because of a lack of data, sensitivity analyses could not be performed for other lag intervals.

The second sensitivity analysis was performed to explore differences among regions. Twenty-six of 31 studies (83.9%) were conducted in Asia (with 20/31 studies [64.5%] from China) and were analyzed separately to assess outcomes from the same region. Sensitivity analysis showed that NO₂ and hemorrhagic stroke showed a weaker and statistically insignificant association (RR, 1.0091; 95% CI, 0.9955-1.0229; $P = 0.19$; Supplemental Fig. S18). CO OR risk outcomes remained unchanged and significant. Additionally, O₃ was associated with an increased risk of hemorrhagic stroke within the included Asian studies, as per a 10-ppb increase in exposure (RR, 1.0579; 95% CI, 1.0122-1.1057), and was statistically significant ($P = 0.01$; Supplemental Fig. S19). Not enough available data supported meaningful sensitivity analyses of other regions.

Table 2. Summary of risk outcomes for air pollution and hemorrhagic stroke

Air pollutant	Exposure interval	Risk value (95% CI)
PM _{2.5}	10 µg/m ³	RR, 1.0002 (0.9985-1.0018) OR, 1.0172 (0.9918-1.0431)
PM ₁₀	10 µg/m ³	RR, 1.0008 (0.9990-1.0025) OR, 1.0076 (0.9957-1.0196)
NO ₂	10 µg/m ³	RR, 1.0142 (1.0000-1.0286)* OR, 0.9853 (0.9685-1.0024)
CO	10 ppb 10 µg/m ³	OR, 1.0634 (0.9623-1.1751) RR, 1.0001 (0.9997-1.0005) OR, 1.0008 (1.0000-1.0016)*
SO ₂	10 ppb 10 µg/m ³	OR, 1.0017 (0.9988-1.0046) RR, 1.0060 (0.9983-1.0138) OR, 0.9982 (0.9805-1.0163)
O ₃	10 ppb 10 µg/m ³	OR, 0.9928 (0.8924-1.1046) RR, 1.0017 (0.9975-1.0060) OR, 1.0024 (0.9814-1.0239)
	10 ppb	OR, 1.0131 (0.9776-1.0499)

CI, confidence interval; CO, carbon monoxide; NO₂, nitrogen dioxide; O₃, ozone; OR, odds ratio; PM_{2.5}, particulate matter ≤ 2.5µm in diameter; PM₁₀, particulate matter ≤ 10µm in diameter; ppb, parts per billion; RR, risk ratio; SO₂, sulfur dioxide.

* $P \leq 0.05$.




Figure 2. Relative risk of hemorrhagic stroke after a 10 µg/m³ increase in nitrogen dioxide exposure.

The third sensitivity analysis was performed to explore differences in study design. Twenty-two of 31 studies (71.0%) had a case-crossover design and were analyzed separately to isolate studies with the same study design. Sensitivity analysis showed that NO₂ and hemorrhagic stroke had a weaker and statistically insignificant association (RR, 1.0024; 95% CI, 0.9958-1.0090; $P = 0.47$; [Supplemental Fig. S20](#)). OR risk outcomes for CO remained unchanged and significant. Not enough available data supported meaningful sensitivity analyses of other study designs.

Discussion

This study showed that short-term exposure to several ambient air pollutants is associated with an increased risk of developing hemorrhagic stroke. Risk measures of RR and OR were the outcomes of interest and they were stratified in this study because of varying literature on their interchangeability.²⁰ Statistically significant associations were found for NO₂ and CO. NO₂ was linked to a 1.42% relative increase in the risk of hemorrhagic stroke when risk was expressed per 10 µg/m³ increment (RR, 1.0142; 95% CI, 1.0000-1.0286; $P = 0.05$). When risk was expressed as an OR per 10 µg/m³ increase in CO, a statistically significant correlation was observed (OR, 1.0008; 95% CI, 1.0000-1.0016; $P = 0.05$). This might seem to be a small increase in risk, but it has large implications because it is derived from population-level data and hence affects millions of people. Additionally, considering that geographical areas can have exposure differences of several increments around the world rather than the standardized increment of 10 µg/m³, it is appreciable that the risk of hemorrhagic stroke can starkly increase accordingly.

In contrast, across RR and OR scales, as well as across 10 µg/m³ and 10 ppb exposure increments, particulate pollutants (PM_{2.5} and PM₁₀), SO₂, and O₃ consistently showed

very small effect sizes and confidence intervals crossed the null. Thus, altogether these results imply that NO₂ and CO might be more pertinent short-term triggers for hemorrhagic stroke compared with others.

Our findings are consistent with and further expand on reported patterns within previous literature on air pollutants and their correlation with cerebrovascular incidents.^{43,47,55} Historically, the short-term effects of air pollution have been more transparent and consistent for total and ischemic stroke than for hemorrhagic stroke. However, the biological plausibility for hemorrhagic stroke might differ because hemorrhagic events are driven primarily by vessel rupture rather than thrombosis.⁵⁶ Accordingly, any effect of air pollutants on hemorrhagic stroke might be mediated more indirectly through acute blood pressure changes, vascular inflammation, endothelial dysfunction, and impaired vascular reactivity in susceptible vessels. In that context, Liu et al.⁴³ reported that NO₂ was the only pollutant, across 14 Chinese cities, that was significantly linked to admissions for hemorrhagic stroke, whereas associations for other pollutants were more erratic and imprecise. This is consistent with our finding of a higher risk per 10 µg/m³ increase in NO₂. Similarly, Li et al.⁴⁷ reported that all 7 pollutants (particulate matter ≤ 1 µm in diameter, PM_{2.5}, PM₁₀, SO₂, NO₂, O₃, and CO) were linked to hospitalizations for ischemic stroke, but only CO was linked to hemorrhagic stroke. This is in line with our pooled estimate, which indicates a statistically significant increase in the risk of hemorrhagic stroke with short-term CO exposure. Moreover, our predominantly null pooled estimates for the risk of hemorrhagic stroke and particulate matter are in line with previous literature. Fisher et al.⁴⁶ reported no evidence of an increased risk of hemorrhagic stroke associated with short-term particulate matter exposure. Likewise, Wu et al.³⁴ and Zhang et al.⁵⁷ reported that particulate pollutants were




Figure 3. Odds ratio of hemorrhagic stroke after a 10 µg/m³ increase in carbon monoxide exposure.

clearly associated with overall or ischemic stroke, but not with hemorrhagic stroke. However, research by Chiu et al.²⁴ showed positive correlations between fine particulate matter and hemorrhagic stroke in lag structures or in single-pollutant models. However, these estimates frequently lost statistical significance when co-pollutants were included. Overall, these studies supplement our review by indicating that short-term exposure to NO₂ and CO has the most consistent adverse effects on hemorrhagic stroke. In contrast, the evidence for particulate matter, SO₂, and O₃ remains inconsistent and context-dependent.

This review has several important limitations. First, the evidence base is primarily composed of observational time-series and case-crossover studies that used regional models or area-level concentrations from fixed-site monitors. These studies do not capture individual time-activity patterns and are likely to introduce nondifferential exposure misclassification, which would bias true effects toward the null. Second, studies differed considerably in terms of covariate adjustments such as temperature, seasonality, and co-pollutants. This limited comparability and prevented the pooling of certain pollutant-metric combinations. Additionally, many studies used data for hospital admissions for risk assessments and did not account for pre-hospital fatal cases. This limits the data in terms of providing a full picture of the adverse effects of short-term pollution exposure. Finally, generalizability to areas with different emission profiles, lower background pollution, and distinct hemorrhagic stroke epidemiology is uncertain, because most studies were conducted in China and other East Asian settings.

To explore these limitations, sensitivity analyses showed that when keeping the lag structure (lag0), region (Asia), or study design (case-crossover) constant, NO₂ RR outcomes no longer appeared significant. This demonstrated that the NO₂ result in the main analysis might have been influenced by these confounders. Although it is important to note that these insignificant results could be due to low power from a low number of studies remaining in the forest plot (for example, only 3 of 7 studies reported lag0 RR NO₂ risk outcomes). More research needs to be done in this area to determine whether this connection is true or not. In contrast, although CO risk outcomes were comparatively smaller than for NO₂, they remained consistent and statistically significant, which suggested that CO was less influenced by these potential confounders. Additionally, O₃ was shown to have an increased risk in hemorrhagic stroke in Asia. However, this result should be interpreted with caution because it is the result of only 2 included Asian studies. More research is warranted on O₃ pollution and hemorrhagic stroke in Asian regions to further test the extent of this association.

Despite the limitations, our findings provide valuable insight. Increases in NO₂ and CO RRs can result in a significant number of hemorrhagic stroke incidents at the population level. This is especially true when the population is already vulnerable because of previous strokes or a high baseline stroke incidence, and when the greater proportion is aging. From a policy standpoint, our findings encourage ongoing efforts to reduce emissions from combustion and traffic through stricter air quality regulations and to further develop and implement cleaner, alternative energy systems. Such interventions are likely to longitudinally help prevent hemorrhagic and ischemic strokes. Future research should prioritize high-quality studies in under-represented areas, improve exposure assessment, and

include subtype-specific analyses that distinguish intracerebral from subarachnoid hemorrhage, alongside comprehensive data on comorbidities and medications.

Conclusions

The results of this meta-analysis showed an association of air pollution exposure with hemorrhagic stroke. More specifically, NO₂ and CO exposures per 10 µg/m³ increment led to increases in hemorrhagic stroke risk; RR, 1.0142 and OR, 1.0008, respectively. Considering the context that the geographical distribution of air pollution can vary by several increments, this finding shows that the risk of hemorrhagic stroke can starkly increase with increased exposure. Because of the large burden of hemorrhagic stroke on cardiovascular health worldwide, it is imperative to consistently examine the effects of air pollution across multiple geographic regions to accurately inform environmental health policy, allocation of economic resources, and preventative measures. These strategies can aid in diminishing the burden of hemorrhagic stroke and other cardiovascular consequences caused by air pollution exposure.

Ethics Statement

This is a systematic review and meta-analysis of articles indexed on Embase, OVID Medline, Web of Science, and PubMed. An institutional review from the Ethics Board was not required.

Patient Consent

The authors confirm that patient consent is not applicable to this article. This systematic review and meta-analysis did not involve direct patient participation or the use of identifiable patient information.

Funding Sources

The authors have no funding sources to declare.

Disclosures

The authors have no conflicts of interest to disclose.

References

1. Chen F, Zhang W, Mfarrej MFB, et al. Breathing in danger: understanding the multifaceted impact of air pollution on health impacts. *Ecotoxicol Environ Saf* 2024;280:116532.
2. Manisalidis I, Stavropoulou E, Stavropoulos A, Bezirtzoglou E. Environmental and health impacts of air pollution: a review. *Front Public Health* 2020;8:14.
3. World Health Organization: Ambient (outdoor) air pollution. Accessed July 08, 2025. Available from: [https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health).
4. World Health Organization: WHO global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Accessed August 08, 2025 Available from: <https://www.who.int/publications/i/item/9789240034228>.
5. Hamanaka RB, Mutlu GM. Particulate matter air pollution: effects on the cardiovascular system. *Front Endocrinol* 2018;9:680.

6. EPA. United States Environmental Protection Agency: Particulate matter (PM) basics. Accessed August 08, 2025. Available from: <https://www.epa.gov/pm-pollution/particulate-matter-pm-basics>.
7. Chen TM, Kushner WG, Gokhale J, Shofer S. Outdoor air pollution: nitrogen dioxide, sulfur dioxide, and carbon monoxide health effects. *Am J Med Sci* 2007;333:249-56.
8. Cui C, Yang R, Chen H, et al. Air toxins disorder the NF- κ B pathway leads to immune disorders and immune diseases in the human health. *Ecotoxicol Environ Saf* 2025;302:118474.
9. Thangavel P, Park D, Lee YC. Recent insights into particulate matter (PM_{2.5})-mediated toxicity in humans: an overview. *Int J Environ Res Public Health* 2022;19:7511.
10. Vogli M, Peters A, Wolf K, et al. Long-term exposure to ambient air pollution and inflammatory response in the KORA study. *Sci Total Environ* 2024;912:169416.
11. Lin Y, Ramanathan G, Zhu Y, et al. Pro-oxidative and proinflammatory effects after traveling from Los Angeles to Beijing. *Circulation* 2019;140:1995-2004.
12. Unnithan AKA, Das JM. Hemorrhagic stroke overview. Accessed October 08, 2025. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK559173/>.
13. Parry-Jones AR, Krishnamurthi R, Ziai WC, et al. World Stroke Organization (WSO): global intracerebral hemorrhage factsheet 2025. *Int J Stroke* 2025;20:145-50.
14. Rajashekar D, Liang JW. Intracerebral hemorrhagic stroke. Accessed October 08, 2025. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK553103/>.
15. Ziu E, Khan Suheb MZ, Mesfin FB. Subarachnoid hemorrhage. Accessed October 08, 2025. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK441958/>.
16. Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet* 2004;364:937-52.
17. Chen L, Luo S, Feng M, et al. Decoding the link between ischemic stroke and myocardial infarction: a pathophysiological and clinical perspective. *Eur J Med Res* 2025;30:984.
18. Seneviratne AN, Miller MR. Air pollution and atherosclerosis. *Atherosclerosis* 2025;406:119240.
19. Toubasi A, Al-Sayegh TN. Short-term exposure to air pollution and ischemic stroke: a systematic review and meta-analysis. *Neurology* 2023;101:e1922-32.
20. George A, Stead TS, Ganti L. What's the risk: differentiating risk ratios, odds ratios, and hazard ratios? *Cureus* 2020;12:e10047.
21. Covidence: The World's #1 Systematic Review Tool. Accessed October 03, 2025. Available from: <https://www.covidence.org/>.
22. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
23. NIH: National Heart, Lung, and Blood Institute. Study quality assessment tools. Accessed January 09, 2025. Available from: <https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools>.
24. Chiu HF, Chang CC, Yang CY. Relationship between hemorrhagic stroke hospitalization and exposure to fine particulate air pollution in Taipei, Taiwan. *J Toxicol Environ Health A* 2014;77:1154-63.
25. Huang Z, Zhou Y, Lu Y, et al. A case-crossover study between fine particulate matter elemental composition and emergency admission with cardiovascular disease. *Acta Cardiol Sin* 2017;33:66-73.
26. Kim SY, Kim JH, Kim YH, et al. Short- and long-term exposure to air pollution increases the risk of stroke. *Int J Stroke* 2022;17:654-60.
27. Shen S, Li X, Yuan C, et al. Association of short-term exposure to sulfur dioxide and hospitalization for ischemic and hemorrhagic stroke in Guangzhou, China. *BMC Public Health* 2020;20:263.
28. Tsai SS, Goggins WB, Chiu HF, Yang CY. Evidence for an association between air pollution and daily stroke admissions in Kaohsiung, Taiwan. *Stroke* 2003;34:2612-6.
29. Zhong H, Shu Z, Zhou Y, et al. Seasonal effect on association between atmospheric pollutants and hospital emergency room visit for stroke. *J Stroke Cerebrovasc Dis* 2018;27:169-76.
30. Guo Y, Xie X, Lei L, et al. Short-term associations between ambient air pollution and stroke hospitalisations: time-series study in Shenzhen, China. *BMJ Open* 2020;10:e032974.
31. Liu X, Li Z, Guo M, et al. Acute effect of particulate matter pollution on hospital admissions for stroke among patients with type 2 diabetes in Beijing, China, from 2014 to 2018. *Ecotoxicol Environ Saf* 2021;217:112201.
32. Liu C, Yu Y, Liu C, et al. Effect of neighbourhood greenness on the association between air pollution and risk of stroke first onset: a case-crossover study in Shandong Province, China. *Int J Hyg Environ Health* 2023;254:114262.
33. Lv X, Shi W, Yuan K, et al. Hourly air pollution exposure and emergency hospital admissions for stroke: a multicenter case-crossover study. *Stroke* 2023;54:3038-45.
34. Wu H, Zhang B, Wei J, et al. Short-term effects of exposure to ambient PM₁, PM_{2.5}, and PM₁₀ on ischemic and hemorrhagic stroke incidence in Shandong Province, China. *Environ Res* 2022;212:113350.
35. Zhao Z, Guo M, Tan P, et al. Effect modification by high-concentration duration in the association between particulate matters and stroke hospital admissions among hypertension patients in Beijing, China. *Air Qual Atmosphere Health* 2023;16:2325-35.
36. Ban J, Wang Q, Ma R, et al. Associations between short-term exposure to PM_{2.5} and stroke incidence and mortality in China: a case-crossover study and estimation of the burden. *Environ Pollut* 2021;268:115743.
37. Jin JQ, Lin GZ, Wu SY, et al. Short-term effects of individual exposure to PM_{2.5} on hospital admissions for myocardial infarction and stroke: a population-based case-crossover study in Guangzhou, China. *Environ Sci Pollut Res* 2023;30:78802-10.
38. Weng L, Li N, Feng T, Zhu R, Zheng ZJ. Short-term association of air pollutant levels and hospital admissions for stroke and effect

- modification by apparent temperature: evidence from Shanghai, China. *Front Public Health* 2021;9:716153.
39. Liu H, Tian Y, Xu Y, et al. Association between ambient air pollution and hospitalization for ischemic and hemorrhagic stroke in China: a multicity case-crossover study. *Environ Pollut* 2017;230:234-41.
40. Czernych R, Kozera G, Badyda AJ, et al. Air pollution increases risk of occurrence of intracerebral haemorrhage but not of subarachnoid haemorrhage: time-series cross-sectional study. *Biomedicines* 2024;12:1562.
41. Han MH, Yi HJ, Kim YS, Kim YS. Effect of seasonal and monthly variation in weather and air pollution factors on stroke incidence in Seoul, Korea. *Stroke* 2015;46:927-35.
42. Han MH, Yi HJ, Ko Y, Kim YS, Lee YJ. Association between hemorrhagic stroke occurrence and meteorological factors and pollutants. *BMC Neurol* 2016;16:59.
43. Liu H, Tian Y, Xu Y, Zhang J. Ambient particulate matter concentrations and hospitalization for stroke in 26 Chinese cities. *Stroke* 2017;48:2052-9.
44. Huang F, Luo Y, Guo Y, et al. Particulate matter and hospital admissions for stroke in Beijing, China: modification effects by ambient temperature. *J Am Heart Assoc* 2016;5:e003437.
45. Sun S, Stewart JD, Eliot MN, et al. Short-term exposure to air pollution and incidence of stroke in the Women's Health Initiative. *Environ Int* 2019;132:105065.
46. Fisher JA, Puett RC, Laden F, et al. Case-crossover analysis of short-term particulate matter exposures and stroke in the health professionals follow-up study. *Environ Int* 2019;124:153-60.
47. Li M, Edgell RC, Wei J, et al. Air pollution and stroke hospitalization in the Beibu Gulf Region of China: a case-crossover analysis. *Ecotoxicol Environ Saf* 2023;255:114814.
48. Villeneuve PJ, Chen L, Stieb D, Rowe BH. Associations between outdoor air pollution and emergency department visits for stroke in Edmonton, Canada. *Eur J Epidemiol* 2006;21:689-700.
49. Wang P, Song W, Li Y, et al. Short-term air pollutants exposure has heterogeneous associations with intracerebral hemorrhage risk: a case-crossover study based on symptom onset time in Chengdu, China. *Atmos Environ* 2022;273:118965.
50. Wang P, Feng W, Luo S, et al. Cleaner outdoor air diminishes the overall risk of intracerebral hemorrhage but brings differential benefits to subpopulations: a time-stratified case-crossover study. *BMC Public Health* 2023;23:1303.
51. Zheng M, Yin Z, Wei J, et al. Submicron particle exposure and stroke hospitalization: an individual-level case-crossover study in Guangzhou, China, 2014-2018. *Sci Total Environ* 2023;886:163988.
52. Xiang H, Mertz KJ, Arena VC, et al. Estimation of short-term effects of air pollution on stroke hospital admissions in Wuhan, China. *PLoS One* 2013;8:e61168.
53. Turin TC, Kita Y, Rumana N, et al. Short-term exposure to air pollution and incidence of stroke and acute myocardial infarction in a Japanese population. *Neuroepidemiology* 2012;38:84-92.
54. Montresor-López JA, Yanosky JD, Mittleman MA, et al. Short-term exposure to ambient ozone and stroke hospital admission: a case-crossover analysis. *J Expo Sci Environ Epidemiol* 2016;26:162-6.
55. Shah ASV, Lee KK, McAllister DA, et al. Short term exposure to air pollution and stroke: systematic review and meta-analysis. *BMJ* 2015;350:h1295.
56. Maida CD, Norrito RL, Rizzica S, Mazzola M, Scarantino ER, Tuttolomondo A. Molecular pathogenesis of ischemic and hemorrhagic strokes: background and therapeutic approaches. *Int J Mol Sci* 2024;25:6297.
57. Zhang R, Liu G, Jiang Y, et al. Acute effects of particulate air pollution on ischemic stroke and hemorrhagic stroke mortality. *Front Neurol* 2018;9:827.

Supplementary Material

To access the supplementary material accompanying this article, visit the online version of the *Canadian Journal of Cardiology* at www.onlinecjc.ca/ and at <https://doi.org/10.1016/j.cjca.2026.05.010>.