

Title	Transcriptome profiling defines a novel regulon modulated by the LysR-type transcriptional regulator MexT in <i>Pseudomonas aeruginosa</i>
Authors	Tian, Zhe-Xian;Fargier, Emilie;Mac Aogáin, Micheál;Adams, Claire;Wang, Yi-Ping;O'Gara, Fergal
Publication date	2009
Original Citation	Tian, Z.-X., Fargier, E., Mac Aogáin, M., Adams, C., Wang, Y.-P. and O'Gara, F. (2009) 'Transcriptome profiling defines a novel regulon modulated by the LysR-type transcriptional regulator MexT in <i>Pseudomonas aeruginosa</i> ', <i>Nucleic Acids Research</i> , 37(22), pp. 7546-7559. doi: 10.1093/nar/gkp828
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkp828 - 10.1093/nar/gkp828
Rights	© 2009, the Authors. Published by Oxford University Press. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by-nc/2.5/uk/), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited. - http://creativecommons.org/licenses/by-nc/2.5/uk/
Download date	2025-06-01 12:31:55
Item downloaded from	https://hdl.handle.net/10468/5030

Transcriptome profiling defines a novel regulon modulated by the LysR-type transcriptional regulator MexT in *Pseudomonas aeruginosa*

Zhe-Xian Tian^{1,2}, Emilie Fargier¹, Micheál Mac Aogáin¹, Claire Adams¹,
Yi-Ping Wang² and Fergal O’Gara^{1,*}

¹BIOMERIT Research Centre, Department of Microbiology, University College Cork, Cork, Ireland and

²National Laboratory of Plant Engineering and Protein Genetic Engineering, College of Life Science, Peking University, Beijing, P.R. China

Received July 22, 2009; Revised and Accepted September 18, 2009

ABSTRACT

The LysR-family regulator MexT modulates the expression of the MexEF-OprN efflux system in the human pathogen *Pseudomonas aeruginosa*. Recently, we demonstrated that MexT regulates certain virulence phenotypes, including the type-three secretion system and early attachment independent of its role in regulating MexEF-OprN. In this study, transcriptome profiling was utilized to investigate the global nature of MexT regulation in *P. aeruginosa* PAO1 and an isogenic *mexEF* mutant. Twelve genes of unknown function were highly induced by overexpressing MexT independent of MexEF-OprN. A well-conserved DNA motif was identified in the upstream regulatory region of nine of these genes and upstream of *mexE*. Reporter fusion analysis demonstrated that the expression of the genes was significantly induced by MexT in *P. aeruginosa* and a heterogenous *Escherichia coli* strain and that the conserved sequence was required for this induction. The conserved DNA motif was further characterized as the MexT binding site by site-directed mutagenesis and electrophoretic mobility shift assays. Genes containing this conserved regulatory sequence were identified across other *Pseudomonas* species, and their expression was activated by MexT. Thus, a novel regulon directly modulated by MexT, that includes but is independent of *mexEF-oprN*, has been identified.

INTRODUCTION

Pseudomonas aeruginosa is a leading cause of hospital-acquired infections and remains the most important

pathogen associated with chronic lung infections in cystic fibrosis patients. *Pseudomonas aeruginosa* is well known for its intrinsic resistance to a wide range of antimicrobial agents and its ability to develop multidrug resistance following antibiotic therapy (1). Resistance-Nodulation-Division (RND) efflux systems are responsible for much of the intrinsic and acquired multidrug resistance in *P. aeruginosa* and genes encoding 12 RND efflux pumps have been identified in its genome. To date, seven—MexAB-OprM, MexCD-OprJ, MexEF-OprN, MexGHI-OpmD, MexJK, MexXY and CzcAB-OpmN—have been functionally characterized and their expression is tightly regulated. Although these efflux pumps can exclude various antimicrobials, most of them are not induced by these substrates in wild-type strains (2). Nevertheless, genes encoding for RND efflux pumps are highly conserved in many living organisms (3) and, recently, increasing attention has been focused on understanding the physiological roles of efflux pumps, that probably precede antibiotics and are relevant to the behaviour of bacteria in their natural ecosystems (3–5). It is now evident that efflux pumps are involved in diverse cellular activities including functions such as the detoxification of intracellular metabolites, bacterial virulence in both animal and plant hosts, cell homeostasis and intercellular signal trafficking. However, little is known about how the expression of these efflux pumps is modulated within innate genetic networks to support these versatile cellular functions in response to changing environments.

MexEF-OprN is a unique RND pump that is modulated by a transcriptional activator, while other RND pumps are modulated by transcriptional repressors in *P. aeruginosa* (2). The expression of *mexEF-oprN* is activated by a LysR-type transcriptional regulator (LTTR) MexT, which is encoded by a gene located just upstream of *mexEF-oprN* in the same orientation in *P. aeruginosa* (6). The MexEF-OprN system is normally

*To whom correspondence should be addressed. Tel: +353 21 4902861; Fax: +353 21 4903101; Email: f.ogara@ucc.ie

quiescent in wild-type cells under normal laboratory conditions, but is highly induced in *nfxC*-type phenotypic mutants. These mutants exhibit increased resistance to chloramphenicol, trimethoprim and fluoroquinolones as well as exhibiting susceptibility to certain β -lactam and aminoglycoside antibiotics (7,8). The underlying mechanisms leading to *nfxC*-type mutants are not fully understood but appear to be multifactorial (9–11). For example, the frequency of isolating *nfxC*-type mutants has been associated with mutations in the *mexT* gene of the parent strain that render the protein active or inactive (9). Indeed, several mutations in the nucleotide sequence of *mexT* were identified in different laboratory PAO1 strains rendering the protein inactive and leading to a low frequency in the isolation of *nfxC*-type mutants. In other PAO1 strains, the *mexT* sequence encoded an active protein leading to a high frequency of *nfxC*-type mutants. In this class of PAO1 strain, the authors suggest that the expression or activity of MexT is suppressed under normal laboratory growth conditions (9). More recently, a deletion in the adjacent *mexS* gene was shown to result in an *nfxC*-type mutation (11). *mexS* encodes a probable oxidoreductase but its role in modulating MexT function is as yet unknown. Nevertheless, in all cases, it appears a functional MexT is crucial for the generation of *nfxC*-type mutants and this function is suppressed under normal growth conditions. Accordingly, and in support of the demand theory of gene regulation whereby the design of molecular control mechanisms is generally dependent on the demand for expression of the regulated genes (12), it is predicted that the expression of *mexEF-oprN* and indeed other possible targets of MexT may be in high demand in natural environments of *P. aeruginosa*.

Besides the *mexEF-oprN* operon, MexT can activate the expression of *mexS*, previously referred to as *qhr*, in *P. aeruginosa* (6). *mexS* is located adjacent to *mexT* and transcribed in the opposite direction, which is a typical arrangement of genes controlled by LTTRs (13). A palindromic DNA sequence has been identified in the promoter region of target genes, to which LTTRs are known to bind. This LTTR box was first identified in *Azorhizobium caulinodans* as an interrupted palindrome with the sequence ATC-N9-GAT, referred to as the 'nod-box' (14). Indeed, a *nod*-box-like sequence in the promoter region of *mexE* has been proposed as the binding site of MexT (6). Interestingly, the promoter region of *mexS* does not contain a clear 'nod-box'.

In addition to antibiotic resistance, overexpression of MexEF-OprN in *nfxC*-type mutants has been linked to reduced levels of homoserine lactone-dependent virulence traits, including pyocyanin, elastase, rhamnolipids and PQS (15) and to reduced expression of type-three secretion effector proteins (16). It was suggested that MexEF-OprN mediates these effects via efflux of cell-signalling intermediates, which ultimately commits the cell to a state of reduced virulence (15,17). Recently, we have observed that MexT downregulates several virulence determinants, such as TTSS gene expression, pyocyanin biosynthesis and early surface attachment, in a MexEF-OprN independent manner in *P. aeruginosa* PAO1 (18). We have suggested that MexT acts as a more global

regulator than previously described and novel target genes may be directly activated by MexT in *P. aeruginosa*. This is supported by the fact that *mexT* is not tightly linked to the *mexEF-oprN* in the genomes of most *Pseudomonas* species, according to *Pseudomonas* Genome Database (www.pseudomonas.com). For instances, *mexT* (PP_2826) and *mexE* (PP_3425) are located in totally different loci in the genome of *P. putida* KT2440 strain.

Elucidating the transcriptional regulatory network linked to MexT will provide a new angle to understand the physiological roles of the MexEF-OprN pump in a broader context. In this study, we demonstrate the existence of a novel regulon incorporating *mexEF-oprN* in *Pseudomonas* species. MexT activates the expression of this novel regulon by binding to a conserved DNA motif in their promoter regions.

MATERIALS AND METHODS

Bacterial strains and culture conditions

The bacterial strains and plasmids used in this study are shown in Table 1. All *Escherichia coli* and *P. aeruginosa* PAO1 strains were routinely grown at 37°C, while *P. fluorescens* PfO-1 and *P. putida* KT2440 strains were grown at 30°C in Luria–Bertani (LB) broth with aeration. When required, antibiotics were added at the following concentrations ($\mu\text{g/ml}$): *E. coli*, ampicillin (Ap, 50), kanamycin (Km, 50), chloramphenicol (Cm, 20), tetracycline (Tc, 10) or gentamicin (Gm, 10); *P. aeruginosa* PAO1, carbenicillin (Cb, 200), gentamicin (Gm, 50), tetracycline (Tc, 50) or streptomycin (Str, 200).

Molecular biology procedures

All routine molecular biology procedures were carried out as described in the standard laboratory manual (19). All DNA primers used in this study are listed in Supplementary Table S1.

Expression profiling experiments

PAO1 (pME6032), PAO1 (pME6032-*mexT*) and PAO1 Δ *mexEF* (pME6032-*mexT*) were grown at 37°C shaking (150 r.p.m.) in 20 ml of LB medium supplemented with tetracycline in 100-ml culture flasks. At an OD₆₀₀ of 0.5, cell cultures were mixed with Qiagen RNeasy Protect Bacteria Reagent (Ratio of 1:2) to stabilize RNA. Total RNA was prepared with an RNeasy mini kit (Qiagen) and contaminating DNA was removed by RNase-free DNase I (Ambion). For each strain, RNA was prepared from three independent batch cultures. cDNA synthesis and hybridization to Affymetrix GeneChip *P. aeruginosa* genome arrays were carried out by a commercial Affymetrix Genechip service supplier (Conway Institute of Biomolecular & Biomedical Research, UCD, Ireland). Array data were normalized using the GC-RMA algorithm, and the data were then further analyzed using the microarray software package Genespring GX 10.0.1 (Agilent). To visualize the microarray data, MA plots (linear regression of log ratio, M, against average

intensity, A) were generated by Genespring GX software, and 3D scatter plots on log₂ based intensity data obtained from three samples were generated by SigmaPlot 11 software (Systat Software Inc.). Genes whose levels of expression were significantly influenced (Fold change ≥ 2 , $P \leq 0.05$) were identified with Genespring GX, using a Benjamini–Hochberg multiple-testing correction and a false detection rate of 5%. The full list of genes whose expression was significantly altered is shown in Supplementary Table S2.

To validate the microarray results, semi-quantitative reverse transcription PCR analysis was carried out using RNA samples prepared from biological replicates not used in the arrays. Two hundred nanograms of RNA samples were reverse transcribed using random primers and AMV reverse transcriptase (Promega). 1/100 aliquot of the reaction mixture was used as template in the subsequent PCR. PCR (20 μ l reaction volume) was performed with the GoTaq[®] Green Master Mix (Promega) with the following conditions: 95°C for 3 min; 30 cycles of 30 s at 95°C; 30 s at 55°C and 30 s at 72°C; 10-min elongation at 72°C. For visualization, 10 μ l of the resulting PCR was subjected to agarose gel electrophoresis and stained with ethidium bromide. The *clpX* gene was used as an internal control to ensure equal amounts of RNA were used in all samples.

Generation of PAO1*nfxC* mutant and PAO1*nfxC* Δ *mexT* strains

PAO1*nfxC* was selected by plating the wild-type strain PAO1 on LB agar plates containing chloramphenicol at 600 μ g/ml, a condition that exclusively selects *nfxC*-type mutants (7). The *mexT* ORF and 500-bp upstream region of this PAO1*nfxC* strain had no sequence alteration(s) compared to the parental PAO1 strain.

In order to construct PAO1*nfxC* Δ *mexT*, a 1-kb SacI–KpnI upstream region including the start codon of *mexT*, and a 1-kb KpnI–BamHI downstream region containing the stop codon of *mexT*, were PCR-amplified and linked together. The resulting 2-kb fragment was cloned into the suicide plasmid pEX18Tc digested with SacI and BamHI. A 1.1-kb KpnI fragment containing the *FRT* gentamicin-resistance (Gm) cassette from plasmid pPS856 (20) was then inserted in-between the flanking regions on the plasmid. The *mexT* of PAO1*nfxC* was then replaced with the plasmid as described by Hoang *et al.* (20). The Gm-resistance sequence in the chromosome was removed by introducing plasmid pFLP2, which carries the Fip recombinase gene (20). Correct insertion in the constructed mutant was verified by PCR using primers bind to flanking chromosomal regions of the fragments cloned in pEX18Tc.

Construction of the promoter-*lacZ* reporter gene fusions and β -galactosidase assays

The promoter region of each putative MexT target gene was PCR-amplified and TA-cloned into pCR2.1-TOPO (Invitrogen). The site-directed mutagenesis of the conserved DNA motif was performed by the protocol described previously (21). Once confirmed by sequencing,

the promoter regions were subcloned into the broad-host low-copy-number plasmid pMP190 (Table 1). The resulting plasmids were introduced into *P. aeruginosa* strains by conjugal transfer from *E. coli* donor strain ST18. ST18 is a *hemA* mutant of S17-1 λ pir strain (Table 1), and can be easily selected against as its growth is strictly dependent on the presence of 5-amino-levulinic acid (5-ALA), even in LB medium (22). Therefore, 50 μ g/ml of 5-ALA was added to the medium during conjugation and omitted when selecting for trans-conjugants. Since PAO1 containing pME6032-*mexT* and PAO1*nfxC* strains have a high chloramphenicol resistance due to the *nfxC* phenotype, all trans-conjugant *P. aeruginosa* cells containing pMP190-derived plasmids were selected on LB agar supplemented with 200 μ g/ml streptomycin. The plasmid transformation into *E. coli* strains in this study was done by routine chemical competent cell protocol (19).

For β -galactosidase assays, cells were grown overnight in LB broth supplemented with appropriate antibiotics, then 1:50 diluted into 10 ml fresh medium in 100-ml culture flasks and incubated at 37°C shaking at 150 r.p.m. Cells were recovered at logarithmic growth phase (OD₆₀₀ 0.5–1.2), and β -galactosidase assays were performed as described by Miller (23). Data are the mean of two independent experiments with triplicate samples.

Purification of His₆-MexT

The plasmids used to overexpress the C-terminal His₆-tagged MexT proteins were constructed by PCR-amplifying the MexT coding sequence and cloned into NcoI–XhoI sites of pET28a. The plasmid harbouring the functional *mexT* from PAO1 was designated as pETmexTH6C, while the plasmid harbouring the non-functional *mexT* due to a mutation of the 39th amino acid Ala to Val was designated as pETmexTm1H6C (Table 1). The plasmids were transformed into the *E. coli* expression host strain BL21-CodonPlus(DE3)-RPL (Merck) and grown at 37°C with vigorous shaking in 200 ml of LB medium containing kanamycin (50 μ g/ml) to an OD₆₀₀ of 1. The cells were then induced with 1 mM IPTG and allowed to express for 1 h at 37°C. The cells were harvested by centrifugation and stored overnight at –20°C. The pellet was resuspended in CelLytic[™] B II buffer (Sigma[®]) (10 ml per gram cell paste) with 5 μ g/ml DNase and 200 μ l per gram cell paste of Protease Inhibitor Cocktail (Sigma[®]). The soluble proteins were extracted according to manufacturer's instructions (Sigma[®], CelLytic[™] B II, Bacterial Cell Lysis Extraction Reagent). The protein extract were applied to a Poly-Prep[®] Chromatography Column (Bio-Rad) containing 1 volume of HIS-Select[™] Nickel Affinity Gel (Sigma[®]). The resin was previously washed with 2 vol of deionized sterile water and equilibrated with 2 vol of Wash buffer (HEPES 100 mM, pH 7.5, 1 mM Imidazole). After all the protein extract was loaded, the column was washed twice with 2 vol of Wash buffer. The His-tag proteins were eluted with 1 vol of Elution buffer (HEPES 100 mM, pH 7.5, 500 mM Imidazole) and their

Table 1. Strains and vectors used in this study

Genotype or phenotype		Source
<i>Pseudomonas aeruginosa</i>		
PAO1 wild-type	Lab stock	
PAO1 <i>AmexEF</i>	<i>mexE-mexF</i> deletion mutant of PAO1	(18)
PAO1 <i>nfxC</i>	<i>mexEF-oprN</i> overexpressing mutant of PAO1	This work
PAO1 <i>nfxCAmexT</i>	<i>mexT</i> deletion mutant of PAO1 <i>nfxC</i>	This work
<i>Escherichia coli</i>		
DH5 α	F- ϕ 80 <i>lacZAM15</i> (<i>AlacZYA-argF</i>)U169 <i>deoR recA1 endA1 hsdR17</i> (r_k^- , m_k^-) <i>phoA supE44 thi-1 gyrA96 relA1Δ</i>	Invitrogen
ST18	<i>hemA</i> λ pir lysogen of S17-1: <i>pro</i> , <i>Res</i> ⁻ , <i>Mob</i> ⁺ , <i>RecA</i> ⁻ derivative of <i>E. coli</i> 294 with Rp4-2 (Tc::mu) (Km::Tn7) in the chromosome	(22)
BL21-CodonPlus (DE3)-RIPL	protein expression host	Merck
<i>Pseudomonas fluorescens</i>		
PfO-1	wild-type	(41)
<i>Pseudomonas putida</i>		
KT2440	mt-2 <i>hsdRI hsdM</i> +	(42)
Vectors		
pCR2.1-TOPO TA	cloning vector, Ap ^r , Km ^r	Invitrogen
pME6032	pVS1-p15A origin, <i>lacI</i> ^q - <i>Ptac</i> expression vector, Tc ^r	(43)
pME6032- <i>mexT</i>	pME6032-derived PAO1 <i>mexT</i> expression vector	(18)
pEX18Tc	<i>oriT</i> ⁺ <i>sacB</i> ⁺ , gene replacement vector, Tc ^r	(20)
pPS856	Gm ^r cassette flanked with FRT sequences, Ap ^r , Gm ^r	(20)
pFLP2	inducible FLP recombinase, Ap ^r	(20)
pMP190	IncQ origin, low-copy-number <i>lacZ</i> fusion vector; Cm ^r Str ^r	(44)
pMP-PAmexEp	pMP190-derived PAO1 <i>mexE</i> promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA1744p	pMP190-derived PAO1 PA1744 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA2759p	pMP190-derived PAO1 PA2759 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA3229p	pMP190-derived PAO1 PA3229 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA4354p	pMP190-derived PAO1 PA4354 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA4623p	pMP190-derived PAO1 PA4623 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA4881p	pMP190-derived PAO1 PA4881 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PA4623mlp	conserved DNA motif mutated version of pMP-PA4623p	This work
pMP-PA4881mlp	conserved DNA motif mutated version of pMP-PA4881p	This work
pMP-Pf012659p	pMP190-derived PfO-1 Pf01_2659 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-Pf013748p	pMP190-derived PfO-1 Pf01_3748 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PP3425p	pMP190-derived <i>P. putida</i> KT2440 PP_3425 promoter- <i>lacZ</i> fusion plasmid	This work
pMP-PP4858p	pMP190-derived <i>P. putida</i> KT2440 PP_4858 promoter- <i>lacZ</i> fusion plasmid	This work
pET28a	T7 promoter-driven His-tag protein expression vector, Km ^r	Novagen
pETmexTH6C	pET28a-derived C-terminal His6-tag PAO1 <i>mexT</i> expression vector	This work
pETmexTmlH6C	pET28a-derived C-terminal His6-tag non-functional <i>mexT</i> expression vector	This work

Ap^r, ampicillin resistance; Km^r, kanamycin resistance; Tc^r, tetracycline resistance; Cm^r, chloramphenicol resistance; Str^r, streptomycin resistance; Gm^r, gentamycin resistance.

homogeneity was verified by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS–PAGE). The protein was then either immediately frozen with 20% of glycerol at –80°C for later use or promptly used for the electrophoretic mobility shift assay (EMSA). The protein concentrations were determined by the Bio-Rad method.

EMSAs

DNA probes of 207 bp containing the putative binding region were generated by PCR using pMP-PA4881p (intact motif) or pMP-4881mlp (mutated motif) as templates using the primer pair fullmexTboxF and fullmexTboxR. The two purified PCR products were 3'-end-labelled with digoxigenin following the manufacturer instruction (Roche Applied Science). The EMSA was carried out using the DIG Gel Shift Kit 2nd Generation (Roche Applied Science) as recommended, with some modifications. Ten fmoles of the DIG-labelled fragment and a range of 0–300 ng of MexT protein were added to the binding reaction. The mixture was allowed to proceed

for 45 min at room temperature. The samples were separated by electrophoresis on 6% native polyacrylamide gels and transferred to Hybond-N blotting membrane (Amersham Life Science) by electroblotting. Protein–DNA complexes were visualized by NBT/BCIP according to manufacturer's instructions (Roche Applied Science). Competition experiments were performed with 200 pmol of unlabelled double-stranded oligonucleotide. The labelled and non-labelled probes were mixed to the binding buffer before adding the protein for the binding reaction.

Identification of a conserved regulatory sequence

For the identification of a conserved regulatory sequence motif in the upstream region of co-regulated genes, the 500-bp sequence upstream of the translational start site of each gene induced by MexT was obtained from the 'www.pseudomonas.com' database. Sequence motifs common to the upstream regions were identified by the online MEME software (http://meme.sdsc.edu/meme4_

1/cgi-bin/meme.cgi) (24) with a motif width range of 5–20 bp and the occurrences of a single motif distributed among the sequences between zero or one per sequence.

RESULTS

Transcriptome profiling of *P. aeruginosa* strains overexpressing MexT

In order to assess the global regulatory nature of the LysR-family transcriptional regulator MexT, independent of its induction of the MexEF-OprN efflux system, whole-genome transcriptome profiling was carried out on PAO1 wild-type and an isogenic *mexEF* mutant overexpressing MexT. In the PAO1 strain used in this study MexT is quiescent under normal laboratory growth conditions. However, when the entire ORF and 600-bp upstream of the *mexT* gene were cloned on plasmid pME6032 (designated as pME6032-*mexT*), MexT was fully functional in *P. aeruginosa* PAO1, increasing the minimum inhibitory concentration (MIC) of chloramphenicol from 32 to 2048 µg/ml, even without IPTG induction of the *tac* promoter on pME6032 (18). Affymetrix *P. aeruginosa* whole-genome microarrays were used to compare the transcription profiles of PAO1 wild-type containing either pME6032-*mexT* or the empty vector control pME6032 and an isogenic *mexEF* deletion mutant strain, PAO1Δ*mexEF*, containing pME6032-*mexT*, during Mid-logarithmic growth conditions. Thus, we identified a potential MexT regulon and differentiated those targets that were dependent or independent of the MexEF-OprN efflux pump.

Genespring GX software was used to analyse the data (Supplementary Table S2) and MA and 3D scatter plots were used to visually compare the overall gene expression profiles of all three samples (Figure 1). Overexpressing *mexT* in PAO1 wild-type (PAO1 *mexT*⁺⁺) increased the expression of 36 genes and decreased the expression of 76 genes compared to PAO1 harbouring the pME6032 empty vector (≥2-fold, *P* ≤ 0.05; see Supplementary Table S2). The most significantly altered genes (≥10-fold) are summarized in Table 2, and include 17 genes upregulated and 23 genes downregulated by overexpressing MexT in wild-type. These genes were also indicated as the most significantly altered genes in the MA plot in Figure 1A. To test if this modulation in gene expression was dependent on the MexEF-OprN pump, the transcriptome profile of PAO1Δ*mexEF* *mexT*⁺⁺ was compared with PAO1 *mexT*⁺⁺. Eighteen of the genes downregulated by MexT in wild-type were upregulated in PAO1Δ*mexEF* *mexT*⁺⁺, indicating their regulation was dependent on *mexEF* expression. In contrast, none of the 17 genes (apart from the *mexEF-oprN* operon) induced by MexT in wild-type was altered in PAO1Δ*mexEF* *mexT*⁺⁺ compared to PAO1 *mexT*⁺⁺ (Table 2), indicating their induction by MexT was independent of *mexEF* expression. This was also evident from the changes in MA plot of PAO1 *mexT*⁺⁺ compared to PAO1Δ*mexEF* *mexT*⁺⁺ whereby only three genes (*mexE*, *mexF* and *oprN*) were indicated as being highly induced, while 18 genes were highly reduced (Figure 1B). The differentiation of

regulatory effects of MexT with regard to their dependency on the MexEF-OprN efflux pump was also evident in the 3D scatter plot (Figure 1C). The expression of the majority of genes was similar among the three samples, forming the main cloud body. In contrast, genes whose expression was highly altered were positioned out of the main cloud body in the 3D scatter plot. These were further categorized into two groups: one incorporating the genes induced by overexpressing MexT independent of MexEF-OprN (in red oval) and the second incorporating the genes downregulated by overexpressing MexT dependent of MexEF-OprN (in green oval). This supports the MA plot analysis and demonstrates that the most significant induction in gene expression by MexT is independent of MexEF-OprN efflux pump, while the most significant reduction by MexT is dependent of MexEF-OprN, in the condition tested. These results support our hypothesis that the genes highly induced by MexT are direct targets of the regulator.

To test the validity of the microarray data, genes whose expression was significantly altered (PA4881, *pqsA* and *lldP*) were selected for semi-quantitative reverse transcription PCR analysis (Figure 2). RNA was prepared from biological replicates not used in the arrays. The housekeeping gene *clpX* was used as the internal control. The transcript level of PA4881 was induced by overexpressing MexT in both PAO1 and PAO1Δ*mexEF* strains but was not detectable in the PAO1 strain containing pME6032. The transcript levels of *pqsA* and *lldP* were reduced by overexpressing MexT in PAO1 but not in PAO1Δ*mexEF* strains. Furthermore, the expression from promoter fusions of a selected number of genes were analysed in the wild-type PAO1 and a heterogeneous *E. coli* background, with and without pME6032-*mexT*, and all genes were highly induced by overexpressing MexT in both backgrounds (Table 3). In all cases, the results confirmed the microarray data.

Due to the limited number and lack of good annotation of the significantly altered genes in the expression profiles (Supplementary Table S2), the pathway mapping through main pathway databases such as KEGG (www.genome.jp/kegg/pathway.html) provided limited information. However, the expression of several operons with known cellular functions was significantly altered by MexT in a *mexEF-oprN*-dependent way. The transcript levels of the *pqs* and *phn* operon genes, encoding enzymes involved in the synthesis of PQS signal molecules, showed a 30- to 220-fold reduction in PAO1 wild-type, when *mexT* was overexpressed and a 23- to 240-fold induction in PAO1Δ*mexEF* *mexT*⁺⁺ relative to PAO1 *mexT*⁺⁺. Similarly, the *hcn* operon genes, encoding hydrogen cyanide synthase, showed a 11- to 52-fold reduction in PAO1 wild-type overexpressing *mexT*, while *hcnA* was upregulated 11-fold in PAO1Δ*mexEF* *mexT*⁺⁺ relative to PAO1 *mexT*⁺⁺. Previously we demonstrated a considerable reduction in *hcnB* expression by overexpressing *mexT* in PAO1 strain but not in PAO1Δ*mexEF* strain using semi-quantitative reverse transcription PCR (18). Our results indicated that *pqs* and *hcn* operons were indirect targets of MexT, as the modulation in expression was dependent on the overexpression of the MexEF-OprN




Figure 1. (A and B) MA-plots showing the relationship between the logarithmic mean signal ratios, M value, and the logarithmic mean signal intensities, A value, are used to spot genes whose expression is significantly altered in (A) PAO1 $mexT^{++}$ compared to PAO1 and (B) PAO1 $mexT^{++}$ compared to PAO1 $\Delta mexEF mexT^{++}$. The spots representing genes altered more than 10-fold are labelled in red. A positive M value indicates increased gene expression, while a negative value indicates reduced expression. (C) 3D scatter-plot projecting the logarithmic mean signal intensities obtained from PAO1, PAO1 $mexT^{++}$ and PAO1 $\Delta mexEF mexT^{++}$ is used to spot genes whose expression was altered by overexpressing $MexT$ dependent or independent of the $MexEF$ -OprN efflux pump. (Left) The orientation of the main cloud body is adjusted towards the reader, genes highly activated by overexpressing $MexT$ independent of the $MexEF$ -OprN efflux pump are grouped in the red oval and genes highly reduced by overexpressing $MexT$ dependent of the $MexEF$ -OprN efflux pump are grouped in the green oval. (Right) The orientation of the main cloud body is adjusted to show the full scale.

Table 2. Genes of expression altered more than 10-fold in response to MexT overexpression

Gene No. ^a	Gene name	PAO1 <i>mexT</i> ⁺⁺ versus PAO1 control ^b	PAO1Δ <i>mexEF</i> <i>mexT</i> ⁺⁺ versus PAO1 <i>mexT</i> ⁺⁺ ^c	Protein description
PA0852	<i>cbpD</i>	−9.4	20	Chitin-binding protein CbpD precursor
PA0996	<i>pqsA</i>	−64	95	Probable coenzyme A ligase
PA0997	<i>pqsB</i>	−51	72	Homologous to β-keto-acyl-acyl-carrier protein synthase
PA0998	<i>pqsC</i>	−219	241	Homologous to β-keto-acyl-acyl-carrier protein synthase
PA0999	<i>pqsD</i>	−72	86	3-Oxoacyl-[acyl-carrier-protein] synthase III
PA1000	<i>pqsE</i>	−87	76	Quinolone signal response protein
PA1001	<i>phnA</i>	−126	95	Anthranilate synthase component I
PA1002	<i>phnB</i>	−34	23	Anthranilate synthase component II
PA1657		−36	<2	Conserved hypothetical protein
PA1658		−12	<2	Conserved hypothetical protein
PA1718	<i>pseE</i>	−12	<2	Type III export protein PseE
PA1744		48	<2	Hypothetical protein
PA1869		−54	11	Probable acyl carrier protein
PA1970		237	<2	Hypothetical protein
PA2193	<i>hcnA</i>	−52	11	Hydrogen cyanide synthase HcnA
PA2194	<i>hcnB</i>	−32	<2	Hydrogen cyanide synthase HcnB
PA2195	<i>hcnC</i>	−11	<2	Hydrogen cyanide synthase HcnC
PA2386	<i>pvdA</i>	<2	10	L-ornithine N5-oxygenase
PA2486		338	<2	Hypothetical protein
PA2491	<i>mexS</i>	75	<2	Probable oxidoreductase
PA2492	<i>mexT</i>	23.3	<2	Transcriptional regulator MexT
PA2493	<i>mexE</i>	2033	−1714	RND multidrug efflux membrane fusion protein MexE precursor
PA2494	<i>mexF</i>	1382	−9.5	RND multidrug efflux transporter MexF
PA2495	<i>oprN</i>	340	−10	Multidrug efflux outer membrane protein OprN precursor
PA2759		72	<2	Hypothetical protein
PA2811		34	<2	Probable permease of ABC-2 transporter
PA2812		12	<2	Probable ATP-binding component of ABC transporter
PA2813		20	<2	Probable glutathione S-transferase
PA3205		11	<2	Hypothetical protein
PA3229		1789	<2	Hypothetical protein
PA3326		−75	26	Probable Clp-family ATP-dependent protease
PA3331		−26	5.0	Cytochrome P450
PA3332		−18	<2	Conserved hypothetical protein
PA3333	<i>fabH2</i>	−13	3.7	3-Oxoacyl-[acyl-carrier-protein] synthase III
PA4141		−32	19	Hypothetical protein
PA4354		11	<2	Conserved hypothetical protein
PA4623		74	<2	Hypothetical protein
PA4770	<i>lldP</i>	−43	46	L-lactate permease
PA4771	<i>lldD</i>	−28	70	L-lactate dehydrogenase
PA4772		−7.0	19	Probable ferredoxin
PA4881		751	<2	Hypothetical protein

^aGene number from the *Pseudomonas* Genome Project (<http://www.pseudomonas.com>).^bFold change in gene expression of PAO1 (pME6032-*mexT*) compared to PAO1 (pME6032); positive value means expression increased and negative value means expression decreased.^cFold change in gene expression of PAO1Δ*mexEF* (pME6032-*mexT*) compared to PAO1 (pME6032-*mexT*).

efflux pump. A reduction in expression of *pqs* and *hcn* operons was also observed previously in the transcriptome profiling of an *nfxC*-type mutant strain overexpressing *mexEF-oprN* (10). The microarray data presented here also suggest that the *lldPD* operon (PA4770-PA4772), PA3326, and the PA3331-PA3334 operon are novel MexEF-OprN-dependent indirect targets of MexT.

The set of genes upregulated by MexT independent of MexEF included *mexS*, previously shown to be activated by MexT in PAO1 (6). The *mexS* gene is located adjacent to *mexT* and divergently transcribed, which is a typical arrangement of genes controlled by LysR-type activators (13). In our transcriptome data, the *mexS* transcript level increased 75-fold in PAO1 overexpressing MexT and was not changed in PAO1Δ*mexEF* *mexT*⁺⁺ relative to PAO1

mexT⁺⁺ (Table 2). These data supported the hypothesis that *mexS* is an MexE-independent target of MexT in *P. aeruginosa* PAO1.

Twelve additional genes were upregulated by overexpressing *mexT* independent of MexEF-OprN (Table 2). Most of these genes were annotated as of unknown function but included genes with probable transporter functions; PA2813-PA2811 encode components of a probable ABC transporter and the PA4354-PA4356 operon encodes a putative repressor, a probable major facilitator superfamily (MFS) transporter and a xenobiotic reductase, respectively. We hypothesized that this set of genes includes direct positively regulated targets of MexT and suggests that MexT may be a key regulator of cellular homeostasis.

Identification of a conserved DNA motif required for MexT regulated expression.

In order to identify possible direct targets of MexT and a potential consensus sequence for MexT binding, the genes that were most highly induced (≥ 10 -fold) by MexT in PAO1 independent of *mexEF* were selected for further analysis (Table 2). In addition to *mexS* and *mexEF-oprN*, the 12 genes showing induction ≥ 10 -fold were PA1744, PA1970, PA2486, PA2759, PA2813-PA2811, PA3205, PA3229, PA4354, PA4623 and PA4881. The DNA sequence of the upstream regulatory region (-500 bp from the ATG translational start site) of each gene (including PA2813 only of the PA2813-PA2811 operon) was aligned using MEME Suite online software

(24). A well-conserved DNA motif ATCA-N₅-GTCGAT-N₄-ACYAT was identified in the upstream regions of 9 of the 12 genes and *mexE* (Figure 3). Interestingly, a clear consensus sequence was not identified in the *mexS* promoter region. The DNA motif overlapped the ATC-N₉-GAT sequence, referred to as a 'nod-box' and proposed as the general motif of the binding sites for LysR-family proteins (13). Previously, this 'nod-box' was proposed to be part of the binding site of MexT in the upstream region of *mexE* (6).

To confirm the transcriptional activation of these putative target genes by MexT, promoter-*lacZ* fusions of selected genes were constructed, transformed into strains of different genetic backgrounds and the level of expression monitored when MexT was overexpressed (Table 3).




Figure 2. Semi-quantitative reverse transcription PCR validation of selected genes identified by microarray analysis as significantly altered by MexT. Total RNA was isolated from PAO1 containing pME6032 (WT vector), PAO1 containing pME6032-*mexT* (WT *mexT*) or PAO1Δ*mexEF* containing pME6032-*mexT* (ΔEF *mexT*). The identity of the genes targeted in each set of PCRs is indicated at the bottom of the figure.

Table 3. The regulatory effect of MexT on the expression of the promoter-*lacZ* fusions of putative target genes

Strain (plasmid) Promoter- <i>lacZ</i>	PAO1 (pME6032)	PAO1 (pME6032- <i>mexT</i>)	PAO1 <i>nfxC</i>	PAO1 <i>nfxCAmexT</i>	<i>E. coli</i> DH5a (pME6032) ^a	<i>E. coli</i> DH5a (pME6032- <i>mexT</i>) ^a
pMP-PAmexEp	45 ± 1	3991 ± 912	954 ± 21	33 ± 2	9 ± 1	230 ± 1
pMP-PA1744p	120 ± 32	2801 ± 143	NA ^b	NA	7 ± 3	7520 ± 849
pMP-PA2759p	52 ± 5	896 ± 104	NA	NA	2 ± 1	4147 ± 666
pMP-PA3229p	76 ± 19	4852 ± 120	NA	NA	4 ± 1	1148 ± 96
pMP-PA4354p	824 ± 83	3385 ± 462	NA	NA	3 ± 1	143 ± 1
pMP-PA4623p	23 ± 3	4302 ± 30	1714 ± 86	29 ± 4	41 ± 4	6479 ± 724
pMP-PA4881p	34 ± 11	1217 ± 91	716 ± 43	20 ± 1	8 ± 2	7783 ± 465
Mutated DNA motif:						
pMP-PA4623m1p	30 ± 10	39 ± 6	25 ± 2	26 ± 1	21 ± 19	18 ± 12
pMP-PA4881m1p	38 ± 9	12 ± 5	22 ± 3	20 ± 2	13 ± 3	12 ± 1
Homologous genes across <i>Pseudomonas</i> species:						
pMP-Pfl2659p	NA	NA	NA	NA	9 ± 1	801 ± 15
pMP-Pfl3748p	NA	NA	NA	NA	7 ± 1	10 474 ± 202
pMP-PP3425p	NA	NA	NA	NA	4 ± 1	2231 ± 168
pMP-PP4858p	NA	NA	NA	NA	2 ± 1	3288 ± 43

^a IPTG (1 mM final conc.) was added in the medium.

^b NA, not assayed.




Figure 3. Alignment of the predicted conserved DNA motif in the upstream regulatory regions of *P. aeruginosa* genes highly induced by MexT (A) and sequence logo for the conserved DNA motif reflecting position-specific probability matrixes (B) deduced from MEME software analysis. Nucleotides with high probability ($\geq 70\%$) are highlighted by red. The number of nucleotides between the conserved DNA motif and the start codon ATG is shown in brackets.

In the wild-type PAO1 strains, the expression level of all the promoters containing the newly deduced conserved DNA motif was highly induced by co-transforming pME6032-*mexT*, compared to the empty vector control (Table 3). As MexEF-OprN is normally quiescent in wild-type cells, but highly induced in *nfxC*-type mutants (7), we investigated if the genes were also induced in an *nfxC*-type mutant background. Therefore, an *nfxC*-type mutant was generated by plating the wild-type strain PAO1 on LB agar plates containing chloramphenicol at 600 $\mu\text{g}/\text{ml}$, a condition previously shown to select *nfxC*-type mutants (7). A selection of promoter-*lacZ* fusion constructs were transformed into the *nfxC* mutant and high levels of expression were observed from all constructs tested, including *mexE* (Table 3). When the *mexT* gene was deleted in this *nfxC*-type mutant the expression of these genes was attenuated (Table 3). We also tested other *nfxC*-type mutant strains from different sources, and all of them showed high induction of PA4881 (data not shown). These results indicated that all the genes with the conserved DNA motif were highly induced in the *nfxC*-type mutant strains and thus, induced by MexT in physiologically relevant conditions.

To further investigate if these genes are direct targets of MexT, we assessed if MexT could activate their expression in a heterogenous *E. coli* background. The promoter-*lacZ* fusion plasmids and pME6032-*mexT* or pME6032 were co-transformed into the *E. coli* DH5 α strains. The *mexT* gene cloned in pME6032-*mexT* was highly expressed in *P. aeruginosa* PAO1 without IPTG induction (Table 2). However, in *E. coli*, the native *mexT* promoter was not induced and IPTG induction was essential for full activation (data not shown). Therefore, a saturated concentration of IPTG (1 mM) was used to induce *mexT* expression from pME6032-*mexT* in *E. coli*. As shown in Table 3, low expression levels of the putative MexT target promoters, including *mexE* was observed in the *E. coli* strains containing empty vector pME6032. In contrast, when *mexT*

was overexpressed, the expression of the promoters were significantly induced (Table 3). These results indicate that MexT may directly activate the expression of the newly identified target genes in *P. aeruginosa*. This was further supported by the observation that an overexpressing *mexT* construct, cloned from a different PAO1 wild-type available in the laboratory collection, and which contained a single base-pair mutation that resulted in an alanine to valine (A to V) change in the 39th residue of the MexT protein, failed to induce the expression of these promoter fusions in either PAO1 or *E. coli* DH5 α backgrounds (data not shown). According to the Pfam database alignment (25), the 39th residue is located in the helix–turn–helix motif of MexT, the DNA-binding motif of the protein. Indeed our results suggested that a mutation in this domain rendered the protein non-functional. While other inactivating mutations in the *mexT* gene have been reported (9) this single base pair mutation has not been previously identified in wild-type strains and appears to be a novel mechanism of inactivating *mexT*. Subsequently, to confirm that the conserved DNA motif in the promoter region was required for MexT activation of the putative target genes, site-directed mutagenesis was carried out to mutate GTCGAT to GTCTGT in the DNA motif of PA4623 and PA4881 promoters. This alteration in the DNA motif completely attenuated activation of the promoters by MexT in the wild-type PAO1, PAO1*nfxC* and the heterogenous *E. coli* DH5 α (Table 3). Thus, the conserved DNA motif is required for the induced expression of these genes by MexT.

MexT protein binds to the conserved DNA motif in target promoters

In order to verify if the MexT protein directly binds the conserved DNA motif, a plasmid expressing C-terminal His-tagged MexT protein was constructed. This tagged MexT protein activated the expression of PA4881 promoter-*lacZ* fusion in *E. coli*, indicating the protein was functional (data not shown). EMSAs were performed to detect binding of MexT to two 207-bp DNA fragments containing the unaltered or mutated PA4881 promoter region. The MexT protein clearly bound to the DNA fragments containing the intact conserved DNA motif (Figure 4A), but not to the DNA fragments containing the mutated conserved DNA motif (Figure 4B). The specificity of MexT binding to its DNA target was confirmed by a competition assay with an excess of unlabeled target DNA (Supplementary Figure S1). These results demonstrated, for the first time, that MexT protein bound to the conserved DNA motif and that a mutation in two of the most conserved nucleotides in the motif abolished the binding of the MexT protein.

To further investigate MexT binding to this conserved motif, the residue 39 A to V mutated non-functional MexT was purified using the His-tag expression plasmid previously used for the wild-type MexT protein. EMSAs were performed to detect binding of the mutated MexT protein to the conserved DNA motif and no binding was detected (Figure 4C). These results demonstrated that this




Figure 4. Purified MexT protein binds to the conserved DNA motif in the promoter region of PA4881. EMSAs were carried out applying 0 ng, 37.5 ng, 75 ng, 150 ng and 300 ng (lanes Ø, 1, 2, 3 and 4) of purified His6-tagged functional (A and B) or non functional (C) MexT proteins to 10 fmol of DIG-labelled DNA target containing the intact conserved DNA motif (A and C), or the mutated DNA target with the conserved DNA motif disrupted (B). Arrow indicates the free DNA probes and the band shifts of the MexT–DNA complex.

particular mutation in the helix–turn–helix motif abolished the binding of MexT protein to the target DNA.

MexT activates the expression of target genes across other *Pseudomonas* species

Orthologues of the *mexEF-oprN* operon can be found in all *Pseudomonas* genomes sequenced to date. The *mexT* gene is located immediately upstream and transcribed in the same direction as *mexEF-oprN* in all *P. aeruginosa* strains and *P. stutzeri* A1501 strain but not in other species. For instance, in the genome of *P. fluorescens* PfO-1 *mexT* (Pfl01_2666) and *mexE* (Pfl01_2659) are separated by six open reading frames (ORFs) and in the genome of *P. putida* KT2440 *mexT* (PP_2826) and *mexE* (PP_3425) are located in totally different loci. To date, it has not been shown if the expression of these *mexE* genes is activated by MexT. In order to investigate this, promoter-*lacZ* fusions of Pfl01_2659 and PP_2826 were constructed and transformed into *E. coli* strains containing pME6032-*mexT*. The expression of both *mexE* fusions was highly induced by MexT in *E. coli* (Table 3).

Two newly identified MexT target genes of particular note were PA4881 and PA4623, which encode hypothetical proteins of 113 amino acids and 129 amino acids, respectively (*Pseudomonas* Genome Database). PA4881 and PA4623 actually encode homologous proteins, each containing four tandem repeat motifs of average 23 amino acids (Figure 5A and B). Genes encoding tandem repeat proteins are widely distributed in the genomes of different organisms (26). Indeed, the genomes of various *Pseudomonas* species harbour a number of genes encoding tandem repeat proteins, which share some conserved residues with PA4881 and PA4623 encoded proteins (Figure 5 and Supplementary Figure S2). Among them, several genes contain the conserved MexT-binding motif in their upstream regulatory regions (Figure 5C–F). To address the question if these genes can be induced by MexT, promoter-*lacZ* fusions of two tandem repeat proteins, *P. fluorescens* PfO-1 Pfl01_3748 and *P. putida* KT2440 PP_4858 were constructed and transformed into *E. coli* strains containing pME6032-*mexT*. The expression of both promoter fusions was highly induced by MexT in *E. coli* (Table 3). These data suggest that MexT activates the expression of a novel regulon across *Pseudomonas* species by binding to a

conserved DNA motif in the promoter regions of target genes.

DISCUSSION

Identification of a novel MexT regulon

Previously, MexT was known to be a transcriptional activator of *mexEF-oprN*, an operon encoding a multidrug efflux pump in *P. aeruginosa* (6). Based on our observation that MexT modulated certain virulence factors in a *mexEF-oprN*-independent manner (18), we hypothesized that MexT has a wider function than previously thought and plays a more global regulatory role in *P. aeruginosa*. To test this hypothesis, we aimed to identify and characterize other regulatory targets in *P. aeruginosa*. Here, we present a genomics-based strategy for uncovering novel genes regulated by MexT in *P. aeruginosa* PAO1. The transcriptome profiling allowed the genes regulated by MexT to be divided into those that were indirectly modulated via MexEF-OprN efflux and those that were regulated by MexT independent of this efflux pump. The MexEF-dependent operons identified were all previously shown to be controlled by quorum sensing signalling (27–29). This correlates with the previous report that overexpression of MexEF-OprN *per se* in *nfxC*-type mutant strains caused a reduction of virulence determinants controlled by cell-to-cell signalling, possibly due to decreased intracellular PQS levels resulting from the efflux of either PQS or a PQS precursor (3,15).

Our main interest, however, was in the observation that in addition to *mexE* and *mexS*, 12 genes were highly induced by MexT independent of MexEF-OprN (Table 2). When the upstream regulatory region of these genes was compared, a well-conserved DNA motif was identified (Figure 3). The conserved motif contained the previously described LysR regulator ‘*nod-box*’, ATC-N9-GAT, but also conserved nucleotides immediately 5’ of the GAT and an additional conserved sequence 4 nucleotides 3’ of the GAT, consisting of ACT/CAT. This sequence was highly conserved in most of the promoters but with some changes in *mexE*, PA1744 and PA1970. The motif was confirmed experimentally as a MexT-binding site (Figure 4) and that the GA of the ‘*nod-box*’ was essential for MexT binding under the conditions tested. Interestingly, when a *P. aeruginosa* genome-wide search (<http://bioinfo.hku.hk/GenoList/index>).

A PA4623 *P. aeruginosa* PAO1

>upstream regulatory DNA sequence
 GCACTATTTCAAACGTCGATGAATACTATGGCGCGCTTCGTTTT
 CCAATCGAGTTTCCCGAACAGAGAATGGCCCCACGTAACGAATAC
 CTGGAGCCACTGCCATG

>amino acid sequence
 1 MKRPLILALATLTANAAFAAPRVESIALRGAQQL
 37 IASDGSDRTLRLRKTQAPRVQEERR
 63 VAESGSEQTIDRLHRDMTSREAR
 86 LAENGSEQTVDRHLHKDMGRVDLR
 109 LADNGSERTVDRHLHREMRPPA

C Pf101_3748 *P. fluorescens* Pf0-1

>upstream regulatory DNA sequence
 CAAACCATCATCGGCGTCGATGTTACCATCACGTCATGAAGTT
 CTCTTAAAAAATCCCGGCGTAACATTGGCTCCATACCAACAAA
 CAACACCCCGGAGCACAAAATG

>amino acid sequence
 1 MKRQVILSIALSVLAFAAKPAHTM
 28 IAEGGSDRLIERR
 41 VAEGGSDRLIERR
 54 VAEGGSDRLIERR
 67 VAEGGSDRLIERR
 80 VAEGGSDRLIERR
 93 VAEGGSDRLIERR
 106 VAEGGSDRLIERR
 119 VAEGGSDRLIERR
 132 VAEGGSDRLIERRVA

E PSEEN4907 *P. entomophila* L48

>upstream regulatory DNA sequence
 TCCAGTATCAATCCCGTCGATTGTACCATATCCCGATGAACCTT
 CTTTCTGGCGTTACCGAGCGTAGCATTGGCTCCGTACCCACTTTC
 CCAGCCCCCAGAGGAGACCGAGATG

>amino acid sequence
 1 MKRHLLSLTLVLAANAFALPAEDQHLSEAERSSAATVS
 41 QPLNT
 46 LAEGGAERLQERAGR
 61 LAEGGSERLLERNNR
 76 VAEGGSDRLIQRNDR
 91 VAEGGSDRLIQRNDR
 106 VAEGGSDRLIQRNDR
 121 VAEGGSDRLIQRNDR
 136 VAEGGADRLNERNNR
 151 VAEGGSDRLVELSRVS

B PA4881 *P. aeruginosa* PAO1

>upstream regulatory DNA sequence
 ACATAATCACCCATGTCGATAGACACTATCATCGCACTTCGTTTT
 TAAATCGACTTATCCGAAGTAAATGACTCCAACGCAATCGACA
 ACCCACTCGGAGTACCGTCATG

>amino acid sequence
 1 MKASLILGLALATLTANAA
 20 FAADGSSRTIDRLHKDMSSVELR
 43 IADNGSEQTIDRLHKDMTKAEAR
 66 LADNGSEQTVDRHLHKDMTKAEAR
 89 LADNGSEQTVDRHLHRSMSRFDLRTA

D PFL_0611 *P. fluorescens* Pf-5

>upstream regulatory DNA sequence
 AAACAATCAATTCCGTTGATTATTACCATTTGTGTTTATGAACCTT
 TATATCGATTATAGCGGCGGTAAACATTAGCTCCGTACCCAATTTAT
 GCCCTACCAAGGAGCAACCCATC

>amino acid sequence
 1 MKRQLLLSLALSVAANAFALPAEQATPQVKASHSVFTQ
 41 T
 42 LAEGGSDRLIERNK
 56 VAADGYDRTPQQQS
 70 VAEDGYDRTPQQQN
 84 VAEDGYDRTPQQQN
 98 VAEDGYDRTPQQQN
 112 VAEGGRDRLEEKQ
 125 LVEDGYSRTPQQQT
 139 VAEGGADRLAERHQAAS

F PP_4858 *P. putita* KT2440

>upstream regulatory DNA sequence
 TCCTCTATCAATATGGTCGATTGTACCATACCCGATGAACCTT
 CTCTCTGGCGTTACGCGGCGTAGCATTGGCTTCGTACCCACTTTC
 CAGCCCCGAGAGGAGACCGAAATG

>amino acid sequence
 1 MKRHLLTLTLVLAANAFALPASEQHLTSEARSSAAEIAQ
 41 PLNT
 45 VAEGGSDRLIERSGR
 60 VAEGGSDRLIERSGR
 75 VAEGGSDRLIERSGR
 90 VAEGGSDRLIERSGR
 105 VAEGGSDRLIERSGR
 120 VAEGGSDRLIEQAGR
 135 VAEGGSDRLIEQAGR
 150 VAEGGSDRLVELSRVS

Figure 5. Upstream regulatory DNA sequences and amino acid sequences of PA4623 (A) and PA4881 (B) and tandem repeat proteins in other *Pseudomonas* (C–F). Locus IDs are from the *Pseudomonas* Genome Project (<http://www.pseudomonas.com>). The conserved DNA motif is underlined and the start codon ATG is boxed. The amino acid sequence of each protein is organized showing alignment of tandem repeats with the position number of the first amino acid residue at the beginning of each line.

pl?database=aerulist) was performed to identify genes that carried a similar sequence motif in their upstream regions, a number of genes were identified that were not altered by MexT in the array analysis. This information warrants further investigation and may be valuable in follow on studies of the MexT-binding site itself and of physiological conditions that may influence binding.

Two of the MexT regulated genes identified in the array analysis do not contain the consensus sequence; PA3205, which was induced just 11-fold, but more interestingly, *mexS*, the divergently transcribed gene previously shown to be induced by MexT (6) does not even contain the core LysR ATC-N9-GAT consensus motif. Furthermore, no binding of the MexT protein to the *mexS* promoter region was detected by EMSA under the conditions used for PA4881 promoter binding (data not shown). This was

surprising as, although it is possible that *mexS* is indirectly regulated by MexT, available data suggest that *mexS* should be a direct target of MexT. Firstly, the genomic arrangement of *mexS-mexT* is a typical arrangement of a LysR-type regulator and its target gene, and it is well conserved across *Pseudomonas* species. Secondly, in addition to its induction by MexT in *P. aeruginosa*, *mexS* can be activated by MexT in a heterogeneous *E. coli* background (6). This anomaly warrants further study, not least to investigate the kinetics of MexT binding under different experimental conditions and to investigate the possibility that MexT may also bind to a modified binding site under different conditions. It has been shown that different co-factors can modify the conformation of the same LysR-type regulator protein and therefore the protein could have different targets

according to one or either conformation (13). To date the co-factor(s) of MexT has not been identified but may indeed influence its binding to target genes.

It has also been shown that a LysR-type regulator can be an activator or a repressor for different target genes (13). The transcriptome data revealed that a number of genes were downregulated >2-fold by MexT independent of *mexEF* expression (Supplementary Table S2). Interestingly, one of these was a gene (*pscE*) encoding a type-three secretion protein. Previously, we demonstrated that MexT exerted a negative effect on virulence traits, including type-three secretion, independent of MexEF-OprN, when grown under TTSS inducing conditions (18). Furthermore, a LysR-type regulator YtxR has been shown to have a global regulatory role in *Yersinia enterocolitica*, including direct suppression of the transcriptional expression of TTSS (30). However, none of the downregulated genes contained an ATC-N9-GAT DNA motif and it remains to be clarified whether they are direct or indirect targets of MexT regulation.

Nevertheless, a novel regulon including *mexEF-oprN* directly activated by MexT has been identified and this is an important step towards understanding the global regulatory role of MexT and the physiological role of MexEF-OprN multidrug efflux pump as a component of this regulon in *P. aeruginosa*.

Possible cellular function of the novel MexT regulon

In addition to *mexEF-oprN*, the novel MexT regulon identified in this study contains two additional putative operons (PA2813-PA2812-PA2811 and PA4354-PA4355-*xenB*) (Table 2 and Supplementary Table S2). PA2813-PA2812-PA2811 encodes a probable glutathione *S*-transferase, a probable ATP-binding component of ABC transporter, and a probable permease of ABC-2 transporter, respectively. Glutathione *S*-transferases constitute a large family of enzymes, which catalyze the addition of glutathione to endogenous or xenobiotic, often toxic electrophilic chemicals. In *Pseudomonas* species, multiple glutathione *S*-transferases are specifically involved in mineralization of recalcitrant compounds, while glutathione *S*-transferase was known to be involved in detoxification in eukaryotes (31). The putative operon PA4354-PA4355-*xenB* encodes an ArsR-type transcriptional regulator, a probable major facilitator superfamily (MFS) transporter and a xenobiotic reductase, respectively. The ArsR-type transcriptional regulators are prokaryotic metallo-regulatory transcriptional repressors. They repress or derepress the expression of operons involved in the detoxification of di- and multivalent heavy metal ions in the absence or presence of these toxics (32). The MFS is the largest characterized family of transporters, capable of transporting various substrates, such as sugars, polyols, drugs, neurotransmitters, amino acids, peptides, and inorganic anions (33). The xenobiotic reductase is a member of the Old Yellow Enzyme family (flavoprotein oxidoreductases) and catalyze the degradation of explosive compounds such as nitroglycerin, 2,4,6-trinitrotoluene and hexahydro-1,3,5-trinitro-1,3,5-triazine in *Pseudomonas* species (34–36). From the compilation of these possible

functions, the MexT regulon is likely involved in a broad-spectrum detoxification or nutrient-scavenging in *P. aeruginosa*.

The MexT regulon also contains six genes (PA1744, PA1970, PA2759, PA3229, PA4623 and PA4881) encoding small proteins (70–130 amino acids) of unknown function but with either type I or type II export signal peptides (37). This suggested that MexT may induce a family of cell envelope or secreted proteins. Of particular note are PA4623 and PA4881, two homologue genes encoding tandem repeat proteins (Figure 5A and B). While the function of these proteins is unknown, recently, it was reported that *Ehrlichia chaffeensis*, an obligately intracellular bacterium, exploits the host cell by a secreted, differentially expressed, tandem repeat protein interacting with multiple host proteins associated with cell signalling, transcriptional regulation and vesicle trafficking (38). Although there is no primary sequence similarity between *P. aeruginosa* tandem repeat proteins encoded by PA4881 and PA4623 and this *E. chaffeensis* tandem repeat protein, it is tempting to speculate that these *P. aeruginosa* tandem repeat proteins may be involved in the interaction with host proteins and warrants further investigation. Furthermore, PA4623 and PA4881 as well as PA3229 and *mexEF-oprN* were shown to be induced in *P. aeruginosa*, after a 12-h interaction with primary normal human airway epithelial cells (39). The role of these proteins in host-cell interactions remains to be elucidated but the report that *nfxC*-type mutants were readily recovered from an experimental model of rat pneumonia in the absence of antibiotic selection (40) indicates that there is some advantage to MexT regulon expression in vivo.

This is supported by the fact that many *Pseudomonas* species have a number of genes encoding tandem repeat proteins, which contain the identical conserved MexT binding motif in their upstream regulatory regions (Figure 5) and can be activated by MexT (Table 3). Moreover, the component genes of the MexT regulon are widely distributed on the genome of *P. aeruginosa* PAO1. These facts strongly suggest that the MexT regulon may serve primary functions in *Pseudomonas* species and understanding their function warrants further investigation. This may also provide new insights into the physiological role of MexEF-OprN pump in *Pseudomonas* species as the pump may function in an integrated way with other components in the MexT regulon.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

We are grateful to Dr Max Schobert for providing the *E. coli* strain ST18. We acknowledge Hazel O'Connor and Marlies Mooij for their scientific contributions, Pat Higgins for his technical assistance and Jerry Reen for his comments on the manuscript.

FUNDING

The European Commission (MTKD-CT-2006-042062; O36314); the Science Foundation of Ireland (SFI 07/IN.1/B948; 08/RFP/GEN1295); the Department of Agriculture and Food (DAF RSF 06 321; DAF RSF 06 377); Irish Research Council for Science, Engineering and Technology (IRCSET) (05/EDIV/FP107); the Health Research Board (RP/2006/271; RP/2007/290); the Environmental Protection Agency (EPA 2006-PhD-S-21; 2008-PhD-S-2); and the Natural Science Foundation of China (to Y.-P.W.) (30830005). Funding for open access charge: Science Foundation Ireland Principal Investigator grant: SFI 07IN.1/B948.

Conflict of interest statement. None declared.

REFERENCES

- Poole, K. and Srikumar, R. (2001) Multidrug efflux in *Pseudomonas aeruginosa*: components, mechanisms and clinical significance. *Curr. Top. Med. Chem.*, **1**, 59–71.
- Schweizer, H.P. (2003) Efflux as a mechanism of resistance to antimicrobials in *Pseudomonas aeruginosa* and related bacteria: unanswered questions. *Genet. Mol. Res.*, **2**, 48–62.
- Martinez, J.L., Sánchez, M.B., Martínez-Solano, L., Hernandez, A., Garmendia, L., Fajardo, A. and Alvarez-Ortega, C. (2009) Functional role of bacterial multidrug efflux pumps in microbial natural ecosystems. *FEMS. Microbiol. Rev.*, **33**, 430–449.
- Piddock, L.J. (2006) Multidrug-resistance efflux pumps – not just for resistance. *Nat. Rev. Microbiol.*, **4**, 629–636.
- Martinez, J.L. (2009) The role of natural environments in the evolution of resistance traits in pathogenic bacteria. *Proc. Biol. Sci.*, **276**, 2521–2530.
- Köhler, T., Epp, S.F., Curty, L.K. and Pechere, J.C. (1999) Characterization of MexT, the regulator of the MexE-MexF-OprN multidrug efflux system of *Pseudomonas aeruginosa*. *J. Bacteriol.*, **181**, 6300–6305.
- Köhler, T., Michéa-Hamzehpour, M., Henze, U., Gotoh, N., Curty, L.K. and Pechère, J.C. (1997) Characterization of MexE-MexF-OprN, a positively regulated multidrug efflux system of *Pseudomonas aeruginosa*. *Mol. Microbiol.*, **23**, 345–354.
- Köhler, T., Michéa-Hamzehpour, M., Plesiat, P., Kahr, A.L. and Pechère, J.C. (1997) Differential selection of multidrug efflux systems by quinolones in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.*, **41**, 2540–2543.
- Maseda, H., Saito, K., Nakajima, A. and Nakae, T. (2000) Variation of the *mexT* gene, a regulator of the MexEF-OprN efflux pump expression in wild-type strains of *Pseudomonas aeruginosa*. *FEMS Microbiol. Lett.*, **192**, 107–112.
- Ramsey, M.M. and Whiteley, M. (2004) *Pseudomonas aeruginosa* attachment and biofilm development in dynamic environments. *Mol. Microbiol.*, **53**, 1075–1087.
- Sobel, M.L., Neshat, S. and Poole, K. (2005) Mutations in PA2491 (*mexS*) promote MexT-dependent *mexEF-oprN* expression and multidrug resistance in a clinical strain of *Pseudomonas aeruginosa*. *J. Bacteriol.*, **187**, 1246–1253.
- Savageau, M.A. (1977) Design of molecular control mechanisms and the demand for gene expression. *Proc. Natl Acad. Sci. USA*, **74**, 5647–5651.
- Maddocks, S.E. and Oyston, P.C.F. (2008) Structure and function of the LysR-type transcriptional regulator (LTTR) family proteins. *Microbiology*, **154**, 3609–3623.
- Goethals, K., Van Montagu, M. and Holsters, M. (1992) Conserved motifs in a divergent nod box of *Azorhizobium caulinodans* ORS571 reveal a common structure in promoters regulated by LysR-type proteins. *Proc. Natl Acad. Sci. USA*, **89**, 1646–1650.
- Köhler, T., van Delden, C., Curty, L.K., Hamzehpour, M.M. and Pechère, J.C. (2001) Overexpression of the MexEF-OprN multidrug efflux system affects cell-to-cell signaling in *Pseudomonas aeruginosa*. *J. Bacteriol.*, **183**, 5213–5222.
- Linares, J.F., Lopez, J.A., Camafeita, E., Albar, J.P., Rojo, F. and Martinez, J.L. (2005) Overexpression of the multidrug efflux pumps MexCD-OprJ and MexEF-OprN is associated with a reduction of type III secretion in *Pseudomonas aeruginosa*. *J. Bacteriol.*, **187**, 1384–1391.
- Cosson, P., Zulianello, L., Join-Lambert, O., Faurisson, F., Gebbie, L., Benghezal, M., Van Delden, C., Curty, L.K. and Köhler, T. (2002) *Pseudomonas aeruginosa* virulence analyzed in a *Dictyostelium discoideum* host system. *J. Bacteriol.*, **184**, 3027–3033.
- Tian, Z.-X., Mac Aogáin, M., O'Connor, H.F., Fargier, E., Mooij, M.J., Adams, C., Wang, Y.-P. and O'Gara, F. (2009) MexT modulates virulence determinants in *Pseudomonas aeruginosa* independent of the MexEF-OprN efflux pump. *Microb. Pathog.*, **21**, 440–446.
- Sambrook, J. and Russell, D.W. (2001) *Molecular Cloning: A Laboratory Manual*, 3rd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Hoang, T.T., Karkhoff-Schweizer, R.R., Kutchma, A.J. and Schweizer, H.P. (1998) A broad-host-range Flp-FRT recombination system for site-specific excision of chromosomally-located DNA sequences: application for isolation of unmarked *Pseudomonas aeruginosa* mutants. *Gene*, **212**, 77–86.
- Fisher, C.L. and Pei, G.K. (1997) Modification of a PCR-based site-directed mutagenesis method. *Biotechniques*, **23**, 570–574.
- Thoma, S. and Schobert, M. (2009) An improved *Escherichia coli* donor strain for diparental mating. *FEMS Microbiol. Lett.*, **294**, 127–132.
- Miller, J.H. (1972) *Experiments in Molecular Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Bailey, T.L. and Elkan, C. (1994) Fitting a mixture model by expectation maximization to discover motifs in biopolymers. *Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology*. AAAI Press, Menlo Park, CA, pp. 28–36.
- Finn, R.D., Tate, J., Misty, J., Coghill, P.C., Sammut, J.S., Hotz, H.R., Ceric, G., Forslund, K., Eddy, S.R., Sonnenhammer, E.L. et al. (2008) The Pfam protein families database. *Nucleic Acids Res.*, **36**, D281–D288.
- Katti, M.V., Sami-Subbu, R., Ranjekar, P.K. and Gupta, V.S. (2000) Amino acid repeat patterns in protein sequences: their diversity and structural-functional implications. *Protein Sci.*, **9**, 1203–1209.
- Schuster, M., Lostroh, C.P., Ogi, T. and Greenberg, E.P. (2003) Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. *J. Bacteriol.*, **185**, 2066–2079.
- Wagner, V.E., Bushnell, D., Passador, L., Brooks, A.I. and Iglewski, B.H. (2003) Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: effects of growth phase and environment. *J. Bacteriol.*, **185**, 2080–2095.
- Wolfgang, M.C., Jyot, J., Goodman, A.L., Ramphal, R. and Lory, S. (2004) *Pseudomonas aeruginosa* regulates flagellin expression as part of a global response to airway fluid from cystic fibrosis patients. *Proc. Natl Acad. Sci. USA*, **101**, 6664–6668.
- Alxer-DiPerte, G.L., Hinchliffe, S.J., Wren, B.W. and Darwin, A.J. (2009) YtxR acts as an overriding transcriptional off switch for the *Yersinia enterocolitica* Ysc-Yop type 3 secretion system. *J. Bacteriol.*, **191**, 514–524.
- Vuilleumier, S. and Pagni, M. (2002) The elusive roles of bacterial glutathione S-transferases: new lessons from genomes. *Appl. Microbiol. Biotechnol.*, **58**, 138–146.
- Busenlehner, L.S., Pennella, M.A. and Giedroc, D.P. (2003) The SmtB/ArsR family of metalloregulatory transcriptional repressors: Structural insights into prokaryotic metal resistance. *FEMS Microbiol. Rev.*, **27**, 131–143.
- Flumana, N. and Bibi, E. (2009) Bacterial multidrug transport through the lens of the major facilitator superfamily. *Biochim. Biophys. Acta*, **1794**, 738–747.
- Blehert, D.S., Fox, B.G. and Chambliss, G.H. (1999) Cloning and sequence analysis of two *Pseudomonas* flavoprotein xenobiotic reductases. *J. Bacteriol.*, **181**, 6254–6263.
- Pak, J.W., Knoke, K.L., Noguera, D.R., Fox, B.G. and Chambliss, G.H. (2000) Transformation of 2, 4, 6-trinitrotoluene

- by purified xenobiotic reductase B from *Pseudomonas fluorescens* I-C. *Appl. Environ. Microbiol.*, **66**, 4742–4750.
36. Fuller, M.E., McClay, K., Hawari, J., Paquet, L., Malone, T.E., Fox, B.G. and Steffan, R.J. (2009) Transformation of RDX and other energetic compounds by xenobiotic reductases XenA and XenB. *Appl. Microbiol. Biotechnol.*, **84**, 535–544.
 37. Lewenza, S., Gardy, J.L., Brinkman, F.S. and Hancock, R.E. (2005) Genome-wide identification of *Pseudomonas aeruginosa* exported proteins using a consensus computational strategy combined with a laboratory-based PhoA fusion screen. *Genome Res.*, **15**, 321–329.
 38. Wakeel, A., Kuriakose, J.A. and McBride, J.W. (2009) An *Ehrlichia chaffeensis* tandem repeat protein interacts with multiple host targets involved in cell signalling, transcriptional regulation, and vesicle trafficking. *Infect. Immun.*, **77**, 1734–1745.
 39. Frisk, A., Schurr, J.R., Wang, G., Bertucci, D.C., Marrero, L., Hwang, S.H., Hassett, D.J. and Schurr, M.J. (2004) Transcriptome analysis of *Pseudomonas aeruginosa* after interaction with human airway epithelial cells. *Infect. Immun.*, **72**, 5433–5438.
 40. Join-Lambert, O.F., Michéa-Hamzehpour, M., Köhler, T., Chau, F., Faurisson, F., Dautrey, S., Vissuzaine, C., Carbon, C. and Pechère, J. (2001) Differential selection of multidrug efflux mutants by trovafloxacin and ciprofloxacin in an experimental model of *Pseudomonas aeruginosa* acute pneumonia in rats. *Antimicrob. Agents Chemother.*, **45**, 571–576.
 41. deFlaun, M.F., Tanzer, A.S., McAteer, A.L., Marshall, B. and Levy, S.B. (1990) Development of an adhesion assay and characterization of an adhesion-deficient mutant of *Pseudomonas fluorescens*. *Appl. Environ. Microbiol.*, **56**, 112–119.
 42. Franklin, F.C., Bagdasarian, M., Bagdasarian, M.M. and Timmis, K.N. (1981) Molecular and functional analysis of the TOL plasmid pWWO from *Pseudomonas putida* and cloning of genes for the entire regulated aromatic ring meta cleavage pathway. *Proc. Natl Acad. Sci. USA*, **78**, 7458–7462.
 43. Heeb, S., Itoh, Y., Nishijyo, T., Schnider, U., Keel, C., Wade, J., Walsh, U., O’Gara, F. and Haas, D. (2000) Small, stable shuttle vectors based on the minimal pVS1 replicon for use in gram-negative, plant-associated bacteria. *J. Mol. Plant Microb. Interact.*, **13**, 232–237.
 44. Spaink, H.P., Okker, R.J.H., Wijffelman, C.A., Pees, E. and Lugtenberg, B.J.J. (1987) Promoters in the nodulation region of the *Rhizobium leguminosarum* Sym plasmid pRL1J1. *Plant Mol. Biol.*, **9**, 27–39.