CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Appropriate use of elective coronary angiography in patients with suspected stable coronary artery disease
Authors	Hanrahan, Michael T.;O'Flynn, Anne Marie;McHugh, Sheena M.;Kearney, Peter;Kearney, Patricia M.
Publication date	2019-01-05
Original Citation	Hanrahan, M. T., O'Flynn, A. M., McHugh, S. M., Kearney, P., Kearney, P. M. (2019) 'Appropriate use of elective coronary angiography in patients with suspected stable coronary artery disease', Irish Journal of Medical Science, pp. 1-7. doi:10.1007/s11845-018-1953-4
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1007/s11845-018-1953-4
Rights	© 2019, Royal Academy of Medicine in Ireland. Published by Springer Nature Switzerland AG. This is a post-peer-review, pre-copyedit version of an article published in the Irish Journal of Medical Science. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11845-018-1953-4
Download date	2026-11-15 17:55:11
Item downloaded from	https://hdl.handle.net/10468/7432

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Table 2. Appropriateness Categories					
Indication #	**Indication**	**Total**	**Inappropriate**	**Uncertain**	**Appropriate**
			N (%)	**N (%)**	**N (%)**
		N = 259	**54 (21%)**	**97 (37%)**	**108 (42%)**
	No non-invasive testing: Asymptomatic				
5	Asymptomatic: Low SCORE risk	3	3	-	-
6	Asymptomatic: Intermediate SCORE risk	2	2	-	-
7	Asymptomatic: High SCORE risk	1	-	1	-
	No non-invasive testing: Symptomatic				
9	Symptomatic: Intermediate pretest probability	9	-	9	-
10	Symptomatic: High pretest probability	29	-	-	29
	Non-invasive testing				
11	Low-risk findings (e.g., DTS ≥5)	54	49	5	-
12	Intermediate-risk findings (e.g., DTS 4 to -10)	78	-	78	-
13	High-risk findings (e.g., DTS ≤-11)	10	-	-	10
14	Other high-risk findings (ST-segment elevation, hypotension with exercise, ventricular tachycardia, prolonged ST-segment depression)	67	-	-	67
24	Newly recognized LV systolic dysfunction (i.e., LVEF ≤40%) with an unknown etiology	2	-	1	1
25	Newly recognized LV systolic dysfunction (i.e., LVEF 41% to 49%) with an unknown etiology	2	-	2	0
26	New regional wall motion abnormality with an unknown etiology and normal LV systolic function	1	-	1	0
33	Coronary CTA: Lesion ≥50% non-left main	1	-	0	1