

Title	Deletion of TLX and social isolation impairs exercise-induced neurogenesis in the adolescent hippocampus
Authors	Kozareva, Danka A.;O'Leary, Olivia F.;Cryan, John F.;Nolan, Yvonne M.
Publication date	2017-10-03
Original Citation	Kozareva, D. A., O'Leary, O. F., Cryan, J. F. and Nolan, Y. M. 'Deletion of TLX and social isolation impairs exercise-induced neurogenesis in the adolescent hippocampus', Hippocampus, In Press. doi: 10.1002/hipo.22805
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1002/hipo.22805
Rights	© 2017 John Wiley & Sons, Inc.This is the peer reviewed version of the following article: Kozareva, D. A., O'Leary, O. F., Cryan, J. F. and Nolan, Y. M., Deletion of Tlx and social isolation impairs exercise-induced neurogenesis in the adolescent hippocampus. Hippocampus. Accepted Author Manuscript. which has been published in final form at http://dx.doi.org/10.1002/hipo.22805. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Download date	2025-08-25 19:11:21
Item downloaded from	https://hdl.handle.net/10468/4846



University College Cork, Ireland Coláiste na hOllscoile Corcaigh

Deletion of Tlx and social isolation impairs exercise-induced neurogenesis in the adolescent hippocampus Danka A. Kozareva, Olivia F. O'Leary, John F. Cryan and Yvonne M. Nolan\* 🔎 Department of Anatomy and Neuroscience, University College Cork, Ireland APC Microbiome Institute, University College Cork, Ireland To whom correspondence should be addressed: \*Yvonne M. Nolan, Department of Anatomy and Neuroscience, University College Cork, Ireland

Tel: (353) 21-420 5476

Fax: (353) 21-427 3518

email: y.nolan@ucc.ie

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/hipo.22805

Abstract

Adolescence is a sensitive period of neurodevelopment during which life experiences and the surrounding environment can have profound effects on the brain. Neurogenesis is a neurodevelopmental process of generating functional neurons from neural stem cells. Hippocampal neurogenesis occurs throughout the lifespan and has been shown to play a role in learning, memory and in mood regulation. In adulthood it is influenced by extrinsic environmental factors such as exercise and stress. Intrinsic factors that regulate hippocampal neurogenesis include the orphan nuclear receptor TLX (Nr2e1) which is primarily expressed in the neurogenic niches of the brain. While mechanisms regulating adult hippocampal neurogenesis have been widely studied, less is known on how hippocampal neurogenesis is affected during adolescence. Thus, the aim of this study was to investigate the influence of both TLX and isolation stress on exercise-induced increases in neurogenesis in running and sedentary conditions during adolescence. Single- (i.e. isolation stress) wild type and Nr2e1-/or pair-housed wild type mice were housed in sedentary conditions or allowed free access to running wheels for 3 weeks during the adolescent period. A reduction of neuronal survival was evident in mice lacking TLX, and exercise did not increase hippocampal neurogenesis in these Nr2e1<sup>-/-</sup> mice. This suggests that TLX is necessary for the pro-neurogenic effects of exercise during adolescence. Interestingly, although social isolation during adolescence did not affect hippocampal neurogenesis, it prevented an exercise-induced increase in neurogenesis in the ventral hippocampus. Together these data demonstrate the importance of intrinsic and extrinsic factors in promoting an exercise-induced increase in neurogenesis at this key point in life.

Key words: adult neurogenesis, adolescence, stress, exercise, TLX

John Wiley & Sons

This article is protected by copyright. All rights reserved.

Adolescence is a critical phase of development associated with plasticity-driven organization of neural circuits in the hippocampus, prefrontal cortex and amygdala (Pattwell et al., 2011; Selemon, 2013). It is also a key period for susceptibility to stress and the emergence of neurobiological disorders such as schizophrenia, depression and anxiety (Green and Nolan, 2014; Hueston et al., 2017; O'Connor and Cryan, 2014; Paus et al., 2008). By approximately postnatal day (P) 30 in rodents, dentate gyrus (DG) formation, cerebellar neurogenesis and myelogenesis are completed, and neurogenesis (the birth of new neurons) is restricted to the niches of the brain where the process persists through adulthood – the subgranular zone (SGZ) of the DG of the hippocampus and the subventricular zone (Lemasson et al., 2005; Li et al., 2009).

Hippocampal neurogenesis has been widely studied in the adult brain and is known to be regulated by several extrinsic and intrinsic factors (Aimone et al., 2014; Gregoire et al., 2014). For example, extrinsic factors such as stress and exercise have been shown to decrease or increase adult hippocampal neurogenesis, respectively (Levone et al., 2015; van Praag et al., 1999a). However, comparatively fewer studies have investigated the impact of these extrinsic factors on hippocampal neurogenesis in the adolescent brain (Abel and Rissman, 2013; Kirshenbaum et al., 2014; Wei et al., 2011). In adult rodents, exercise has been shown to enhance learning and memory (Creer et al., 2010; Marlatt et al., 2012; van Praag et al., 1999a), protect against stress-induced depression and anxiety-like behaviours (Duman et al., 2008; Grippo et al., 2014) and protect against cognitive deficits in neurodegenerative disorders (Barbour et al., 2007; Cotman et al., 2007; Ryan and Nolan, 2016). How exercise can facilitate processes as diverse as spatial learning and memory, anxiety and responses to stress is not yet clear. However, accumulating evidence suggests that the hippocampus is functionally segregated along its dorsoventral axis in rodents such that the dorsal

hippocampus (dHi) plays a predominant role in spatial learning and memory while the ventral hippocampus (vHi) plays a predominant role in anxiety and the stress response (Bannerman et al., 2004; Fanselow and Dong, 2010). Similarly, there is an emerging view that neurogenesis might also be similarly functionally segregated along this axis (O'Leary and Cryan, 2014; Tanti and Belzung, 2013). Intrinsic factors that control hippocampal neurogenesis include the orphan nuclear receptor TLX (Nr2e1) which is primarily expressed in the neurogenic niches of the postnatal brain and has been implicated as an important regulator of neural stem cells by maintaining them in their proliferative and non-differentiated state (Roy et al., 2004; Shi et al., 2004). In a spontaneous deletion mouse model, adult TLX knock out (Nr2e1<sup>-/-</sup>) mice display altered neurogenesis and synaptic plasticity, as well as an impairment of dendritic structures in the DG. These mice also present with an aggressive phenotype and display cognitive impairments in hippocampal-dependent tasks (Christie et al., 2006; Young et al., 2002). Interestingly, some of these effects are apparent in adolescence (O'Leary et al., 2016a; O'Leary et al., 2016b).

Taken together, while mechanisms regulating adult hippocampal neurogenesis have been thoroughly investigated, less is known about how neurogenesis is affected during the adolescent period. Moreover, the impact of facilitators and impeders of neurogenesis, such as exercise and stress, on the adolescent brain has yet to be established. This is surprising given that adolescence is a critical period for the maturation of neurons as well as a time during which profound social and physiological change occurs. Thus, the aim of this study was to investigate (1) the impact of exercise during adolescence on hippocampal neurogenesis; (2) the role of the intrinsic factor TLX on exercise-induced changes in hippocampal neurogenesis and (3) whether social isolation stress influences exercise-induced changes in hippocampal neurogenesis. Given the growing evidence of segregated effects on neurogenesis along the

dorsoventral axis of the hippocampus, we examined the impact of exercise, TLX, and social isolation on neurogenesis in the dorsal versus ventral hippocampus.

This animal study was conducted in strict compliance with the European Directive 2010/63/EU, and under an authorization issued by the Health Products Regulatory Authority Ireland and approved by the Animal Ethics Committee of University College Cork (UCC). Breeding pairs of Nr2e1<sup>-/-</sup> mice exhibiting a spontaneous deletion of TLX were kindly provided by Prof. Elizabeth Simpson (University of British Colombia) and were generated as previously described (Wong et al., 2010). Male Nr2e1<sup>-/-</sup> mice and wild type (WT) littermate controls on a BL6129 background were singly-housed upon weaning and given ad libitum access to food and water under a 12-12 h light/dark cycle. At 4 weeks of age (P28), mice received 4 x bromodeoxyuridine (BrdU; 75mg/kg, i.p., Sigma Cat# B5002) injections at 2hour intervals to label newly-born cells. Half of the animals from each genotype were then housed with free access to a running wheel for 3 weeks (Med Associates Inc Cat# ENV-044), thus there were 4 experimental groups (WT sedentary, WT running, Nr2e1<sup>-/-</sup> sedentary; Nr2e1<sup>-/-</sup> running; n = 5-6 per group; Fig. 1). Male mice were collected from 11 litters in total. For welfare reasons  $Nr2e1^{-/-}$  mice have to be singly housed due to their aggressive phenotype (Young et al., 2002). Thus, their corresponding wild type littermates were also singly housed. Since single housing is a social isolation stressor, we sought to examine whether this chronic stress influenced the effect of exercise on hippocampal neurogenesis in WT mice (Fig. 1).

Three weeks following BrdU administration and initiation of exercise, mice (P49) were anesthetized with Euthasol (1.0 mL/kg, i.p.) and transcardially perfused with phosphate buffered saline (PBS) solution followed by 4.0% paraformaldehyde (PFA). Brains were postfixed overnight, cryoprotected in 30% sucrose and subsequently flash frozen. Coronal

John Wiley & Sons

This article is protected by copyright. All rights reserved.

sections (40 µm) through the entire hippocampus were collected onto slides in a 1:6 series and surviving cells and surviving newly born neurons were identified using immunohistochemistry for BrdU alone, and colocalization of BrdU and the neuronal marker NeuN, respectively. Briefly, sections were incubated in HCl (2M; 37°C, 45 mins), renatured in 0.1M sodium tetraborate (pH 8.5) and then blocked in 3% normal donkey serum (NDS; Sigma Cat# D9663). Sections were incubated with rat anti-BrdU antibody (Abcam Cat# AB6326; 1:250; overnight at 4°C), followed by AlexaFluor594 donkey anti-rat (Abcam Cat# ab150156; 1:500; 2 hours at room temperature) and mouse anti-NeuN (Millipore Cat# MAB377; 1:100; overnight at 4°C). Sections were then incubated in AlexaFluor488 donkey anti-mouse (Abcam Cat# ab150105 1:500; 2 hours at room temperature). To assess cell death, immunohistochemistry for the apoptosis marker caspase-3 was performed. Briefly, sections were incubated in  $H_2O_2$  (in 1% methanol, 40 mins at room temperature) and then blocked in 10% normal goat serum (NGS; Sigma Cat# G9023). Sections were incubated with rabbit anti-active caspase-3 antibody (Promega Cat# G7481; 1:250; overnight at 4°C), followed by sequential incubations with the streptavidin-biotin immunoenzymatic antigen detection system (Abcam, Cat# ab64261). Images were obtained using an Olympus BX533 upright microscope coupled to an Olympus DP72 camera and Olympus FV1000 scanning laser confocal system (BioSciences Imaging Centre, UCC). Immunofluorescent Z-stack images with a 4.4 µm step size were collected using a 10x objective, while DAB staining was analysed at 20x magnification with bright field. Systematic random sampling was employed through the whole DG by counting the cells on both hemispheres of each section in 1:6 series (240 µm apart). Cell quantification was performed using the image processing software package, ImageJ v 1.43m. All cell numbers are expressed as an average per section. The dorsal DG was defined as AP: - 0.94 to AP: - 2.30 and the ventral DG as AP: - 2.46 to AP: -3.80 as described previously (O'Leary et al., 2012; Paizanis et al., 2010).

Exercise significantly increased the number of surviving new cells in the whole DG of WT mice (exercise effect:  $F_{1,19} = 7.915$ , p = 0.011), but not in Nr2e1<sup>-/-</sup> mice (exercise x genotype) interaction:  $F_{1, 19} = 6.747$ , p = 0.018; Fig. 2A). Furthermore, both sedentary and running Nr2e1<sup>-/-</sup> mice exhibited a reduction in the number of surviving new cells compared to their WT littermates (genotype effect:  $F_{1, 19} = 244.002$ , p < 0.001; Fig. 2A). These changes in cell survival were driven by an exercise-induced increase in the number of surviving new neurons (neurogenesis) in the DG of WT mice (exercise effect:  $F_{1,19} = 6.024$ , p = 0.024), but not in Nr2e1<sup>-/-</sup> mice (exercise x genotype interaction:  $F_{1,19} = 5.704$ , p = 0.027; Fig. 2B). Additionally, sedentary Nr2e1<sup>-/-</sup> mice also exhibited reduced hippocampal neurogenesis compared to their WT littermates (genotype effect:  $F_{1,19} = 130.096$ , p < 0.001; Fig. 2B). Upon subdivision of the DG into dorsal (dDG) and ventral (vDG) regions, this genotype effect was apparent in both subregions for both new cell (BrdU+) and new neuron (BrdU+NeuN+) survival (BrdU+: dDG: F<sub>1,19</sub> = 98.723, p < 0.001; vDG: F<sub>1,19</sub> = 92.748, p < 0.001; Fig 2A; BrdU+NeuN+: dDG:  $F_{1.19} = 69.635$ , p < 0.001; vDG:  $F_{1.19} = 88.056$ , p < 0.001; Fig. 2B). However, the exercise-induced increase in new cell survival and neurogenesis was only apparent in the dDG (exercise x genotype interaction: BrdU+:  $F_{1,19} =$ 4.078, p = 0.058; **BrdU+NeuN+:**  $F_{1,19} = 4.502$ ; p = 0.047) but not in the vDG of WT mice (Fig. 2). No difference was observed in the percentage of surviving new cells that adopted a neuronal fate in the DG of WT animals; almost all BrdU+ cells were immunopositive for NeuN and there was no effect of exercise on cells adopting a neuronal fate. In the DG of Nr2e1-/- mice, however, a significantly smaller percentage of surviving cells matured into neurons in both sedentary and exercise conditions (genotype effect: F1,19 = 31.530, p <

0.001; % BrdU+NeuN+ cells/BrdU+ cells: WT sedentary: M = 70.156, SD = 4.065; WT

running: M = 74.398, SD = 9.338; Nr2e1-/- sedentary: M = 53.839, SD = 9.130; Nr2e1-/running: M = 51.151, SD = 9.257).

Since exercise increased neurogenesis in the dorsal but not ventral DG of WT mice, we decided to investigate whether the stress of this social isolation during adolescence could explain the lack of effect of exercise-induced increases in neurogenesis in the vDG. Thus we compared the effect of running on cell survival and hippocampal neurogenesis in singlehoused compared to pair-housed WT mice (Fig. 3). Two way ANOVA revealed that exercise significantly increased cell survival in paired- but not single-housed mice in the whole DG, and this effect persisted in both dDG and vDG (exercise x stress interaction: total DG: F<sub>1,17</sub> = 7.420, p = 0.014; dDG: F<sub>1,17</sub> = 5.911, p = 0.026; vDG: F<sub>1,17</sub> = 5.825, p = 0.027; Fig. 3A). Interestingly, exercise significantly increased neurogenesis of both paired and single-housed mice who had access to a running wheel for three weeks compared to their sedentary counterparts (exercise effect: DG:  $F_{1,17} = 34.054$ ; p < 0.001). However, the exercise-induced increase in neurogenesis was significantly lower in single-housed animals than in pair-housed animals (exercise x stress interaction: DG:  $F_{1,17} = 6.866$ ; p = 0.018). Moreover, upon analysis of the dorsal and ventral regions of the DG, we found that although exercise increased neurogenesis in the dDG in both pair-housed and single-housed mice (dDG:  $F_{1,17} = 29.350$ ; p < 0.001; Fig. 3B), this effect was attenuated by the stress of single housing (exercise x stress interaction dDG:  $F_{1,17} = 5.239$ , p = 0.035). Additionally, the exercise-induced increases in neurogenesis observed in the vDG of pair-housed mice (exercise effect: vDG:  $F_{1,17} = 15.421$ ; p = 0.001) was prevented in singly housed mice (exercise x stress interaction: vDG:  $F_{1,17} = 6.933$ , p = 0.017; Fig. 3B).

There was a trend towards a reduction of caspase-3+ cells in the vDG of Nr2e1<sup>-/-</sup> mice with access to a running wheel, compared to their sedentary and WT running counterparts (exercise x genotype interaction:  $F_{1,12} = 3.8$ , p = 0.077; **Fig. 4A**). There was a significant reduction in the number of apoptotic cells in the vDG but not dDG or whole DG in pair-housed mice with access to a running wheel compared to their single-housed or pair-housed sedentary littermates (exercise x stress interaction: vDG:  $F_{1,12} = 5.770$ , p < 0.05; **Fig. 4B**).

Our results indicate that the well-known pro-neurogenic effect of exercise observed during adulthood (van Praag et al., 1999a; van Praag et al., 1999b) also occurs in adolescent male mice. This finding corroborates the effects of exercise on cognitive function observed in adolescent rats (Hopkins et al., 2011). Mice exposed to exercise during adolescence have increased hippocampal levels of pro-neurogenic brain-derived neurotrophic factor (BDNF) (Gallego et al., 2015) and enhanced expression of synaptic plasticity genes (Abel and Rissman, 2013). Exercise has also been shown to rescue alcohol-induced deficits in cell proliferation in adolescent rats (Helfer et al., 2009). Interestingly, a physical skills training task has recently been reported to increase cell survival in the DG of adolescent rats (DiFeo and Shors, 2017). However, training on this type of tasks involves physical exercise and learning, both of which increase neurogenesis and so a definitive conclusion on the effects of exercise alone during adolescence on hippocampal neurogenesis cannot be determined from this study. The present report is the first to demonstrate that voluntary running increases the survival of newly born cells and neurons in the DG of adolescent mice.

In the absence of TLX, a key regulator of adult neurogenesis, we found no pro-neurogenic effect of exercise. It has been previously shown that deletion of TLX leads to significant reduction in adult neurogenesis, synaptic plasticity and impaired dendritic structure in the DG

of adult mice (Christie et al., 2006). Similarly, we show reduced survival of newborn cells and neurons in the DG of sedentary Nr2e1<sup>-/-</sup> adolescent mice, an effect which was not mitigated by exercise. This positions TLX as a regulator of exercise-induced increases in neurogenesis during adolescence. Further studies will determine whether TLX mitigates the pro-neurogenic effects of exercise in adulthood and indeed throughout the lifespan. We have previously shown that a lack of TLX is associated with impairments in hippocampal-related cognitive and anxiety behaviours during adolescence. Specifically, adolescent but not adult Nr2e1<sup>4<sup>-</sup></sup> mice showed deficits in spatial working memory, contextual fear conditioning and cued fear conditioning (O'Leary et al., 2016a). The fact that a lack of TLX expression from birth through adulthood did not persistently induce the same impairments during adolescence and adulthood may point towards compensatory mechanisms occurring past the adolescent period, which ameliorate to some degree the deficits caused by deletion of TLX, independent of neurogenesis. It is worth noting that in contrast to our data with TLX deficiency, ablation of neurogenesis through irradiation, another model of reduced hippocampal neurogenesis, has been shown to be sensitive to the rescue effects of exercise (Clark et al., 2008; Naylor et al., 2008). However, the effects of adolescent hippocampal irradiation on neuronal survival remain unknown.

We observed an exercise-induced increase in neurogenesis, but not cell survival, in the DG of singly-housed adolescent mice which is consistent with previous reports using adult mice (Dostes et al., 2016; Gregoire et al., 2014; Kannangara et al., 2009). It is worth noting however that single housing has been shown to blunt cell proliferation in running rats (Leasure and Decker, 2009; Stranahan et al., 2006) but these differences have not been reconciled in adolescent rodents. To date, the effect of exercise and housing conditions on neurogenesis across the septo-temporal axis of the hippocampus has not been examined. This

#### John Wiley & Sons

This article is protected by copyright. All rights reserved.

is interesting in light of the emerging view that the ventral hippocampus may be the predominant sub-region involved in stress responses (Bannerman et al., 2004; O'Leary and Cryan, 2014; Tanti and Belzung, 2013). Hence, the conflict in findings on hippocampal neurogenesis from group-housed and single-housed rodents may be a function of the differential effects of exercise on the dorsal and ventral hippocampus. Our results demonstrate that social isolation prevents the pro-neurogenic effects of exercise in the vDG but not the dDG. Specifically, exercise increased the survival of new neurons in the dDG of both single- and pair-housed mice, but only in the vDG of pair-housed mice. In addition, the exercise-induced increase in neurogenesis in the dDG of both single and pair-housed mice was significantly attenuated by single housing. Importantly, the studies to date report on exercise-induced increases in neurogenesis in the whole DG, which may explain why studies employing single-housed mice consistently replicate the pro-neurogenic effects of exercise (Mustroph et al., 2012) observed in group-housed mice (van Praag et al., 1999a). Here we show that social isolation during adolescence acts as differential regulator of exercise across the distinct anatomical regions of the DG and propose that neurogenesis in both the dDG and vDG should be taken into consideration when investigating the role of hippocampal neurogenesis in exercise and stress-induced changes in behaviour. Nonetheless, the effects reported here may be specific to the adolescent period. Thus an examination of the effect of social isolation stress during other time periods of the lifespan on any potential exerciseinduced changes in neurogenesis in the subregions of the DG is warranted in future studies.

The mechanisms underlying the differential effect of social isolation stress on exerciseinduced changes in neurogenesis in the dDG and vDG during adolescence remain unclear. It has been reported that unpredictable chronic mild stress in adult mice preferentially decreased the survival of new neurons in the vDG (Elizalde et al., 2010; Tanti et al., 2012), supporting

This article is protected by copyright. All rights reserved.

the view that neurogenesis in the ventral pole of the DG may be more susceptible to the effects of stress. Interestingly, environmental enrichment, which includes exercise, promoted neurogenesis only in the dorsal hippocampus (Tanti et al., 2012). The effects of stress on neurogenesis are thought to be mediated by the glucocorticoid (GR) and mineralocorticoid (MR) receptors (Saaltink and Vreugdenhil, 2014), which are highly expressed in the hippocampus of rodents (Montaron et al., 2003). However, there are some preliminary and inconclusive findings regarding the difference in expression levels of the GR receptor in the dDG and vDG (Lin et al., 2012; Robertson et al., 2005), while MR receptor expression has been shown to be more concentrated in the vDG, at least in the rat brain (Robertson et al., 2005). Notwithstanding, both receptors have been shown to have a distinct activation pattern across the septo-temporal axis of the DG in response to acute stress (Caudal et al., 2014; Dorey et al., 2012). Moreover, a recent report has demonstrated that exercise increased GR expression in the hippocampus in single-housed but not pair-housed adult mice (Pan-Vazquez et al., 2015). Whether adolescent social isolation stress can affect the pro-neurogenic effect of exercise through the differential expression and activation of GR and MR in the vDG remains to be investigated. Another potential vDG-mediated mechanism underlying the attenuation of exercise-induced increase in neurogenesis by stress is through changes in the pro-neurogenic plasticity molecule BDNF (Berchtold et al., 2005; Chen and Russo-Neustadt, 2009; Ploughman et al., 2007; Tang et al., 2008). This is due to the fact that a stressful spatial navigation task has previously been shown to differentially affect the expression of BDNF in the dorsal (increased expression) and ventral (decreased expression) subregions of the DG (Hawley et al., 2012). The N-methyl-D-aspartate (NMDA) receptor activation pathway has also been implicated in both stress- and exercise-related changes in hippocampal neurogenesis. Specifically, exercise induced a robust increase in the activation of NMDA receptor albeit in cortical mouse tissue (Dietrich et al., 2005) and NMDA receptors have been

reported to operate downstream of the stress hormone, corticosterone to regulate hippocampal neurogenesis (Cameron et al., 1998). Given that there is a lower density of binding sites for NMDA receptors in the vDG compared to dDG (Pandis et al., 2006), it could be speculated that the two stimuli (isolation stress and exercise) compete for activation of the same pathway. Finally, isolation stress and exercise may also differentially impact upon cell death in the dorsal and ventral DG. In the current study, we show a significant decrease in the number of apoptotic cells in the vDG of pair housed mice with access to a running wheel. This may reflect a protective mechanism of exercise against cell death, which is attenuated by isolation-induced stress. However, it is important to note that the apoptosis measured here accounted for all cells in the vDG, hence we cannot conclude that the effects of isolation stress and exercise on apoptosis are specific to newly generated neurons.

It is surprising that we found no differences in the survival of newborn neurons of either area of the DG in single- and pair- housed sedentary adolescent mice. Studies conducted during adulthood have reported that social isolation results in anxiety- and depression-like behaviours in mice along with a reduction in levels of neuroplasticity genes (Berry et al., 2012; Ieraci et al., 2016). Social isolation during adolescence in non-human primates (marmosets) also impaired hippocampal neurogenesis in time-dependent manner (Cinini et al., 2014). Moreover, social isolation during adulthood has been shown to delay the proneurogenic effects of exercise in rats (Stranahan et al., 2006). Interestingly, evidence from the Pereira lab suggests that an enriched environment is necessary to promote neurogenesis in single- housed adult mice (Monteiro et al., 2014). It is possible that in the absence of other external stimuli, social condition does not affect neurogenesis during the adolescent period, possibly due to the high basal rate of neurogenesis that occurs during adolescence compared to adulthood (He and Crews, 2007). Alternatively, the mouse strain used in the current study

(generated on a BL6129 background) may have been a confounding factor by potentially limiting our ability to detect any downregulation of new neurons. In several independent studies examining the role of genetic influence on the baseline level of hippocampal neurogenesis, the B6129SF1 and 129Sv were among the strains showing the lowest levels of newborn neurons (Clark et al., 2011; Kempermann et al., 1997; Merritt and Rhodes, 2015).

In conclusion, our results demonstrate that social isolation stress during adolescence attenuates an exercise-induced increase in neurogenesis. We show that this effect is most pronounced in the ventral hippocampus, a brain sub-region which plays a predominant role in anxiety and in regulating the stress response. Adolescence is a critical period for susceptibility to stress-related disorders as well as a time during which remodeling of hippocampal connectivity, including neurogenesis occurs. Thus the impact of stress during adolescence on hippocampal neurogenesis and associated behaviours may be particularly potent. We also show that TLX is necessary for the pro-neurogenic effects of exercise during adolescence and have previously shown that the role of TLX in anxiety-related behaviours is most apparent during adolescence. TLX is thus an important intrinsic regulator of exerciseinduced changes in neurogenesis and may be a key target in understanding the interaction between positive and negative modifiable lifestyle factors such as stress and exercise on hippocampal neurogenesis and associated behaviours during adolescence.

### Acknowledgements

This work was funded by Science Foundation Ireland (SFI) under grant number SFI/IA/1537. We thank Professor Elizabeth Simpson, University of British Columbia, Vancouver, Canada for kindly providing us with Nr2e1 mice.

# References

Abel JL, Rissman EF. 2013. Running-induced epigenetic and gene expression changes in the adolescent brain. International journal of developmental neuroscience : the official journal of the International Society for Developmental Neuroscience 31(6):382-90.

Aimone JB, Li Y, Lee SW, Clemenson GD, Deng W, Gage FH. 2014. Regulation and function of adult neurogenesis: from genes to cognition. Physiological reviews 94(4):991-1026.

Bannerman DM, Rawlins JN, McHugh SB, Deacon RM, Yee BK, Bast T, Zhang WN, Pothuizen HH, Feldon J. 2004. Regional dissociations within the hippocampusmemory and anxiety. Neuroscience and biobehavioral reviews 28(3):273-83.

Barbour KA, Edenfield TM, Blumenthal JA. 2007. Exercise as a treatment for depression and other psychiatric disorders: a review. Journal of cardiopulmonary rehabilitation and prevention 27(6):359-67.

Berchtold NC, Chinn G, Chou M, Kesslak JP, Cotman CW. 2005. Exercise primes a molecular memory for brain-derived neurotrophic factor protein induction in the rat hippocampus. Neuroscience 133(3):853-61.

Berry A, Bellisario V, Capoccia S, Tirassa P, Calza A, Alleva E, Cirulli F. 2012. Social deprivation stress is a triggering factor for the emergence of anxiety- and depression-like behaviours and leads to reduced brain BDNF levels in C57BL/6J mice. Psychoneuroendocrinology 37(6):762-72.

Cameron HA, Tanapat P, Gould E. 1998. Adrenal steroids and N-methyl-D-aspartate receptor activation regulate neurogenesis in the dentate gyrus of adult rats through a common pathway. Neuroscience 82(2):349-54.

Caudal D, Jay TM, Godsil BP. 2014. Behavioral stress induces regionally-distinct shifts of brain mineralocorticoid and glucocorticoid receptor levels. Front Behav Neurosci 8:19.

Chen MJ, Russo-Neustadt AA. 2009. Running exercise-induced up-regulation of hippocampal brain-derived neurotrophic factor is CREB-dependent. Hippocampus 19(10):962-72.

Christie BR, Li AM, Redila VA, Booth H, Wong BK, Eadie BD, Ernst C, Simpson EM. 2006. Deletion of the nuclear receptor Nr2e1 impairs synaptic plasticity and dendritic structure in the mouse dentate gyrus. Neuroscience 137(3):1031-7. Cinini SM, Barnabe GF, Galvao-Coelho N, de Medeiros MA, Perez-Mendes P, Sousa MB, Covolan L, Mello LE. 2014. Social isolation disrupts hippocampal neurogenesis in young non-human primates. Front Neurosci 8:45.

- Clark PJ, Brzezinska WJ, Thomas MW, Ryzhenko NA, Toshkov SA, Rhodes JS. 2008. Intact neurogenesis is required for benefits of exercise on spatial memory but not motor performance or contextual fear conditioning in C57BL/6J mice. Neuroscience 155(4):1048-58.
- Clark PJ, Kohman RA, Miller DS, Bhattacharya TK, Brzezinska WJ, Rhodes JS. 2011. Genetic influences on exercise-induced adult hippocampal neurogenesis across 12 divergent mouse strains. Genes Brain Behav 10(3):345-53.
- Cotman CW, Berchtold NC, Christie LA. 2007. Exercise builds brain health: key roles of growth factor cascades and inflammation. Trends in neurosciences 30(9):464-72.
- Creer DJ, Romberg C, Saksida LM, van Praag H, Bussey TJ. 2010. Running enhances spatial pattern separation in mice. Proceedings of the National Academy of Sciences of the United States of America 107(5):2367-72.
- Dietrich MO, Mantese CE, Porciuncula LO, Ghisleni G, Vinade L, Souza DO, Portela LV. 2005. Exercise affects glutamate receptors in postsynaptic densities from cortical mice brain. Brain Res 1065(1-2):20-5.
- DiFeo G, Shors TJ. 2017. Mental and physical skill training increases neurogenesis via cell survival in the adolescent hippocampus. Brain research 1654(Pt B):95-101.
- Dorey R, Pierard C, Chauveau F, David V, Beracochea D. 2012. Stress-induced memory retrieval impairments: different time-course involvement of corticosterone and glucocorticoid receptors in dorsal and ventral hippocampus.
  - Neuropsychopharmacology 37(13):2870-80.
- Dostes S, Dubreucq S, Ladeveze E, Marsicano G, Abrous DN, Chaouloff F, Koehl M. 2016. Running per se stimulates the dendritic arbor of newborn dentate granule cells in mouse hippocampus in a duration-dependent manner. Hippocampus 26(3):282-8.
- Duman CH, Schlesinger L, Russell DS, Duman RS. 2008. Voluntary exercise produces antidepressant and anxiolytic behavioral effects in mice. Brain research 1199:148-58.
- Elizalde N, Garcia-Garcia AL, Totterdell S, Gendive N, Venzala E, Ramirez MJ, Del Rio J, Tordera RM. 2010. Sustained stress-induced changes in mice as a model for chronic depression. Psychopharmacology (Berl) 210(3):393-406.
- Fanselow MS, Dong HW. 2010. Are the dorsal and ventral hippocampus functionally distinct structures? Neuron 65(1):7-19.

- Gallego X, Cox RJ, Funk E, Foster RA, Ehringer MA. 2015. Voluntary exercise decreases ethanol preference and consumption in C57BL/6 adolescent mice: sex differences and hippocampal BDNF expression. Physiology & behavior 138:28-36.
- Green HF, Nolan YM. 2014. Inflammation and the developing brain: consequences for hippocampal neurogenesis and behavior. Neuroscience and biobehavioral reviews 40:20-34.
- Gregoire CA, Bonenfant D, Le Nguyen A, Aumont A, Fernandes KJ. 2014. Untangling the influences of voluntary running, environmental complexity, social housing and stress on adult hippocampal neurogenesis. PloS one 9(1):e86237.
- Grippo AJ, Ihm E, Wardwell J, McNeal N, Scotti MA, Moenk DA, Chandler DL, LaRocca MA, Preihs K. 2014. The effects of environmental enrichment on depressive and anxiety-relevant behaviors in socially isolated prairie voles. Psychosomatic medicine 76(4):277-84.
- Hawley DF, Morch K, Christie BR, Leasure JL. 2012. Differential response of hippocampal subregions to stress and learning. PLoS One 7(12):e53126.
- He J, Crews FT. 2007. Neurogenesis decreases during brain maturation from adolescence to adulthood. Pharmacology, biochemistry, and behavior 86(2):327-33.
- Helfer JL, Goodlett CR, Greenough WT, Klintsova AY. 2009. The effects of exercise on adolescent hippocampal neurogenesis in a rat model of binge alcohol exposure during the brain growth spurt. Brain research 1294:1-11.
- Hopkins ME, Nitecki R, Bucci DJ. 2011. Physical exercise during adolescence versus adulthood: differential effects on object recognition memory and brain-derived neurotrophic factor levels. Neuroscience 194:84-94.
- Hueston CM, Cryan JF, Nolan YM. 2017. Stress and adolescent hippocampal neurogenesis: diet and exercise as cognitive modulators. Transl Psychiatry 7(4):e1081.
- Ieraci A, Mallei A, Popoli M. 2016. Social Isolation Stress Induces Anxious-Depressive-Like Behavior and Alterations of Neuroplasticity-Related Genes in Adult Male Mice. Neural plasticity 2016:6212983.
- Kannangara TS, Webber A, Gil-Mohapel J, Christie BR. 2009. Stress differentially regulates the effects of voluntary exercise on cell proliferation in the dentate gyrus of mice. Hippocampus 19(10):889-97.
- Kempermann G, Kuhn HG, Gage FH. 1997. Genetic influence on neurogenesis in the dentate gyrus of adult mice. Proc Natl Acad Sci U S A 94(19):10409-14.

Kirshenbaum GS, Lieberman SR, Briner TJ, Leonardo ED, Dranovsky A. 2014. Adolescent but not adult-born neurons are critical for susceptibility to chronic social defeat. Frontiers in behavioral neuroscience 8:289.

Leasure JL, Decker L. 2009. Social isolation prevents exercise-induced proliferation of hippocampal progenitor cells in female rats. Hippocampus 19(10):907-12.

- Lemasson M, Saghatelyan A, Olivo-Marin JC, Lledo PM. 2005. Neonatal and adult neurogenesis provide two distinct populations of newborn neurons to the mouse olfactory bulb. The Journal of neuroscience : the official journal of the Society for Neuroscience 25(29):6816-25.
- Levone BR, Cryan JF, O'Leary OF. 2015. Role of adult hippocampal neurogenesis in stress resilience. Neurobiology of stress 1:147-55.
- Li Y, Mu Y, Gage FH. 2009. Development of neural circuits in the adult hippocampus. Current topics in developmental biology 87:149-74.
- Lin YL, Lin SY, Wang S. 2012. Prenatal lipopolysaccharide exposure increases anxiety-like behaviors and enhances stress-induced corticosterone responses in adult rats. Brain Behav Immun 26(3):459-68.
- Marlatt MW, Potter MC, Lucassen PJ, van Praag H. 2012. Running throughout middle-age improves memory function, hippocampal neurogenesis, and BDNF levels in female C57BL/6J mice. Developmental neurobiology 72(6):943-52.
- Merritt JR, Rhodes JS. 2015. Mouse genetic differences in voluntary wheel running, adult hippocampal neurogenesis and learning on the multi-strain-adapted plus water maze. Behav Brain Res 280:62-71.

Montaron MF, Piazza PV, Aurousseau C, Urani A, Le Moal M, Abrous DN. 2003.

Implication of corticosteroid receptors in the regulation of hippocampal structural plasticity. Eur J Neurosci 18(11):3105-11.

- Monteiro BM, Moreira FA, Massensini AR, Moraes MF, Pereira GS. 2014. Enriched environment increases neurogenesis and improves social memory persistence in socially isolated adult mice. Hippocampus 24(2):239-48.
- Mustroph ML, Chen S, Desai SC, Cay EB, DeYoung EK, Rhodes JS. 2012. Aerobic exercise is the critical variable in an enriched environment that increases hippocampal neurogenesis and water maze learning in male C57BL/6J mice. Neuroscience 219:62-71.
- Naylor AS, Bull C, Nilsson MK, Zhu C, Bjork-Eriksson T, Eriksson PS, Blomgren K, Kuhn HG. 2008. Voluntary running rescues adult hippocampal neurogenesis after

Page 19 of 27

### Hippocampus

- irradiation of the young mouse brain. Proceedings of the National Academy of Sciences of the United States of America 105(38):14632-7.
- O'Connor RM, Cryan JF. 2014. Adolescent brain vulnerability and psychopathology through the generations: role of diet and dopamine. Biological psychiatry 75(1):4-6.
- O'Leary JD, Kozareva DA, Hueston CM, O'Leary OF, Cryan JF, Nolan YM. 2016a. The nuclear receptor Tlx regulates motor, cognitive and anxiety-related behaviours during adolescence and adulthood. Behavioural Brain Research 306:36-47.
- O'Leary JD, O'Leary OF, Cryan JF, Nolan YM. 2016b. Regulation of behaviour by the nuclear receptor TLX. Genes, brain, and behavior.
- O'Leary OF, Cryan JF. 2014. A ventral view on antidepressant action: roles for adult hippocampal neurogenesis along the dorsoventral axis. Trends in pharmacological sciences 35(12):675-87.
- O'Leary OF, O'Connor RM, Cryan JF. 2012. Lithium-induced effects on adult hippocampal neurogenesis are topographically segregated along the dorso-ventral axis of stressed mice. Neuropharmacology 62(1):247-55.
- Paizanis E, Renoir T, Lelievre V, Saurini F, Melfort M, Gabriel C, Barden N, Mocaer E, Hamon M, Lanfumey L. 2010. Behavioural and neuroplastic effects of the newgeneration antidepressant agomelatine compared to fluoxetine in glucocorticoid receptor-impaired mice. The international journal of neuropsychopharmacology 13(6):759-74.
- Pan-Vazquez A, Rye N, Ameri M, McSparron B, Smallwood G, Bickerdyke J, Rathbone A, Dajas-Bailador F, Toledo-Rodriguez M. 2015. Impact of voluntary exercise and housing conditions on hippocampal glucocorticoid receptor, miR-124 and anxiety.
   Mol Brain 8:40.
- Pandis C, Sotiriou E, Kouvaras E, Asprodini E, Papatheodoropoulos C, Angelatou F. 2006. Differential expression of NMDA and AMPA receptor subunits in rat dorsal and ventral hippocampus. Neuroscience 140(1):163-75.
- Pattwell SS, Bath KG, Casey BJ, Ninan I, Lee FS. 2011. Selective early-acquired fear memories undergo temporary suppression during adolescence. Proceedings of the National Academy of Sciences of the United States of America 108(3):1182-7.
- Paus T, Keshavan M, Giedd JN. 2008. Why do many psychiatric disorders emerge during adolescence? Nature reviews. Neuroscience 9(12):947-57.

Ploughman M, Granter-Button S, Chernenko G, Attwood Z, Tucker BA, Mearow KM, Corbett D. 2007. Exercise intensity influences the temporal profile of growth factors involved in neuronal plasticity following focal ischemia. Brain Res 1150:207-16.

- Robertson DA, Beattie JE, Reid IC, Balfour DJ. 2005. Regulation of corticosteroid receptors in the rat brain: the role of serotonin and stress. Eur J Neurosci 21(6):1511-20.
- Roy K, Kuznicki K, Wu Q, Sun Z, Bock D, Schutz G, Vranich N, Monaghan AP. 2004. The Tlx gene regulates the timing of neurogenesis in the cortex. The Journal of neuroscience : the official journal of the Society for Neuroscience 24(38):8333-45.
- Ryan SM, Nolan YM. 2016. Neuroinflammation negatively affects adult hippocampal neurogenesis and cognition: can exercise compensate? Neuroscience and biobehavioral reviews 61:121-31.
- Saaltink DJ, Vreugdenhil E. 2014. Stress, glucocorticoid receptors, and adult neurogenesis: a balance between excitation and inhibition? Cell Mol Life Sci 71(13):2499-515.
- Selemon LD. 2013. A role for synaptic plasticity in the adolescent development of executive function. Translational psychiatry 3:e238.
- Shi Y, Chichung Lie D, Taupin P, Nakashima K, Ray J, Yu RT, Gage FH, Evans RM. 2004.
  Expression and function of orphan nuclear receptor TLX in adult neural stem cells.
  Nature 427(6969):78-83.
- Stranahan AM, Khalil D, Gould E. 2006. Social isolation delays the positive effects of running on adult neurogenesis. Nature neuroscience 9(4):526-33.
- Tang SW, Chu E, Hui T, Helmeste D, Law C. 2008. Influence of exercise on serum brainderived neurotrophic factor concentrations in healthy human subjects. Neurosci Lett 431(1):62-5.
- Tanti A, Belzung C. 2013. Neurogenesis along the septo-temporal axis of the hippocampus: are depression and the action of antidepressants region-specific? Neuroscience 252:234-52.
- Tanti A, Rainer Q, Minier F, Surget A, Belzung C. 2012. Differential environmental regulation of neurogenesis along the septo-temporal axis of the hippocampus. Neuropharmacology 63(3):374-84.
- van Praag H, Christie BR, Sejnowski TJ, Gage FH. 1999a. Running enhances neurogenesis, learning, and long-term potentiation in mice. Proceedings of the National Academy of Sciences of the United States of America 96(23):13427-31.
- van Praag H, Kempermann G, Gage FH. 1999b. Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. Nature neuroscience 2(3):266-70.

- Wei L, Meaney MJ, Duman RS, Kaffman A. 2011. Affiliative behavior requires juvenile, but not adult neurogenesis. The Journal of neuroscience : the official journal of the Society for Neuroscience 31(40):14335-45.
- Wong BK, Hossain SM, Trinh E, Ottmann GA, Budaghzadeh S, Zheng QY, Simpson EM. 2010. Hyperactivity, startle reactivity and cell-proliferation deficits are resistant to chronic lithium treatment in adult Nr2e1(frc/frc) mice. Genes, brain, and behavior 9(7):681-94.
- Young KA, Berry ML, Mahaffey CL, Saionz JR, Hawes NL, Chang B, Zheng QY, Smith RS, Bronson RT, Nelson RJ and others. 2002. Fierce: a new mouse deletion of Nr2e1; violent behaviour and ocular abnormalities are background-dependent. Behavioural Brain Research 132(2):145-158.

Acc

### **Figure Captions**

## Figure 1: Experimental design.

Outline of the experimental groups and timeline illustrating the duration of the experiment.

### Figure 2: Nr2e1 is necessary for the pro-neurogenic effect of exercise to occur.

Mean number of BrdU+ (**A**) and BrdU+NeuN+ (**B**) cells per section in the whole, dorsal and ventral hippocampus of singly-housed wild type or Nr2e1<sup>-/-</sup> adolescent mice with or without access to running wheels. Data are expressed as mean  $\pm$  SEM. \*\*\* p < 0.001 compared to WT counterparts; ++ p < 0.01, + p < 0.05 compared to WT sedentary mice; (Two-way ANOVA, Fisher's LSD), n = 5-6. Representative confocal images through the DG from WT sedentary (**C**) and running (**D**) and Nr2e1<sup>-/-</sup> sedentary (**E**) and running (**F**) mice. Immunohistochemical staining shows BrdU+ (red), NeuN+ (green) and BrdU+NeuN+ (orange) cells at 10X magnification. Scale bar = 100 µm. Higher magnification images depict immunopositive cells in the DG for BrdU (**C**' – **F**'), NeuN (**C**'' – **F**'') and merged channels (**C**''' – **F**'''). Scale bar = 25 µm.

Figure 3: Differential modulation of neurogenesis by isolation stress and exercise across the septo-temporal axis of the DG in adolescent mice.

Mean number of BrdU+ (**A**) and BrdU+NeuN+ (**B**) cells per section in the whole, dorsal and ventral hippocampus of singly- or pair- housed adolescent mice with or without access to running wheels. Data are expressed as mean  $\pm$  SEM. \* p < 0.05, \*\*\* p < 0.001 compared to single- or pair- housed sedentary counterparts; + p < 0.05, ++ p < 0.01 compared to pair-housed running mice; (Two-way ANOVA, Fisher's LSD), n = 4-6. Representative confocal images of coronal sections through the dDG and vDG immunohistochemically stained with BrdU (red) and NeuN (green) from pair-housed sedentary (dDG: **C**, vDG: **G**), pair-housed

running (dDG: **D**, vDG: **H**), single-housed sedentary (dDG: **E**, vDG: **I**) and single-housed running (dDG: **F**, vDG: **J**) mice. Images were taken at 10X magnification. Scale bar = 100  $\mu$ m. Higher magnification images depict immunopositive cells in the DG for BrdU (dDG: **C'** - **F'**, vDG: **G'** - **J'**), NeuN (dDG: **C''** - **F''**, vDG: **G''** - **J''**) and merged channels (dDG: **C'''** - **F''''**, vDG: **G'''** - **J'''**). Scale bar = 25  $\mu$ m.

Figure 4: Differential modulation of apoptosis by isolation stress and exercise across the septotemporal axis of the DG in WT but not Nr2e1<sup>-/-</sup> adolescent mice.

Mean number of active caspase-3+ cells per section in whole, dorsal and ventral hippocampus of singly-housed wild type or Nr2e1<sup>-/-</sup> adolescent mice (**A**) and singly- or pair-housed wild type adolescent mice with or without access to running wheels (**B**). Data are expressed as mean  $\pm$  SEM. \* p < 0.05, compared to single- or pair-housed sedentary counterparts; + p < 0.05, compared to single-housed running mice; (Two-way ANOVA, Fisher's LSD), n = 4. Representative bright field images of coronal sections through the hippocampus immunocytochemically stained with active caspase-3 (dark brown; black arrows) from a wild type control (**C**) and Nr2e1<sup>-/-</sup> (**D**) mouse. Images were taken at 20X magnification. Scale bar = 100 µm.



Figure 1: Experimental design.

321x184mm (300 x 300 DPI)

Accepter



Figure 2: Nr2e1 is necessary for the pro-neurogenic effect of exercise to occur.

336x191mm (300 x 300 DPI)

John Wiley & Sons



Figure 3: Differential modulation of neurogenesis by isolation stress and exercise across the septo-temporal axis of the DG in adolescent mice.

345x273mm (300 x 300 DPI)

John Wiley & Sons



Figure 4: Differential modulation of apoptosis by isolation stress and exercise across the septotemporal axis of the DG in WT but not Nr2e1-/- adolescent mice.

338x190mm (96 x 96 DPI)

Accepte