

	7			
Title	Changing antiepilepsy drug-prescribing trends in women with epilepsy in the UK and Ireland and the impact on major congenital malformations			
Authors	Kinney, Michael O.;Morrow, James;Patterson, Chris C.;Campbell, Ellen;Russell, Aline;Smithson, W. Henry;Parsons, Linda;Morrison, Patrick J.;Bromley, Rebecca;Liggan, Brenda;Delanty, Norman;Irwin, Beth;Hunt, Stephen J.;Craig, John J.			
Publication date	2018-04-16			
Original Citation	Kinney, M. O., Morrow, J., Patterson, C.C., Campbell, E., Russell, A., Smithson, W. H., Parsons, L., Morrison, P. J., Bromley, R., Liggan, B., Delanty, N., Irwin, B., Hunt, S. J. and Craig, J. J. (2018) 'Changing antiepilepsy drug-prescribing trends in women with epilepsy in the UK and Ireland and the impact on major congenital malformations', Journal of Neurology, Neuroscience and Psychiatry. doi:10.1136/jnnp-2017-317368			
Type of publication	Article (peer-reviewed)			
Link to publisher's version	10.1136/jnnp-2017-317368			
Rights	© 2018, Article authors (or their employer(s) unless otherwise stated in the text of the article). This article has been accepted for publication in Journal of Neurology, Neuroscience and Psychiatry, 2018, following peer review, and the Version of Record can be accessed online at https://dx.doi.org/ 10.1136/jnnp-2017-317368. Reuse of this manuscript version is permitted strictly pursuant to the terms of the CC-BY-NC-ND 4.0 license: http://creativecommons.org/licenses/by-nc-nd/4.0/ - https://creativecommons.org/licenses/by-nc-nd/4.0/			
Download date	2024-04-25 07:21:50			
Item downloaded from	https://hdl.handle.net/10468/6404			



## **Data Supplement**

		Monotherapy	Polytherapy	No AED exposures
Total number of exposures		6785	1858	604
Outcomes	Live births	6470	1711	585
	Spontaneous abortions	224	115	9
	Stillbirths/Intrauterine deaths	42	15	2
	Induced abortion	49	17	8
	Neonatal deaths	21	7	2
	Pregnancy losses with informative outcomes	27	13	2
	Total major congenital malformations	211	94	15
	Total Informative outcomes	6476	1717	585
	MCM rate (95% CI)*	3.3 (2.8-3.7) %	5.5 (4.5-6.7) %	2.6 (1.4-4.2) %

<sup>\*</sup> Malformation rates were calculated as [total major congenital malformations (which includes pregnancy losses with MCM i.e. informative outcomes] ÷ [total informative outcomes (i.e. all births where a complete assessment could be made of the presence of MCM and if a uninformative outcome (pregnancy loss in this context) occurred, these pregnancies were excluded from the denominator]

Abbreviations: CI; Confidence interval, AED; Anti-epileptic drugs, MCM; Major congenital malformation

<u>Supplementary Table 1</u>: Cumulative data showing frequency of exposure to different treatments, with associated outcome and malformation data through to August 2016.