

Title	Swallow-breathing coordination during incremental ascent to altitude
Authors	Huff, Alyssa;Day, Trevor A.;English, Mason;Reed, Mitchell D.;Zouboules, Shaelynn;Saran, Gurkarn;Leacy, Jack K.;Mann, Carli;Peltonen, Joel D. B.;O'Halloran, Ken D.;Sherpa, Mingma T.;Pitts, Teresa
Publication date	2019
Original Citation	Huff, A., Day, T. A., English, M., Reed, M. D., Zouboules, S., Saran, G., Leacy, J. K., Mann, C., Peltonen, J. D., O'Halloran, K. D. and Sherpa, M. T. (2019) 'Swallow-breathing coordination during incremental ascent to altitude', Respiratory Physiology and Neurobiology, 265, pp.121-126. https://doi.org/10.1016/ j.resp.2018.06.005
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.resp.2018.06.005
Rights	© 2019, Elsevier Ltd. All rights reserved. This manuscript version is made available under the CC BY-NC-ND 4.0 license https:// creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2025-07-16 09:13:55
Item downloaded from	https://hdl.handle.net/10468/15597



University College Cork, Ireland Coláiste na hOllscoile Corcaigh



HHS Public Access

Respir Physiol Neurobiol. Author manuscript; available in PMC 2019 July 01.

Published in final edited form as:

Author manuscript

Respir Physiol Neurobiol. 2019 July ; 265: 121-126. doi:10.1016/j.resp.2018.06.005.

Swallow-breathing coordination during incremental ascent to altitude

Alyssa Huff^{1,2}, Trevor A. Day³, Mason English², Mitchell D. Reed², Shaelynn Zouboules³, Gurkarn Saran³, Jack K. Leacy⁴, Carli Mann³, Joel DB. Peltonen³, Ken D. O'Halloran⁴, Mingma T. Sherpa⁵, and Teresa Pitts²

¹Department of Physiology, University of Louisville, Louisville, KY, USA ²Department of Neurological Surgery, Kentucky Spinal Cord Injury Research Center, University of Louisville, Louisville, KY, USA ³Department of Biology, Faculty of Science and Technology, Mount Royal University, Calgary, Alberta, Canada ⁴Department of Physiology, School of Medicine, University College Cork, Cork, Ireland ⁵Kunde Hospital, Khunde, Solukhumbu, Nepal

Abstract

Swallow and breathing are highly coordinated behaviors reliant on shared anatomical space and neural pathways. Incremental ascent to high altitudes results in hypoxia/hypocapnic conditions altering respiratory drive, however it is not known whether these changes also alter swallow. We examined the effect of incremental ascent (1,045m, 3,440m and 4,371m) on swallow motor pattern and swallow-breathing coordination in seven healthy adults. Submental surface electromyograms (sEMG) and spirometry were used to evaluate swallow triggered by saliva and water infusion. Swallow-breathing phase preference was different between swallows initiated by saliva versus water. With ascent, saliva swallows changed to a dominate pattern of occurrence during the transition from inspiration to expiration. Additionally, water swallows demonstrated a significant decrease in submental sEMG duration and a shift in submental activity to earlier in the apnea period, especially at 4,371m. Our results suggest that there are changes in swallow-breathing coordination and swallow production that likely increase airway protection with incremental ascent to high altitude. The adaptive changes in swallow were likely due to the exposure to hypoxia and hypocapnia, along with airway irritation.

Keywords

High altitude ascent; Electromyograms; Airflow; Apnea duration; Airway protection; Swallow coordination

Corresponding Author: Teresa Pitts, PhD, Department of Neurological Surgery, Kentucky Spinal Cord Injury Research Center, School of Medicine, University of Louisville, 511 S Floyd Street, Louisville, KY 40202. Conflict of Interest: None of the authors have a conflict of interest to report.

1. Introduction

Swallow and breathing are highly coordinated airway protective behaviors. Swallow is a multi-phase event, however the pharyngeal phase presents the highest risk for aspiration (Paydarfar et al., 1995). During the pharyngeal phase, supra-laryngeal/hyoid musculature moves the larynx superiorly and anteriorly resulting in closure of the airway and a functional apnea (German et al., 2009; Wheeler Hegland et al., 2011; Wheeler Hegland et al., 2009).

The expiratory phase of breathing is the preferred phase for swallow to occur, likely due to the limited inspiratory airflow (Martin-Harris et al., 2003). The central mechanism is thought to be due to interactions of breathing and swallow pattern generators (Dick et al., 1993; Miller, 1982), however this preference can be modified by peripheral feedback (Pitts et al., 2015b) and disease (Brodsky et al., 2010; Leslie et al., 2002; Troche et al., 2011). Specifically, alterations in respiratory mechanics due to chronic obstructive pulmonary disease (Nagami et al., 2017; Pinto et al., 2017) and/or upper abdominal laparotomy can shift swallow occurrences to inspiration, potentially increasing risk of aspiration (Pitts et al., 2015b). Additionally, there is also limited evidence that alterations in blood gasses (i.e., oxygen $[O_2]$ and carbon dioxide $[CO_2]$) can also increase the likelihood that swallow will occur during inspiration (D'Angelo et al., 2014), (Ghannouchi et al., 2013).

Incremental ascent to high altitudes (>2,000m) produces hypoxia (low O₂) induced hyperventilation, resulting in hypocapnia (low CO₂) (Huang et al., 1984; Weil, 1986). As climbers acclimatize to high altitude they can reach a new "steady-state chemoreflex drive" in which balance is achieved between hypoxia and hypocapnia, while ventilation parameters can return to near baseline conditions (Bruce CD, 2018; Pfoh et al., 2017). Additionally, healthy individuals that are not acclimatized to high altitude conditions can have changes in pulmonary mechanics due to interstitial pulmonary edema, which can be accompanied with accumulation of fluid within and around the airway walls (Cremona et al., 2002; Pratali et al., 2010; Schoene et al., 1988). Early symptoms such as shortness of breath and cough are often overlooked leading to mortality (Dunin-Bell and Boyle, 2009).

Due to the significant coordination necessary for swallow and breathing, it is likely that conditions which significantly alter respiratory drive and mechanics would also affect swallow production and swallow-breathing coordination. We hypothesized that with incremental ascent to high altitude there would be a decrease in swallow duration, and a shift in swallow phase preference to inspiration.

2. Methods

Ethics and Participant Recruitment

This study abided by the Canadian Government Tri-Council policy on research ethics with human participants (TCPS2) and the Declaration of Helsinki, except for registration in a database. Ethical approval was received in advance through Mount Royal University Human Research Ethics Board (Protocol 100012) and was harmonized with the Nepal Health Research Council (Protocol 109–2017). Participants were recruited via email

correspondence or direct verbal communication, and provided written, voluntary, informed and ongoing consent.

Ten participants were recruited for the study, while only seven (two males, five females) completed the study. One participant voluntarily withdrew from the study during ascent, another was excluded following baseline data acquisition due to a persistent cough and a third was excluded due to complications with data acquisition. Exclusion criteria included facial hair, as electrodes were unable to effectively adhere to skin, and health status (e.g., persistent cough, severe altitude illness). No pre-existing medical conditions were reported by any participants. Participants avoided rigorous exercise for at least 12 hours prior to data collection.

Incremental ascent to high altitude

Baseline measurements were recorded at 1,045m (Calgary) prior to the departure to Nepal. Following arrival in Kathmandu (1,400m), participants spent up to 3 days in Kathmandu before flying to Lukla (2,860m) where the trek to high altitude commenced (Figure 1). Consecutive measurements were obtained on rest days at 3,440m (Namche; day 3 at altitude) and 4,371m (Pheriche; day 5 at altitude) on every second day following arrival in Lukla (Figure 1), following one night sleep at each respective altitude.

Data Collection

Data acquisition was performed using an analog to digital data acquisition system [Powerlab/16SP ML880; AD Instruments (ADI), Colorado Springs, CO, USA], and data was collected, archived and analyzed offline using commercially available software (LabChart Pro software version 8) and a personal laptop computer. Surface electromyogram (sEMG) (ADI MLA2503 & ADI FE132) electrodes were placed approximately 3 cm posterior to the mental region of the mandible, on each side of the midline, capturing the submental complex. The grounding electrode was placed inferior to the participant's left clavicle. Voluntary swallow was performed in advance to ensure an adequate electrical signal through the sEMG electrodes.

A pneumotachometer (800L flow head; Series 3813; Hans Rudolph Inc.) and spirometer amplifier (ADI ML141) were used to monitor respiratory variables using a mouthpiece and nose-clip. Calibration of the flow head was performed with a 3L calibration syringe before data acquisition in each participant. Respiratory flow (L/s) was measured directly by the pneumotachograph. Inspired volume (V_{TI}; L) and respiratory frequency (f_R ; min⁻¹) were derived from respiratory flow. The product of V_{TI} and f_R was used to determine instantaneous minute ventilation (\dot{V}_I ; L/min). The pressure of end-tidal P_{ET}CO₂ was

measured using a portable, calibrated capnograph (Masimo EMMA, Danderyd, Sweden) with a personal mouthpiece and nose clip and peripheral oxygen saturation (SpO₂) was measured with a portable finger pulse oximeter (Masimo SET® Rad-5, Danderyd, Sweden). Electrocardiography (ECG; ADI MLA2503 & ADI FE132; lead II configuration) was utilized to derive instantaneous heart rate (HR; 1/R-R Interval in min⁻¹). The protocol was carried out with participants sitting comfortably in a dark, quiet room with ear plugs and eyes closed. Resting ventilation at each altitude was analyzed from a one-minute

representative period near the end of a 10-min baseline period, whereas $P_{ET}CO_2$ and SpO_2 measures were obtained after stability was achieved.

Swallow stimulation

1. Swallows produced during the baseline respiratory data via normal saliva collection in the mouth, termed *saliva swallows*.

2. *Water swallows* were trigged via water delivery from a 250 mL wash bottle (Nalgene 2089–0008 Narrow-Mouth Economy Bottle; Thermo Scientific, Waltham, MA, USA) inserted approximately 5 cm into the participant's mouth, lateral to the pneumotachometer mouthpiece. The wash bottle was positioned by each participant to ensure comfort with the water delivery. The infusion protocol began by recording a thirty-second baseline with all instrumentation in place. Following this baseline, water was infused at ~1 mL/second for 30 seconds into the participants' mouths. Finally, a 30 second washout was conducted after all instrumentation remaining in place. In all instances, participants were instructed before the introduction of water to swallow normally as needed.

Statistical Analysis

Data was analyzed from seven participants (5 female and 2 male) ages 19–23 at 1,045m (Calgary), 3,440m (Namche; day 3 at altitude), and 4,371m (Pheriche; day 7 at altitude) (Figure 1). All results were expressed as means \pm standard deviation (SD) using SPSS software (IBM).

To examine changes in swallow phase preference the following designations were used for respiratory phase: A) transition from inspiration to expiration (In-Ex); within expiration (Ex-Ex); transition from expiration to inspiration (Ex-In); and within inspiration (In-In). Then the following assigned coding system was used with In-Ex = 1; Ex-Ex = 2; Ex-In = 3; and In-In = 4 to categorize where each swallow occurred (Table 1). Finally, Wilcoxon signed ranks tests were run to determine changes across swallow-type and altitude, as we have previously used (Pitts et al., 2015b).

Swallow apnea duration was measured as the period of zero airflow in the event of a swallow (Figure 2). The apnea duration then was divided into three sub-phases: a) preswallow apnea, b) duration of submental sEMG, and c) post-swallow apnea (Figure 2). Preswallow apnea began at the time of zero airflow before the submental activation. Submental sEMG duration was measured as the activation and inactivation of submental sEMG. Postswallow apnea was measured as the zero airflow after the inactivation of submental complex (Figure 2). Additionally respiratory rate, heart rate, mean arterial pressure (MAP), \dot{v}_I , SpO₂, P_{ET}CO₂ and steady-state chemoreflex drive (SS-CD) were measured. The SS-CD was computed by calculating a stimulus index (SI; P_{ET}CO₂/SpO₂), and then comparing minute ventilation against SI (Bruce CD, 2018; Pfoh et al., 2017). A repeated measures ANOVA was used to determine differences in swallow motor pattern and respiratory parameters across the three elevations with significance at *p* 0.05, and if significance was met the LSD post-hoc test was used. A *p* 0.07 was designated as "approaching significance".

3. Results

Swallow was present during baseline respiratory measurements (saliva swallows), and reliably elicited with infusion of water in all subjects (water swallows). A total of 379 swallows (122 saliva and 257 water) were analyzed across the three altitudes (142 at 1,045m; 121 at 3,440m; and 116 at 4,371m).

Swallow-breathing coordination

Table 1 reports percent of swallow occurrences across each respiratory phase/transition. Water swallows had a strong In-Ex phase preference (69–79%) which was maintained through the ascent protocol. For saliva swallows at 1,045m only 43% occurred during In-Ex [significantly different than water (Z = -3.3, p < 0.001)], but this shifted at 3,440m with 76% of swallows occurring during In-Ex [significantly different than 1,045m (Z = -3.3, p < 0.001)]. At the highest altitude 4,371m the percent of swallows which occurred during the In-Ex transition reduced to 55% (p = 0.07). Interestingly, at 1,045m 21% of saliva swallows occurred during inspiration (In-In), which reduced to 6% at 3,440m and at 4,371m none occurred. In contrast <6% of water swallows occurred during inspiration (Table 1).

Change of swallow motor pattern with increasing altitude

Figure 3 demonstrates changes in pre-swallow apnea, submental duration, and post swallow apnea plotted by subjects across the three altitude locations. For swallows elicited by water, the average submental duration (ms) approached significance $[1170 \pm 539, 1038 \pm 218, \text{ and} 710 \pm 227 \text{ respectively } (F_{2, 12} = 4.19, p = 0.07)]$. As elevation increased pre-swallow apnea duration (ms) significantly decreased $[-256 \pm 236, -115 \pm 99, \text{ and} -5 \pm 172 \text{ respectively}$ ($F_{2, 12} = 4.218, p = 0.06$)], and post-swallow apnea duration (ms) significantly increased [56 \pm 109, 111 \pm 171, and 241 \pm 218 ($F_{2, 12} = 6.137, p < 0.05$)] (Table 2 and Figure 2 and 3). Of note, pre-swallow submental sEMG activity was seen during swallows at each elevation and of each type (Figure 2). For saliva swallows there was no significant change in submental sEMG and apnea duration, or swallow frequency (Table 2).

Breathing related variables

Table 2 also illustrates resting minute ventilation (V_I), the pressure of end-tidal $P_{ET}CO_2$,

peripheral oxygen saturation (SpO₂), stimulus index (SI) and measurement of steady-state chemoreflex drive (SS-CD) during incremental ascent to high altitude. All variables changed in predictable ways with incremental ascent. Heart rate [81.6 ± 9.5 , 97.8 ± 7.9 , and 93.5 ± 5.8 respectively (F_{2,12} =10.29, p < 0.05)], MAP [90.4 ± 8.4 , 96.0 ± 6.5 , and 99.1 ± 9.2 respectively (F_{2,12} = 11.88, p < 0.05)] and SS-CD significantly *increased* as altitude increased [36.8 ± 8.5 , 49.3 ± 12.7 , and 58.7 ± 19.5 respectively (F_{2,12} = 7.41, p < 0.05)]. SpO₂ [96.2 ± 1.0 , 88.1 ± 2.3 , and 83.3 ± 5.3 respectively (F_{2,12} = 37.44, p < 0.001)] and P_{ET}CO₂ [31.1 ± 4.2 , 25.9 ± 2.7 , and 21.3 ± 2.3 respectively (F_{2,12} = 31.61, p = 0.001)] significantly *decreased* as altitude increased. Additionally, respiratory rate and instantaneous minute ventilation remained stable across all elevations (Table 2).

4. Discussion

This is the first evidence of a significant change in swallow-breathing coordination as well as swallow production during incremental ascent to high altitude. There was a significant change in swallow phase preference comparing saliva to water swallows during baseline and approached significance at the highest elevation (4,371m). This was due to a shift in the dominance of the In-Ex pattern seen during water swallows and at 3,440m for saliva swallows. Additionally, in the water trials there was a significant increase in the post-swallow apnea period and a decrease (approaching significance) in the submental duration and pre-swallow apnea, while the overall swallow apnea duration did not change.

Phase Preference

Swallow phase preference has been intensely studied in humans (Martin-Harris, 2008; Martin-Harris et al., 2008; Martin-Harris et al., 2003; Martin-Harris and McFarland, 2013; Pratali et al., 2010; Wheeler Hegland et al., 2011; Wheeler Hegland et al., 2009), as well as in cats (Dick et al., 1993; Pitts et al., 2015a; Pitts et al., 2013; Pitts et al., 2015b), goats (Bonis et al., 2011; Feroah et al., 2002a; Feroah et al., 2002b), and rats (Saito et al., 2002a, b). However, all the peripheral stimulations and/or central mechanisms which regulate their interactions are not entirely understood. In the present study there was *not* a strong expiratory phase preference (~80%) which is observed in single swallow studies in which a 5 or 10 mL bolus is placed in the mouth (Wheeler Hegland et al., 2009). Saliva swallows (probably most akin to the typical single swallow task) demonstrated only 9% occurred during expiration, with 43% occurring in the transition of In-Ex, and of great interest is that 21% of these swallows occurred during inspiration (Table 1).

The dominance of In-Ex preference may be due in part to the mouthpiece which forces an "open mouth" swallow. It has been shown that muscle spindle afferents, in the masseter muscle, increase in discharge frequency during active opening of the jaw (Taylor et al., 1997). It has also been shown that input of muscle spindle afferents influence other central pattern generators [i.e. locomotion (Pearson, 1995)], and has been speculated that muscle spindle afferents influence mastication CPG output (Kolta et al., 1990; Lund, 2011). This information allows speculation that position of the jaw, indicated by proprioception of muscle spindle afferents can modulate the interaction between the swallow and breathing CPGs.

These changes could also be related to the effects of hypoxia and/or hypocapnia on swallow. Although there are limited studies, there are also conflicting results. In mice an increase in swallow frequency was reported (Khurana and Thach, 1996), no change in rat (Ghannouchi et al., 2013), and a decrease in the cat (Nishino et al., 1986). Hypoxia has also been studied in nonnutritive swallow in newborn lambs which showed a decrease in frequency during quiet sleep (Duvareille et al., 2007). Interestingly, hypercapnia shifts swallows towards In and Ex-In (D'Angelo et al., 2014) while we found that hypocapnia with hypoxia shifts swallow toward In-Ex. In light of the present data, further studies may need to investigate swallow-breathing coordination not only with variation of respiratory drive but swallow drive as well. We speculate that the water trials increased swallow excitability, which likely altered and stabilized its relationship with breathing.

Swallow motor pattern

In contrast to the swallow-breathing coordination data, the largest changes in the swallow motor pattern with ascent were on the water swallows, with a 39% decrease in the submental duration (Figure 2–3) at the highest altitude (compared to Calgary). This effect has been demonstrated in cats when swallow was coordinated with cough (airway irritation discussed below) (Leow et al., 2006); however we could find no study demonstrating a decrease in submental sEMG in healthy adults when using a mechanical/cold stimulus on the back of the mouth (Sciortino et al., 2003) or altering oral stimulation with taste (Leow et al., 2006).

To protect the airway during the pharyngeal phase of swallow the vocal folds must be adducted (zero flow; swallow apnea) during the laryngeal exposure to the bolus (Butler et al., 2004; Chi-Fishman and Sonies, 2000; Ding et al., 2003; Kijima et al., 1999; Martin-Harris et al., 2003; Martin et al., 1994; Paydarfar et al., 1995; Wheeler Hegland et al., 2011). In a review by Martin-Harris (2008), she stated that increases in the timing from the onset of the submental activity to the apnea period is related to significant clinical risk for aspiration. Evidence of this has been demonstrated in patients with Parkinson's disease with dysphagia (Ertekin, 2014). Based on this current data, we speculate that the decrease in submental sEMG and the shift in its activity to closer to the start of the swallow apnea period could increase airway protection. Of note, Ertekin and colleagues (Ertekin, 2014; Gürgör et al., 2013) demonstrated an activation of the submental complex during the pre-swallow respiratory phase that is likely related to infusion of water into the mouth (termed foreburst). Figure 2 demonstrates the difference between swallow-related and pre-swallow submental activity.

Airway Irritation

Exposure to high altitude conditions is also associated with airway irritation from dry air and insensible water loss, which results in a chronic cough (Freer, 2004). The most common diagnosis in the Nepal Himalaya is "Khumbu cough", also known as "high altitude hack" (Freer, 2004), thought to be caused by dry air, sub-zero temperatures, dust, and exposure to yak dung stoves in the lodges (Linoby et al., 2013). There is evidence that dry air increases airway responsiveness (Van Oostdam et al., 1986), and prolonged exposure results in an inflammatory response, desquamation of the epithelium, and edema of submucosa (Florey et al., 1932). While each subject did have evidence of coughing across the recording period, none were actively coughing during the measurement period. It is possible that activation of irritant receptors can alter swallow production without cough as a presenting feature.

Respiratory Drive

The changes in swallow and swallow-breathing coordination were also accompanied by changes/adaption of the chemoreflexes driving breathing. It is known that these reflexes become more dynamic as individuals acclimatize to their respective environment (Pfoh et al., 2017) (Steinback and Poulin). To asses this adaptation, Pfoh and colleagues (2017) created an index of steady-state chemoreflex drive (SS-CD), taking into account resting ventilation indexed against the overall contributions of both low O₂ and low CO₂ during exposure to hypoxia. Based on the magnitude of this index the significant change in the SS-

CD from 1,045m to 3,440m is evidence of respiratory acclimatization in our participants [see also (Huang et al., 1984)].

Blood levels of O₂ and CO₂ are maintained in part by central (brainstem) and peripheral (carotid body) chemoreceptors. Central chemoreceptors, located throughout brainstem, detect PCO₂/[H⁺] accumulation (Guyenet and Bayliss, 2015). Peripheral chemoreceptors located bilaterally within carotid bodies detect rapid changes in both O₂ and CO₂ synergistically (Fitzgerald and Parks, 1971; Lahiri and DeLaney, 1975; López-Barneo et al., 2016). A primary location for integrating these signals is in the nucleus tractus solitarius (NTS) (Jordan and Spyer, 1986; Paton et al., 2001). Due to the overlap in sensory integration in the NTS for breathing and swallow (Jean, 1984, 2001), this may be a site of shared central excitability which affects both respiratory and swallow central pattern generators.

Clinical Implications

Altitude exposure has inherent risks with 1–2% experiencing high altitude pulmonary edema (HAPE) (Houston, 1960; Hultgren, 1969; Schoene et al., 1986), a form of high altitude sickness, and of those 65% are diagnosed with a concomitant respiratory infection (most commonly pneumonia) (Leshem et al., 2008). It would be of interest to know if climbers with pneumonia display the same adaptations in swallow, especially in light of our knowledge of pneumonia rates with dysphagia.

5. Conclusions

Our results suggest that there are changes in swallow-breathing coordination and swallow motor production that increase airway protection with incremental ascent to high altitude. In conclusion, we suspect the adaptive changes in swallow were likely due to the exposure to superimposed hypoxia and hypocapnia, along with the increased airway irritation.

Acknowledgements:

R00-HL 111215, The Kentucky Spinal Cord and Head Injury Trust, The Commonwealth of Kentucky Challenge for Excellence, Natural Sciences and Engineering Research Council of Canada Discovery grant (RGPIN-2016–04915). We also want to thank the guides and research subjects we are grateful for your participation in this project.

References

- Bonis J, Neumueller S, Marshall B, Krause K, Qian B, Pan L, Hodges M, Forster H, 2011 The effects of lesions in the dorsolateral pons on the coordination of swallowing and breathing in awake goats. Respiratory Physiology and Neurobiology 175, 272–282. [PubMed: 21145433]
- Brodsky MB, McFarland DH, Dozier TS, Blair J, Ayers C, Michel Y, Gillespie MB, Day TA, Martin-Harris B, 2010 Respiratory-swallow phase patterns and their relationship to swallowing impairment in patients treated for oropharyngeal cancer. Head Neck 32, 481–489. [PubMed: 19780056]
- Bruce CD SG, Pfoh JR, Leacy JK, Zouboules SM, Peltonen JDB, Linares AM, Chiew AE, O'Halloran KD, Sherpa MT, Day TA, 2018 What is the Point of The Peak? Assessing Steady-State Chemoreflex Drive in High Altitude Field Studies. Adv Exp Med Biol. In Press.
- Butler SG, Postma GN, Fischer E, 2004 Effects of viscosity, taste, and bolus volume on swallowing apnea duration of normal adults. Otolaryngology Head and Neck Surgery 131, 860. [PubMed: 15577781]
- Chi-Fishman G, Sonies BC, 2000 Motor strategy in rapid sequential swallowing: new insights. Journal of Speech, Language and Hearing Research 43, 1481.

- Cremona G, Asnaghi R, Baderna P, Brunetto A, Brutsaert T, Cavallaro C, Clark TM, Cogo A, Donis R, Lanfranchi P, 2002 Pulmonary extravascular fluid accumulation in recreational climbers: a prospective study. The Lancet 359, 303–309.
- D'Angelo OM, Diaz-Gil D, Nunn D, Simons JC, Gianatasio C, Mueller N, Meyer MJ, Pierce E, Rosow C, Eikermann M, 2014 Anesthesia and increased hypercarbic drive impair the coordination between breathing and swallowing. Anesthesiology: The Journal of the American Society of Anesthesiologists 121, 1175–1183.
- Dick T, Oku Y, Romaniuk J, Cherniack N, 1993 Interaction between central pattern generators for breathing and swallowing in the cat. The Journal of Physiology 465, 715. [PubMed: 8229859]
- Ding R, Logemann JA, Larson CR, Rademaker AW, 2003 The effects of taste and consistency on swallow physiology in younger and older healthy individuals: a surface electromyographic study. Journal of Speech, Language and Hearing Research 46, 977.
- Dunin-Bell O, Boyle S, 2009 Secondary prevention of HAPE in a Mount Everest summiteer. High altitude medicine & biology 10, 293–296. [PubMed: 19775220]
- Duvareille C, Lafrance M, Samson N, St-Hilaire M, Pladys P, Micheau P, Bournival V, Langlois C, Praud J-P, 2007 Effects of hypoxia and hypercapnia on nonnutritive swallowing in newborn lambs. Journal of Applied Physiology 103, 1180–1188. [PubMed: 17626833]
- Ertekin C, 2014 Electrophysiological Evaluation of Oropharyngeal Dysphagia in Parkinson's Disease. Journal of Movement Disorders 7, 31–56. [PubMed: 25360228]
- Feroah TR, Forster H, Fuentes CG, Lang IM, Beste D, Martino P, Pan L, Rice T, 2002a Effects of spontaneous swallows on breathing in awake goats. Journal of Applied Physiology 92, 1923–1935. [PubMed: 11960942]
- Feroah TR, Forster H, Fuentes CG, Wenninger J, Martino P, Hodges M, Pan L, Rice T, 2002b Contributions from rostral medullary nuclei to coordination of swallowing and breathing in awake goats. Journal of Applied Physiology 93, 581–591. [PubMed: 12133868]
- Fitzgerald RS, Parks DC, 1971 Effect of hypoxia on carotid chemoreceptor response to carbon dioxide in cats. Respiration physiology 12, 218–229. [PubMed: 4327983]
- Florey H, Carleton H, Wells A, 1932 Mucus secretion in the trachea. British journal of experimental pathology 13, 269.
- Freer L, 2004 Descriptive report of experience designing and staffing the first-ever medical clinic at Mt. Everest base camp, 2003. High altitude medicine & biology 5, 89–90. [PubMed: 15072722]
- German RZ, Crompton AW, Thexton AJ, 2009 Integration of the reflex pharyngeal swallow into rhythmic oral activity in a neurologically intact pig model. J Neurophysiol 102, 1017–1025. [PubMed: 19515957]
- Ghannouchi I, Duclos C, Marie J, Verin E, 2013 Modification in swallowing and ventilation coordination during hypercapnia, hypoxia, and tachypnea in unrestrained animals. Neurogastroenterology & Motility 25, 308. [PubMed: 23278913]
- Gürgör N, Arıcı , Incesu TK, Seçil Y, Tokuço lu F, Ertekin C, 2013 An electrophysiological study of the sequential water swallowing. Journal of Electromyography and Kinesiology 23, 619–626. [PubMed: 23466271]
- Guyenet PG, Bayliss DA, 2015 Neural control of breathing and CO2 homeostasis. Neuron 87, 946–961. [PubMed: 26335642]
- Houston CS, 1960 Acute pulmonary edema of high altitude. New England Journal of Medicine 263, 478–480. [PubMed: 14403413]
- Huang S, Alexander J, Grover R, Maher J, McCullough R, McCullough R, Moore L, Sampson J, Weil J, Reeves J, 1984 Hypocapnia and sustained hypoxia blunt ventilation on arrival at high altitude. Journal of Applied Physiology 56, 602–606. [PubMed: 6423588]
- Hultgren HN, 1969 High altitude pulmonary edema. Biomedicine of high terrestrial elevations, 131–141.
- Jean A, 1984 Control of the central swallowing program by inputs from the peripheral receptors. A review. Journal of the autonomic nervous system 10, 225–233. [PubMed: 6384335]
- Jean A, 2001 Brain stem control of swallowing: neuronal network and cellular mechanisms. Physiological Review 81, 929–969.

- Jordan D, Spyer KM, 1986 Brainstem integration of cardiovascular and pulmonary afferent activity. Prog Brain Res 67, 295–314. [PubMed: 3823479]
- Khurana A, Thach BT, 1996 Effects of upper airway stimulation on swallowing, gasping, and autoresuscitation in hypoxic mice. Journal of applied physiology (Bethesda, Md. : 1985) 80, 472–477.
- Kijima M, Isono S, Nishino T, 1999 Coordination of swallowing and phases of respiration during added respiratory loads in awake subjects. Am J Respir Crit Care Med 159, 1898–1902. [PubMed: 10351937]
- Kolta A, Lund J, Rossignol S, 1990 Modulation of activity of spindle afferents recorded in trigeminal mesencephalic nucleus of rabbit during fictive mastication. Journal of neurophysiology 64, 1067– 1076. [PubMed: 2258735]
- Lahiri S, DeLaney R, 1975 Stimulus interaction in the responses of carotid body chemoreceptor single afferent fibers. Respiration physiology 24, 249–266. [PubMed: 242049]
- Leow L, Huckabee ML, Sharma S, Tooley T, 2006 The influence of taste on swallowing apnea, oral preparation time, and duration and amplitude of submental muscle contraction. Chemical senses 32, 119–128. [PubMed: 17071940]
- Leshem E, Pandey P, Shlim DR, Hiramatsu K, Sidi Y, Schwartz E, 2008 Clinical features of patients with severe altitude illness in Nepal. Journal of travel medicine 15, 315–322. [PubMed: 19006504]
- Leslie P, Drinnan MJ, Ford GA, Wilson JA, 2002 Swallow respiration patterns in dysphagic patients following acute stroke. Dysphagia 17, 202–207. [PubMed: 12140646]
- Linoby, A.F., Nias, M.A., Ahmad, B.E., Zaki, S., Canda, R., Sariman, H., Azam, Z., Amat, A., 2013. Physiological Responses and Adaptations to Exposure from Moderate to Extreme Altitude: A Case Study of the Youngest Malaysian Climber to Scale Mt. Everest.
- López-Barneo J, González-Rodríguez P, Gao L, Fernández-Agüera MC, Pardal R, Ortega-Sáenz P, 2016 Oxygen sensing by the carotid body: mechanisms and role in adaptation to hypoxia. American Journal of Physiology-Cell Physiology 310, C629–C642. [PubMed: 26764048]
- Lund JP, (2011). Chapter 15 Chew before you swallow, in: Gossard JP, Dubuc R, Kolta A (Eds.), Progress in Brain Research. Elsevier, pp. 219–228.
- Martin-Harris B, 2008 Clinical implications of respiratory–swallowing interactions. Current opinion in otolaryngology & head and neck surgery 16, 194–199. [PubMed: 18475070]
- Martin-Harris B, Brodsky MB, Michel Y, Castell DO, Schleicher M, Sandidge J, Maxwell R, Blair J, 2008 MBS measurement tool for swallow impairment—MBSImp: establishing a standard. Dysphagia 23, 392–405. [PubMed: 18855050]
- Martin-Harris B, Brodsky MB, Price CC, Michel Y, Walters B, 2003 Temporal coordination of pharyngeal and laryngeal dynamics with breathing during swallowing: single liquid swallows. Journal of Applied Physiology 94, 1735. [PubMed: 12506044]
- Martin-Harris B, McFarland DH, (2013). Coordination of Deglutition and Respiration, Principles of Deglutition. Springer, pp. 25–34.
- Martin B, Logemann J, Shaker R, Dodds W, 1994 Coordination between respiration and swallowing: respiratory phase relationships and temporal integration. Journal of Applied Physiology 76, 714. [PubMed: 8175582]
- Miller AJ, 1982 Deglutition. Physiological Review 62, 129-184.
- Nagami S, Oku Y, Yagi N, Sato S, Uozumi R, Morita S, Yamagata Y, Kayashita J, Tanimura K, Sato A, Takahashi R, Muro S, 2017 Breathing–swallowing discoordination is associated with frequent exacerbations of COPD. BMJ Open Respiratory Research 4, e000202.
- Nishino T, Kohchi T, Honda Y, Shirahata M, Yonezawa T, 1986 Differences in the effects of hypercapnia and hypoxia on the swallowing reflex in cats. BJA: British Journal of Anaesthesia 58, 903–908. [PubMed: 3089244]
- Paton JF, Deuchars J, Li YW, Kasparov S, 2001 Properties of solitary tract neurones responding to peripheral arterial chemoreceptors. Neuroscience 105, 231–248. [PubMed: 11483315]
- Paydarfar D, Gilbert RJ, Poppel CS, Nassab PF, 1995 Respiratory phase resetting and airflow changes induced by swallowing in humans. The Journal of physiology 483, 273–288. [PubMed: 7776238]
- Pearson KG, 1995 Proprioceptive regulation of locomotion. Current Opinion in Neurobiology 5, 786– 791. [PubMed: 8805415]

- Pfoh JR, Steinback CD, Berg ERV, Bruce CD, Day TA, 2017 Assessing chemoreflexes and oxygenation in the context of acute hypoxia: Implications for field studies. Respiratory physiology & neurobiology 246, 67–75. [PubMed: 28757365]
- Pinto CF, Balasubramanium RK, Acharya V, 2017 Nasal airflow monitoring during swallowing: Evidences for respiratory-swallowing incoordination in individuals with chronic obstructive pulmonary disease. Lung India : official organ of Indian Chest Society 34, 247–250. [PubMed: 28474650]
- Pitts T, Gayagoy A, Rose M, Poliacek I, Condrey J, Musslewhite M, Shen T, Davenport P, Bolser D, 2015a Suppression of Abdominal Motor Activity during Swallowing in Cats and Humans. PloS one 10, e0128245–e0128245. [PubMed: 26020240]
- Pitts T, Rose MJ, Mortensen AN, Polia ek I, Sapienza CM, Lindsey BG, Morris KF, Davenport PW, Bolser DC, 2013 Coordination of cough and swallow: a meta-behavioral response to aspiration. Respiratory physiology & neurobiology 189, 543–551. [PubMed: 23998999]
- Pitts T, Rose MJ, Poliacek I, Condrey J, Davenport PW, Bolser DC, 2015b Effect of laparotomy on the swallow-breathing relationship in the cat. Lung 193, 129–133. [PubMed: 25331536]
- Pratali L, Cavana M, Sicari R, Picano E, 2010 Frequent subclinical high-altitude pulmonary edema detected by chest sonography as ultrasound lung comets in recreational climbers. Critical care medicine 38, 1818–1823. [PubMed: 20562696]
- Saito Y, Ezure K, Tanaka I, 2002a Difference between hypoglossal and phrenic activities during lung inflation and swallowing in the rat. J Physiol 544, 183–193. [PubMed: 12356891]
- Saito Y, Ezure K, Tanaka I, 2002b Swallowing-related activities of respiratory and non-respiratory neurons in the nucleus of solitary tract in the rat. Journal of Physiology 540, 1047–1060. [PubMed: 11986389]
- Schoene RB, Hackett PH, Henderson WR, Sage EH, Chow M, Roach RC, Mills WJ, Martin TR, 1986 High-altitude pulmonary edema. Jama 256, 63–69. [PubMed: 3012134]
- Schoene RB, Swenson ER, Pizzo CJ, Hackett PH, Roach RC, Mills WJ Jr, Henderson W Jr, Martin T, 1988 The lung at high altitude: bronchoalveolar lavage in acute mountain sickness and pulmonary edema. Journal of Applied Physiology 64, 2605–2613. [PubMed: 3403445]
- Sciortino KF, Liss JM, Case JL, Gerritsen KGM, Katz RC, 2003 Effects of Mechanical, Cold, Gustatory, and Combined Stimulation to the Human Anterior Faucial Pillars. Dysphagia 18, 16– 26. [PubMed: 12497192]
- Steinback CD, Poulin MJ, 2007 Ventilatory responses to isocapnic and poikilocapnic hypoxia in humans. Respir Physiol Neurobiol 155, 104–113. [PubMed: 16815106]
- Taylor A, Hidaka O, Durbaba R, Ellaway PH, 1997 Fusimotor influence on jaw muscle spindle activity during swallowing-related movements in the cat. J Physiol 503 (Pt 1), 157–167. [PubMed: 9288683]
- Troche MS, Huebner I, Rosenbek JC, Okun MS, Sapienza CM, 2011 Respiratory-Swallowing Coordination and Swallowing Safety in Patients with Parkinson's Disease. Dysphagia 26, 218– 224. [PubMed: 20623304]
- Van Oostdam J, Walker D, Knudson K, Dirks P, Dahlby R, Hogg J, 1986 Effect of breathing dry air on structure and function of airways. Journal of Applied Physiology 61, 312–317. [PubMed: 3733618]
- Weil JV, 1986 Ventilatory control at high altitude. Comprehensive Physiology.
- Wheeler Hegland K, Huber JE, Pitts T, Davenport PW, Sapienza CM, 2011 Lung Volume Measured During Sequential Swallowing in Healthy Young Adults. Journal of Speech, Language, and Hearing Research 54, 777.
- Wheeler Hegland KM, Huber JE, Pitts T, Sapienza CM, 2009 Lung volume during swallowing: single bolus swallows in healthy young adults. Journal of Speech, Language, and Hearing Research 52, 178.



Figure 1.

Timeline of travel, ascent, and recording locations. The (\bigstar) represents where data was collected, and (\clubsuit) indicates flights.



Figure 2.

Example of submental sEMG and airflow from the same participant from Calgary (1,045m) and Pheriche (4,371m) during the water swallow protocol. B to C marks the swallow apnea period. A to B is the pre-swallow submental activity, A to D is the submental duration and C to D is the post-swallow apnea period. At 4,371m, there was a significant increase in the post-swallow apnea as well as a decrease submental duration. The "foreburst" is activity related to water being introduced to the oral cavity.

Huff et al.



Figure 3.

Scatter plot of duration measures (pre-swallow, submental and post-swallow) for each subjects across the recording locations for the saliva (A) and water (B) swallow tasks. Repeated measures ANOVA *p*-value reported for each dependent measure, and gray line represents group mean.

Table 1.

Percent of swallow occurrence during breathing across the three levels of ascent.

In-Ex Ex-Ex Ex-In In-In Saliva Swallow - <td< th=""><th></th><th></th><th></th><th></th><th></th></td<>					
Saliva Swallow 1,045m 43 9 27 21 3,440m 76 12 6 6 4,371m 55 15 30 0 Water Swallow 1,045m 79 9 9 2 3,440m 69 14 11 6 4,371m 76 11 11 2		In-Ex	Ex-Ex	Ex-In	In-In
1,045m 43 9 27 21 3,440m 76 12 6 6 4,371m 55 15 30 0 Water Swallow 1,045m 79 9 9 2 3,440m 69 14 11 6 4,371m 76 11 11 2	Saliva Swallow				
3,440m 76 12 6 6 4,371m 55 15 30 0 Water Swallow 1,045m 79 9 9 2 3,440m 69 14 11 6 4,371m 76 11 11 2	1,045m	43	9	27	21
4,371m 55 15 30 0 Water Swallow 1,045m 79 9 9 2 3,440m 69 14 11 6 4,371m 76 11 11 2	3,440m	76	12	6	6
Water Swallow 1,045m 79 9 9 2 3,440m 69 14 11 6 4,371m 76 11 11 2	4,371m	55	15	30	0
1,045m 79 9 9 2 3,440m 69 14 11 6 4,371m 76 11 11 2	Water Swallow				
3,440m 69 14 11 6 4,371m 76 11 11 2	1,045m	79	9	9	2
4,371m 76 11 11 2	3,440m	69	14	11	6
	4,371m	76	11	11	2

Author Manuscript

Table 2.

different elevations are shown in this table. Resting respiratory rate (RR), mean arterial pressure (MAP), peripheral O₂ saturation (SpO₂), end tidal CO₂ Means, standard deviations (SD), and *p*-values comparing ventilatory, cardiac and acclimation values, as well as saliva and water swallows at the three pressure ($P_{ET}CO_2$), instantaneous minute ventilation (\dot{V}_I), and steady-state chemoreflex drive (SS-CD) are recorded. Submental (swallow) duration, swallow apnea duration, pre-swallow apnea and post-swallow apnea (Figure 2) are recorded in both saliva and water conditions. Figure 3 displays swallow data by participant.

	Calgary	7 (1,((45m)	Namo	he (3	,440m)	Pher	iche	(4,371m)	and and a
	Mean	+I	SD	Mean	+I	SD	Mean	+I	SD	<i>p</i> -values
Resting RR (min ⁻¹)	14.0	+1	6.0	15.2	+1	3.9	15.5	+1	5.2	0.42
Resting Heart Rate	81.6	+1	9.5	97.8	+1	7.9 ***	93.5	+1	5.8 **	0.004
MAP (mm Hg)	90.4	+I	8.4	96.0	+I	6.5 **	99.1	+I	9.2 ***	0.004
SpO ₂ (%)	96.2	+I	1.0	88.1	+I	2.3 ***	83.3	+I	5.3 ***††	<0.001
$P_{ET}CO_2$ (Torr)	31.1	+1	4.2	25.9	+1	2.7 **	21.3	+1	2.3 ***†††	0.001
\dot{V}_I (L/min)	11.9	+1	2.7	14.2	+1	2.6	14.7	+1	3.8	0.07
SS-CD (V _I /SI)	36.8	+1	8.5	49.3	+1	12.7 *	58.7	+1	$19.5 \ ^{**7}$	0.02
Saliva Swallow Data										
Submental Duration (ms)	1480	+1	804	1070	+1	490	1015	+1	457	0.35
Swallow Apnea Duration (ms)	1088	+1	433	1121	+1	253	1010	+1	374	0.77
Pre-Swallow Apnea (ms)	-296	+1	220	-183	+1	113	-141	+I	265	0.45
Post-Swallow Apnea (ms)	66-	+1	363	233	+1	385	174	+1	101	0.30
Water Swallow Data										
Submental Duration (ms)	1170	+1	539	1038	+1	218	710	+I	227 *†††	0.07
Swallow Apnea Duration (ms)	973	+1	398	1030	+1	165	946	+1	285	0.60
Pre-Swallow Apnea (ms)	-256	+1	236	-115	+1	66	-5	+1	126 *	0.06
Post-Swallow Apnea (ms)	56	+1	109	111	+1	171	241	+1	$218 \ ^{**/}$	0.02
Reported <i>p</i> -values are for repeated	d measure	s one	way Al	VOVA an	d sig	nificant val	ues are b e	olded	-	

Respir Physiol Neurobiol. Author manuscript; available in PMC 2019 July 01.

* Significant difference from Calgary *p 0.06,

** p<0.05

Author Manuscript

Author Manuscript

 † Significant difference between Namche and Pheriche † p 0.06 †† $p_{-}0.05$ ††† $p_{-}0.01$