CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

| | |
|---|---|
| Title | Habitat use of culturally distinct Galápagos sperm whale Physeter macrocephalus clans |
| Authors | Eguiguren, Ana;Pirotta, Enrico;Cantor, Maurício;Rendell, Luke;Whitehead, Hal |
| Publication date | 2019-01-17 |
| Original Citation | Eguiguren, A., Pirotta, E., Cantor, M., Rendell, L. and Whitehead, H. (2019) 'Habitat use of culturally distinct Galápagos sperm whale Physeter macrocephalus clans', Marine Ecology Progress Series, 609, pp. 257-270. doi:10.3354/meps12822 |
| Type of publication | Article (peer-reviewed) |
| Link to publisher's version | https://www.int-res.com/abstracts/meps/v609/p257-270/ - 10.3354/meps12822 |
| Rights | © 2019, Inter-Research. All rights reserved. |
| Download date | 2026-11-03 11:47:41 |
| Item downloaded from | https://hdl.handle.net/10468/7517 |

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

# Habitat use of culturally distinct Galápagos sperm whale (*Physeter macrocephalus*) clans

Running page head: Sperm whale clan habitat

Ana Eguiguren[1,2]*, Enrico Pirotta[3,4], Maurício Cantor[5,6], Luke Rendell[7], Hal Whitehead[1]

[1]Department of Biology, Dalhousie University, Oxford Street, Halifax, Nova Scotia B3H 4J1 Canada

[2]Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito, PC 17-1200-841 Ecuador

[3]Department of Mathematics and Statistics, Washington State University, Vancouver, WA 98686 USA

[4]School of Biological, Earth and Environmental Sciences,University College, Cork, T12K8AF, Ireland

[5]Departamento de Ecologia e Zoologia, Universidade Federal de Santa Catarina, Florianópolis, SC 88040-970 Brazil

[6]Centro de Estudos do Mar, Universidade Federal do Paraná, Pontal do Sul, PR 83255-976 Brazil

[7]Center for Social Learning and Cognitive Evolution, and Sea Mammal Research Unit, University of St Andrews, St Andrews, Fife, KY16 9TH UK

*Corresponding author email: anaeguibur@gmail.com

## Abstract

Ecological niche is traditionally defined at the species level, but individual niches can vary considerably within species. Research on intra-specific niche variation has been focused on intrinsic drivers. However, differential transmission of socially learned behaviours can also lead to intra-specific niche variation. In sperm whales (*Physeter macrocephalus*), social transmission of information is thought to generate culturally distinct clans, which at times occur sympatrically. Clans have distinct dialects, foraging success rates, and movement patterns, but whether the niches of clan members are also different remains unknown. We evaluated the differences in habitat use of clans off the Galápagos Islands, using data collected over 63 encounters between 1985 and 2014. During encounters, we recorded geographic positions, determined clan identity through analysis of group vocalizations and individual associations, and used topographical and oceanographic variables as proxies of sperm whale prey distribution. We used logistic Generalized Additive Models, fitted with Generalized Estimating Equations to account for spatiotemporal autocorrelation, to predict clan identity as a function of the environment descriptors. Oceanographic variables marginally contributed to differentiating clans. Clan identity could be predicted almost entirely based on geographic location. This fine-scale, within-region spatial partitioning likely derives from whales preferring areas where members of their clans occur over temporal scales of a few months to a few years. By identifying differences in clans' space use, we have uncovered another level of sperm whale life that is likely influenced by their cultural nature.

**Key words:** habitat preference, cetacean, culture, GAM, GEE, Galápagos

## Introduction

Traditionally, ecological niche and habitat use have been defined at the species level (Hutchinson 1957, Leibold 1995). However, mounting evidence for individuals of the same population having low niche overlap reminds us that conspecifics are not always ecologically equivalent (Bolnick et al. 2003). To date, most of the theoretical work on individual niche variation has focused on intrinsic sources of variation, such as morphological, physiological, and ontogenic traits (Van Valen 1965, Roughgarden 1972, Svanbäck & Persson 2004). Less attention has been given to social learning as a mechanism for individual niche variation (but see Galef 1976; Laland et al. 2000; Slagsvold and Wiebe 2007; Sargeant and Mann 2009).

When behavioural traits are socially learned and shared among groups of individuals, there is culture (Boyd & Richerson 1996, Laland & Hoppitt 2003). Culture, as so defined, can play an important role in the divergence of resource and space use among individuals, especially in species in which foraging strategies and habitat selection are socially transmitted (e.g. Laland & Galef, 2009; Whitehead & Rendell, 2014). Notable cases include apes and monkeys that learn to use different tools to exploit nuts and termites (McGrew et al. 1979, Boesch et al. 1994, Whiten et al. 1999, van Schaik et al. 2003, Ottoni & Izar 2008), birds that learn about feeding areas and prey sizes from their parents' choices (Slagsvold & Wiebe 2011), female mountain sheep retaining the home ranges of their social groups (Geist 1971), dolphins using the same foraging tactics and areas of their mothers and/or peers (Mann & Patterson 2013, Cantor et al. 2018), and sea otters using foraging tools to meet their matrilineally transmitted dietary preferences (Estes et al. 2003). These and other foraging techniques and habitat use patterns are socially acquired behavioural traits that result in different resource use patterns, and so reduce trophic niche

64 overlap among subsets of individuals within the same population (Jaeggi et al. 2010, Slagsvold
65 & Wiebe 2011, Allen et al. 2013).

66 However, it is not always straightforward to disentangle culture from other underlying causes of
67 foraging behaviour variation. Both genetic and ecological factors are explanatory candidates for
68 behavioural divergence, especially in allopatric populations (e.g. Laland and Galef 2009; Koops
69 et al. 2013). One way to overcome this issue is excluding all sources of non-cultural behavioural
70 variation (Whiten et al. 1999), but this has proved problematic (Laland & Janik 2006).
71 Alternatively, by studying resource-use variation among sympatric groups of genetically-similar
72 individuals, one can account for such environmental and genetic mechanisms. Two particularly
73 well-known marine examples are killer whales (*Orcinus orca*) and Indo-Pacific bottlenose
74 dolphins (*Tursiops* sp.). Mammal-eating and fish-eating killer whales use the same waters off
75 British Columbia but feed exclusively on very different prey (Ford et al. 1998). Off Shark Bay,
76 Australia, part of a bottlenose dolphin population uses marine sponges as tools to forage on the
77 seafloor for prey that are hard to access otherwise, leading to distinct social communities of
78 "sponging" and "non-sponging" dolphins that coexist in the same habitat (Mann et al. 2012).
79 Neither case can be explained by genetic variation alone (Krützen et al. 2005, Mann et al. 2012,
80 Riesch et al. 2012).

81 Over much wider spatial scales, there is the case of sympatric cultural divergence among female
82 sperm whales (*Physeter macrocephalus*) into clans. While males lead mostly solitary lives in
83 high latitudes, females and immatures live in tightly-knit social units, containing few matrilines,
84 in tropical and subtropical waters (Best 1979, Christal et al. 1998). Social units form temporary
85 larger groups (Whitehead et al. 1991), but they do so with other units with which they share a
86 large proportion of their acoustic repertoire, thus delineating a higher social level: the *vocal clan*

87 (Rendell & Whitehead 2003, Whitehead et al. 2012, Gero et al. 2016). Sperm whale clans of the
88 Eastern Tropical Pacific are genetically indistinct (Rendell et al. 2012) and sympatric (Rendell &
89 Whitehead 2003). Members of different clans can encounter one another easily, in theory.
90 However, they not only maintain distinct vocal dialects over time (Rendell & Whitehead 2005),
91 but also differ in movement and social behaviour, reproductive and foraging success, and diet
92 composition (Whitehead & Rendell 2004, Marcoux 2005, Marcoux et al. 2007a, Cantor &
93 Whitehead 2015). These divergences suggest that sperm whales belonging to culturally distinct
94 but sympatric clans may use different habitats, but this has not yet been studied directly.

95 Understanding sperm whale niche is hampered by logistical constraints. Their trophic niche, for
96 instance, is known only indirectly. Sperm whales seem to primarily prey on cephalopods, but
97 since they live offshore and feed at great depths (Papastavrou et al. 1989), observations of
98 predation are rare. Moreover, analyses of stomach contents and defecation yield contrasting
99 results regarding the species consumed (see Clarke et al. 1988, Clarke & Paliza 2001, Smith &
100 Whitehead 2000). While many bathypelagic squid have overlapping ranges and niches
101 (Nigmatullin et al. 2001), different age and size classes within single species have different
102 distributions and dietary preferences (Nigmatullin et al. 2001; Markaida 2006). On the other
103 hand, the habitat component of sperm whale niche can be assessed via the environmental
104 variables that influence the distribution of the cephalopods they prey upon (Jaquet & Whitehead
105 1996), such as bottom topography and oceanographic variables that are related to upwelling
106 processes and increased productivity (Jaquet & Whitehead 1996, Pirotta et al. 2011, Wong &
107 Whitehead 2014).

108 Here, we evaluate whether sympatric sperm whale clans differ in habitat use by investigating the
109 spatial, oceanographic, and topographic characteristics of the waters they occupy off the

Galápagos Islands. Specifically, we compared the relative habitat use of two vocal clans that were particularly common in the area in the 1980's (Rendell & Whitehead 2003), and of two other clans that have recently replaced them in the 2010's (Cantor et al. 2016).

## Methods

### *Field Methods*

We studied sperm whales off the Galápagos Archipelago (93º-88ºW; 3ºN-3ºS) aboard dedicated research sailboats (10-12m) between January and June, in 1987, 1989, 2013, and 2014 (Table 1). We searched for whales acoustically, monitoring hydrophones that could detect sperm whale clicks up to about 7 kilometres away every 15-60 minutes (Whitehead 2003). During daylight hours, we also searched for whales visually within a range of 0.2 to 2.0 km, depending on weather conditions. Upon finding a group of sperm whales, we approached it cautiously to photograph their flukes for individual identification (Arnbom 1987). We refer to the periods during which we had continuous (within less than 6 hours) visual and/or acoustic contact with the same group of females as *encounters*.

Groups of females and immatures (identified based on body size and behaviour; Whitehead 2003) were followed for as long as possible, during which time the vessel's geographic location was recorded. Until 1993, positions were estimated by interpolation from SATNAV fixes at least every 3 hours; after 1993, positions were recorded every 1-5 min using GPS (as in Whitehead and Rendell 2004). Vessel positions were used as indicators of the whales' locations, which, given the range of acoustic detection, could be up to 7 kilometres away from the vessel.

### *Clan identification*

We assigned clan identity to groups of female and immature sperm whales based on the similarity of their communication sounds, called codas (see Rendell and Whitehead 2003; Cantor et al. 2016). A clan was considered a collection of groups of sperm whales that shared an identifiable part of their coda repertoires (see Rendell & Whitehead 2003). At least four vocal clans were commonly sighted around Galápagos (Rendell & Whitehead 2003, Cantor et al. 2016): *Regular* (typically producing regularly-spaced clicks); *Plus-One* (typical codas with an extended pause before the last click), *Short* (typical codas with fewer than 5 clicks), and *Four-Plus* clan (typical codas with a basis of 4 regular clicks).

We assigned clan memberships to all groups of whales that were photo-identified together and had their acoustic repertoire sufficiently sampled (see Rendell & Whitehead 2003, Cantor et al. 2016). Geographic positions within a day were assigned to a corresponding clan because: 1) typically only one group of whales was tracked per day; 2) whales of the same group belong to the same clan; 3) groups from different clans are typically found some days apart (Whitehead & Rendell 2004). However, in four multiple-day encounters, more than one clan was identified, likely due to the replacement of the tracked group by one of another clan during the night. Since we could not determine the time the new group of whales was found, for these encounters, we used only geographic positions that were recorded in daylight (06:00-18:00), during which photo-identifications were available (see Whitehead and Rendell 2004).

*Environmental descriptors*

As topographical variables, we used depth from the General Bathymetric Chart of the Oceans (http://www.gebco.net/data_and_products/gridded_bathymetry_data/) and percentage of slope incline, calculated with Spatial Analysis tools in ArcGIS. As oceanographic variables, we used relative mean sea surface temperature (relSST) as a proxy for upwelling and standard deviation

154 of SST (sdSST) as a proxy for frontal activity from the Pathfinder Version 5.0 & 5.1 dataset
155 collected by the Advanced Very High Resolution Radiometer (AVHRR) and processed by the
156 NOAA National Oceanographic Data Center for 1980's data points, and Aqua-MODIS satellite
157 images distributed by the NOAA CoastWatch Program and NASA's Goddard Space Flight
158 Center for 2010's data points (see Griffin 1999; Praca et al. 2009; Pirotta et al. 2011). We
159 calculated relSST as the difference between SST at a geographic position and the mean SST over
160 the entire Galápagos region (defined as 93º-88ºW; 2ºN-2ºS for the 1980's period and 93º-88ºW;
161 1.5ºN-2ºS for the 2010's period) for the corresponding month We also considered chlorophyll-a
162 concentration (Chla) as a measure of primary productivity for the 2013-2014 survey period,
163 which was not available for the earlier studies. We obtained these data from NOAA CoastWatch
164 Program Aqua MODIS satellite images. Since the sperm whales' cephalopod prey are
165 themselves predatory, there is an expected temporal lag of about 3-4 months between primary
166 productivity peaks and increases in cephalopod biomass (see Jaquet 1996; Pirotta et al. 2011).
167 Thus, we considered the monthly Chla concentration averaged over the three months prior to the
168 encounter date. We note that while relSST, sdSST, and Chla reflect processes that affect primary
169 productivity at the surface, these values may not reflect high productivity hundreds of metres
170 below the surface, which is where sperm whale prey is found (Volkov & Moroz 1977; Pierce et
171 al. 2008). However, an association between surface and subsurface waters is suggested by the
172 significant correlation between sperm whale feeding success and surface conditions (Smith &
173 Whitehead 1996). Finally, we used latitude and longitude to account for spatial variation
174 unexplained by oceanographic and topographical variables.

175 We linked values of depth and slope to geographic positions using the raster package in R (R
176 Core Team 2016). We obtained SST and Chla values for each geographic position using the

177 rerddapXtracto R package (Mendelssohn 2016). Topographic and oceanographic variables were
178 extracted at 0.10° resolution, to reflect the distances over which sperm whales could be detected
179 visually and acoustically. Oceanographic variables were weekly averages. In the case of Chla,
180 we used the monthly mean averaged over three months, starting from three months prior to
181 recorded geographic positions. During analysis, we found that models fitted using environmental
182 variables extracted at coarser spatial and temporal scales did not produce substantially different
183 results (Supplement 1)

184 *Modelling differences in habitat use*

185 To examine whether the different clans of sperm whales had different habitat use patterns, we
186 used logistic Generalized Additive Models (GAMs) and Generalized Estimating Equations
187 (GEEs) in which oceanographic and topographic variables were used as predictors of clan
188 identity (following Pirotta et al. 2011). We used GEEs to account for spatiotemporal
189 autocorrelation expected from our continuous method of data collection (Pirotta et al. 2011).
190 This method has previously been used in ecological studies when data were sequentially
191 collected or when measurements were gathered repeatedly from a group of individuals (Dormann
192 et al. 2007, Pirotta et al. 2011, Pirotta et al. 2014, Scott-Hayward et al. 2015). Specifically,
193 sequential data points are grouped into independent blocks and a correlation structure is fitted
194 within blocks (Liang & Zeger 1986). We used a working independence model, which is
195 preferred when the true nature of the correlation is unknown (Liang & Zeger 1986, McDonald
196 1993, Pan 2001). This approach returns more realistic estimates of uncertainty compared with a
197 standard GAM to account for the observed degree of autocorrelation within blocks, but
198 parameter estimates are not affected.

We analyzed data collected in the 1980's and in the 2010's separately, because different clans were sighted during each of these periods (Table 1; see also Cantor et al. 2016): predominantly *Plus-One* and *Regular* in the former; *Short* and *Four-Plus* in the latter. For the 1980's analysis, we included only sightings with *Plus-One* and *Regular* clans as there was only one encounter with each of the *Short* and *Four-Plus* clans over this period (Table 1). We binarized records in each period (i.e. assigning "0" to one clan, "1" to the other). We used individual geographic positions as our unit of analysis and encounters with single clans as the blocking variable, because each encounter represented one group of whales. All locations within each encounter were included within a block. Autocorrelation function (ACF) plots of residuals from individual encounters for the final models (see below) rapidly converged to zero, indicating that encounter was an appropriate blocking variable (Scott-Hayward et al. 2013; See Figs. S1 & S2 in Supplement 2). We tested whether latitude and longitude were best entered as linear terms or cubic spline smooths (see below), while other variables were treated as linear terms, because we assumed that relationships between habitat use and oceanographic and topographic variables would be monotonic.

Habitat use can be influenced by behavioral states in cetacean species (Cañadas & Hammond 2008; Palacios et al. 2013) but we did not include behavioural information in our analyses. Sperm whales have two very distinct behavioural states – they forage for about 75% of the time and socialize during the rest (Whitehead and Weilgart 1991). While socializing, sperm whales tend to move slowly and in more variable directions (Whitehead and Weilgart 1991), so that at the spatial scales of this study (>10 km) positions collected during socializing would not be much different, if at all, from those recorded at the end and beginning of the foraging bouts

221 respectively preceding and following the period of socialising. Therefore, in this case, habitat use
222 records will largely be determined by foraging behaviour.

223 We subsampled or interpolated geographic positions so that they were available approximately
224 every hour and retained only geographic positions collected in areas that were sufficiently
225 surveyed during both study periods (see Supplement 3 for further details). To identify and avoid
226 collinearity, we calculated correlation coefficients for all pairs of explanatory variables (Tables
227 S1-2 in Supplement 4). When variables were collinear ($|r| > 0.4$), we fit alternative initial models
228 that included only uncorrelated variables.

229 *Model selection*

230 To select the most parsimonious combination of uncorrelated variables and the best form (linear
231 or smooth) in which latitude and longitude should be included, we used the quasi-likelihood
232 under independence model criterion (QIC)—an adaptation of Akaike's information criterion
233 (AIC) for GEEs (Pan 2001, Cui & Qian 2007) available in the MuMIn R package (Barton,
234 2016). First, we fitted alternative initial models using uncorrelated predictors, in which latitude
235 and longitude were entered as either linear terms or cubic splines, and then used QIC to select the
236 best shape at which these should be entered. Next, we used backwards stepwise selection to
237 determine which variables to include.

238 We also fitted null models that included only latitude and longitude, aiming to capture variation
239 in relative habitat preferences that could not be accounted for by any of the oceanographic or
240 topographic variables available and investigate the degree to which oceanographic and
241 topographic variables retained in the model improved predictive ability. All explanatory
242 variables were standardised by subtracting the mean and dividing by the standard deviation.

243 *Prediction maps*

244 To examine the spatial distribution of predicted probabilities of encountering a given clan, we
245 produced prediction maps for each study period within areas where whales were found, using the
246 final models (see Supplement 5). We also generated maps of predicted probabilities under the
247 null models for each study period. To identify regions where predictions from the final and the
248 null model differed the most, we generated a mean difference raster. Specifically, for each study
249 period, we obtained the absolute difference between the calculated probabilities generated from
250 the final best model for each year and those calculated through the null model, and averaged
251 annual differences to create a single raster.

252 *Validation*

253 To validate the final models, we analyzed the following three aspects of predictive performance.
254 First, we used goodness of fit (GOF)—a measure of how well the final models fit the data—by
255 generating confusion matrices to assess the models' accuracy in predicting the data used to fit
256 models (Fielding & Bell 1997). To build confusion matrices, we estimated the predicted
257 probability that locations during encounters indicated a given clan. We transformed predicted
258 probability values into a binary assignment using a cut-off that maximized the distance between
259 the Receiver Operating Characteristic (ROC) curve and a 1:1 line using the ROCR package in R
260 (Fielding & Bell 1997, Sing et al. 2005). Second, we used leave-one-out cross validation (LOO)
261 to quantify how accurately a model predicted clan identity for an encounter when that encounter
262 was iteratively removed from the data used to fit the model. In each encounter, we calculated the
263 percentage of geographic positions for which clan identity was correctly assigned (Hastie et al.
264 2009). Finally, we used external cross-validation, i.e. assessed how accurately models predicted
265 clan identity in data that were not used in the model fitting and selection process. We calculated

the accuracy in predicting clan identity for whales found in 1985 for the 1980's models, and for whales found in the western region during 2013 and 2014 for the 2010's models. For each study period, we compared these three aspects of performance of the final models to those of corresponding null models.

## Results

### *1980's period*

We analyzed 596 geographic positions collected between 1987 and 1989. Of these, 168 positions were collected while following the *Plus-One* clan whales and 479 while following *Regular* clan whales. Most encounters occurred in the west and northwest of the archipelago (Fig. 1a), and lasted between an hour and 6 days, averaging 1.6 days (SD = 1.4 days). We fitted two alternative initial models (Table S1 in Supplement 6). Our final model included latitude and longitude as cubic splines and slope and weekly sdSST as linear terms (GOF = 85.2%) (Table S2 in Supplement 6).

Most of the variation among the clans was explained by geographic variables. Whales of the *Plus-One* clan were more likely to be found north of 0.25°N, although uncertainty in predicting clan identity in that region was high (Fig. 2a-i). This is consistent with the observed latitudinal distributions of the *Plus-One* and *Regular* clans north of the Equator, but not with their distributions in the southern limits of the study region where only *Plus-One* clan whales were found (Fig. 3a-i). *Plus-One* whales were also found predominantly in more western waters, but uncertainty in predicting clan identity increased east of the archipelago (91º W; Fig. 2a-ii). This was consistent with the observed distribution of *Plus-One* whales throughout study years, which was restricted to areas west of 91.5º W, and with the distribution of *Regular* clan whales, which

288 occurred throughout the longitudinal range of sperm whale distribution (Fig. 3a-ii). High
289 uncertainty in predicting clan identity in the east likely resulted from the small number of
290 encounters that occurred in that area (Fig. 3a-ii). Although our final model included slope and
291 weekly sdSST (Figs. 2a-iii-iv), response curves did not reflect the observed slope, and sdSST at
292 which the clans were found (Figs. 3a-iii-iv).

293 The predominant effects of geographic variables in differentiating clan identity were also
294 apparent from the similarity between predictive maps generated using the final model and the
295 null model (Figs. 4a-i, ii). These two models predicted identical clan distributions in areas both
296 close to and far from the Galápagos Islands, where there was little spatial overlap among the
297 *Plus-One* and *Regular* clans, but more dissimilar distributions in regions of higher spatial overlap
298 between the clans (Fig. 4a-iii).

299 The inclusion of oceanographic and topographic variables in the final model did not significantly
300 improve the goodness of fit or the average predictive accuracy through LOO cross-validation in
301 comparison to the null model (Fig. 5). Moreover, the inclusion of these variables did not improve
302 the null model's poor ability to predict the clan identity of whales found in 1985 (Fig. 5).

303 *2010's period*

304 Between 2013 and 2014, we analyzed 370 geographic positions to the south of the Galápagos
305 Islands (Fig. 1b). Of these, 226 positions were collected while following the *Short* clan whales
306 and 144 while following *Four-Plus* clan whales. Encounters lasted between 1 hour and 8 days,
307 and averaged 1.3 days (SD = 2.3 days). We fitted six initial candidate models (Table S3 in
308 Supplement 6). The best final model included latitude and longitude as cubic splines, and weekly
309 relSST and sdSST (Table S4 in Supplement 6; GOF = 87 %).

The variation in clan distribution during this period was explained by geographic and oceanographic variables. We found that *Four-Plus* whales were most likely to occur at around 2.2 and 1.8°S, and least likely to occur over latitudinal ranges between these values (Fig. 2b-i). *Four-Plus* whales were also more likely to occur east of 90.5°W, but uncertainty in predicting clan identity was high further west, where there was only one encounter (with *Short* clan whales; Fig. 2b-ii). This predicted geographic distribution reproduced the observed distribution of clans during the 2010's study period (Fig. 3b-i & ii). *Four-Plus* whales were also more likely to occur in areas of higher weekly relSST (Fig. 2b-iii), and lower weekly sdSST (Fig. 2b-iv). The modelled relationships between weekly relSST and sdSST and clan identity were consistent with the oceanographic conditions measured during the 2010's study period (Figs. 3b-iii, iv). However, we note that the relSST mean is skewed towards lower temperatures by an encounter with *Short* clan whales that consistently covered colder waters.

The importance of oceanographic variables in differentiating the habitat of *Four-Plus* and *Short* clans was illustrated by the different prediction maps yielded by the final model and null models (Fig. 4b-i, ii). While both the full and null models generated identical probabilities in the easternmost region where only *Short* clan whales were encountered, they differed greatly over the regions where both clans overlapped (Fig. 4b-iii)

However, while modelled differences in the oceanographic conditions over which *Four-Plus* and *Short* clans occurred were consistent with observed differences in habitat use between *Four-Plus* and *Short* clans, models that included oceanographic variables performed worse in terms of LOO than the null model (Fig. 5b). The same was true regarding performance measured through external cross-validation (Fig. 5b). Further, the performance measured through LOO and external cross-validation of both null and full model was poor overall (<50%; Fig. 5b).

## Discussion

We found that culturally distinct sperm whale clans that are sympatric at the regional scale, around the Galápagos Archipelago, vary considerably in fine-scale habitat use, delineated by spatial partitioning and, to a lesser degree, by oceanographic characteristics. In the 1980's, whales from the *Regular* and *Plus-One* clan used different geographical locations, while in the 2010's, *Four-Plus* and *Short* clan whales used waters with different oceanographic features. In the following sections, we discuss how the sociality of this species may influence its space use patterns via social transmission of habitat preferences and foraging behaviours.

### *Spatial partitioning*

We found sperm whale clans used different areas around the Galápagos Archipelago. In the 1980's *Plus-One* whales were more common in offshore western waters than *Regular* clans whales—consistent with previous findings (Whitehead & Rendell 2004). In the 2010's period, only the *Four-Plus* clan occurred west of the archipelago and, in the southern region, the areas of overlap with the *Short* clan were limited.

Previous analysis has shown that, over days up to a few weeks, areas on the scale at which we can survey from a small vessel are predominantly occupied by groups of whales of a single clan (Whitehead & Rendell 2004). Social units may group to forage together. Individuals may benefit from eavesdropping on group members' echolocation clicks and locate prey more easily, or use other social information on prey location (Whitehead 1989, Whitehead et al. 1991). At daily to weekly scales, we hypothesise that social units could benefit from remaining in an area where other clan members are found and/or avoiding areas dominated by social units of other clans. In this sense, the distribution of sperm whales could be affected by the distributions of fellow clan

355 members as well as by where members of other clans. The reactions of sperm whales to
356 encounters with other clans have not been documented, but active avoidance of members of
357 different cultural entities has been proposed for *transient* and *resident* killer whales (Bigg 1979,
358 Baird & Dill 1995). We note, however, that because these killer whale ecotypes have very
359 different diets, social avoidance could be entangled with different spatial use driven by prey
360 distribution, whereas diet differences are likely much subtler among sperm whale clans
361 (Marcoux et al. 2007), making social avoidance more evident.

362 We found that the spatial partitioning among sperm whale clans over few days and weeks was
363 consistent throughout the months over at least two years. This was most remarkable in the
364 1980's, during which the overall distribution of the clans was maintained despite variation in
365 environmental conditions and sperm whale feeding success between 1987—a strong el Niño
366 year— and 1989—a normal year (Whitehead & Rendell 2004). During the 1987 El Niño,
367 temperatures were 4°C higher than in 1989 (Whitehead & Rendell 2004). Increased temperatures
368 during El Niño events are associated with decreased marine production, which affects the fitness
369 of species across taxa (Trillmich & Dellinger 1991, Boersma 1998, Schaeffer et al. 2008, Wolff
370 et al. 2012). Feeding rates of both *Regular* and *Plus-One* sperm whales were significantly lower
371 in 1987 than in 1989 (Whitehead and Rendell 2004). While there is no direct information on
372 sperm whale prey abundance off the Galápagos Islands, decline in the biomass of the squid
373 *Dosidicus gigas*, an important prey of sperm whales in the region (Clarke et al. 1988, Clarke &
374 Paliza 2001) has been documented across the eastern Pacific during strong El Niño years (Taipe
375 et al. 1991, Markaida 2006). The distribution of clans remained relatively constant across two
376 highly different years, suggesting that site fidelity over the annual temporal scale may be
377 maintained if social units rely on the presence of other clan members as a cue for habitat

selection. Thus, while sperm whale clans are often described as sympatric at a regional scale—for example, around the Galápagos Archipelago, off the Coast of Chile, and in the Caribbean (Gero et al. 2016; Rendell & Whitehead 2003)—spatial partitioning was apparent at a finer spatial scale (less than 10 km).

Studies that span greater temporal and spatial scales indicate however that clan-specific habitat use patterns become diluted. Our study focused on a window of up to three years around the Galápagos and was restricted to the months between January-June, which are mostly representative of the warm season. This represents a snapshot of a female sperm whale's lifespan—60 to 70 years (Rice 1989)—and a portion of the home range of such nomadic animals—at least 2000 km across the Eastern Pacific (Whitehead et al. 2008, Mizroch & Rice 2013, Cantor et al. 2016). But throughout the decades, the clan composition in the Galápagos Islands shifted abruptly from being dominated by the *Regular* and *Plus-One* clans in the 1980's, to the *Regular* clan in the 1990's, and to the *Short* and *Four-Plus* clans in the 2010's (Cantor et al. 2016). This shift may have resulted from movements triggered by environmental changes and fluctuation in prey availability over large scales (Cantor et al. 2016, 2017). Additionally, patterns of habitat use for the same clans in other areas were less discrete (Whitehead & Rendell 2004). Off the Chilean coast in the year 2000, *Regular*, *Short,* and *Plus-One* clans ranges overlapped more than off the Galápagos (Whitehead & Rendell 2004). Movement patterns of *Regular* clan whales off Chile were also significantly more convoluted than those of *Regular* clan whales off the Galápagos (Whitehead & Rendell 2004).

*Oceanographic variation*

Whether oceanographic conditions drive variation in clan space use remains uncertain. During the 1980's, oceanographic variables did not contribute to discriminating the space use of *Plus-*

*One* and *Regular* clans. However, three lines of evidence suggest that oceanic conditions were different in the areas occupied by the *Plus One* and *Regular* clans. First, the relative species composition of sperm whale diet varied regionally, as described by the analysis of fecal samples off the Galápagos Islands (Smith & Whitehead 2000). Second, *Regular* clan whales in this period had a higher carbon-13 isotope signature compared to *Plus-One* clan whales (Marcoux et al. 2007b). Higher C-13 signatures are characteristic of less turbulent habitats, and have been suggested to reflect the difference in oceanic flow conditions between the more inshore habitat of the *Regular* clan and the oceanic habitat of *Plus-One* clan whales (France 1995, Marcoux et al. 2007a). And third, *Regular* and *Plus-One* clan whales had significantly different movement patterns and foraging success rates during this period (Whitehead & Rendell 2004). Thus, different conditions between the areas in which the clans were found could have existed but may have not captured by the oceanographic variables we included in the present analysis. However, it remains uncertain whether observed behavioural differences in *Regular* and *Plus-One* clans were a consequence of different habitat conditions or if these behaviours caused different habitat selection patterns among the clans (Whitehead & Rendell 2004).

In the 2010's, *Four-Plus* clan whales were found in warmer waters and areas of higher variation in SST than *Short* clan whales. These differences may have arisen if these clans were directly tracking different environmental cues to find their prey or if the prey they preferred was found in association with different environmental conditions. Alternatively, these differences might also be a by-product of the spatial segregation described above. In addition, these patterns were described based on a limited number of unevenly represented encounters and models that captured these patterns performed poorly through cross-validation (although they fit well to the

423 data). Thus, our sample may not be sufficient to accurately represent the habitat of the *Short* and
424 *Four-Plus* clans during this period.

425 Some of the uncertainty in characterizing the habitat of the clans arises from the difficulty in
426 measuring sperm whales' habitat accurately, and is further confounded by the lack of detailed
427 information on diving behaviour. Although the oceanographic and topographic variables we used
428 are valid proxies for the distribution of sperm whale prey (Jaquet & Whitehead 1996, Pirotta et
429 al. 2011, Wong & Whitehead 2014), they do not equate to their presence, abundance or quality.
430 Furthermore, our measurements of oceanographic variables describe surface conditions. It is
431 uncertain the degree to which indicators of upwelling or frontal activity at the sea surface
432 represent those in deeper waters, because these features can be displaced or dissipated at greater
433 depths (Jaquet 1996). Our inclusion of mostly surface-level oceanographic variables also likely
434 explains the small contribution that these variables had in predicting clan identity. Recent
435 advances in echosounding technology used to measure composition, biomass, and movements of
436 bathypelagic squid offer a promising way to better characterize the fine-scale habitat of sperm
437 whales (Benoit-Bird et al. 2015, Benoit-Bird et al. 2017). Additionally, we aimed to identify
438 differences in niche traits among the clans but did not evaluate the possibility of niche width
439 varying among the clans, which has been found among killer whale ecotypes (Foote et al. 2009).
440 Thus, our decision to study only linear differences in habitat-use patterns may have restricted our
441 ability to find non-monotonic contrasts in the oceanographic conditions where clans were found.

442 **Conclusions**

443 Our study reveals fine-scale spatial partitioning among clans around the Galápagos Islands that
444 suggests another layer of complexity in the cultural lives of sperm whales. We show that clans
445 differ in fine-scale space use, in addition to vocal repertoire (Rendell & Whitehead 2003),

movement patterns (Whitehead & Rendell 2004), fitness (Marcoux et al. 2007a), diet (Marcoux et al. 2007b) and social behaviour (Cantor & Whitehead 2015). Taken together, these findings suggest the niche of sperm whale clans is constructed on the basis of both social and environmental information, both of which interact over different spatial and temporal scales (see also Boyd and Richerson 1988; Whitehead 2007; van der Post and Hogeweg 2009). The potential ability of sperm whales to balance socially acquired traditions with environmental cues likely plays a part in their ecological success in such a highly dynamic, mesopelagic environment (see also Laland et al. 2000; Whitehead 2007).

To further understand clan-specific niches of sperm whales, future studies should collect spatial data from other regions of the eastern Tropical Pacific and couple them with detailed diving data using tag technologies and direct measurements of prey availability through echosounding devices (Watwood et al. 2006, Benoit-Bird et al. 2015, Benoit-Bird et al. 2017). Combining such large- and fine-scale spatial data will help clarify whether clans have consistently different foraging strategies or if these behaviours are a response to varying environmental conditions.

## Ethics statement

Field procedures for approaching, photographing, and recording sperm whales were approved by the Committee on Laboratory Animals of Dalhousie University.

## Acknowledgements

We are grateful for all volunteer crewmembers for their hard work at sea, to G. Merlen and F. Félix for help with logistics, and to the Ministerio de Defensa Nacional, Ministerio del Ambiente, and Dirección del Parque Nacional Galápagos for research permits. We thank all

those who help processing data in the lab. We also thank Roy Mendelssohn for his help with satellite data, and Ari Friedlaender, Daniel Palacios, Marie Auger-Méthe, and Cindy Staicer for insightful comments on the manuscripts. AE thanks the contribution from the Dalhousie University Faculty of Graduate Studies, Nova Scotia Graduate Scholarship, and the Patrick F. Lett Graduate Students' Assistance Bursary; MC was funded by Conselho Nacional de Desenvolvimento Científico e Tecnológico (202581/2011-0, 153797/2016-9) and the Killam Trusts; LR was supported by the Marine Alliance for Science and Technology for Scotland (MASTs) pooling initiative and their support is gratefully acknowledged. MASTs is funded by the Scottish Funding Council (grant reference HR09011) and contributing institutions; HW was funded by Natural Sciences and Engineering Research Council of Canada, the National Geographic Society, the International Whaling Commission, the Whale and Dolphin Conservation Society, Cetacean Society International and the Green Island Foundation.

**Literature cited**

Allen J, Weinrich M, Hoppitt W, Rendell L (2013) Network-based diffusion analysis reveals cultural transmission of lobtail feeding in humpback whales. Science 340:485–488

Arnbom T (1987) Individual identification of sperm whales. Rep Int Whaling Comm 37:201–204

Baird RW, Dill LM (1995) Occurrence and behaviour of transient killer whales: seasonal and pod-specific variability, foraging behaviour, and prey handling. Can J Zool 73:1300–1311

Barton K (2016) MuMIn: Multi-model inference. R package version 1.15.6. Available at: CRAN.R-project.org/package=MuMIn

490 Benoit-Bird KJ, Moline MA, Southall BL (2015) Deep mapping of teuthivorous whales and their
491 prey fields. Report under contract to the Department of Defense Strategic Environmental
492 Research and Development Program; OMB No. 0704-0188
493 Benoit-Bird KJ, Moline MA, Southall BL (2017) Prey in oceanic sound scattering layers
494 organize to get a little help from their friends. Limnol Oceanogr 73:2788–2798
495 Best PB (1979) Social organization in sperm whales, *Physeter macrocephalus*. In: Norris KS
496 (ed) Behavior of marine animals, Vol. 3. University of California Press, Berkeley
497 Bigg MA (1979) Interaction between pods of killer whales off British Columbia and
498 Washington. In: Third biennial conference on the biology of marine mammals: 3
499 Bivand RS, Keitt T, Rowlingson B (2016). rdgal: bindings for the geospatial data abstraction
500 library. R package version 1.2-5. Available at: CRAN.R-project.org/package=rdgal
501 Bivand RS, Pebesma EJ, Gomez-Rubio V (2013). Applied spatial data analysis with R, Second
502 Edition. Springer, New York
503 Block WM, Brennan LA (1993) The habitat concept in ornithology: theory and applications. In
504 Power DM (ed) Current ornithology, Vol. 11. Plenum Press, New York, p 35-91
505 Boesch C, Marchesi P, Marchesi N, Fruth B, Joulian F (1994) Is nut cracking in wild
506 chimpanzees a cultural behavior? J Hum Evol 26:325–338
507 Bolnick DI, Svanbäck R, Fordyce JA, Yang LH, Davis JM, Hulsey CD, Forister ML (2003) The
508 ecology of individuals: incidence and implications of individual specialization. Am Nat
509 161:1–28
510 Boyd R, Richerson PJ (1988) An evolutionary model of social learning: the effects of spatial and
511 temporal variation. In: Zentall TR, Galef BG (eds) Social learning: psychological and
512 biological perspectives. Lawrence Erlbaum Associates, Hillsdale, p 29–48

513 Boyd R, Richerson PJ (1996) Why culture is common, but cultural evolution is rare. Proc Br
514 Acad 88:77–93
515 Cantor M, Eguiguren A, Merlen G, Whitehead H (2017) Galápagos sperm whales (*Physeter*
516 *macrocephalus*): waxing and waning over three decades. Can J Zool, 95: 645–652
517 Cantor M, Simões-Lopes PC, Daura-Jorge, FG (2018) Spatial consequences for dolphins
518 specialized in foraging with fishermen. Anim Behav 139: 19-27
519 Cantor M, Whitehead H (2015) How does social behavior differ among sperm whale clans? Mar
520 Mamm Sci 31:1275–1290
521 Cantor M, Whitehead H, Gero S, Rendell L (2016) Cultural turnover among Galápagos sperm
522 whales. R Soc Open Sci 3:160615
523 Cañadas A, Hammond P (2008) Abundance and habitat preferences of the short-beaked common
524 dolphin *Delphinus delphis* in the southwestern Mediterranean: implications for
525 conservation. Endanger Species Res 4: 309-331
526 Christal J, Whitehead H, Lettevall E (1998) Sperm whale social units: variation and change. Can
527 J Zool 76:1431–1440
528 Clarke R, Paliza O (2001) The food of sperm whales in the southeast pacific. Mar Mamm Sci
529 17:427–429
530 Clarke R, Paliza O, Aguayo A (1988) Sperm whales of the southeast Pacific, Part IV: Fatness,
531 food and feeding. In: Pilleri G (ed) Investigations on Cetacea, Vol. XXI. Privately
532 published by G. Pilleri, Berne, p 53–195
533 Cui J, Qian G (2007) Selection of working correlation structure and best model in GEE analyses
534 of longitudinal data. Comm Stat Simul Comput 36:987–996

535 Dormann CF, McPherson JM, Araújo MB, Bivand R, Bolliger J, Carl G, Davies RG, Hirzel A,
536 Jetz W, Kissling WD, Kühn I, Ohlemüller R, Peres-Neto PR, Reineking B, Schröder B,
537 Schurr FM, Wilson R (2007) Methods to account for spatial autocorrelation in the
538 analysis of species distributional data: a review. Ecography 30: 609-628
539 Estes JA, Riedman ML, Staedler MM, Tinker MT, Lyon BE (2003) Individual variation in prey
540 selection by sea otters: patterns, causes and implications. J Anim Ecol 72:144–155
541 Fielding AH, Bell JF (1997) A review of methods for the assessment of prediction errors in
542 conservation presence/absence models. Environ Conserv 24:38–49
543 Foote AD, Newton J, Piertney SB, Willerslev E, Gilbert MTP (2009) Ecological, morphological
544 and genetic divergence of sympatric North Atlantic killer whale populations. Mol Ecol
545 18:5207–5217
546 Ford JK, Ellis GM, Barrett-Lennard LG, Morton AB, Palm RS, Balcomb III KC (1998) Dietary
547 specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal
548 British Columbia and adjacent waters. Can J Zool 76:1456–1471
549 France RL (1995) Carbon-13 enrichment in benthic compared to planktonic algae: foodweb
550 implications. Mar Ecol Prog Ser 124:307–312
551 Galef BG (1976) Social transmission of acquired behavior: a discussion of tradition and social
552 learning in vertebrates. Adv Stud Behav 3:77–100
553 Geist V (1971) Sheep society and home range formation. In: Mountain Sheep. The University of
554 Chicago Press, Chicago
555 Gero S, Whitehead H, Rendell L (2016) Individual, unit and vocal clan level identity cues in
556 sperm whale codas. R Soc Open Sci 3:150372
557 Griffin RB (1999) Sperm whale distributions and community ecology associated with a warm-

core ring off Georges Bank. Mar Mamm Sci 15:33–51

Grinell J (1971) The niche-relationships of the California thrasher. Auk 34:427–433

Hall, LS, Krausman PR, Morrison ML (1997) The habitat concept and a plea for standard terminology. Wildl Soc Bull 25:173-182

Hastie T, Tibshirani R, Friedman J (2009) The Elements of Statistical Learning, Second Edition. Springer Science & Business Media, New York

Hijmans RJ (2016) raster: geographic data analysis and modeling. R package version 2.5-8. Available at: CRAN.R.project.org/package=raster

Hutchinson GE (1957) Concluding remarks. Cold Spring Harb Symp Quant Biol 22:415–427

Jaeggi AV., Dunkel LP, Noordwijk MA van, Wich SA, Sura AA, van Schaik CP (2010) Social learning of diet and foraging skills by wild immature Bornean orangutans: implications for culture. Am J Primatol 72:62–71

Jaquet N (1996) How spatial and temporal scales influence understanding of sperm whale distribution: a review. Mammal Rev 26:51–65

Jaquet N, Whitehead H (1996) Scale-dependent correlation of sperm whale distribution with environmental features and productivity in the South Pacific. Mar Ecol Prog Ser 135:1–9

Kanwisher JW, Ridgway SH (1983) The physiological ecology of whales and porpoises. Sci Am 248:110–120

Kawakami T (1980) A review of sperm whale food. Sci Rep Whales Res Inst 32:199–218

Koops K, McGrew WC, Matsuzawa T (2013) Ecology of culture: do environmental factors influence foraging tool use in wild chimpanzees, *Pan troglodytes verus*? Anim Behav 85:175–185

Krützen M, Mann J, Heithaus MR, Connor RC, Bejder L, Sherwin WB (2005) Cultural

transmission of tool use in bottlenose dolphins. Proc Natl Acad Sci U S A 102:8939–43

Laland KN, Galef BG (2009) The question of animal culture. Harvard University Press, Cambridge

Laland KN, Janik VM (2006) The animal cultures debate. Trends Ecol Evol 21:542-547

Laland KN, Odling-Smee J, Feldman MW (2000) Niche construction, biological evolution, and cultural change. Behav Brain Sci 23: 131-175

Leibold MA (1995) The niche concept revisited: mechanistic models and community context. Ecology 76:1371–1382

Liang KY, Zeger SL (1986) Longitudinal data analysis using generalized linear models. Biometrika 73:13–22

Mann J, Patterson EM (2013) Tool use by aquatic animals. Philos Trans R Soc Lond, B 368, 20120424

Mann J, Stanton MA, Patterson EM, Bienenstock EJ, Singh LO (2012) Social networks reveal cultural behaviour in tool-using using dolphins. Nat Commun 3:980

Marcoux M (2005) Vocalizations, diet and fitness among acoustic clans of sperm whales (*Physeter macrocephalus*). PhD dissertation, Dalhousie University, Halifax, NS

Marcoux M, Rendell L, Whitehead H (2007a) Indications of fitness differences among vocal clans of sperm whales. Behav Ecol Sociobiol 61:1093–1098

Marcoux M, Whitehead H, Rendell L (2007b) Sperm whale feeding variation by location, year, social group and clan: evidence from stable isotopes. Mar Ecol Prog Ser 333:309–314

Markaida U (2006) Food and feeding of jumbo squid *Dosidicus gigas* in the Gulf of California and adjacent waters after the 1997-98 El Niño event. Fish Res 79: 16-27

McDonald BW (1993) Estimating logistic regression parameters for bivariate binary data. JR

604 Stat Soc Ser B Stat Methodol 55:629–642
605 McGrew WC, Tutin CEG, Baldwin PJ (1979) Chimpanzees, tools, and termites: cross-cultural
606 comparisons of Senegal, Tanzania, and Rio Muni. J R Anthropol Inst 14:185–214
607 Mizroch SA, Rice DW (2013) Ocean nomads: distribution and movements of sperm whales in
608 the North Pacific shown by whaling data and Discovery marks. Mar Mamm Sci
609 29:136–165
610 Mendelssohn R (2016) rerddapXtracto: extracts environmental data from ERD's ERDDAP web
611 service. R package version 0.1.0. Available at: github.com/rmendels/rerddapXtracto
612 Nigmatullin CM, Nesis KN, Arkhipkin AI (2001) A review of the biology of the jumbo squid
613 *Dosidicus gigas* (Cephalopoda: Ommastrephidae). Fish Res 54: 9-19
614 Ottoni EB, Izar P (2008) Capuchin monkey tool use: Overview and implications. Evol Anthropol
615 17:171–178
616 Palacios D, Baumgartner M, Laidre K, Gregr E (2013) Beyond correlation: integrating
617 environmentally and behaviourally mediated processes in models of marine mammal
618 distributions. Endanger Species Res 22: 191-203
619 Pan W (2001) Akaike's information criterion in generalized estimating equations. Biometrics
620 57:120–125
621 Papastavrou V, Smith SC, Whitehead H (1989) Diving behaviour of the sperm whale, *Physeter*
622 *macrocephalus*, off the Galapagos Islands. Can J Zool 67:839–846
623 Pirotta E, Matthiopoulos J, MacKenzie M, Scott-Hayward L, Rendell L (2011) Modelling sperm
624 whale habitat preference: a novel approach combining transect and follow data. Mar Ecol
625 Prog Ser 436:257–272
626 Pirotta E, Thompson PM, Miller PI, Brookes KL, Cheney B, Barton TR, Graham IM, Lusseau D

627 (2014) Scale-dependent foraging ecology of a marine top predator modelled using
628 passive acoustic data. Funct Ecol 28:206–217
629 Pebesma EJ, Bivand RS (2005) Classes and methods for spatial data in R. R News 5:9-13
630 Praca E, Gannier A, Das K, Laran S (2009) Modelling the habitat suitability of cetaceans:
631 example of the sperm whale in the northwestern Mediterranean Sea. Deep Res I 56:648–
632 657
633 R Development Core Team (20016). R: A language and environment for statistical computing. R
634 Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
635 http://www.R-project.org.
636 Rendell L, Mesnick SL, Dalebout ML, Burtenshaw J, Whitehead H (2011) Can genetic
637 differences explain vocal dialect variation in sperm whales, *Physeter macrocephalus*?
638 Behav Genet 42:332–343
639 Rendell L, Whitehead H (2003) Vocal clans in sperm whales (*Physeter macrocephalus*). Proc
640 R Soc B 270:225–231
641 Rendell L, Whitehead H (2005) Spatial and temporal variation in sperm whale coda
642 vocalizations: stable usage and local dialects. Anim Behav 70:191–198
643 Rice DW (1989) Sperm whale *Physeter macrocephalus* Linnaeus, 1758. In: Ridgway SH,
644 Harrison R (eds) Handbook of marine mammals, Vol 4. Academic Press, London
645 Riesch R, Barrett-Lennard LG, Ellis GM, Ford JKB, Deecke VB (2012) Cultural traditions and
646 the evolution of reproductive isolation: ecological speciation in killer whales? Biol J Linn
647 Soc 106:1–17
648 Roughgarden J (1972) Evolution of niche width. Am Nat 106:683–718
649 Sargeant BL, Mann J (2009) From social learning to culture: intrapopulation variation in

650 bottlenose dolphins. In: Laland KN, Galef B (eds) The Question of Animal Culture.
651 Harvard University Press, Cambridge, p 152–173
652 Scott-Hayward L, Mackenzie ML, Ashe E, Williams R (2015) Modelling killer whale feeding
653 behaviour using a spatially adaptive Complex Region Spatial Smoother (CReSS) and
654 Generalized Estimating Equations (GEEs). J Agric Biol Environ Stat 20: 305-322
655 Scott-Hayward L, Oedekoven C, Mackenzie ML, Walker CG, Rextad E (2013) User guide for
656 the MRSea package: statistical modelling of bird and cetacean distributions in offshore
657 renewables development areas. University of St. Andrews contract for Marine Scotland;
658 SB9 (CR/2012/05)
659 Sing T, Sander O, Beerenwinkel N, Lengauer T (2005) ROCR: Visualizing classifier
660 performance in R. Bioinformatics 21:3940–3941
661 Slagsvold T, Wiebe KL (2007) Learning the ecological niche. Proc R Soc B 274:19–23
662 Slagsvold T, Wiebe KL (2011) Social learning in birds and its role in shaping a foraging niche.
663 Philos Trans R Soc Lond B Biol 366:969–977
664 Smith SC, Whitehead H (2000) The diet of Galápagos sperm whales *Physeter macrocephalus* as
665 indicated by fecal sample analysis. Mar Mamm Sci 16:315–325
666 Svanbäck R, Persson L (2004) Individual diet specialization, niche width and population
667 dynamics: implications for trophic polymorphisms. J Anim Ecol 73:973–982
668 Taipe A, Yamashiro C, Rojas P, Roque C (2001) Distribution and concentrations of jumbo flying
669 squid (*Dosidicus gigas*) off the Peruvian coast between 1991 and 1999. Fish Res 54:21–
670 32
671 van der Post DJ, Hogeweg P (2009) Cultural inheritance and diversification of diet in variable
672 environments. Anim Behav 78:155–166

673 van Schaik CP, Ancrenaz M, Borgen G, Galdikas B, Knott CD, Singleton I, Suzuki A, Utami SS,
674 Merrill M (2003) Orangutan cultures and the evolution of material culture. Science
675 299:102–105
676 Van Valen L (1965) Morphological variation and width of ecological niche. Am Nat 99:377–390
677 Volkov AF, Moroz IF (1977) Oceanological conditions of the distribution of cetacean in the
678 Eastern Tropical part of the Pacific Ocean. Rep Int Whal Comn 27: 186–188
679 Watwood SL, Miller PJO, Johnson M, Madsen PT, Tyack PL (2006) Deep-diving foraging
680 behaviour of sperm whales (*Physeter macrocephalus*). J Anim Ecol 75:814–825
681 Whitehead H (1989) Formations of foraging sperm whales, *Physeter macrocephalus*, off the
682 Galapágos Islands. Can J Zool 67:2131–2140
683 Whitehead H (2003) Sperm Whales, Social Evolution in the Ocean. The University of Chicago
684 Press, London
685 Whitehead H (2007) Learning, climate and the evolution of cultural capacity. J Theor Biol
686 245:341–350
687 Whitehead H, Antunes R, Gero S, Wong SNP, Engelhaupt D, Rendell L (2012) Multilevel
688 societies of female sperm whales (*Physeter macrocephalus*) in the Atlantic and Pacific:
689 why are they so different? Int J Primatol 33:1142–1164
690 Whitehead H, Coakes A, Jaquet N, Lusseau S (2008) Movements of sperm whales in the tropical
691 Pacific. Mar Ecol Prog Ser 361:291–300
692 Whitehead H, Rendell L (2004) Movements, habitat use and feedings success of cultural clans of
693 South Pacific sperm whales. J Anim Ecol 73:190–196
694 Whitehead H, Rendell L (2014) The cultural lives of whales and dolphins. University of Chicago
695 Press, Chicago

696 Whitehead H, Waters S, Lyrholm T (1991) Social organization of female sperm whales and their
697 offspring: constant companions and casual acquaintances. Behav Ecol Sociobiol 29:385–
698 389
699 Whitehead H, Weilgart L (1991) Patterns of visually observable behaviour and vocalizations in
700 groups of female sperm whales. Behaviour 118: 275-296
701 Whiten A, Goodall J, McGrew WC, Nishida T, Reynolds V, Sugiyama Y, Tutin CEG,
702 Wrangham RW, Boesch C (1999) Cultures in chimpanzees. Nature 399:682–685
703 Wong SNP, Whitehead H (2014) Seasonal occurrence of sperm whales (*Physeter*
704 *macrocephalus*) around Kelvin Seamount in the Sargasso Sea in relation to
705 oceanographic processes. Deep Sea Res I 91:10–16
706


## Tables

Table 1. Summary of time spent following female and juvenile sperm whales during the 1980's and 2010's surveys off the Galápagos Islands. Encounters were defined as consecutive geographic positions that were assigned to the same clan and occurred within < 6 hours of each other.

| Year | Surveyed period | Days spent following whales | Encounters with females and immatures[a] | *Regular* clan encounters | *Plus-One* clan encounters | *Short* clan encounters | *Four-Plus* clan encounters |
|---|---|---|---|---|---|---|---|
| 1985[c] | Jan. 18 – Apr. 22 | 29 | 12 | 10 | 1 | 1 | 0 |
| 1987 | Jan. 2 – Jun. 30 | 51 | 21 | 12 | 7 | 1 | 0 |
| 1989 | Apr. 4 – May 22 | 32 | 16 | 10 | 3 | 0 | 1 |
| 2013 (Southern)[b] | Apr. 9 – Apr. 12 | 4 | 9 | 0 | 0 | 3 | 2 |
| 2013[c] (Western)[b] | Jan. 3 –Feb. 21 | 10 | 2 | 0 | 0 | 0 | 2 |
| 2014 (Southern)[b] | Jan. 23 – May 22 | 24 | 11 | 0 | 0 | 2 | 3 |
| 2014[c] (Western)[b] | Jan. 13 – Feb. 10 | 2 | 1 | 0 | 0 | 0 | 1 |
| | Total | 152 | 72 | 32 | 11 | 7 | 9 |

a. Encounter number includes encounters for which clan identity was not assigned, which is why this number does not always equal the sum of encounters with each of the clans
b. Southern regions consist of areas south of 1.3°S and Western regions are north of 1.3°S (Fig. 1)
c. Data from these survey periods were used for external cross-validation only

716 **Figures**




717

718 Figure 1. Geographic positions in (a) 1987 and 1989 of *Plus-One* and *Regular* clan sperm whales, and (b)
719 in 2013 and 2013 of *Four-Plus and Short* clan sperm whales off the Galápagos Islands. The southern
720 region that was included in the 2010's period is delineated by the dashed rectangle. A section of South
721 and Central America is shown for reference





Figure 2. Partial plots of $\log_e$(odds) of female and juvenile sperm whales found off the Galápagos Islands belonging to (a) the *Plus-One* clan in the 1980's study period and (b) the *Four-Plus* clan in the 2010's study period. (a) In the 1980's, clan identity = *Plus-One* is modelled as function of (a-i) latitude, (a-ii) longitude, (a-iii) slope incline, (a-iv) weekly standard deviation of SST (sdSST). (b) In the 2010's, clan identity = *Four-Plus* is modelled as a function of (b-ii) latitude, (b-ii) longitude, (b-iii) weekly relSST, and (b-iv) weekly sdSST. Grey lines represent 95% confidence intervals.






Figure 3. Bean-plots of observed geographic and oceanographic variables by clan; (a) shows the 1980's distribution of variables in which *Plus-One* and *Regular* clan whales were found off the Galápagos Islands: (a-i) latitude, (a-ii) longitude, (a-iii) slope incline, and (a-iv) weekly standard deviation of sea surface temperature (sdSST); (b) shows the 2010's distribution of variables in which *Four-Plus* and *Short* clan sperm whales were found: (b-i) latitude (b-ii) longitude, (b-iii) weekly relSST, and (b-iv) weekly sdSST.




738

739 Figure 4. Predicted probability of sperm whales belonging to different clans off the Galápagos Islands
740 mapped at 0.12° resolution. (a) sperm whales of the *Plus-One* and *Regular* clans in 1987 and1989 as a
741 function of (a-i) a full model, (a-ii) a null model (latitude and longitude only), and (a-iii) absolute
742 difference between the full and null models. (b) sperm whales of the *Short* and *Four-Plus* clans in 2013
743 and 2014 as a function of (b-i) a full model, (b-ii), a null model (latitude and longitude only), and (b-iii)
744 the absolute difference between the full and null models.

745

746

747

748





Figure 5. Predictive accuracy (%) of null models (fit with latitude and longitude only) and full models of clan identity of sperm whales off the Galápagos Islands in the 1980's (1987 and 1989), and 2010's (2013 and 2014). Predictive accuracy was measured through leave-one-out (LOO) and external cross-validation. Standard errors are shown for LOO accuracy.