

Title	Next-generation food research: Use of meta-omic approaches for characterizing microbial communities along the food chain
Authors	Yap, Min;Ercolini, Danilo;Álvarez-Ordóñez, Avelino;O'Toole, Paul W.;O'Sullivan, Orla;Cotter, Paul D.
Publication date	2021-10-22
Original Citation	Yap, M., Ercolini, D., Álvarez-Ordóñez, A., O'Toole, P. W., O'Sullivan, O. and Cotter, P. D. (2021) 'Next-generation food research: Use of meta-omic approaches for characterizing microbial communities along the food chain', Annual Review of Food Science and Technology,13(1), pp. 361-384. doi: 10.1146/annurev-food-052720-010751
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1146/annurev-food-052720-010751
Rights	© 2022, Annual Reviews. All rights reserved. Posted with permission from the Annual Review of Food Science and Technology, Volume 13 © Annual Reviews, http://www.annualreviews.org .
Download date	2025-05-12 19:34:35
Item downloaded from	https://hdl.handle.net/10468/13119

Next-Generation Food Research: Use of Meta-Omic Approaches for Characterizing Microbial Communities Along the Food Chain

Min Yap^{1,2}, Danilo Ercolini^{3,4}, Avelino Álvarez-Ordóñez^{5,6}, Paul W O'Toole^{2,7}, Orla O'Sullivan^{1,7,8}, Paul D. Cotter^{1,7,8*}

¹Teagasc Food Research Centre, Moorepark, Fermoy, County Cork, P61C996, Ireland

²School of Microbiology, University College Cork, County Cork, T12YT20, Ireland

³Department of Agricultural Sciences, University of Naples Federico II, Portici, 80055, Italy

⁴Task Force on Microbiome Studies, University of Naples Federico II, Naples, 80100, Italy

⁵Department of Food Hygiene and Technology, Universidad de León, 24071 León, Spain

⁶Institute of Food Science and Technology, Universidad de León, 24071 León, Spain

⁷APC Microbiome Ireland, University College Cork, County Cork, T12YT20, Ireland

⁸VistaMilk SFI Research Centre, Moorepark, Fermoy, County Cork, P61C996, Ireland

* Corresponding author: paul.cotter@teagasc.ie

Keywords:

food microbiome, food-processing environment, meta-omic approaches, high-throughput sequencing

Abstract

Microorganisms exist along the food chain and impact the quality and safety of foods in both positive and negative ways. Identifying and understanding the behaviour of these microbial communities enables the implementation of preventative or corrective measures in public health and food industry settings. Current culture-dependent microbial analyses are time-consuming and only target specific subsets of microbes. However, the greater use of culture-independent meta-omic approaches has the potential to facilitate a thorough characterisation of the microbial communities along the food chain. Indeed, these methods have shown potential in contributing to outbreak investigation, ensuring food authenticity, assessing the spread of antimicrobial resistance, tracking microbial dynamics during fermentation and processing, and uncovering the factors along the food chain that impact food quality and safety. This review examines the community-based approaches, and particularly the application of sequencing-based meta-omics strategies, for characterizing microbial communities along the food chain.

INTRODUCTION

Microorganisms along the food chain from farm to fork influence food quality and safety. Historically, culture-based techniques have been used extensively to characterise these microbes. However, with the development of molecular methods and high-throughput sequencing technologies, culture-independent techniques have become more relevant to food microbiome analysis. This has resulted in a corresponding shift from the investigation of specific taxa or groups of food microorganisms to a broader community-based analysis (Cocolin and Ercolini 2015). These methods are based on the extraction of nucleic acids, proteins, and/or metabolites, allowing for the detection and characterisation of microbes within an environment without the need for culturing. As the occurrence of and interactions between microorganisms impact the quality and safety of food, a deeper understanding of these processes allows for early interventions to adverse food safety events, ensuring optimal food quality, and identifying the source of desirable or undesirable microorganisms.

Despite being labor-intensive and time-consuming, culture methods are still the methods employed most regularly in the food industry (Dwivedi and Jaykus 2011, Sohler et al. 2014). However, one of the biggest limitations is that these approaches frequently detect only a

fraction of the microbes that are present in the sample as they rely on the isolation and growth of single microbes on culture media whose metabolic and physiological requirements can be reproduced *in vitro*. This not only overlooks the portion of microbes that are viable but not culturable (VBNC) but also fails to consider the relationships within the community of bacteria present in the sample (Cao et al. 2017). Uncultured microorganisms are estimated to account for up to 99% of the microorganisms in many environments, meaning that the use of traditional culture methods causes a gross underestimation of the microbial population (Handelsman 2004). Although this level of underestimation may not be as considerable in food systems, nevertheless, because the microbiomes of food and food processing environments are composed of complex, dynamic microbial communities, meta-omic approaches have the potential to provide a more accurate and greater understanding of these communities.

In this review, the contribution of microorganisms to the quality and safety of food and the traditional approaches to microbial characterisation are briefly described. The main focus of the review is on sequencing-based meta-omic approaches and their contribution to understanding microbial community dynamics in food, food-associated environments and along the food processing microbiome. Other non-sequencing-based meta-omic approaches are also mentioned in brief.

IMPORTANCE OF MICROORGANISMS THROUGH THE FOOD CHAIN

Quality: Flavor, Texture, Fermentation and Spoilage

Food quality is often associated with physical parameters such as pH and moisture content, which can influence the growth and survival of microorganisms within a food and the food chain. Food spoilage is a process or change that renders a product undesirable or unacceptable for consumption, which is impacted by both the food's intrinsic characteristics and the extrinsic environment (Blackburn 2006). It is a complex process, whereby food undergoes biochemical changes, often due to microbial activity according to ecological determinants (Nychas and Panagou 2011). Some common spoilage bacteria include *Pseudomonas* spp., *Shewanella* spp., *Bacillus* spp., *Clostridium* spp., lactic acid bacteria (LAB), and Enterobacteriaceae (Blackburn 2006). Ultimately, different bacteria cause varying quality problems in different types of food, with some examples presented in Table 1.

Furthermore, despite their slower growth rate, yeasts and molds are able to exploit many ecological niches in food systems and can utilise substrates and tolerate extreme conditions that are not possible for bacteria (in't Veld 1996, Petruzzi et al. 2017). Common spoilage yeasts include species of *Zygosaccharomyces* (in high sugar foods), *Saccharomyces* (a cause of gassiness and turbidity in wines), or *Candida* (cause off-flavors in meat and dairy products) and common spoilage molds include *Zygomycetes* (in produce with high water content), *Penicillium* spp. (cause rot in fruits), and *Aspergillus* (in grains, spices and nuts) (Sahu and Bala 2017).

However, microbes can also improve food quality by changing its intrinsic characteristics. This is evident in fermented foods, where the activity of microbes can improve their organoleptic and nutritive qualities in addition to extending shelf life. Fermented food microbes can either be introduced spontaneously (from the raw materials or production or processing environments) or inoculated as starter cultures and, over time, can produce enzymes, volatile compounds, and antimicrobial molecules, such as organic acids, fatty acids, hydrogen peroxide, diacetyl, and bacteriocins, which can help to slow down or prevent the growth of spoilage and pathogenic microbes (Reis et al. 2012). Although the spontaneous introduction of microbes is of specific relevance to this review, here we briefly provide an overview of some of the most important microorganisms in fermented foods in general.

LAB are among the most important microbes in the production of several fermented foods (Hatti-Kaul et al. 2018). This is reflected in the natural adaptation of many LAB to fermented food environments but has been complemented by many years of research to better understand and enhance their contribution to product safety as well as organoleptic, nutritional, and health properties (Leroy and De Vuyst 2004). Different species of LAB have been used in dairy products (cheese and fermented milks), meats (sausage), fish, vegetables (sauerkraut and pickles), soy sauce, cereals (sourdough), and alcoholic beverages (wine) (Leroy and De Vuyst 2004). Another group of bacteria associated with fermentation are acetic acid bacteria, which mainly consist of *Acetobacter* and *Gluconoacetobacter*. This group of bacteria play important roles in coffee, cocoa and vinegar fermentation because of their ability to oxidize carbon substrates (Schwan and Ramos 2014). *Bacillus subtilis* and *Bacillus licheniformis* are important for the industrial-scale fermentation of soybeans as they grow rapidly, resulting in short fermentation times (Schallmey et al. 2004). Yeast can also

play an important role in the production of many fermented foods. *Saccharomyces cerevisiae* is used in alcoholic fermentation, and yeasts are ultimately used in many indigenous fermented foods as they are acid tolerant, able to grow at high temperatures, and are present in many environments (Schwan and Ramos 2014). In Asia, indigenous foods fermented with yeast, such as miso, soy sauce and wines, are commonly consumed (Aidoo et al. 2006).

Safety: Pathogens and Microbial Antagonism

Despite the value of fermented food and other microbes in contributing to food quality and safety, with respect to food safety, microbes are frequently regarded negatively, with foodborne pathogens responsible for foodborne illness and outbreaks across the globe annually. The consumption of contaminated food causes an estimated 4,500 deaths annually in Europe (World Health Organisation 2017). The causative agents of foodborne outbreaks in Europe in 2019 were bacterial pathogens (26.4%), bacterial toxins (19.3%), viruses (10.7%), and parasites and other agents (3.6%), and 40% of reported outbreaks had unknown causative agents (European Food Safety Authority and European Centre for Disease Prevention and Control 2019). Common pathogenic bacteria include *Bacillus cereus*, *Campylobacter jejuni*, *Clostridium botulinum*, *Clostridium perfringens*, *Cronobacter sakazakii*, *Escherichia coli*, *Listeria monocytogenes*, *Salmonella* spp., *Shigella* spp., *Staphylococcus aureus*, *Vibrio* spp., and *Yersinia enterocolitica*. Viruses such as norovirus and hepatitis E as well as parasites, including *Toxoplasma gondii* and *Trichinella spiralis*, are also common causes of outbreaks and have been recently reviewed (Bintsis 2017). There are various ways that pathogenic microorganisms can enter the food chain. They can be inherent to the raw ingredients, or introduced along the processing line via equipment, food handlers or packaging materials, among other routes. Microbial communities can also be present in the form of biofilms, which are microbial communities that adhere to solid surfaces and may contain pathogenic and spoilage species that can persist on surfaces in food-processing facilities (Coughlan et al. 2016). Once attached, these biofilms can be difficult to remove as they are embedded in a polymeric matrix and cells in the biofilm may be resistant to disinfectants or antimicrobials, particularly in mixed-species biofilms (Yuan et al. 2020). Research efforts on control strategies to prevent biofilm formation and remove existing biofilms are ongoing to overcome this challenge in the food industry. Food safety management systems, including hazard analysis and critical control points, and risk assessment principles have been widely implemented to prevent foodborne illnesses and outbreaks and control the spread of

pathogens along the food chain. However, these management systems are reliant on having a thorough understanding of the microorganisms present and the risk they may pose.

Although microbes are often viewed negatively from a food safety perspective, some have been useful in biocontrol or biopreservation. Microbial antagonism has been applied in the food industry through the use of bacteriocins, phages, and more (Jordan et al. 2014). Bacteriocins, which consist of antibacterial peptides, have been used to target spoilage and pathogenic bacteria in food and in turn prolong the shelf life and improve the safety of food (Galvez et al. 2008). Bacteriocins from LAB such as nisin has been approved for use in foods and is most commonly used in foods such as meat, dairy and vegetable products (Jordan et al. 2014). Bacteriophages or phages are virus predators of bacteria that have shown great promise as they are naturally occurring and control for specific pathogenic bacteria without impacting the quality and microbiota of foods (O'Sullivan et al. 2019). Phages have been applied to a range of foods at various stages from farm to fork to eliminate common pathogenic bacteria such as *Campylobacter jejuni*, *Salmonella* spp., *E. coli* O157:H7, and more (Vikram et al. 2020). Additionally, biofilms from some species can aid in improving food safety by outcompeting undesirable bacteria. Some LAB strains were found to exhibit antagonistic properties against unwanted bacteria, act as a natural barrier, and alter biofilm formation of spoilage microbes (Ouali et al. 2014). In food, the microbial community and the interactions between microbes play essential roles in food quality and safety.

TRADITIONAL APPROACHES TO CHARACTERISATION OF MICROBES

As noted above, culture-based assays have historically been used for the detection, enumeration and isolation of viable foodborne pathogens or spoilage microbes in food and environmental samples (Dwivedi and Jaykus 2011). In general, samples are first homogenized and then often undergo enrichment steps (pre-enrichment and selective enrichment), followed by selective or differential plating to distinguish from other microbes present and, finally, confirmation with biochemical, serological, or other methods. Pre-enrichment is used to recover injured cells and dilute inhibitory compounds in food samples, whereas selective enrichment increases the concentration of a target pathogen while suppressing the growth of other microflora (Dwivedi and Jaykus 2011). These conventional methods are inexpensive but time consuming and labour-intensive and can take from 2-3 days to a week for inoculation, isolation and confirmation depending on the targeted

microorganism (Mandal et al. 2011). Furthermore, owing to the possible presence of VBNC bacteria, false negatives may occur, which could mean the unsuccessful detection of pathogens or spoilage bacteria in food.

With the need for more timely detection of bacteria for foodborne outbreak response, rapid methods that replace conventional plating steps with faster immunology or molecular-based approaches have been investigated in depth and adopted more widely in recent years (Wang and Salazar 2016). High levels of sensitivity and specificity are needed for food pathogen detection (Feng 2007). Immunology-based methods like enzyme-linked immunosorbent assay (ELISA) based on the specific binding of antigens with antibodies, have shown potential but their lack of sensitivity and relatively high limit of detection [$10^3 - 10^5$ colony forming units (CFUs)/mL] has meant that enrichment is generally first required before detection. Nevertheless, when ELISA is coupled with nanotechnology, its application for food analysis has achieved greater sensitivity, specificity and stability (Wu et al. 2019).

Compared to traditional culture methods, nucleic acid-based techniques such as polymerase chain reaction (PCR), quantitative PCR (qPCR) and loop-mediated isothermal amplification (LAMP) require shorter time and through multiplexing, can facilitate concurrent detection, real-time monitoring and quantification of multiple microbial targets (Liu et al. 2017, Tao et al. 2020). Development and optimization of each assay is important, as complex food matrices may hinder nucleic acid extraction and contain inhibitors that may interfere with reactions in the assay. Particularly when multiplexing, primer design is crucial, as primer sets require similar annealing temperatures for a successful assay (Wang and Salazar 2016). Unfortunately, these methods, like immunology-based assays, often still require enrichment or concentration steps because of their limit of detection ($10^3 - 10^4$ CFUs/ mL), and as pathogens are often in low concentrations in foods, direct detection is difficult (Ceuppens et al. 2014, Wang and Salazar 2016). Another rapid method used largely in clinical settings is matrix assisted laser desorption ionization-time-of-flight mass spectrometry (MALDI-TOF MS), which analyzes signals from ribosomal proteins after ionization and time of flight detection that are distinctive for each strain, allowing for rapid microbial identification (de Koster and Brul 2016). Although these rapid methods are generally preferable to culture-based approaches, thorough validation is required by industry and regulatory bodies before routine adoption.

Although culture-based and rapid methods are useful in identifying microbes in complex food- or food environment-related samples for public health and commercial purposes, they create an unbalanced emphasis on specific microorganisms and, despite multiplexing, still only capture a small percentage of the microbial community as a whole (Fleet 1999). As microorganisms exist in communities, it is important to study them as such because the growth, survival, and activity of one species or strain may impact or be associated with the presence of another. Furthermore, from a practical perspective, approaches that could theoretically allow the simultaneous identification of all pathogens and spoilage microbes in a sample from the food chain could have a disruptive positive influence.

META-OMIC APPROACHES: COMMUNITY APPROACHES

Advances in technologies have provided the opportunity for faster and superior characterisation of food chain microbiomes, with a shift toward replacing or supplementing culture-dependent methods with culture-independent, molecular-based methods (Sohier et al. 2014). Whole-genome sequencing has successfully complemented culture-dependent methods by providing deeper discrimination of microbial strains than previous typing methods, with regulatory bodies in the United States and Europe now including it as a tool for pathogen typing and antimicrobial resistance (AMR) surveillance (Rantsiou et al. 2018). In *E. coli* O157:H7 outbreak investigations, genome sequencing stood out from other typing methods, providing insights that enabled improved epidemiological case and cluster identification, geographical origin tracking, and information of potential emerging strains (Jenkins et al. 2019).

In contrast, culture-independent methods including the use of DNA sequencing technologies, have enabled identification and characterisation of multiple microbes in foods or along the food chain at the same time while also bypassing the need to culture microbes (Cocolin and Ercolini 2015). Some current community-based approaches are shown in Figure 1.

Sequencing-Based Meta-Omic Approaches: Metagenetics, Metagenomics and Metatranscriptomics

Metagenetics, also known as amplicon sequencing, metataxonomics, metabarcoding and sometimes, 16S metagenomics or 16S rRNA gene sequencing, is a targeted approach that

involves the amplification of marker genes from mixed genomic DNA by PCR, followed by direct sequencing and alignment against a reference database to identify the taxonomic composition of whole microbial communities (Franzosa et al. 2015). The 16S ribosomal RNA (rRNA) gene is most frequently used in the identification of bacteria, as it is universally found in bacteria, and the gene contains nine hypervariable regions, some or all of which can be targeted through amplification and sequencing to identify the corresponding bacterial taxonomy. A similar approach can be applied to fungi, through targeting the 18S or 23S rRNA genes or the internal transcribed spacer (ITS) regions of the rRNA operon.

Shotgun metagenomics, commonly referred to as metagenomics, is the untargeted genomic analysis of a population of microorganisms by sequencing the entire DNA sample extracted from a mixed microbial community (Quince et al. 2017). This method involves fragmentation of the sample DNA, followed by preparation of a library that is sequenced, with the resulting data analysed to provide information on both taxonomic composition and functional potential of the entire microbial community. Due to the untargeted nature of metagenomics, information relating to all categories of microbes, including bacteria, viruses, archaea, and single-celled eukaryotes like fungi, can be derived from the sample (Quince et al. 2017), assuming the DNA extraction method is appropriate. The lack of an amplification step removes the bias that metagenetics may have and has greater sensitivity, enabling taxonomic classification up to the strain level. Another advantage of shotgun metagenomics is the potential for the recovery, if sufficient sequencing depth is applied, of metagenome-assembled genomes (MAGs), which can provide more genomic information, revealing functional and safety-related properties of specific taxa (Bowers et al. 2017), and allow for the investigation of strain-level diversity in food-related microbial species such as LAB (Pasolli et al. 2020).

Metatranscriptomics relates to the untargeted sequencing of total mRNA isolated from a sample, which allows for the identification of transcriptionally active microbes in the sample, and may provide further insights into the potential functional characteristics of the microbial community. This approach reveals the microbes that are viable and, indeed, most active within a community while also enabling a deeper understanding of how microbial communities in complex food microbiomes or food-related environments interact with each other. This approach can also be used to look at *in situ* gene expression in food, collecting information on the metabolic activities potentially related to food fermentation and/or

spoilage that are currently expressed in a food ecosystem. An additional advantage of metatranscriptomics is the ability to detect RNA-based viruses, including foodborne pathogens such as norovirus (Lewis et al. 2020).

Other Meta-Omic Approaches: Metaproteomics and Meta-metabolomics

Other non-sequencing community-based methods, i.e., metaproteomics and meta-metabolomics, also have the potential to be used in food microbiome studies. Metaproteomics is the large-scale study of the entire protein complement produced by microbial communities within a sample at a given time point, which can aid in linking genomic and transcriptomic data to biological function, deepening the understanding of phenotypic changes as conditions change (Soggiu et al. 2016). Metaproteomics provides information on the microbial communities and their abundances and functions, the interactions within the community, the changes in community metabolism and physiology, and the substrate utilization, carbon sources and assimilation pathways of the microbes in the sample (Kleiner 2019). Mass spectrometry is used for metaproteomics and its application in the food microbiome has mainly been in the characterisation of fermented foods such as fermented soybean and cheese (Soggiu et al. 2016, Xie et al. 2019). Meta-metabolomics, however, involves the use of chemistry, biochemistry, and bioinformatics to detect and analyse small weight metabolites in samples and provide insights into microbial phenotypic characteristics. There are two categories of meta-metabolomic analyses: untargeted, which focuses on the detection of as many groups of metabolites as possible, and targeted, which focuses on a specific user-selected group of metabolites under determined conditions (Li et al. 2020). Researchers in this field typically use either mass spectrometry or nuclear magnetic resonance to evaluate food ingredients, quality, safety, authenticity, and traceability (Kim et al. 2016). The integration of metaproteomics and meta-metabolomics with other omic approaches has been used to provide novel insights and link genomic information with phenotypes (Kim et al. 2016, Pinu et al. 2019).

Current Challenges in Sequencing-Based Meta-Omic Approaches

Despite their promise, these sequencing-based approaches have challenges to overcome to achieve wider application. The major challenge for these meta-omic approaches is the lack of standardization, causing variation in results because of the use of different extraction methods, sequencing platforms, databases, and bioinformatics tools. To highlight this point,

we refer to several studies that have found differing conclusions depending on the approaches taken for analysis, thereby highlighting the need to identify those that provide the greatest accuracy and, ultimately, their use in a standardized manner (Lewis et al. 2020, McHugh et al. 2021, Walsh et al. 2018, Yang et al. 2020).

Similarly in terms of analysis, the fact that results are typically presented in terms of relative abundances may lead to misinterpretations, as an increase in the relative abundance of one taxon results in the concurrent decrease of others. For this reason, it is necessary to quantify microbial communities by using complementary methods and efforts to do so have included digital PCR, qPCR, flow cytometry and culture. It is notable that the use of synthetic standards in sequencing has produced varying results for different methods, again highlighting a need for caution during analysis (Galazzo et al. 2020).

Furthermore, for metagenetics, there is a difficulty when endeavoring to compare outcomes across different studies arising because of a lack of consistency relating to the hypervariable region of the 16S rRNA gene targeted (Claesson et al. 2010). Additionally, the focus on one marker gene can cause other issues. In particular, the operon copy number for 16S rRNA genes differs across taxa, which may inadvertently affect quantitative estimation. Single-copy target genes like *recA*, *rpoB*, and *gyrB* have been suggested as alternatives, but their use is limited because of their relevance to specific taxa of bacteria only and/or the absence of databases that are sufficiently populated (Ogier et al. 2019, Poirier et al. 2018)

From the perspective of methodology, extracting DNA/RNA of sufficient concentration and quality is essential for sequencing, which may be challenging in some circumstances, such as from environmental swab samples taken from areas with a low microbial load (De Filippis et al. 2020). Amplification methods such as multiple displacement amplification (MDA) have been applied to generate DNA of sufficient quantities and the inclusion of controls have been investigated to reduce contamination, but these methods can lead to biases (Marine et al. 2014, McHugh et al. 2021).

For both standard metagenetic and metagenomic sequencing, there is no differentiation between DNA extracted from living and dead organisms within a microbiome, which is of key importance with respect to food or food environment microbiome studies. Propidium monoazide (PMA) and the previously more commonly used ethidium monoazide (EMA) are

DNA-binding dyes that selectively bind to accessible DNA present in the matrix, essentially binding to DNA from dead bacteria and other cells and preventing its amplification during library preparation (Nocker et al. 2006). Treatment with PMA before DNA extraction thereby selects for the subsequent sequencing of DNA from viable cells. The successful application of PMA with sequencing allowed the selective analysis of viable cells during milk processing and cheese manufacturing (Erkus et al. 2016, Kable et al. 2019). However, its performance can be influenced by the microbial community and sample biomass (Wang et al. 2021). Research on the application of these dyes with the use of internal standards could provide insights that may allow for quantification of live and dead cells, and optimization of this treatment in different food or environment matrices. Another option is the sequencing of RNA in place of or alongside DNA, as measuring RNA copies targets the active microbial fraction, which allows the differentiation of viable and non-viable microbes (Mira Miralles et al. 2019), although the instability of mRNA can again provide challenges. Further research on the discrimination of live and dead cells is required, particularly for the application of these sequencing-based approaches for food-related samples.

It is also important to note that for metagenetics in particular, the short reads generated by some sequencing platforms, such as those developed by the market leader, Illumina, can be limiting with respect to assigning taxonomy at the species level. Other sequencing platforms that produce longer reads, such as those from Oxford Nanopore Technologies (ONT) or Pacific Biosciences, may address this but lower read accuracy and higher sequencing costs can be issues.

Although shotgun metagenomic sequencing overcomes many of the biases associated within amplicon-based approaches, one of its biggest challenge is the reduced microbial sequencing depth that occurs when randomly sequencing samples that contain high amounts of host DNA. Although most studies remove the reads from host DNA during bioinformatic analysis, a more efficient alternative is to deplete host DNA or enrich microbial DNA through various chemical methods and commercially available kits (Marotz et al. 2018, Yap et al. 2020). Similarly in metatranscriptomics, highly abundant rRNA can result in increasing costs and complex downstream analysis. To overcome this challenge, rRNA depletion or mRNA enrichment strategies before sequencing and/or post-sequencing removal during downstream analysis have been adopted (Shakya et al. 2019).

Additionally, with the use of sequencing technologies, advanced computational power and bioinformatics skills are necessary for their use, which add to the challenges when considering the application of these approaches.

APPLICATIONS OF SEQUENCING-BASED META-OMIC APPROACHES ALONG THE FOOD CHAIN

Public Health Applications

The sequencing-based meta-omic approaches mentioned above have contributed significantly to the study of various diverse microbiomes, including by facilitating significant advances in food microbiome research. From a public health perspective, these meta-omic approaches have provided insights relating to pathogen detection, outbreak investigation, AMR determination and food authenticity and source tracking.

Pathogen detection and outbreak investigation.

Meta-omic approaches are advantageous, as they bypass the need for culturing and enrichment of pathogens from samples before identification and characterisation of putative etiological agents. They also are able to reveal the presence of uncultured and hard to culture microbes, which may be useful in surveillance, source attribution, risk assessment and epidemiological analysis when traditional methods fall short (EFSA Panel on Biological Hazards et al. 2019). Both metagenetic and metagenomic approaches have enabled the detection and characterisation of pathogens in various foods, including vegetables, meat, and dairy products (Aw et al. 2016, McHugh et al. 2018, Mira Miralles et al. 2019, Yang et al. 2016). Metatranscriptomics, although less widely applied because of the challenges in RNA isolation, also has great potential for identifying viable pathogens in food (Yang et al. 2020).

Use of metagenomic approaches can extend beyond the food chain, where metagenomic sequencing of patient stool samples collected during the outbreak in Germany of STEC (Shiga toxin-producing *E. coli*) O104:H4 assisted the recovery of genomes of the outbreak strain (Loman et al. 2013). Moreover, metagenomics is useful when a viral agent is the cause of the outbreak or, in the case of multi strain outbreaks, it is able to discriminate and characterise several strains, allowing them to be distinguished considerably faster than traditional culture-based methods (Buytaers et al. 2020). Compared to metagenetics, which

may be more useful for low biomass samples because of the amplification of the target, metagenomics facilitates more sensitive characterisation to the species level and further investigation of the functional potential of microbes present (Grützke et al. 2019).

Despite the potential of these meta-omic approaches, they are currently not widely used. One reason is the lack of harmonized methods and standardized, accredited workflows/pipelines that would allow consistent detection and characterisation of outbreak-causing agents (EFSA Panel on Biological Hazards et al. 2019). However, the usefulness of metagenomic analyses can be enhanced when they are complemented with further quantitative molecular assays, highlighting their effectiveness in determining pathogen contamination or outbreak events. A big technical challenge that hinders greater adoption of meta-omic techniques as a routine screening tool for pathogens is that these techniques are not always sufficiently sensitive (Leonard et al. 2015, Lewis et al. 2020). With low numbers of pathogenic cells in samples, substantial sequencing depth is required, particularly for shotgun sequencing, as samples contain DNA from other microbes or contaminants such as animal, plant or human DNA (Yang et al. 2016). With sufficient sequencing depth, shotgun metagenomics can be a faster and more valuable tool that provides more information than current conventional workflows, which permit linking food/environment outbreak-related samples with clinical samples (Buytaers et al. 2020, Grützke et al. 2019, Li et al. 2020). Although the complexity of various food matrices can be a challenge, this is not as great an issue for less biologically complex matrices, such as water used in food production or some minimally processed foods (Fernandez-Cassi et al. 2017).

Identification of antimicrobial resistance-encoding genes.

Over the past decades, AMR has been identified as a serious public health threat and because of this, more tools have been published for the detection of genetic determinants of AMR from sequencing data. Although whole-genome sequencing of cultured isolates is usually utilized, metagenomic sequencing shows great potential for monitoring AMR, as it has outperformed culture-based methods in quantifying resistance in swine herds (Munk et al. 2017). Shotgun sequencing has shown success in the monitoring of AMR genes in the environment from farm to slaughter (Noyes et al. 2016, Pitta et al. 2016). It has also been used to understand the association between antimicrobial use and resistance and the effect of processing on the resistome and virulome (Campos Calero et al. 2018, Mencía-Ares et al. 2020, Van Gompel et al. 2019).

As with other metagenomic approaches, sequencing depth and the presence of host DNA should be considered, as they have been found to affect resistome profiling in environmental and food samples (Gweon et al. 2019, Rubiola et al. 2020). Other challenges include the difficulty in assigning ARG to their host species or strains, which may be addressed by sequencing with long-read technology and the choice of reference resistance gene database, where differences were found between gene variants from the same reference sequence from different databases, reiterating the need for comprehensive databases and standardized workflows (Doyle et al. 2020, Slizovskiy et al. 2020). It is also important to note that the AMR data may not always be phenotypically relevant, as these genes might not be expressed or the choice of bioinformatic tools can result in false positives or negatives (Doyle et al. 2020). From the perspective of gene expression, metatranscriptomics can potentially be employed to complement the analysis (Wang et al. 2020). The analysis of the mobilome (all mobile genetic elements of the microbiome) has also been paired with resistome analysis to understand the potential spread of AMR genes and virulence factors through horizontal gene transfer (Slizovskiy et al. 2020).

Food authenticity.

Food fraud is a global issue that has many consequences, including possible health risks, economic losses, and hindering sustainability efforts. Metabarcoding has been used to determine the authenticity and origin of honey, traditional Chinese medicines, fish, and more (Carvalho et al. 2017, Coghlan et al. 2012, Khansaritoreh et al. 2020, Liu et al. 2020). The basic concept is that the microbiome associated with a traditional food is closely linked to the geographical origin and mode of production of the food as the microbes are typical of raw materials and environment. Although there have been some successes, there are challenges associated with using microbiomes as a means of determining the provenance of food. These include the need for the existence of databases containing the components of the expected microbiome of the food and the potential alteration of the microbiome due to storage or processing conditions (Liu et al. 2020). Similar to other meta-omic applications, the reliance on the completeness of reference databases together with the accuracy of food matrix authentication are important to avoid inaccurate conclusions. Haiminen et al. (2019) found both DNA and RNA shotgun sequencing to be accurate untargeted methods for food authentication and contaminant detection, which has been applied by Kamilari et al. (2019) to

characterise Protected Designation of Origin (PDO) cheeses with complementary metabolomics to define product origin differentiating factors.

Other public health-related fields.

Besides the food industry, other fields have also found benefits in the application of community-based microbiome analysis methods. Community-based approaches have contributed to the increasing knowledge of the indigenous microbial community and AMR patterns in both healthcare settings and water systems that have provided evidence for the greater need for surveillance (King et al. 2016, O'Hara et al. 2017, Zhang et al. 2017). In hospital settings, meta-omic approaches have provided clues to the routes of entry and relationships between pathogens and non-pathogens, as well as helped in environmental surveillance to fight hospital-acquired infections and AMR (Comar et al. 2019, Rampelotto et al. 2019). Similarly, when supplemented with other techniques, shotgun metagenomics was effective in uncovering the presence of virulence factors and novel biomarkers of pathogen-related species in drinking water distribution systems (Zhang et al. 2017). Additionally, on an international scale, urban sewage and waste from aircraft flights have been cited as economically and ethically acceptable approaches for continuous global surveillance and prediction of AMR using metagenomics (Hendriksen et al. 2019, Petersen et al. 2015).

Food Industry Applications

Microbial communities exist throughout the food chain and understanding their dynamics and the conditions that promote or hinder their growth would be useful for food safety and quality purposes. Research efforts using meta-omic approaches have looked into foods, food-associated environments, and food-processing steps, as presented in Figure 2, which are elaborated in the following sections.

Foods: fermented and non-fermented.

One of the main applications of community-based approaches is in the study of fermented foods. Previous reviews noted that most of the early studies on fermented foods employed metagenetics to monitor the activity of microorganisms during fermentation (De Filippis et al. 2017). In recent years, more studies have utilized metagenomics and metatranscriptomics to understand the changes in microbial community diversity and activity during fermentation in a broad range of foods, including vegetables, cheeses, and more (De Filippis et al. 2016,

Duru et al. 2018, Jung et al. 2013, Kim et al. 2020, Liu et al. 2020, Pham et al. 2019, Xiao et al. 2020). Metatranscriptomic analysis revealed the changes in gene expression and metabolic properties of LAB during fermentation of vegetables (Jung et al. 2013, Xiao et al. 2020). Likewise from metatranscriptomic analysis of cheese, metabolic interactions within the microbial community, and temperature-driven functional changes during ripening were revealed (De Filippis et al. 2016, Pham et al. 2019). The use of both metagenomic and metatranscriptomic analyses allowed for the detection of active microbes during fermentation and of microbes responsible for biogenic amine production in fermented soy products (Kim et al. 2020, Liu et al. 2020). This parallel approach was also useful in understanding the dynamics of the microbial community during ripening, revealing the impact of temperature on the microbial community and genes expressed (Dugat-Bony et al. 2015, Duru et al. 2018). These are selected examples of studies within the continuously growing pool of research that employ these methods to study the microbial consortia in fermented foods. Unsurprisingly, it has been suggested that multiple meta-omic approaches facilitate the improved, efficient, and sustainable production of fermented foods through detailed functional characterisation of their microbiomes (Chen et al. 2017).

Although the number of studies using meta-omic approaches to study non-fermented foods is considerably lower than that of fermented foods, those that have been completed highlight the great potential of such approaches. Most of these applications have related to the characterisation of food-associated environments or food-processing steps, which are elaborated in the following sections. Other than those studies, there have been promising studies involving the use of community-based methods to screen for spoilage or pathogenic microorganisms. However, because of the complex nature of food samples and the frequently low pathogen abundances, direct sequencing of DNA or RNA of food has, to date, been found to be less sensitive than conventional culture-based or amplicon-based methods (Lewis et al. 2020, Yang et al. 2020). It should also be noted that, even though both short and long-read sequencing technologies have shown promise with respect to accurate classification of microbes to the family and genus levels, not all approaches sufficiently classify to the species or strain level needed for pathogen detection (Grützke et al. 2019). This is sometimes a significant limitation, especially in terms of food safety, where identifying at only the genus level may not be informative enough to understand the actual species present that could cause food safety or quality issues along the food chain. The need for sensitive and specific tests coupled with other challenges prove that these community-based approaches are currently not

applicable at the regulatory compliance level, but with further development that will be the standard be in the future (Yang et al. 2016).

Food-associated environments.

Food-associated environments, from farm to processing facility, have repeatedly been found to impact, both positively and negatively, the final product microbiome.

Environmental factors.

Microorganisms can enter the food chain at a number of different points. This includes the crops and animals from which the foods are sourced/derived as well as environmental factors such as soil, water, farming systems, pests, and climate conditions. Meta-omic approaches have found that factors such as pasture systems, animal housing, airborne dust, irrigation water, and several others can influence the microbiota diversity and composition of food (Allard et al. 2019, Doyle et al. 2017, Wu et al. 2019). Besides diversity and composition, meta-omic approaches used to characterise the resistome reveal that farm environments are potential vehicles for AMR bacteria and genes, originating from dust and animal feces that contribute to AMR spread and worker exposure (Luiken et al. 2020, Noyes et al. 2016). The use of animal waste as fertilizer (manure/wastewater) can also cause the dissemination of AMR bacteria and genes in the environment, which in turn affect the microbiota of crops grown or animals raised on the land (Allard et al. 2019, He et al. 2019). Seasonality is another contributing factor to the microbiota of the animal and plant environment. Seasonal impacts were evident in certain products, like milk and beef, where the use of metagenetics and metagenomics has revealed seasonal variations in the microbiota of final products (Hwang et al. 2020, Kable et al. 2016, McHugh et al. 2020).

Food-processing environments.

Meta-omic techniques have been adopted in the characterisation of several environments involved in the processing of foods such as meat (De Filippis et al. 2013, Hultman et al. 2015, Stellato et al. 2016), dairy (Anvarian et al. 2016, Doyle et al. 2017, Kable et al. 2016), and alcoholic beverages (Bokulich et al. 2015, Bokulich et al. 2013, Wang et al. 2018). One key observation from using meta-omic approaches for such studies is the presence of a resident microbiome that persists within the processing environments and has the potential to affect final food product quality and safety. This was highlighted in a recent review relating

to the use of high-throughput sequencing to characterise the dominant taxa found in both processing environments and food products, which summarized the evidence that the processing environment can act as a reservoir and source of microbial transfer to food (De Filippis et al. 2020). This can be both beneficial and detrimental, with, for example, beneficial effects apparent in fermented food production. In this regard, microbes in the environment were found to contribute positively to the production of fermented vegetables, wine, and Chinese liquor (Bokulich et al. 2013, Einson et al. 2018, Wang et al. 2018). In contrast, spoilage or pathogenic microorganisms have been found on surfaces of various dairy-, meat- and vegetable-processing facilities using different meta-omic approaches (Hultman et al. 2015, McHugh et al. 2020, Pothakos et al. 2015, Stellato et al. 2016, Zwirzitz et al. 2020). For example, *Pseudomonas* spp. was found in drain biofilms in cheese- and salmon-processing plants (Dzieciol et al. 2016, Langsrud et al. 2016) and pathogens like *Staphylococcus* and *Yersinia* were found on surfaces in milk- and meat-processing plants (Hultman et al. 2015, Kable et al. 2019). Indeed, correlation of microbial communities in biofilms, as determined by metagenetics, with environmental factors has been used to track persistence over time, showing that bacterial communities were location-specific in meat- and fish-processing plants (Rodríguez-López et al. 2020). Additionally, microbial co-occurrences of pathogens with other microbes and microbial interactions within complex ecosystems can be evaluated through meta-omic approaches, which may determine patterns that favour or prevent the growth or survival of foodborne pathogens (den Besten et al. 2018, Illegheems et al. 2015). This was investigated through 16S rRNA sequencing that examined interactions between *Listeria* spp. and the microbiome within a food production facility, and identified species that acted as apparent protagonists or antagonists that had impacts on the presence of *L. monocytogenes* within the processing plant (Fox et al. 2014).

Handling can be a potential source of contamination or microbial transfer, whereby microbes can be unknowingly transferred from surfaces to the food product. *Moraxella* spp., a prominent meat-spoilage bacteria, was found on gloves of employees, which were identified as a potential source of contamination using full-length 16S rRNA gene sequencing throughout a pork-processing plant (Zwirzitz et al. 2020). Similarly, handling was identified as a catalyst in the proliferation of spoilage bacteria in beef products after high-throughput sequencing uncovered the origin of spoilage-associated bacteria from carcasses and their persistence in the environment (De Filippis et al. 2013).

Food-processing steps.

Using meta-omic approaches to monitor the changes in food microbiomes during food processing has been useful in understanding the impact of processes on the quality and safety of foods. This has been studied through two approaches. One approach has involved profiling the entire food-processing chain, where samples were taken from the start to the end of the process and meta-omic methods were used to track the changes in microbial community dynamics, which can facilitate the generation of mitigation measures. This whole-chain approach often involves sampling of both food and environmental samples and has highlighted areas where contamination or spoilage can potentially occur; e.g., in meat processing, animal carcasses or hides were identified as possible sources of contamination and measures taken during and after slaughter were found to be key in reducing bacterial load and transmission of AMR genes to meat products (Calero et al. 2020, De Filippis et al. 2013, Noyes et al. 2016, Yang et al. 2016). A similar approach to studying sausage production showed that the emulsification step selected for gram-positive spoilage bacteria (Hultman et al. 2015). Other investigations have highlighted the impact of storage, low temperatures, and equipment on the milk microbiota in dairy processing (Falardeau et al. 2019, Kable et al. 2016, McHugh et al. 2020), whereas in breweries, food contact surfaces were noted as areas that could allow transmission of spoilage bacteria or genes (Bokulich et al. 2015).

The other approach that has been taken when using meta-omic methodologies is process focused, where specific processing steps that are often considered critical points in food safety management systems are examined. Processes such as heat treatment, cold storage, packaging, cleaning, and others have been studied to understand the microbial dynamics during these processes and ensure their efficacy at eliminating or reducing growth of bacteria. Metagenetics used to investigate heat treatments unsurprisingly found a reduction of bacterial abundance and diversity in meatballs and cheese but it also affected the quality of the final products (Kamilari et al. 2020, Li et al. 2021). Similarly, monitoring the ripening processes of cheese using metagenomics and metatranscriptomics has provided a better understanding of the temperature-driven differences in flavor development (De Filippis et al. 2016, Duru et al. 2018), whereas metagenetics, proteomics and complementary physicochemical and sensory analysis revealed the efficacy of high-pressure processing in improving the quality and shelf life of fish fillets and led to the identification of quality markers for further study (Tsironi et al. 2019). For storage in particular, metagenetic and metagenomic analysis revealed cold temperature storage is an area along the processing chain that allowed for the proliferation

and dominance of certain psychrotrophic spoilage microorganisms in meat and dairy (McHugh et al. 2020, Stellato et al. 2016). Monitoring microbial dynamics to understand the effect of storage temperature on the microbial community has been performed using metagenetics coupled with sensory assessment or culture-dependent methods in sausage and fish, which has resulted in the development of models to infer spoilage dynamics and associations of bacterial species during storage (Benson et al. 2014, Zotta et al. 2019). Modified atmosphere packaging (MAP) is currently used to extend the shelf life of various foods like fresh and processed meat and seafood, fruit and vegetables, but optimization of the gas composition is required to keep the product's quality. In the evaluation of MAP for poultry, Wang et al. (2017) identified a shift in the bacterial community compared to other packaging conditions using metagenomics, and Höll et al. (2020) used metatranscriptomics to monitor the regulation responses of two spoilage bacteria to different atmospheric conditions. Similarly, evaluations of shelf life of fish fillets in MAP and vacuum packaging at low temperatures have been performed with metagenetics and sensory analysis or metabolomics to understand the dynamics of spoilage bacteria over time (Jääskeläinen et al. 2019, Sørensen et al. 2020). The efficacy of cleaning and disinfection has been investigated with metagenetics, with evidence of bacterial diversity and abundance altered after cleaning in dairy and pig facilities (Bridier et al. 2019, Dass et al. 2018). Similarly, RNA-based 16S rRNA sequencing showed current cleaning practices with ozonation were effective and caused shifts in potentially active microbiota in meat-processing plants (Botta et al. 2020). In contrast, sanitation in salmon-processing plants, determined by metagenetics, was found to be inadequate as *Pseudomonas* spp. persisted in biofilms on conveyor belts (Langsrud et al. 2016).

Ultimately, sequencing-based meta-omic approaches have been found to be effective tools in identifying microorganisms along the processing chain, and routine implementation can help to uncover the factors that influence microbial population dynamics (McHugh et al. 2020, Zwirzitz et al. 2020). The numerous studies carried out to date show that there is great potential for the use of meta-omic approaches in tracking microbial communities along the food chain.

SUMMARY POINTS

1. Microorganisms are important contributors to the quality and safety of a food product and they exist throughout the whole food chain.
2. As microbes exist in communities, it is valuable to study them as such. Meta-omic approaches bypass the need for culturing and isolating microbes and allow for the greater characterisation of microbial communities.
3. Metagenetics and metagenomics are two sequencing-based meta-omic approaches that are already being used in the characterisation of foods, food-associated environments and food processing microbiomes. Although only a few studies have used metatranscriptomics, results show potential in assessing the dynamics of viable microbes along the food chain.
4. The use of sequencing-based meta-omic approaches shows promise in better characterisation of microbiomes along the food chain and would allow for greater understanding of the factors contributing to food safety and quality. However, standardized workflows/pipelines are necessary to allow for data sharing and comparability and widespread adoption at a regulatory and industry level.

FUTURE ISSUES

1. With increasing adoption of these meta-omic approaches to uncover the microbiome of food and food-related environments, there is a great need for standardized workflows/pipelines for methodology and analysis.
2. Large amounts of data are generated by sequencing. This requires good data management practices and systematic metadata documentation to facilitate data sharing of research outputs. Additionally, bioinformatics expertise for the analysis of the data generated is currently essential to draw accurate and correct interpretations from the sequencing data. Future efforts will need to focus on accurate, automated analytical tools.
3. As substantial parts of the analysis require referencing available databases, the results from sequencing studies are only as good as these databases. Databases are currently compiled mainly from human microbiome studies, as more research has been done in that field, which may result in a bias toward human-related microbes. The ongoing

increase in microbiome studies on food and other fields should correct this imbalance to enable better characterisation of microbiomes.

4. With the further development of assays to overcome the challenges of meta-omic approaches, such as host DNA depletion and the ability to distinguish viable microbes in the microbial community, there will be an even wider application of meta-omic approaches for the characterisation of microbes along the food-processing chain.
5. From metagenomic data, the recovery of MAGs could make way for more single-strain studies that can contribute to a greater understanding of the resident microflora of food environments as well as the strains responsible for fermentation or spoilage in foods. Additionally, increasing the number of studies into the functional properties of microorganisms within food environments using metatranscriptomics or metagenomics with complementary approaches like metabolomics can provide greater insight into the active microorganisms and metabolic pathways involved in processes along the food chain.
6. Portable sequencing devices from ONT have allowed for field/onsite sequencing which has proven to be useful in clinical outbreak investigations and environmental sampling. These portable devices could enable rapid detection of microbiological contaminants or pathogens in food-production or food-processing environments. Although some studies have explored this possibility, further comparisons with other sequencing technologies and platforms are required to determine accuracy and comparability (McHugh et al. 2021, Yang et al. 2020).

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGEMENTS

M.Y. is funded by the Irish Dairy Levy. Research in the Cotter laboratory is funded by Science Foundation Ireland (SFI) under grant number SFI/12/RC/2273 (APC Microbiome Ireland), by SFI together with the Irish Department of Agriculture, Food and the Marine

under grant number SFI/16/RC/3835 (VistaMilk), and by the European Commission under the Horizon 2020 program under grant number 818368 (MASTER).

LITERATURE CITED

- Aidoo KE, Rob Nout M, Sarkar PK. 2006. Occurrence and function of yeasts in Asian indigenous fermented foods. *FEMS Yeast Res.* 6(1):30-39
- Allard SM, Callahan MT, Bui A, Ferelli AMC, Chopyk J et al. 2019. Creek to Table: Tracking fecal indicator bacteria, bacterial pathogens, and total bacterial communities from irrigation water to kale and radish crops. *Sci. Total Environ.* 666:461-71
- Anvarian AH, Cao Y, Srikumar S, Fanning S, Jordan K. 2016. Flow cytometric and 16S sequencing methodologies for monitoring the physiological status of the microbiome in powdered infant formula production. *Front. Microbiol.* 7:968
- Aw TG, Wengert S, Rose JB. 2016. Metagenomic analysis of viruses associated with field-grown and retail lettuce identifies human and animal viruses. *Int. J. Food Microbiol.* 223:50-56
- Benson AK, David JR, Gilbreth SE, Smith G, Nietfeldt J et al. 2014. Microbial successions are associated with changes in chemical profiles of a model refrigerated fresh pork sausage during an 80-day shelf life study. *Appl. Environ. Microbiol.* 80(17):5178-94
- Bintsis T. 2017. Foodborne pathogens. *AIMS microbiology* 3(3):529
- Blackburn CW. 2006. *Food spoilage microorganisms*. Cambridge, UK: Woodhead Publishing
- Bokulich NA, Bergsveinson J, Ziola B, Mills DA. 2015. Mapping microbial ecosystems and spoilage-gene flow in breweries highlights patterns of contamination and resistance. *Elife* 4:e04634
- Bokulich NA, Ohta M, Richardson PM, Mills DA. 2013. Monitoring seasonal changes in winery-resident microbiota. *PLOS ONE* 8(6)
- Botta C, Ferrocino I, Pessione A, Cocolin L, Rantsiou K. 2020. Spatiotemporal Distribution of the Environmental Microbiota in Food Processing Plants as Impacted by Cleaning and Sanitizing Procedures: The Case of Slaughterhouses and Gaseous Ozone. *Appl. Environ. Microbiol.* 86(23)

776 Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D et al. 2017.
777 Minimum information about a single amplified genome (MISAG) and a metagenome-
778 assembled genome (MIMAG) of bacteria and archaea. *Nat. Biotechnol.* 35(8):725-31

779 Bridier A, Le Grandois P, Moreau M-H, Prénom C, Le Roux A et al. 2019. Impact of
780 cleaning and disinfection procedures on microbial ecology and Salmonella
781 antimicrobial resistance in a pig slaughterhouse. *Sci. Rep.* 9(1):1-13

782 Buytaers FE, Saltykova A, Denayer S, Verhaegen B, Vanneste K et al. 2020. A Practical
783 Method to Implement Strain-Level Metagenomics-Based Foodborne Outbreak
784 Investigation and Source Tracking in Routine. *Microorganisms* 8(8):1191

785 Calero GC, Gómez NC, Lerma LL, Benomar N, Knapp CW, Abriouel H. 2020. In silico
786 mapping of microbial communities and stress responses in a porcine slaughterhouse
787 and pork products through its production chain, and the efficacy of HLE disinfectant.
788 *Food Res. Int.* 136:109486

789 Campos Calero G, Caballero Gómez N, Benomar N, Pérez Montoro B, Knapp CW et al.
790 2018. Deciphering resistome and virulome diversity in a porcine slaughterhouse and
791 pork products through its production chain. *Front. Microbiol.* 9:2099

792 Cao Y, Fanning S, Proos S, Jordan K, Srikumar S. 2017. A review on the applications of next
793 generation sequencing technologies as applied to food-related microbiome studies.
794 *Front. Microbiol.* 8:1829

795 Carvalho DC, Palhares RM, Drummond MG, Gadanhho M. 2017. Food metagenomics: Next
796 generation sequencing identifies species mixtures and mislabeling within highly
797 processed cod products. *Food Control* 80:183-86

798 Ceuppens S, Li D, Uyttendaele M, Renault P, Ross P et al. 2014. Molecular methods in food
799 safety microbiology: interpretation and implications of nucleic acid detection.
800 *Comprehensive Reviews in Food Science and Food Safety* 13(4):551-77

801 Chen G, Chen C, Lei Z. 2017. Meta-omics insights in the microbial community profiling and
802 functional characterization of fermented foods. *Trends Food Sci. Technol.* 65:23-31

803 Claesson MJ, Wang Q, O'Sullivan O, Greene-Diniz R, Cole JR et al. 2010. Comparison of
804 two next-generation sequencing technologies for resolving highly complex microbiota
805 composition using tandem variable 16S rRNA gene regions. *Nucleic Acids Res.*
806 38(22):e200-e00

807 Cocolin L, Ercolini D. 2015. Zooming into food-associated microbial consortia: a
808 'cultural' evolution. *Current Opinion in Food Science* 2:43-50

809 Coghlan ML, Haile J, Houston J, Murray DC, White NE et al. 2012. Deep sequencing of
810 plant and animal DNA contained within traditional Chinese medicines reveals legality
811 issues and health safety concerns. *PLoS Genet.* 8(4):e1002657

812 Comar M, D'Accolti M, Cason C, Soffritti I, Campisciano G et al. 2019. Introduction of NGS
813 in environmental surveillance for healthcare-associated infection control.
814 *Microorganisms* 7(12):708

815 Coughlan LM, Cotter PD, Hill C, Alvarez-Ordóñez A. 2016. New weapons to fight old
816 enemies: novel strategies for the (bio) control of bacterial biofilms in the food
817 industry. *Front. Microbiol.* 7:1641

818 Dass SC, Wang B, Stratton JE, Bianchini A, Ababdappa A. 2018. Food processing
819 environment surveillance using amplicon metagenomics: assessing the change in the
820 microbiome of a fluid milk processing facility before and after cleaning. *BAOJ Food*
821 *Sci & Tec* 2(1):12

822 De Filippis F, Genovese A, Ferranti P, Gilbert JA, Ercolini D. 2016. Metatranscriptomics
823 reveals temperature-driven functional changes in microbiome impacting cheese
824 maturation rate. *Sci. Rep.* 6:21871

825 De Filippis F, La Storia A, Villani F, Ercolini D. 2013. Exploring the sources of bacterial
826 spoilers in beefsteaks by culture-independent high-throughput sequencing. *PLOS*
827 *ONE* 8(7)

828 De Filippis F, Parente E, Ercolini D. 2017. Metagenomics insights into food fermentations.
829 *Microb. Biotechnol.* 10(1):91-102

830 De Filippis F, Valentino V, Alvarez-Ordóñez A, Cotter PD, Ercolini D. 2020. Environmental
831 microbiome mapping as a strategy to improve quality and safety in the food industry.
832 *Current Opinion in Food Science*

833 de Koster CG, Brul S. 2016. MALDI-TOF MS identification and tracking of food spoilers
834 and food-borne pathogens. *Current Opinion in Food Science* 10:76-84

835 den Besten HM, Amézquita A, Bover-Cid S, Dagnas S, Ellouze M et al. 2018. Next
836 generation of microbiological risk assessment: potential of omics data for exposure
837 assessment. *Int. J. Food Microbiol.* 287:18-27

838 Doyle CJ, Gleeson D, O'Toole PW, Cotter PD. 2017. Impacts of seasonal housing and teat
839 preparation on raw milk microbiota: a high-throughput sequencing study. *Appl.*
840 *Environ. Microbiol.* 83(2)

841 Doyle RM, O'Sullivan DM, Aller SD, Bruchmann S, Clark T et al. 2020. Discordant
842 bioinformatic predictions of antimicrobial resistance from whole-genome sequencing
843 data of bacterial isolates: An inter-laboratory study. *Microbial genomics* 6(2)

844 Dugat-Bony E, Straub C, Teissandier A, Onésime D, Loux V et al. 2015. Overview of a
845 surface-ripened cheese community functioning by meta-omics analyses. *PLOS ONE*
846 10(4):e0124360

847 Duru IC, Laine P, Andreevskaya M, Paulin L, Kananen S et al. 2018. Metagenomic and
848 metatranscriptomic analysis of the microbial community in Swiss-type Maasdam
849 cheese during ripening. *Int. J. Food Microbiol.* 281:10-22

850 Dwivedi HP, Jaykus L-A. 2011. Detection of pathogens in foods: the current state-of-the-art
851 and future directions. *Crit. Rev. Microbiol.* 37(1):40-63

852 Dzieciol M, Schornsteiner E, Muhterem-Uyar M, Stessl B, Wagner M, Schmitz-Esser S.
853 2016. Bacterial diversity of floor drain biofilms and drain waters in a *Listeria*
854 monocytogenes contaminated food processing environment. *Int. J. Food Microbiol.*
855 223:33-40

856 EFSA Panel on Biological Hazards, Koutsoumanis K, Allende A, Alvarez-Ordóñez A,
857 Bolton D et al. 2019. Whole genome sequencing and metagenomics for outbreak
858 investigation, source attribution and risk assessment of food-borne microorganisms.
859 *EFSA Journal* 17(12):e05898

860 Einson JE, Rani A, You X, Rodriguez AA, Randell CL et al. 2018. A vegetable fermentation
861 facility hosts distinct microbiomes reflecting the production environment. *Appl.*
862 *Environ. Microbiol.* 84(22):e01680-18

863 Erkus O, de Jager VC, Geene RT, van Alen-Boerrigter I, Hazelwood L et al. 2016. Use of
864 propidium monoazide for selective profiling of viable microbial cells during Gouda
865 cheese ripening. *Int. J. Food Microbiol.* 228:1-9

866 European Food Safety Authority, European Centre for Disease Prevention and Control. 2019.
867 The European Union one health 2018 zoonoses report. *EFSA Journal* 17(12):e05926

868 Falardeau J, Keeney K, Trmčić A, Kitts D, Wang S. 2019. Farm-to-fork profiling of bacterial
869 communities associated with an artisan cheese production facility. *Food Microbiol.*
870 83:48-58

871 Feng P. 2007. Rapid methods for the detection of foodborne pathogens: current and next-
872 generation technologies. In *Food Microbiology: Fundamentals and Frontiers, Third*
873 *Edition*, pp. 911-34: American Society of Microbiology

874 Fernandez-Cassi X, Timoneda N, Gonzales-Gustavson E, Abril J, Bofill-Mas S, Girones R.
 875 2017. A metagenomic assessment of viral contamination on fresh parsley plants
 876 irrigated with fecally tainted river water. *Int. J. Food Microbiol.* 257:80-90
 877 Fleet GH. 1999. Microorganisms in food ecosystems. *Int. J. Food Microbiol.* 50(1-1):101-17
 878 Fleet GH. 2011. Yeast spoilage of foods and beverages. In *The yeasts*, pp. 53-63: Elsevier
 879 Fox EM, Solomon K, Moore JE, Wall PG, Fanning S. 2014. Phylogenetic profiles of in-
 880 house microflora in drains at a food production facility: comparison and biocontrol
 881 implications of *Listeria*-positive and-negative bacterial populations. *Appl. Environ.*
 882 *Microbiol.* 80(11):3369-74
 883 Franzosa EA, Hsu T, Sirota-Madi A, Shafquat A, Abu-Ali G et al. 2015. Sequencing and
 884 beyond: integrating molecular'omics' for microbial community profiling. *Nat. Rev.*
 885 *Microbiol.* 13(6):360-72
 886 Galazzo G, van Best N, Benedikter B, Janssen K, Bervoets L et al. 2020. How to count our
 887 microbes? The effect of different quantitative microbiome profiling approaches.
 888 *Front. Cell. Infect. Microbiol.* 10: 403. doi: 10.3389/fcimb
 889 Galvez A, López RL, Abriouel H, Valdivia E, Omar NB. 2008. Application of bacteriocins in
 890 the control of foodborne pathogenic and spoilage bacteria. *Crit. Rev. Biotechnol.*
 891 28(2):125-52
 892 Grützke J, Malorny B, Hammerl JA, Busch A, Tausch SH et al. 2019. Fishing in the soup—
 893 Pathogen detection in food safety using Metabarcoding and Metagenomic sequencing.
 894 *Front. Microbiol.* 10:1805
 895 Gweon HS, Shaw LP, Swann J, De Maio N, AbuOun M et al. 2019. The impact of
 896 sequencing depth on the inferred taxonomic composition and AMR gene content of
 897 metagenomic samples. *Environmental Microbiome* 14(1):1-15
 898 Haiminen N, Edlund S, Chambliss D, Kunitomi M, Weimer BC et al. 2019. Food
 899 authentication from shotgun sequencing reads with an application on high protein
 900 powders. *NPJ science of food* 3(1):1-11
 901 Handelsman J. 2004. Metagenomics: application of genomics to uncultured microorganisms.
 902 *Microbiol. Mol. Biol. Rev.* 68(4):669-85
 903 Hatti-Kaul R, Chen L, Dishisha T, Enshasy HE. 2018. Lactic acid bacteria: From starter
 904 cultures to producers of chemicals. *FEMS Microbiol. Lett.* 365(20):fny213
 905 He L-Y, He L-K, Liu Y-S, Zhang M, Zhao J-L et al. 2019. Microbial diversity and antibiotic
 906 resistome in swine farm environments. *Sci. Total Environ.* 685:197-207

907 Hendriksen RS, Munk P, Njage P, Van Bunnik B, McNally L et al. 2019. Global monitoring
908 of antimicrobial resistance based on metagenomics analyses of urban sewage. *Nature*
909 *communications* 10(1):1124

910 Höll L, Hilgarth M, Geissler AJ, Behr J, Vogel RF. 2020. Metatranscriptomic analysis of
911 modified atmosphere packaged poultry meat enables prediction of *Brochothrix*
912 *thermosphacta* and *Carnobacterium divergens* in situ metabolism. *Arch. Microbiol.*
913 202:1945-55

914 Hultman J, Rahkila R, Ali J, Rousu J, Björkroth KJ. 2015. Meat processing plant microbiome
915 and contamination patterns of cold-tolerant bacteria causing food safety and spoilage
916 risks in the manufacture of vacuum-packaged cooked sausages. *Appl. Environ.*
917 *Microbiol.* 81(20):7088-97

918 Hwang BK, Choi H, Choi SH, Kim B-S. 2020. Analysis of Microbiota Structure and
919 Potential Functions Influencing Spoilage of Fresh Beef Meat. *Front. Microbiol.*
920 11:1657

921 Illegghems K, Weckx S, De Vuyst L. 2015. Applying meta-pathway analyses through
922 metagenomics to identify the functional properties of the major bacterial communities
923 of a single spontaneous cocoa bean fermentation process sample. *Food Microbiol.*
924 50:54-63

925 in't Veld JHH. 1996. Microbial and biochemical spoilage of foods: an overview. *Int. J. Food*
926 *Microbiol.* 33(1):1-18

927 Jääskeläinen E, Jakobsen LM, Hultman J, Eggers N, Bertram HC, Björkroth J. 2019.
928 Metabolomics and bacterial diversity of packaged yellowfin tuna (*Thunnus albacares*)
929 and salmon (*Salmo salar*) show fish species-specific spoilage development during
930 chilled storage. *Int. J. Food Microbiol.* 293:44-52

931 Jenkins C, Dallman TJ, Grant KA. 2019. Impact of whole genome sequencing on the
932 investigation of food-borne outbreaks of Shiga toxin-producing *Escherichia coli*
933 serogroup O157: H7, England, 2013 to 2017. *EuroSurveillance* 24(4):1800346

934 Jordan K, Dalmaso M, Zentek J, Mader A, Bruggeman G et al. 2014. Microbes versus
935 microbes: control of pathogens in the food chain. *J. Sci. Food Agric.* 94(15):3079-89

936 Jung JY, Lee SH, Jin HM, Hahn Y, Madsen EL, Jeon CO. 2013. Metatranscriptomic analysis
937 of lactic acid bacterial gene expression during kimchi fermentation. *Int. J. Food*
938 *Microbiol.* 163(2-3):171-79

939 Kable ME, Srisengfa Y, Laird M, Zaragoza J, McLeod J et al. 2016. The core and seasonal
940 microbiota of raw bovine milk in tanker trucks and the impact of transfer to a milk
941 processing facility. *MBio* 7(4):e00836-16

942 Kable ME, Srisengfa Y, Xue Z, Coates LC, Marco ML. 2019. Viable and total bacterial
943 populations undergo equipment-and time-dependent shifts during milk processing.
944 *Appl. Environ. Microbiol.* 85(13):e00270-19

945 Kamilari E, Anagnostopoulos DA, Papademas P, Kamilaris A, Tsaltas D. 2020.
946 Characterizing halloumi cheese's bacterial communities through metagenomic
947 analysis. *LWT*:109298

948 Kamilari E, Tomazou M, Antoniadis A, Tsaltas D. 2019. High throughput sequencing
949 technologies as a new toolbox for deep analysis, characterization and potentially
950 authentication of protection designation of origin cheeses? *International journal of*
951 *food science*

952 Khansaritoreh E, Salmaki Y, Ramezani E, Azirani TA, Keller A et al. 2020. Employing DNA
953 metabarcoding to determine the geographical origin of honey. *Heliyon* 6(11):e05596

954 Kim KH, Chun BH, Kim J, Jeon CO. 2020. Identification of biogenic amine-producing
955 microbes during fermentation of ganjang, a Korean traditional soy sauce, through
956 metagenomic and metatranscriptomic analyses. *Food Control* 121:107681

957 Kim S, Kim J, Yun EJ, Kim KH. 2016. Food metabolomics: from farm to human. *Curr.*
958 *Opin. Biotechnol.* 37:16-23

959 King P, Pham LK, Waltz S, Sphar D, Yamamoto RT et al. 2016. Longitudinal metagenomic
960 analysis of hospital air identifies clinically relevant microbes. *PLOS ONE* 11(8)

961 Kleiner M. 2019. Metaproteomics: much more than measuring gene expression in microbial
962 communities. *MSystems* 4(3):e00115-19

963 Langsrud S, Moen B, Møretrø T, Løype M, Heir E. 2016. Microbial dynamics in mixed
964 culture biofilms of bacteria surviving sanitation of conveyor belts in salmon-
965 processing plants. *J. Appl. Microbiol.* 120(2):366-78

966 Leonard SR, Mammel MK, Lacher DW, Elkins CA. 2015. Application of metagenomic
967 sequencing to food safety: detection of Shiga toxin-producing *Escherichia coli* on
968 fresh bagged spinach. *Appl. Environ. Microbiol.* 81(23):8183-91

969 Leroy F, De Vuyst L. 2004. Lactic acid bacteria as functional starter cultures for the food
970 fermentation industry. *Trends Food Sci. Technol.* 15(2):67-78

971 Lewis E, Hudson J, Cook N, Barnes J, Haynes E. 2020. Next-generation sequencing as a
972 screening tool for foodborne pathogens in fresh produce. *J. Microbiol.*
973 *Methods*:105840

974 Li R, Wang C, Zhou G, Li C, Ye K. 2021. The effects of thermal treatment on the bacterial
975 community and quality characteristics of meatballs during storage. *Food Science &*
976 *Nutrition* 9(1):564-73

977 Li S, Mann DA, Zhang S, Qi Y, Meinersmann RJ, Deng X. 2020. Microbiome-Informed
978 Food Safety and Quality: Longitudinal Consistency and Cross-Sectional
979 Distinctiveness of Retail Chicken Breast Microbiomes. *Msystems* 5(5)

980 Liu N, Zou D, Dong D, Yang Z, Ao D et al. 2017. Development of a multiplex loop-mediated
981 isothermal amplification method for the simultaneous detection of *Salmonella* spp.
982 and *Vibrio parahaemolyticus*. *Sci. Rep.* 7(1):1-7

983 Liu X-F, Liu C-J, Zeng X-Q, Zhang H-Y, Luo Y-Y, Li X-R. 2020. Metagenomic and
984 metatranscriptomic analysis of the microbial community structure and metabolic
985 potential of fermented soybean in Yunnan Province. *Food Science and Technology*
986 40(1):18-25

987 Liu X, Teixeira JS, Ner S, Ma KV, Petronella N et al. 2020. Exploring the potential of the
988 microbiome as a marker of the geographic origin of fresh seafood. *Front. Microbiol.*
989 11

990 Loman NJ, Constantinidou C, Christner M, Rohde H, Chan JZ-M et al. 2013. A culture-
991 independent sequence-based metagenomics approach to the investigation of an
992 outbreak of Shiga-toxigenic *Escherichia coli* O104: H4. *JAMA* 309(14):1502-10

993 Luiken RE, Van Gompel L, Bossers A, Munk P, Joosten P et al. 2020. Farm dust resistomes
994 and bacterial microbiomes in European poultry and pig farms. *Environ. Int.*
995 143:105971

996 Mandal PK, Biswas AK, Choi K, Pal UK. 2011. Methods for rapid detection of foodborne
997 pathogens: an overview. *American Journal of Food Technology* 6:87-102

998 Marine R, McCarren C, Vorrasane V, Nasko D, Crowgey E et al. 2014. Caught in the middle
999 with multiple displacement amplification: the myth of pooling for avoiding multiple
1000 displacement amplification bias in a metagenome. *Microbiome* 2(1):3

1001 Marotz CA, Sanders JG, Zuniga C, Zaramela LS, Knight R, Zengler K. 2018. Improving
1002 saliva shotgun metagenomics by chemical host DNA depletion. *Microbiome* 6(1):42

1003 McHugh AJ, Feehily C, Fenelon MA, Gleeson D, Hill C, Cotter PD. 2020. Tracking the
1004 Dairy Microbiota from Farm Bulk Tank to Skimmed Milk Powder. *Msystems* 5(2)

- McHugh AJ, Feehily C, Tobin JT, Fenelon MA, Hill C, Cotter PD. 2018. Mesophilic sporeformers identified in whey powder by using shotgun metagenomic sequencing. *Appl. Environ. Microbiol.* 84(20):e01305-18
- McHugh AJ, Yap M, Crispie F, Feehily C, Hill C, Cotter PD. 2021. Microbiome-based environmental monitoring of a dairy processing facility highlights the challenges associated with low microbial-load samples. *npj Science of Food* 5(4)
- Mencía-Ares O, Cabrera-Rubio R, Cobo-Díaz JF, Álvarez-Ordóñez A, Gómez-García M et al. 2020. Antimicrobial use and production system shape the fecal, environmental, and slurry resistomes of pig farms. *Microbiome* 8(1):1-17
- Mira Miralles M, Maestre-Carballea L, Lluésma-Gomez M, Martínez-García M. 2019. High-throughput 16S rRNA sequencing to assess potentially active bacteria and foodborne pathogens: a case example in ready-to-eat food. *Foods* 8(10):480
- Munk P, Andersen VD, de Knecht L, Jensen MS, Knudsen BE et al. 2017. A sampling and metagenomic sequencing-based methodology for monitoring antimicrobial resistance in swine herds. *J. Antimicrob. Chemother.* 72(2):385-92
- Nocker A, Cheung C-Y, Camper AK. 2006. Comparison of propidium monoazide with ethidium monoazide for differentiation of live vs. dead bacteria by selective removal of DNA from dead cells. *J. Microbiol. Methods* 67(2):310-20
- Nogarol C, Acutis P, Bianchi D, Maurella C, Peletto S et al. 2013. Molecular characterization of *Pseudomonas fluorescens* isolates involved in the Italian “blue mozzarella” event. *J. Food Prot.* 76(3):500-04
- Noyes NR, Yang X, Linke LM, Magnuson RJ, Dettenwanger A et al. 2016. Resistome diversity in cattle and the environment decreases during beef production. *Elife* 5:e13195
- Nychas GE, Panagou E. 2011. Microbiological spoilage of foods and beverages. In *Food and Beverage Stability and Shelf Life*, ed. Kilcast D, Subarmaniam P, pp. 3-28. Cambridge, UK: Woodhead Publishing
- O'Sullivan L, Bolton D, McAuliffe O, Coffey A. 2019. Bacteriophages in food applications: From foe to friend. *Annu. Rev. Food Sci. Technol.* 10:151-72
- O'Hara NB, Reed HJ, Afshinnekoo E, Harvin D, Caplan N et al. 2017. Metagenomic characterization of ambulances across the USA. *Microbiome* 5(1):125
- Ogier J-C, Pages S, Galan M, Barret M, Gaudriault S. 2019. rpoB, a promising marker for analyzing the diversity of bacterial communities by amplicon sequencing. *BMC Microbiol.* 19(1):171

- Ouali FA, Al Kassaa I, Cudennec B, Abdallah M, Bendali F et al. 2014. Identification of lactobacilli with inhibitory effect on biofilm formation by pathogenic bacteria on stainless steel surfaces. *Int. J. Food Microbiol.* 191:116-24
- Pasolli E, De Filippis F, Mauriello IE, Cumbo F, Walsh AM et al. 2020. Large-scale genome-wide analysis links lactic acid bacteria from food with the gut microbiome. *Nature communications* 11(1):1-12
- Peirson MD, Guan TY, Holley RA. 2003. Thermal resistances and lactate and diacetate sensitivities of bacteria causing bologna discolouration. *Int. J. Food Microbiol.* 86(3):223-30
- Petersen TN, Rasmussen S, Hasman H, Carøe C, Bælum J et al. 2015. Meta-genomic analysis of toilet waste from long distance flights; a step towards global surveillance of infectious diseases and antimicrobial resistance. *Sci. Rep.* 5:11444
- Petruzzi L, Corbo MR, Sinigaglia M, Bevilacqua A. 2017. Microbial spoilage of foods: Fundamentals. In *The microbiological quality of food*, pp. 1-21: Elsevier
- Pham N-P, Landaud S, Lieben P, Bonnarme P, Monnet C. 2019. Transcription Profiling Reveals Cooperative Metabolic Interactions in a Microbial Cheese-Ripening Community Composed of *Debaryomyces hansenii*, *Brevibacterium aurantiacum*, and *Hafnia alvei*. *Front. Microbiol.* 10:1901
- Pinu FR, Beale DJ, Paten AM, Kouremenos K, Swarup S et al. 2019. Systems biology and multi-omics integration: viewpoints from the metabolomics research community. *Metabolites* 9(4):76
- Pitta DW, Dou Z, Kumar S, Indugu N, Toth JD et al. 2016. Metagenomic evidence of the prevalence and distribution patterns of antimicrobial resistance genes in dairy agroecosystems. *Foodborne Pathog. Dis.* 13(6):296-302
- Poirier S, Rue O, Peguilhan R, Coeuret G, Zagorec M et al. 2018. Deciphering intra-species bacterial diversity of meat and seafood spoilage microbiota using gyrB amplicon sequencing: A comparative analysis with 16S rDNA V3-V4 amplicon sequencing. *PLOS ONE* 13(9):e0204629
- Pothakos V, Stellato G, Ercolini D, Devlieghere F. 2015. Processing environment and ingredients are both sources of *Leuconostoc gelidum*, which emerges as a major spoiler in ready-to-eat meals. *Appl. Environ. Microbiol.* 81(10):3529-41
- Quince C, Walker AW, Simpson JT, Loman NJ, Segata N. 2017. Shotgun metagenomics, from sampling to analysis. *Nat. Biotechnol.* 35(9):833

1072 Rampelotto PH, Sereia AF, de Oliveira LFV, Margis R. 2019. Exploring the hospital
1073 microbiome by high-resolution 16S rRNA profiling. *Int. J. Mol. Sci.* 20(12):3099

1074 Rantsiou K, Kathariou S, Winkler A, Skandamis P, Saint-Cyr MJ et al. 2018. Next generation
1075 microbiological risk assessment: opportunities of whole genome sequencing (WGS)
1076 for foodborne pathogen surveillance, source tracking and risk assessment. *Int. J. Food*
1077 *Microbiol.* 287:3-9

1078 Reis J, Paula A, Casarotti S, Penna A. 2012. Lactic acid bacteria antimicrobial compounds:
1079 characteristics and applications. *Food Engineering Reviews* 4(2):124-40

1080 Rodríguez-López P, Rodríguez-Herrera JJ, Cabo ML. 2020. Tracking bacteriome variation
1081 over time in *Listeria monocytogenes*-positive foci in food industry. *Int. J. Food*
1082 *Microbiol.* 315:108439

1083 Rubiola S, Chiesa F, Dalmaso A, Di Ciccio P, Civera T. 2020. Detection of Antimicrobial
1084 Resistance Genes in the Milk Production Environment: Impact of Host DNA and
1085 Sequencing Depth. *Front. Microbiol.* 11:1983

1086 Sahu M, Bala S. 2017. Food processing, food spoilage and their prevention: an overview. *Int.*
1087 *J. Life. Sci. Scienti. Res* 3(1):753-59

1088 Schallmeyer M, Singh A, Ward OP. 2004. Developments in the use of *Bacillus* species for
1089 industrial production. *Can. J. Microbiol.* 50(1):1-17

1090 Schwan RF, Ramos CL. 2014. Role of microbes and their diversity in fermented foods. In
1091 *Beneficial microbes in fermented and functional foods*, ed. Ravishankar Rai V, Bai
1092 JA, pp. 524-47. Boca Raton, FL: CRC Press

1093 Shakya M, Lo C-C, Chain PS. 2019. Advances and challenges in Metatranscriptomic
1094 analysis. *Frontiers in Genetics* 10:904

1095 Slizovskiy IB, Mukherjee K, Dean CJ, Boucher C, Noyes NR. 2020. Mobilization of
1096 antibiotic resistance: are current approaches for colocalizing resistomes and
1097 mobilomes useful? *Front. Microbiol.* 11:1376

1098 Soggiu A, Piras C, Mortera SL, Alloggio I, Urbani A et al. 2016. Unravelling the effect of
1099 clostridia spores and lysozyme on microbiota dynamics in Grana Padano cheese: a
1100 metaproteomics approach. *J. Proteomics* 147:21-27

1101 Sohier D, Pavan S, Riou A, Combrisson J, Postollec F. 2014. Evolution of microbiological
1102 analytical methods for dairy industry needs. *Front. Microbiol.* 5:16

1103 Sørensen JS, Bøknæs N, Mejlholm O, Dalgaard P. 2020. Superchilling in combination with
1104 modified atmosphere packaging resulted in long shelf-life and limited microbial

1105 growth in Atlantic cod (*Gadus morhua* L.) from capture-based-aquaculture in
 1106 Greenland. *Food Microbiol.* 88:103405

1107 Stellato G, La Stora A, De Filippis F, Borriello G, Villani F, Ercolini D. 2016. Overlap of
 1108 spoilage-associated microbiota between meat and the meat processing environment in
 1109 small-scale and large-scale retail distributions. *Appl. Environ. Microbiol.*
 1110 82(13):4045-54

1111 Stohr V, Joffraud J-J, Cardinal M, Leroi F. 2001. Spoilage potential and sensory profile
 1112 associated with bacteria isolated from cold-smoked salmon. *Food Res. Int.* 34(9):797-
 1113 806

1114 Tao J, Liu W, Ding W, Han R, Shen Q et al. 2020. A multiplex PCR assay with a common
 1115 primer for the detection of eleven foodborne pathogens. *J. Food Sci.* 85(3):744-54

1116 Tsironi T, Anjos L, Pinto PI, Dimopoulos G, Santos S et al. 2019. High pressure processing
 1117 of European sea bass (*Dicentrarchus labrax*) fillets and tools for flesh quality and
 1118 shelf life monitoring. *J. Food Eng.* 262:83-91

1119 Van Gompel L, Luiken RE, Sarrazin S, Munk P, Knudsen BE et al. 2019. The antimicrobial
 1120 resistome in relation to antimicrobial use and biosecurity in pig farming, a
 1121 metagenome-wide association study in nine European countries. *J. Antimicrob.*
 1122 *Chemother.* 74(4):865-76

1123 Vikram A, Woolston J, Sulakvelidze A. 2020. Phage Biocontrol Applications in Food
 1124 Production and Processing. *Bacterial Viruses: Exploitation for Biocontrol and*
 1125 *Therapeutics*:283-334

1126 Walsh AM, Crispie F, O'Sullivan O, Finnegan L, Claesson MJ, Cotter PD. 2018. Species
 1127 classifier choice is a key consideration when analysing low-complexity food
 1128 microbiome data. *Microbiome* 6(1):50

1129 Wang H, Zhang X, Wang G, Jia K, Xu X, Zhou G. 2017. Bacterial community and spoilage
 1130 profiles shift in response to packaging in yellow-feather broiler, a highly popular meat
 1131 in Asia. *Front. Microbiol.* 8:2588

1132 Wang X, Du H, Zhang Y, Xu Y. 2018. Environmental microbiota drives microbial succession
 1133 and metabolic profiles during Chinese liquor fermentation. *Appl. Environ. Microbiol.*
 1134 84(4):e02369-17

1135 Wang Y, Hu Y, Liu F, Cao J, Lv N et al. 2020. Integrated metagenomic and
 1136 metatranscriptomic profiling reveals differentially expressed resistomes in human,
 1137 chicken, and pig gut microbiomes. *Environ. Int.* 138:105649

1138 Wang Y, Salazar JK. 2016. Culture-independent rapid detection methods for bacterial
 1139 pathogens and toxins in food matrices. *Comprehensive Reviews in Food Science and*
 1140 *Food Safety* 15(1):183-205

1141 Wang Y, Yan Y, Thompson KN, Bae S, Accorsi EK et al. 2021. Whole microbial community
 1142 viability is not quantitatively reflected by propidium monoazide sequencing approach.
 1143 *Microbiome* 9(1):1-13

1144 World Health Organisation. 2017. The burden of foodborne diseases in the WHO European
 1145 Region. pp. 48. Copenhagen, Denmark: WHO Regional Office for Europe

1146 Wu H, Nguyen QD, Tran TT, Tang MT, Tsuruta T, Nishino N. 2019. Rumen fluid, feces,
 1147 milk, water, feed, airborne dust, and bedding microbiota in dairy farms managed by
 1148 automatic milking systems. *Animal Science Journal* 90(3):445-52

1149 Wu L, Li G, Xu X, Zhu L, Huang R, Chen X. 2019. Application of nano-ELISA in food
 1150 analysis: recent advances and challenges. *TrAC Trends in Analytical Chemistry*
 1151 113:140-56

1152 Xiao Y, Huang T, Xu Y, Peng Z, Liu Z et al. 2020. Metatranscriptomics reveals the gene
 1153 functions and metabolic properties of the major microbial community during Chinese
 1154 Sichuan Paocai fermentation. *Food Microbiol.*:103573

1155 Xie M, Wu J, An F, Yue X, Tao D et al. 2019. An integrated metagenomic/metaproteomic
 1156 investigation of microbiota in dajiang-meju, a traditional fermented soybean product
 1157 in Northeast China. *Food Res. Int.* 115:414-24

1158 Yang M, Cousineau A, Liu X, Luo Y, Sun D et al. 2020. Direct metatranscriptome RNA-seq
 1159 and multiplex RT-PCR amplicon sequencing on Nanopore MinION—promising
 1160 strategies for multiplex identification of viable pathogens in food. *Front. Microbiol.*
 1161 11:514

1162 Yang X, Noyes NR, Doster E, Martin JN, Linke LM et al. 2016. Use of metagenomic
 1163 shotgun sequencing technology to detect foodborne pathogens within the microbiome
 1164 of the beef production chain. *Appl. Environ. Microbiol.* 82(8):2433-43

1165 Yap M, Feehily C, Walsh CJ, Fenelon M, Murphy EF et al. 2020. Evaluation of methods for
 1166 the reduction of contaminating host reads when performing shotgun metagenomic
 1167 sequencing of the milk microbiome. *Sci. Rep.* 10(1):1-11

1168 Yuan L, Hansen MF, Røder HL, Wang N, Burmølle M, He G. 2020. Mixed-species biofilms
 1169 in the food industry: current knowledge and novel control strategies. *Crit. Rev. Food*
 1170 *Sci. Nutr.* 60(13):2277-93


1171 Zhang Y, Kitajima M, Whittle AJ, Liu W-T. 2017. Benefits of genomic insights and
1172 CRISPR-Cas signatures to monitor potential pathogens across drinking water
1173 production and distribution systems. *Front. Microbiol.* 8:2036
1174 Zotta T, Parente E, Ianniello RG, De Filippis F, Ricciardi A. 2019. Dynamics of bacterial
1175 communities and interaction networks in thawed fish fillets during chilled storage in
1176 air. *Int. J. Food Microbiol.* 293:102-13
1177 Zwirzitz B, Wetzels SU, Dixon ED, Stessl B, Zaiser A et al. 2020. The sources and
1178 transmission routes of microbial populations throughout a meat processing facility.
1179 *NPJ biofilms and microbiomes* 6(1):1-12
1180

1181 **TABLES**

1182 Table 1. Some classical examples of types of food spoilage that can be caused by different organisms.

Spoilage characteristic	Spoilage organism in food type	References
Off-odor and off-flavors	<i>Pseudomonas</i> spp., <i>Carnobacterium</i> spp., <i>Serratia</i> spp., <i>Leuconostoc</i> spp. and <i>Brochothrix thermosphacta</i> produce off-odors and off-flavors in meat, fish and poultry <i>Shewanella putrefaciens</i> causes rancid and sulfurous odors and <i>Aeromonas</i> spp. produces a sour flavor in smoked salmon Various Enterobacteriaceae cause off-odors and off-flavors in preserved seafood products <i>Citrobacter</i> and <i>Proteus</i> have been found to cause off-odors in poultry <i>Candida</i> spp. and <i>Kluyveromyces</i> spp. cause off-odors and flavors in fermented dairy products.	Blackburn (2006), Stohr et al. (2001), Fleet (2011)
Changes in texture	<i>Pseudomonas</i> spp. cause meat and poultry to become slimy/mushy due to the action of degradative enzymes LAB can cause poor texture in cheese <i>Bacillus</i> spp. are able to cause ropiness in breads and bakery products <i>Clostridium</i> spp. and <i>Bacillus</i> spp. cause softening in vegetables and fruit <i>Erwinia</i> and <i>Penicillium</i> spp. cause soft rots in vegetables, leading to a mushy texture. The texture of cheese and yogurts is altered by <i>Candida</i> spp. and <i>Kluyveromyces</i> spp.	Blackburn (2006), Nychas and Panagou (2011), Fleet (2011)
Discolouration	<i>Pseudomonas fluorescens</i> is able to cause blue coloration in cheese <i>Carnobacterium viridans</i> causes green discoloration in cooked cured sausage	Nogarol et al. (2013), Peirson et al. (2003)
Gas formation	<i>Clostridium</i> spp. cause gas formation resulting in bloating in canned or vacuum-packed goods and late blowing defects in cheese Enterobacteriaceae is responsible for gas production in salad products <i>Saccharomyces</i> causes gassiness in wines Several yeast species cause swelling in juice packets	Petruzzi et al. (2017), Sahu and Bala (2017)

1183



1185
1186
1187
1188
1189
1190
1191
1192

Figure 1. Current meta-omic approaches used in microbiome research. Sequencing-based approaches include Metagenetics, Metagenomics and Metatranscriptomics and other community-based methods include Metaproteomics and Meta-metabolomics, which are currently being used in human, environment, and food microbiome studies.




Figure 2. Current applications of meta-omic sequencing-based approaches along the food chain. Metagenetics, Metagenomics and Metatranscriptomics have been used in studies investigating the microbial community of food, food processing steps and food-associated environments.