

Title	Short-term consumption of a high-fat diet increases host susceptibility to <i>Listeria monocytogenes</i> infection
Authors	Las Heras, Vanessa;Clooney, Adam G.;Ryan, Feargal J.;Cabrera-Rubio, Raul;Casey, Pat G.;Hueston, Cara M.;Pinheiro, Jorge;Rudkin, Justine K.;Melgar, Silvia;Cotter, Paul D.;Hill, Colin;Gahan, Cormac G.
Publication date	2019
Original Citation	Las Heras, V., Clooney, A. G., Ryan, F. J., Cabrera-Rubio, R., Casey, P. G., Hueston, C. M., Pinheiro, J., Rudkin, J. K., Melgar, S., Cotter, P. D., Hill, C. and Gahan, C. G. M. (2019) 'Short-term consumption of a high-fat diet increases host susceptibility to <i>Listeria monocytogenes</i> infection', <i>Microbiome</i> , 7(1) (12pp). DOI: 10.1186/s40168-019-0621-x
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-019-0621-x - 10.1186/s40168-019-0621-x
Rights	© The Author(s). 2019. © The Author(s). 2019Open AccessThis article is distributed under the terms of the Creative Commons Attribution 4.0International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver(http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated. - https://creativecommons.org/licenses/by/4.0/
Download date	2024-03-29 10:13:53
Item downloaded from	https://hdl.handle.net/10468/7680



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Supplemental data, Las Heras et al *Fig S4*.




Figure S4. β -diversity metrics. **A.** Changes in Firmicutes to Bacteroidetes bacterial ratio of abundance during dietary shifts (D13 and D16). **B and C.** Changes in total bacterial abundance between low and high-fat diets at the family level. A volcano plot showing the fold change between high and low fat diets at (B. D13 and C. D16). Each point represents an Operational Taxonomic Unit (OTU). The X axis represents in the log2 of the fold change while the Y axis is the negative log10 of DESeq2 P values adjusted for multiple testing using the False Discovery Rate method. Points to the right of the plot with positive log2FoldChange values represent bacterial taxa with increased abundance in the mice on the high fat diet relative to the mice on the low fat and those with negative log2FoldChange values represent bacterial taxa with increased abundance in the low fat diet relative to the high fat diet. The horizontal black line represents the cut off for statistical significance, an adjusted p-value of 0.05.