


Title	Detailed mapping of Bifidobacterium strain transmission from mother to infant via a dual culture-based and metagenomic approach
Authors	Feehily, Conor;O'Neill, Ian J.;Walsh, Calum J.;Moore, Rebecca L.;Killeen, Sarah Louise;Geraghty, Aisling A.;Lawton, Elaine M.;Byrne, David F.;Sanchez-Gallardo, Rocio;Nori, Sai Ravi Chandra;Nielsen, Ida Busch;Wortmann, Esther;Matthews, Elizabeth;O'Flaherty, Roisin;Rudd, Pauline M.;Groeger, David S.;Shanahan, Fergus;Salдова, Radka;McAuliffe, Fionnuala M.;van Sinderen, Douwe;Cotter, Paul D.
Publication date	2023-01-01
Original Citation	Feehily, C., O'Neill, I.J., Walsh, C.J., Moore, R.L., Killeen, S.L., Geraghty, A.A., Lawton, E.M., Byrne, D., Sanchez-Gallardo, R., Nori, S.R.C., Nielsen, I.B., Wortmann, E., Matthews, E., O'Flaherty, R., Rudd, P.M., Groeger, D., Shanahan, F., Salдова, R., McAuliffe, F.M., Van Sinderen, D. and Cotter, P.D. (2023) 'Detailed mapping of Bifidobacterium strain transmission from mother to infant via a dual culture-based and metagenomic approach', Nature Communications, 14(1), 3015 (14pp). doi: 10.1038/s41467-023-38694-0
Type of publication	Article (peer-reviewed);journal-article
Link to publisher's version	10.1038/s41467-023-38694-0
Rights	© The Author(s) 2023. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license,

	visit http://creativecommons.org/licenses/by/4.0/ - https://creativecommons.org/licenses/by/4.0/
Download date	2026-10-12 05:27:27
Item downloaded from	https://hdl.handle.net/10468/15305






UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh



Supplementary Figure 1. a Heatmap representing the top 10 species of each sample type, whereby increased relative abundance for that species goes from yellow to purple on the heat gradient. Each colour bar represents a distinct metadata feature. Samples are clustered by Bray-Curtis, and slices are split by timepoint. **b** Alpha diversity of both maternal (16W $n=121$, 34W $n=129$, 1M $n=118$) and infant (Del $n=53$, 1W $n=45$, 1M $n=118$) stool samples. Boxplots highlight the middle 50% of data with the median relative abundance shown in the central line. The whiskers extend to show the range of values. Asterisks indicate significant differences from two-sided t -tests rooted to 16-weeks maternal sample. **c** Number of total *Bifidobacterium* isolated from each sample type split by timepoint. **d** envfit covariate analysis revealing the effect size of 22 different covariates on the microbiota beta diversity of each sample type. Significant associations are denoted with a black asterisk. Infant male/female split is 62/70, respectively. **e** Total number of metagenome assembled genomes (MAGS) recovered from each sample site. **f** Summary statistics for all sequenced *Bifidobacterium* genomes.



Supplementary Figure 2. a Transmitted *Bifidobacterium* strains identified by strain isolation.

Transmitted strains are highlighted by green colour strips in each plot.

b StrainPhlAn3-

determined normalised phylogenetic distance of transmitted trains ($n=27$) identified by strain

isolation. Boxplots highlight the middle 50% of data with the median relative abundance


shown in the central line. The whiskers extend to show the range of values. **c** SNP distance of

strains between ($n=41,129$) and within dyads with significant difference ($p = 1.1 \times 10^{-12}$)


denoted by asterisk determined by two-sided t -test. Boxplots highlight the middle 50% of data

with the median relative abundance shown in the central line. The whiskers extend to show


the range of values.



Supplementary Figure 3. a Stratification of all strains transferred by each combination of test method used in this study. **b** nMDS plot of the infant stool microbiome at 1-month, red arrows indicate the direction of effect of significant covariates as determined by envfit. **c** Comparison of the relative abundance within the infant 1-month stool of species of which strains were found to transfer, split by samples that displayed any transfer event. Boxplots highlight the middle 50% of data with the median relative abundance shown in the central line. The whiskers extend to show the range of values. Asterisks indicate significant difference of Bonferroni corrected p -value < 0.05 . **d** Heatmap of all species above 1% relative abundance in at least a single 1-month infant stool sample. Top annotations indicate samples where a transfer of one of the significant covariates from (b) are present.



(b)



Supplementary Figure 4. a Density plots indicating the number of detected strain sharing as determined by different normalised phylogenetic distance thresholds (horizontal dashed lines). **b** Midpoint rooted phylogenetic tree of alignment of SNP regions of *B. breve* strains from this study and 92 publicly available *B. breve* genomes.