

Title	Perceptions of adolescents and young adults with allergy and/or asthma and their parents on EAACI guideline recommendations about transitional care: A European survey
Authors	Khaleva, Ekaterina;Knibb, Rebecca;DunnGalvin, Audrey;Vazquez-Ortiz, Marta;Comberiati, Pasquale;Alviani, Cherry;Garriga-Baraut, Teresa;Gowland, M. Hazel;Gore, Claudia;Angier, Elizabeth;Blumchen, Katharina;Duca, Bettina;Hox, Valérie;Jensen, Britt;Mortz, Charlotte G.;Pite, Helena;Pfaar, Oliver;Santos, Alexandra F.;Sanchez-Garcia, Silvia;Timmermans, Frans;Roberts, Graham
Publication date	2021-09-26
Original Citation	Khaleva, E., Knibb, R., DunnGalvin, A., Vazquez-Ortiz, M., Comberiati, P., Alviani, C., Garriga-Baraut, T., Gowland, M. H., Gore, C., Angier, E., Blumchen, K., Duca, B., Hox, V., Jensen, B., Mortz, C. G., Pite, H., Pfaar, O., Santos, A. F., Sanchez-Garcia, S., Timmermans, F. and Roberts, G. (2021) 'Perceptions of adolescents and young adults with allergy and/or asthma and their parents on EAACI guideline recommendations about transitional care: A European survey', Allergy. doi: 10.1111/all.15109
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1111/all.15109
Rights	© 2021, John Wiley & Sons, Inc. This is the peer reviewed version of the following article: Khaleva, E., Knibb, R., DunnGalvin, A., Vazquez-Ortiz, M., Comberiati, P., Alviani, C., Garriga-Baraut, T., Gowland, M. H., Gore, C., Angier, E., Blumchen, K., Duca, B., Hox, V., Jensen, B., Mortz, C. G., Pite, H., Pfaar, O., Santos, A. F., Sanchez-Garcia, S., Timmermans, F. and Roberts, G. (2021) 'Perceptions of adolescents and young adults with allergy and/or asthma and their parents on EAACI guideline recommendations about transitional care: A European survey', Allergy, doi: 10.1111/all.15109, which has been published in final form at https://doi.org/10.1111/all.15109 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Download date	2025-04-25 22:53:24
Item downloaded from	https://hdl.handle.net/10468/12048



DR. EKATERINA KHALEVA (Orcid ID : 0000-0002-2220-7745)

DR. REBECCA CLAIRE KNIBB (Orcid ID : 0000-0001-5561-0904)

DR. AUDREY DUNNGALVIN (Orcid ID : 0000-0002-1540-3959)

DR. CHERRY ALVIANI (Orcid ID : 0000-0003-1527-0495)

DR. CHARLOTTE G. MORTZ (Orcid ID : 0000-0001-8710-0829)

DR. HELENA PITE (Orcid ID : 0000-0002-7300-928X)

PROF. OLIVER PFAAR (Orcid ID : 0000-0003-4374-9639)

DR. ALEXANDRA SANTOS (Orcid ID : 0000-0002-7805-1436)

PROF. GRAHAM C ROBERTS (Orcid ID : 0000-0003-2252-1248)

Article type : Position Paper

Perceptions of adolescents and young adults with allergy and/or asthma and their parents on EAACI guideline recommendations about transitional care: A European survey.

Manuscript Acceptance Date: 08-Aug-2021

Ekaterina Khaleva¹, Rebecca Knibb², Audrey DunnGalvin^{3,4}, Marta Vazquez-Ortiz⁵, Pasquale Comberiati^{6,7}, Cherry Alviani^{1,8}, Teresa Garriga-Baraut⁹, M Hazel Gowland¹⁰, Claudia Gore^{5,11}, Elizabeth Angier¹², Katharina Blumchen¹³, Bettina Duca^{5,11}, Valérie Hox¹⁴, Britt Jensen¹⁵, Charlotte G Mortz¹⁵, Helena Pite¹⁶, Oliver Pfaar¹⁷, Alexandra F. Santos¹⁸⁻²⁰, Silvia Sanchez-Garcia²¹, Frans Timmermans²², Graham Roberts^{1,8,23}

Affiliations:

1. Faculty of Medicine, University of Southampton, Southampton, UK
2. School of Psychology, College of Health and Life Sciences, Aston University, Birmingham, UK

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ALL.15109](https://doi.org/10.1111/ALL.15109)

This article is protected by copyright. All rights reserved

3. Applied Psychology and Paediatrics and Child Health, University College Cork, Cork, Ireland
4. Paediatrics and Child Infectious Diseases, First Moscow State Medical University, Russia.
5. Section of Inflammation, Repair and Development, National Heart and Lung Institute, Imperial College London. United Kingdom
6. Department of Clinical and Experimental Medicine, Section of Paediatrics, University of Pisa, 56126 Pisa, Italy
7. Department of Clinical Immunology and Allergology, I.M. Sechenov First Moscow State Medical University, 119991 Moscow, Russia.
8. The David Hide Asthma and Allergy Research Centre, St Mary's Hospital, Isle of Wight, UK
9. Unitat d'Allergologia Pediàtrica, Hospital Universitari Vall d'Hebron, Barcelona, Spain. Grup d'Investigació "Creixement i Desenvolupament", Institut de Recerca de l'Hospital Universitari Vall d'Hebron (VHIR), Barcelona, Spain.
10. Allergy Action, St Albans, UK
11. Department of Paediatrics, Imperial College Healthcare NHS Trust, London, UK
12. Primary Care and Population Sciences, University of Southampton, Southampton, UK
13. Department of Paediatric and Adolescent Medicine, Paediatric Pneumology, Allergology and Cystic Fibrosis, University Hospital Frankfurt, Frankfurt am Main, Germany
14. Department of Otorhinolaryngology, Head and Neck Surgery, University Hospitals Saint-Luc, Brussels, Belgium.
15. Department of Dermatology and Allergy Centre, Odense Research Centre for Anaphylaxis (ORCA), Odense University Hospital, Klovevænget 15, DK-5000 Odense C, Denmark
16. Allergy Center, CUF Descobertas Hospital and CUF Tejo Hospital; CEDOC, Chronic Diseases Research Center, NOVA Medical School/Faculdade de Ciências Médicas, Universidade Nova de Lisboa, Lisbon, Portugal
17. Department of Otorhinolaryngology, Head and Neck Surgery, Section of Rhinology and Allergy, University Hospital Marburg, Philipps-Universität Marburg, Marburg, Germany
18. Department of Women and Children's Health (Paediatric Allergy), School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, United Kingdom
19. Peter Gorer Department of Immunobiology, School of Immunology and Microbial Sciences, King's College London, London, United Kingdom
20. Children's Allergy Service, Guy's and St Thomas' Hospital, London, United Kingdom; Asthma UK Centre in Allergic Mechanisms of Asthma, London, United Kingdom
21. Allergy Department, Hospital Infantil Universitario del Niño Jesús, Madrid, Spain
22. Nederlands Anafylaxis Netwerk – European Anaphylaxis Taskforce, Dordrecht, The Netherlands
23. NIHR Southampton Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust, Southampton, UK

Address for correspondence:

Professor Graham Roberts, Paediatric Allergy and Respiratory Medicine, University Child Health (MP803), University Hospital, Southampton NHS Foundation Trust, Tremona Road, Southampton SO16 6YD, UK.

Tel.: +44 (0) 2380796160 E-mail: g.c.roberts@soton.ac.uk

Short title: Feedback on recommendations about transitional care in allergy and asthma.

ACKNOWLEDGEMENTS

We would like to acknowledge the support of EAACI in running the survey. We would like to thank Millie Basu, Lea Blum, Anna Borushko, Davide Caimmi, Antoine Deschildre, Amandine Divaret-Chauveau, Anastasiya Kolotilina, Sarah Lyons, Sonia Martin-Pereda, Maura Pedrini and Lea Tammen, for assisting in generating different language versions of the adolescents/young adults and parent survey. We would like to thank the following for distributing the survey: Federasma e Allergie Onlus; Associazione Respiriamo Insieme; Associazione Laziale Asma e Malattie Allergiche (ALAMA); Associazione Liguri Allergici (ALA); Associazione Asma Sardegna Onlus; Cibo Amico Associazione per le Allergie Alimentari e l'Anafilassi; Associazione Nazionale Dermatite Atopica (ANDeA); Associazione Food Allergy Italia; APS Chiara Ribechini; Uniamo Federazione Italiana Malattie Rare onlus, Ari-aaa3 ONLUS; Società Italiana di Allergologia e Immunologia Pediatrica (SIAIP); Società Italiana di Allergologia; Asma e Immunologia Clinica (SIAAIC); Montserrat Alvaro, Jaime Lozano, Joan Bartra, Fundació Humans, Fundació Lovexair, Edryx Healthcare; patients and parents from the pediatric allergy unit of the University Hospital Vall d'Hebron, Barcelona; patients and parents from Centre Mèdic del Berguedà and Gabinet Mèdic del Berguedà, Berga, Barcelona; Institut Vescomtat de Cabrera, Hostalric; Institut Blanca d'Anjou, El Perelló; Societat Catalana d'Al·lergologia i Immunologia Clínica (SCAIC); Sociedad Española de Inmunología Clínica y Alergología Pediátricas (SEICAP); Sociedad Española de Alergología e Inmunología Clínica (SEAIC); Lara Pimenta; Mário Morais Almeida; Andre Moreira (EAACI SPC Chair), Portuguese Association of Asthmatics; Deutscher Allergie- und Asthmabund (DAAD) (Sabine Schnadt),

Mönchengladbach, Germany; Daniel Munblit; Maria Kardakova; 'Allergomoms', 'АтД по-доказательному' (Anna Taranova); Danish patient organization; Asthma Allergy Denmark; Nederlands Anafylaxis Netwerk; Allergy UK (Carla Jones); Anaphylaxis Campaign (Lynne Regent); Hilary Allen; Aideen Byrne; John Fitzsimons; Antoine Deschildre; Amandine Divaret-Chauveau; "Association Asthme et Allergies" (Christine Rolland); "AFPRAL" (Pascale Couratier); Elena Bradatan; Virginie Doyen; and Swiss patient organization and all the patients and parents who completed the survey. Finally, we would also like to thank the EAACI Executive Committee for their helpful comments and suggestions.

ABSTRACT

Background. The European Academy of Allergy and Clinical Immunology has developed a guideline to provide evidence-based recommendations for healthcare professionals to support the transitional care of adolescents and young adults (AYA) with allergy and/or asthma. The goal of this work was to ensure that the draft recommendations are also important for patients.

Methods. We surveyed patients aged 11-25 years with allergy and/or asthma and their parents across Europe between 17th February and 16th March 2020. The multilingual survey was distributed through national allergy and asthma patient organisations in Europe as well as through social media.

Results. A total of 1210 responses from 24 European countries were collected. There were 415 (34.3%) AYA and 795 (65.7%) parents. The majority of AYA (72.3%) and parents (81.9%) were female. Patients had a history of asthma (61.1%), allergic rhinoconjunctivitis (54.1%), food allergy (53.8%), atopic eczema (42.6%), and anaphylaxis (28.8%). All recommendations achieved the median score of either 'important' or 'very important'. The least supported recommendations were the use of joint clinics with both paediatric and adult physicians attending and the use of web-based or mobile technologies for communication with the AYA. The most supported recommendation was checking that the AYA is knowledgeable and compliant with their prescribed medication. Qualitative analysis revealed conditional approval for some recommendations.

Conclusions. There was agreement from patients and parents on the importance of the draft recommendations on transitional care for adolescents and young adults with allergy and/or asthma

and their parents. The recommendations now need to be implemented into clinical practice across Europe.

Key words: adolescents, allergy, asthma, transition, survey, young adults

Word count: 2586 words; **Tables:** 2; **Figures:** 3.

INTRODUCTION

Transition has been defined as an 'active and evolving process that addresses the medical, psychosocial, and educational needs of young people as they prepare to move from child- to adult-centered health care' ¹. Several guidelines on general transitional care have been published by the European Academy of Paediatrics², Canadian Association of Paediatric health centres³, American Academy of Pediatrics⁴, and National Institute for Health and Care Excellence⁵. Disease specific guidelines are also available to optimise transition for adolescent and young adults (AYA) with different long-term conditions⁶⁻⁸. Transition programmes have shown significant improvements in patient care during transition leading to low rates of loss of follow up, high scores for AYA's satisfaction with transition and self-efficacy in managing their disease⁹.

Recently, the European Academy of Allergy and Clinical Immunology (EAACI) published the first European guideline on the effective transition of adolescents and young adults (AYA) with allergy and/or asthma¹⁰. As part of the guideline development process, an online survey was conducted to ensure the draft recommendations developed by the EAACI Task Force (TF) are important for AYA with allergy and asthma and their parents or carers (hereafter referred to as "parents") across Europe. Involving AYA in refining the European recommendations is an essential step because these are the patients who need to be transitioned and adapt to a new type of care. Understanding what is important to them helps to facilitate a smooth transition, ensure treatment adherence and subsequently improve health care outcomes. Parents also play an important role in preparing and supporting adolescents during transition to becoming independent adults. Parents must make a challenging transition themselves from taking full responsibility for their adolescent's healthcare to their child self-managing their disease and becoming a competent patient. It has been shown that assessment of parental perceptions on the transition of AYA with congenital heart defects offered insights into how transition planning could be optimised¹¹.

By obtaining the perspective of AYA's and their parents on EAACI draft recommendations for transitional care, we aimed 1) to evaluate the importance of each recommendation independently for each group and 2) to identify additional factors that need to be included in the 'other considerations' section of recommendations.

METHODS

Study design

A quantitative, online, cross-sectional survey was conducted. The survey (available in the supplementary materials) was based on recommendations developed by the members of the EAACI Adolescent and Young Adult Task Force (TF) who come from a range of disciplinary and clinical backgrounds, including allergists (specialists and subspecialists), general practitioners, paediatricians, dermatologists, otolaryngologists, adult physicians, nurses, psychologists and patient representatives. Recommendations are based on the results of systematic reviews on the challenges of AYA with allergic conditions¹², interventions for these patients¹³ as well as generic transition recommendations from evidence-based guidelines over the last 5 years. In addition, three rounds of a Delphi survey were conducted among TF members in order to achieve a consensus for D level recommendations¹⁰.

Participants and data collection

We invited AYA, aged 11-25 with allergy and asthma and their parents across Europe who were able to read English, Dutch, Danish, German, Spanish, Portuguese, Italian, French or Russian, to participate in the survey. The survey was distributed through national allergy and asthma patient organisations in Europe (UK, The Netherlands, Italy, Portugal, Spain, Ireland, Germany, Russia, Denmark, France) who then disseminated the link to the survey in SurveyMonkey among their members. In addition, the survey was advertised on social media (e.g. Facebook, Twitter). Before accessing the questionnaire, potential respondents were informed about the survey's purpose, the organisations conducting the survey and the average time required to complete. Responses were collected between 17th February and 16th March 2020.

The questionnaire

The anonymised survey included 24 questions divided between two parts: 5 questions about demographic information (AYA or parent, age, gender, country, allergic diseases) and 19 questions about the level of importance of each recommendation. Recommendations were divided into five groups, namely 1) generic advice, 2) treatment of allergy and asthma, 3) self-management, 4)

psychological issues and help, 5) support from family, friends and others. Participants were asked to rank the level of importance for each recommendation using a 5-point scale: 1 "Not important," 2 "Slightly Important," 3 Fairly important", 4 "Important," and 5 "Very Important," plus a "No opinion" option was available. An average score of at least 2 was set as the level for acceptance of the recommendation. Options for free-text responses were provided. The wording of the questionnaire was carefully checked to ensure it could be understood by this lay audience, translated into eight languages (English, German, Spanish, Portuguese, Italian, Danish, Dutch, French and Russian) and back translated into English to ensure validity and accuracy. Reading age was tested prior to dissemination of the survey to ensure clarity and understanding. A group of target participants also tested the time required to complete the survey, which ranged from 15 to 20 minutes.

Qualitative analysis

Qualitative data analysis was used to summarise free-text comments from AYA and parents. All comments from AYA and parents were analysed separately. Comments in languages other than English were translated by one TF member and checked by a second TF member. Braun and Clarke's steps for thematic analysis were used for analysis¹⁴. Each comment was coded and the codes were then combined into themes. In order to ensure that responder views were correctly interpreted, each comment and its code was reviewed in duplicate by patient representatives, clinicians and psychologists from the EAACI TF (EK, GR, CA, RCK, ADG, PC, T. G-B, MHG). Any discrepancies were resolved through discussion and, if necessary, a third reviewer (M.V-O) was consulted. Codes were then split based on the direction of the content: supportive, conditionally supportive or non-supportive (Supplementary materials).

Statistical analysis

All data were collected and analysed using SPSS software version 25.0. Descriptive statistics were used to describe respondent characteristics. Medians with lower and upper quartiles are presented for continuous variables given the distribution of the data. Frequency tables with percentages are provided for categorical variables.

A minimum of 50 responders per country was required for the comparison of data between countries. The comparisons were performed by using the Kruskal Wallis test. Summary tables were used to represent the results. Data were considered significant if statistical tests produce a p-value of <0.05.

RESULTS

Respondent demographics and characteristics

Among 1425 received responses, 215 (16%) were excluded as they were either incomplete or came from outside Europe. A total of 1210 responses from 24 European countries were analysed. The most

common countries of residence were Italy (20.7%), Portugal (17.0%), France (15.6%) and Russia (15.4%). There were 415 (34.3%) AYA and 795 (65.7%) parents. The majority of AYA (72.3%) and parents (81.9%) were female. Patients had a history of asthma (61.1%), allergic rhinoconjunctivitis (54.1%), food allergy (53.8%), eczema (42.6%), anaphylaxis (28.8%). Most AYA had one (29.7%) or two (26.9%) allergic conditions but 22.0% had more than four allergic comorbidities. Respondents' characteristics are listed in Table 1.

Importance of the recommendations on transitional care

All recommendations were (at minimum) scored as "important" (median score of 4 on the 5-point scale) (Table 2). The most supported recommendation was that of checking that the AYA is knowledgeable and compliant with their prescribed medication, with 68% of AYA and 77.7% of parents reporting this as "very important". The least supported recommendations related to the use of joint clinics with both paediatric and adult physicians attending (only 27.7% AYA and 38.1% parents reported this as "very important") and the use of web-based mobile technologies for communication with the AYA (only 27.0% AYA and 35.6% of parents reported this as 'very important'). In addition to overall data, a sensitivity analysis showed that the results were similar in individual countries with more than 50 responses (Table S1).

Of the 1356 free-text comments received, 978 (72.1%) were made by parents. Summary of the feedback with all themes from AYA and parents may be found in the supplementary materials (Tables S2-S19). Figure 1 (and Figure S1) show the key 'supportive, conditionally supportive and not supportive comments for generic recommendations, Figure 2 presents comments on treatment and self-management of allergy, skin disease and asthma and comments on psychological issues and help, and support from family, friends and others are found in Figure 3. In general, AYA's and parents' feedback was similar (Tables S2-S19). There were some notable exceptions such as parents were more likely to comment that transition should start earlier in adolescence (Table S2).

Almost all comments were supportive with a few conditional ones which were used to improve the recommendations. For example, respondents suggested that a personal action plan should be developed not only when AYA is about to be transferred to the adults' department, but should instead be utilised at an early stage before adolescence. In addition, it was suggested that the action plan should be regularly reviewed by all HCP involved in the individual's care. With regards to motivational interviewing, participants proposed that active involvement of the AYA in discussion would improve self-management of asthma and quality of life. Adolescents suggested that conversation about possible stressful life events that may impact disease control, should be explored in a very sensitive way, and that psychological interventions using cognitive behavioural therapy to improve adherence, self-management and symptom control are likely to be most effective for specific patients, for example, those struggling to self-manage their condition. Professionals who support those with difficulties, such

as psychologists, should have expertise in allergy and asthma. The family should be enrolled early in the transition process, however the shift from parent to self-management should be done gradually to support AYA empowerment. Telling friends may be difficult for some and AYA suggested that they should be supported and encouraged to start by talking to a few close friends. Parents added that there should be a balance between autonomy and safety and recognised that it may be challenging for them to have to reduce control of their child's life.

Key negative comments related to the suggestion that adolescents should learn about self-management before 11 years of age; that the use of aids to improve adherence might undermine AYA taking responsibility for their care; and some felt that AYA might not be happy to share the details of their allergic diagnoses with their friends.

DISCUSSION

The findings from this pan-European survey supports the value of draft guideline recommendations on the transitional care developed by the EAACI Adolescent and Young Adult Task Force for both AYA with allergic conditions and their parents. Conducting this survey in different European countries and health systems demonstrated that the recommendations are understood by AYA/parents and remain relevant. The rich qualitative data set of almost 1400 comments, and the thematic analysis of the responses to the open-ended questions provided additional insights on their opinions about each recommendation. From this survey we were able to further refine the recommendations based on the range of supportive, conditionally supportive, and minimal number of non-supportive free-text responses.

This survey contributes to the growing literature on the benefits of patient-centred care and understanding of patient perceptions and views for healthcare decision-making¹⁵. For example, patient-reported experience measures have been used to guide quality improvement across different settings¹⁶. Moreover, a set of considerations to engage AYA in research have been proposed¹⁷ and several guidelines on transitional care involved this age group and their parents in the development of guideline recommendations^{6,18}.

Survey participants recognised the value of actively monitoring adherence to treatment throughout transition, but did not consider an overlap in care between paediatric and adult HCP to be particularly important. This finding may be because some countries do not have separate paediatric and adult specialists (allergists, dermatologists, respiratory physicians) and care is mostly led by general practitioners or specialists who treat patients of all ages. In contrast, AYA with inflammatory bowel disease rated joint consultation with both HCP as one of the most valuable features of the transition care program¹⁹. Other methods of communication between HCP and AYA such as web-based, mobile technologies were also considered less important. Research indicates that AYA generally

prefer using web-based methods of communications²⁰ and are less interested in using social media sites for communication with HCP due to privacy concerns²¹. These two less popular recommendations align with other guidelines, which emphasise the importance of suitable use of technology for interaction between patients and HCP^{1,5,8} and joint consultations^{8,22}.

Limitations inherent in the survey method impacted our ability to investigate possible reasons for our findings. Further qualitative interviews might identify why AYA do not fully support joint care by adult and paediatric HCP or the use of technologies for communication with HCP. Some respondents reported that self-management training should start before 11 years of age. We would agree that is appropriate for some AYA in line with developmentally appropriate healthcare.²³ Some AYA find technological solutions, such as an alarm on a mobile phone, to be very helpful to improve their treatment adherence.²⁴ Some AYA reported concerns about what their friends might think if they shared the details of their allergic diagnoses with them, although others reported this as a positive experience with AYA receiving considerable support from their close friends.²⁵

Strengths and limitations of the survey

Although the survey covered most European countries, it may not represent the opinions of AYA and their parents in countries not surveyed. Technical limitations meant that comments were not included for the second recommendation; however ranking did allow us to assess importance in both groups. The representativeness of the survey is likely to be high given the large number of responses from patients with a range of allergic conditions, and their parents. Furthermore, 29.7% had one and 22% more than 4 allergic comorbidities. To the best of our knowledge, this survey is the first of its kind to evaluate draft recommendations by patients and their parents through a pan-European survey design and, the high number of free-text comments analysed in this survey allowed for more in-depth understanding of the patient perspective on recommendations concerning transition and transfer of AYA with allergy and/or asthma from paediatric to adult services.

Implications

There are several important implications arising from the findings of this survey. Firstly recommendations on transitional care are important for AYA with allergy and/or asthma and therefore, should be implemented within clinical practice across Europe. Secondly, qualitative analysis of comments can help to refine recommendations by allowing for the provision of additional information from the patient perspective.

All generic as well as allergy specific recommendations on adherence, self-management, support and psychological help received positive feedback from both AYA with allergies and/or asthma and their parents in Europe. Recommendations are intended to be useful, practical, facilitate local teams

to work together and promote transition and transfer where necessary. We would like to emphasise that the application of these recommendations and the effectiveness of the healthcare for adolescents and young adults with allergies and asthma should be reevaluated through regular audits¹⁰. Assessment should involve AYA and families, as well as policy makers, researchers and government agencies²⁶. It is hoped that harmonisation of the transition process and practice will improve psychological and physical outcomes as well as the quality of life of these patients.

CONCLUSIONS

This pan-European survey showed that adolescents and young adults with allergy and/or asthma and their parents find draft EAACI recommendations on transitional care for these conditions important or very important. Qualitative analysis of responses to open-ended questions confirmed the value of these recommendations and provided additional information from a patient perspective. Next steps should include implementation of recommendations into clinical practice taking into account differences between European countries in how transition may be organised.

CONFLICTS OF INTEREST

GR and RK report research funding from Asthma UK and National Institutes of Health Research into the challenge associated with asthma during adolescents. FT reports being a parent of a young adult with food allergy. None of the other authors have anything to disclose.

CONTRIBUTIONS

Survey concept and design: GR, EK, M.V.-O. Statistical analysis and interpretation of data: EK, GR. Qualitative analysis: EK, GR, CA, RCK, AG, PC, T.G-B, MHG. Drafting of the original manuscript: EK,GR. Reviewed and edited the manuscript: GR, M.V.-O., EA, KB, RCK, PC, CA, BD, CGM, AD, CG, VH, BJ, HP, AFS, T.G-B, SSG, MHG, FT, OF. All authors provided critical review of the manuscript and approved the final version. Obtained funding, GR, M.V.-O.

REFERENCES

1. Vajro P, Fischler B, Burra P, et al. The Health Care Transition of Youth With Liver Disease Into the Adult Health System: Position Paper From ESPGHAN and EASL. *J Pediatr Gastroenterol Nutr.* 2018;66(6):976-990.

2. Mazur A, Dembinski L, Schrier L, Hadjipanayis A, Michaud PA. European Academy of Paediatric consensus statement on successful transition from paediatric to adult care for adolescents with chronic conditions. *Acta Paediatr.* 2017;106(8):1354-1357.
3. Velazquez JR, Atzin AJ, Garcia M, Teran LM, Perez-Rodriguez M. Increased frequency of CD56-CD16+/-NK cells in exacerbated asthma respiratory disease-A suggestive role as biomarker. *Allergy: European Journal of Allergy and Clinical Immunology.* 2017;72 (Supplement 103):660-661.
4. White PH, Cooley WC, Transitions Clinical Report Authoring G, American Academy Of P, American Academy Of Family P, American College Of P. Supporting the Health Care Transition From Adolescence to Adulthood in the Medical Home. *Pediatrics.* 2018;142(5):11.
5. Komalla V, Allam VSRR, Kwok PCL, et al. A phospholipid-based formulation for the treatment of airway inflammation in chronic respiratory diseases. *European Journal of Pharmaceutics and Biopharmaceutics.* 2020;157:47-58.
6. Foster HE, Minden K, Clemente D, et al. EULAR/PReS standards and recommendations for the transitional care of young people with juvenile-onset rheumatic diseases. *Ann Rheum Dis.* 2017;76(4):639-646.
7. Ludvigsson JF, Agreus L, Ciacci C, et al. Transition from childhood to adulthood in coeliac disease: the Prague consensus report. *Gut.* 2016;65(8):1242-1251.
8. Brooks AJ, Smith PJ, Cohen R, et al. UK guideline on transition of adolescent and young persons with chronic digestive diseases from paediatric to adult care. *Gut.* 2017;66(6):988-1000.
9. Walter M, Kamphuis S, van Pelt P, de Vroed A, Hazes JMW. Successful implementation of a clinical transition pathway for adolescents with juvenile-onset rheumatic and musculoskeletal diseases. *Pediatr Rheumatol Online J.* 2018;16(1):50.
10. Roberts G, Vazquez-Ortiz M, Knibb R, et al. EAACI Guidelines on the effective transition of adolescents and young adults with allergy and asthma. *Allergy.* 2020;75(11):2734-2752.
11. Gaydos LM, Sommerhalter K, Raskind-Hood C, et al. Health Care Transition Perceptions Among Parents of Adolescents with Congenital Heart Defects in Georgia and New York. *Pediatr Cardiol.* 2020;41(6):1220-1230.
12. Vazquez-Ortiz M, Angier E, Blumchen K, et al. Understanding the challenges faced by adolescents and young adults with allergic conditions: a systematic review. *Allergy.* 2020;Aug;75(8):1850-1880.
13. Knibb RC, Alviani C, Garriga-Baraut T, et al. The effectiveness of interventions to improve self-management for adolescents and young adults with allergic conditions: a systematic review. *Allergy.* 2020;Aug;75(8):1881-1898.

14. Braun V, Clarke V. Using thematic analysis in psychology. *Qualitative Research in Psychology*. 2006;3(2):77-101.
15. Rand L, Dunn M, Slade I, Upadhyaya S, Sheehan M. Understanding and using patient experiences as evidence in healthcare priority setting. *Cost Effectiveness and Resource Allocation*. 2019;17(1):20.
16. Gleeson H, Calderon A, Swami V, Deighton J, Wolpert M, Edbrooke-Childs J. Systematic review of approaches to using patient experience data for quality improvement in healthcare settings. *BMJ Open*. 2016;6(8):e011907.
17. Lolocono Merves M, Rodgers CR, Silver EJ, Sclafane JH, Bauman LJ. Engaging and sustaining adolescents in community-based participatory research: structuring a youth-friendly community-based participatory research environment. *Fam Community Health*. 2015;38(1):22-32.
18. Le Roux E, Mellerio H, Jacquin P, et al. Practical generic guidelines for paediatric-to-adult transition for adolescents with chronic disease. *Eur J Public Health*. 2019;29(3):442-448.
19. Blazquez ML, Hernani MT, Calatayud GA, et al. Transition Care Program of Adolescents With Inflammatory Bowel Disease: Perceptions From A Survey for Patients. *Inflamm Bowel Dis*. 2019;25(3):e22-e23.
20. Low JK, Manias E. Use of Technology-Based Tools to Support Adolescents and Young Adults With Chronic Disease: Systematic Review and Meta-Analysis. *JMIR Mhealth Uhealth*. 2019;7(7):e12042.
21. Applebaum MA, Lawson EF, von Scheven E. Perception of transition readiness and preferences for use of technology in transition programs: teens' ideas for the future. *Int J Adolesc Med Health*. 2013;25(2):119-125.
22. Italian Society of Paediatric Gastroenterology H, Nutrition IAOHG, Endoscopists ISoEISoG, et al. Transition of gastroenterological patients from paediatric to adult care: A position statement by the Italian Societies of Gastroenterology. *Dig Liver Dis*. 2015;47(9):734-740.
23. Farre A, Wood V, Rapley T, Parr JR, Reape D, McDonagh JE. Developmentally appropriate healthcare for young people: a scoping study. *Arch Dis Child*. 2015;100(2):144-151.
24. Holley S, Walker D, Knibb R, et al. Barriers and facilitators to self-management of asthma in adolescents: An interview study to inform development of a novel intervention. *Clin Exp Allergy*. 2018;48(8):944-956.
25. Macadam C, Barnett J, Roberts G, et al. What factors affect the carriage of epinephrine auto-injectors by teenagers? *Clin Transl Allergy*. 2012;2(1):3.
26. Canadian Association of Paediatric Health Centres. A Guideline for Transition From Paediatric to Adult Health Care for Youth with Special Health Care Needs: A National Approach. <https://ken.childrehealthcarecanada.ca/xwiki/bin/download/Transitioning+from+Paediatric+to+Adult>

Table 1. Demographics of survey responders.

All responders (n=1210)	Number (%) of responders
Adolescents	415 (34.3)
Male	115 (27.7)
Female	300 (72.3)
Parents	795 (65.7)
Male	144 (18.1)
Female	651 (81.9)

Type of allergic disease	
Asthma	739 (61.1)
Allergic rhinoconjunctivitis	655 (54.1)
Food allergy	651 (53.8)
Eczema	515 (42.6)
Anaphylaxis	348 (28.8)
Urticaria	248 (20.5)
Drug allergy	132 (10.9)
Venom allergy	68 (5.6)
Number of allergic diseases*	
1	359 (29.7)
2	326 (26.9)
3	257 (21.2)
≥4	266 (22.0)
Countries	
Italy	251 (20.7)
Portugal	206 (17.0)
France	189 (15.6)
Russia	186 (15.4)
United Kingdom	103 (8.5)
Spain	71 (5.9)
Denmark	54 (4.5)
Netherlands	46 (3.8)
Germany	26 (2.1)
Ireland	25 (2.1)
Others**	53 (4.4)

* Anaphylaxis is not included as a separate condition. **: Bulgaria n=1 (0.1 %), Greece n=1 (0.1%), Luxembourg n=1 (0.1 %), Poland n=1 (0.1 %), Romania n=1 (0.1 %), Georgia n=1 (0.1 %), Azerbaijan n=1 (0.1 %), Turkey n=1 (0.1 %), Switzerland n=6 (0.5%), Kazakhstan n=3 (0.2%), Belarus n=6 (0.5%), Ukraine n=15 (1.2%), Belgium n=15 (1.2%).

Table 2. The level of importance for each recommendation as regarded by participants .

Recommendation	AYA n=415 Median (IQR)	Parents n=795 Median (IQR)
Generic advice		
1. Young people should start to learn how to manage their asthma, allergies and skin disease when they are about 11-13 years old.	5 (4,5)	5 (4,5)
2. It is important to think about:		
Making sure that clinics have a plan/special document about how to care for young people as they grow up.	5 (4,5)	5 (4,5)
Telling the young person and their parents or carers about how the plan will work and how it will change as they grow up.	5 (4,5)	5 (4,5)
Using a checklist to find out when the young person is ready to take more responsibilities for their asthma, allergy and skin disease as they grow up.	4 (4,5)	5 (4,5)
Checking that the young person is able to and takes the medicines they have been given.	5 (4,5)	5 (5,5)
If the young person has to move across from a children's clinic to one for adults, it would be helpful for them to see both children's and adult's doctors in one clinic transiently while they get used to the change.	4 (3,4)	4 (3,5)
The doctors and nurses caring for young people in children's and adult clinics should have regular meetings to discuss their care.	4 (3,4)	4 (4,5)
3. Doctors and nurses could use web-based and other mobile technologies such as texts or skype to communicate with the young person.	4 (3,4)	4 (3,5)
4. It may be helpful for doctors and nurses to talk to young people about how their asthma, allergies and skin disease may affect their social life (e.g. when being with friends or family), education and career plans.	5 (4,5)	5 (4,5)
5. Doctors, nurses and other medical staff should have special training to help young people with asthma, skin symptoms and allergies.	5 (4,5)	5 (4,5)
6. There should be regular checks of how well the clinic works to make sure it is	4 (4,5)	5 (4,5)

effective and helpful for young people.		
Treatment of allergy, skin disease and asthma		
7. The doctors and nurses should try to make the young people's treatment easy to follow.	4 (4,5)	5 (4,5)
8. Phone reminders, apps and other methods may be useful to help young people to remember their treatment and take more responsibility for looking after their asthma, skin disease and allergies.	4 (4,5)	4 (4,5)
Self-management of allergy, skin disease and asthma		
9. A personal action plan covering what to do would help young people manage their asthma, skin disease or allergies.	4 (4,5)	5 (4,5)
10. It would be helpful during the hospital visit to focus on issues and ways to manage asthma, skin disease and allergies where the young person is less confident.	4 (4,5)	5 (4,5)
11. Young people and their families might want guidance from doctors and nurses on how to manage their asthma, skin disease or allergies when the young person is at social events (e.g. sports, celebration, holidays).	4 (3,5)	5 (4,5)
12. Young people could learn from other young people with asthma, skin disease and allergies about how to manage their life.	4 (3,5)	4 (4,5)
13. Doctors or nurses should have conversations with young people designed to strengthen their motivation and commitment to improving their asthma management.	4 (4,5)	5 (4,5)
Psychological issues and help		
14. Doctors and nurses should look out for young people who feel anxious or depressed as these may affect their asthma, skin disease and allergies.	5 (4,5)	5 (4,5)
15. Doctors and nurses should find out if young people have experienced stressful events (such as parents' divorce or bullying) which may affect their asthma, skin disease and allergies.	4 (4,5)	4 (4,5)
16. A psychologist may be able to help young people to manage their asthma, skin disease and allergies better.	4 (3,4)	4 (4,5)
Support from family, friends and others		
17. Families should be encouraged to support young people as they start to manage their asthma, skin disease and allergies.	5 (4,5)	5 (4,5)
18. Young people should be encouraged to let their friends know about their asthma,	5 (4,5)	5 (4,5)

skin disease and allergies and how they can help in an emergency.		
19. Clinics should recommend reliable websites and other useful sources of information about asthma, skin disease and allergies to young people.	4 (4,5)	4 (4,5)

IQR, interquartile range. Potential responses were 1 “Not important,” 2 “Slightly Important,” 3 Fairly important”, 4 “Important,” and 5 “Very Important,” plus a “No opinion” option was available. AYA, adolescents and young adults.

Figure legends.

Figure 1. Summary of feedback on generic recommendations


AAI, adrenaline autoinjectors; AYA, adolescents and young adults; GP, general practitioner; P, parents; HCP, healthcare professionals. The thematic map includes themes where the total number of comments for each theme ≥ 11 . Information in brackets specifies which group (AYA and/or P) has more than 11 comments in each theme. If none of the groups reported ≥ 11 comments in the theme but it has ≥ 11 total number of comments this theme is also included.

Figure 2. Summary of feedback on adherence and self-management recommendations

AAI, adrenaline autoinjectors; AYA, adolescents and young adults; GP, general practitioner; P, parents; HCP, healthcare professionals. The thematic map includes themes where the total number of comments for each theme ≥ 11 . Information in brackets specifies which group (AYA and/or P) has more than 11 comments in each theme. If none of the groups reported ≥ 11 comments in the theme but it has ≥ 11 total number of comments this theme is also included.

Figure 3. Summary of feedback on psychology and support recommendations

AAI, adrenaline autoinjectors; AYA, adolescents and young adults; GP, general practitioner; P, parents; HCP, healthcare professionals. The thematic map includes themes where the total number of comments for each theme ≥ 11 . Information in brackets specifies which group (AYA and/or P) has more than 11 comments in each theme. If none of the groups reported ≥ 11 comments in the theme but it has ≥ 11 total number of comments this theme is also included.








Figure 2. Summary of feedback on adherence and self-management recommendations





all_15109_f4.jpg