

Title	Response rates to questionnaire-based studies in the contemporary dental literature: A systematic review
Authors	Al Khalaf, Khaleel;O'Dowling Keane, Shane;da Mata, Cristiane;McGillycuddy, Catherine T;Chadwick, Barbara L;Lynch, Christopher D
Publication date	2022-09-15
Original Citation	Al Khalaf, K., O'Dowling Keane, S., da Mata, C., McGillycuddy, C. T., Chadwick, B. L. and Lynch, C. D. (2022) 'Response rates to questionnaire-based studies in the contemporary dental literature: A systematic review', Journal of Dentistry, 126, 104284 (7pp). doi: 10.1016/j.jdent.2022.104284
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.jdent.2022.104284
Rights	© 2022, The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/) - https:// creativecommons.org/licenses/by/4.0/
Download date	2025-05-16 06:21:41
Item downloaded from	https://hdl.handle.net/10468/13718



University College Cork, Ireland Coláiste na hOllscoile Corcaigh Contents lists available at ScienceDirect



Journal of Dentistry



Response rates to questionnaire-based studies in the contemporary dental literature: A systematic review



Khaleel Al Khalaf^{a,b,*}, Shane O'Dowling Keane^a, Cristiane da Mata^a, Catherine T. McGillycuddy^c, Barbara L. Chadwick^d, Christopher D. Lynch^a

^a Restorative Dentistry Department, University Dental School & Hospital, University College Cork, Wilton, Co. Cork, Ireland

^b Prosthodontics Department, Dammam Specialist Dental Centre, Dammam Medical Complex, Dammam, Saudi Arabia

^c General Dental Practitioner, Tralee, Co. Kerry, Ireland

^d School of Dentistry, Cardiff University, Heath Park, Cardiff, United Kingdom

ABSTRACT

Objectives: This systematic review aimed to investigate what is a reasonable response rate for dental questionnaire-based studies in recent literature and to assess the factors that affect the response rates.

Methods: We used MEDLINE/PubMed to search the dental literature of 2021 (January-October). Two reviewers independently assessed studies eligibility and extracted data using standardized electronic extraction form.

Results: One hundred and seventy-two studies were eligible, of these a total of 149 response rates were reported from 133 studies, whereas the remaining 39 studies were excluded as they did not report response rates. The median response rate across the included studies was 77% (mean = 70.8%). We found significant negative correlation between the response rate and the actual number of distributed questionnaires (sample size) (r = -0.4127; P < 0.001). We also found an association between the response rate and the area of distribution, e.g., national or international (P = 0.0012). However, a wide variation was observed in the quality of information reported within this review and we did not find clear evidence of association between the response rate and other variables such as questionnaire piloting, number of questions in the questionnaire and the journal impact factor.

Conclusions: The findings of this systematic review confirm the association between the response rate and the sample size, where the response rate increases when the sample size less than 300 participants. In addition, a higher response rate could be achieved when the study conducted within the same institution (e.g., university). *Significance*: Questionnaire-based research can provide answers to several questions that could not be answered by other types of research related to the field of dentistry, dental health practitioners and students' attitudes and behaviours and more. Questionnaire-based publications can effectively contribute to dental research; thus, dental journals should consider development of a minimum set of guidelines in the reporting of questionnaire-based manuscripts.

1. Introduction

Questionnaire-based research are widely published within the dental literature. This methodology can offer an objective methods of collecting information about people's knowledge, beliefs, attitudes, and behavior [1]. Questionnaires can be used as the sole research instrument e.g., cross-sectional surveys or within clinical trials and epidemiological studies [2]. However, concern exists in relation to what represents a 'reasonable' response rate – i.e., being of sufficient magnitude to reliably answer the questions posed, while excluding non-responder bias. Therefore a questionnaire should aim to obtain as representative a range of responses as possible and thereby provide reliable and valid answers to the research questions posed [3].

A brief review of the dental literature reveals a wide range of response rates – from as low as 7% on the assessment of the knowledge

and attitudes of western Australian dental health practitioners towards identifying and reporting child abuse [4] to as high as 100% on the assessment of the diagnostic skills of general dentists in different types of orthodontic malocclusions [5]. Thus, questionnaire response rates have become one of the "most controversial issues" which may affect the journal acceptance and publication of the questionnaire-based researches [6].

In 1997, Tan and Burke conducted a review that included a sample of 77 articles from four dental journals during the period 1989–1992. Their aim was to investigate the range and factors affecting response rates for mailed distributed questionnaire-based studies [7]. The authors found that 64% is the average response rate for the investigated questionnaire-based studies. Additionally, they suggested that questionnaire subject, incentives offered and length of the questionnaire may influence response rates [7]. However, since then, research methods,

https://doi.org/10.1016/j.jdent.2022.104284

Received 11 August 2022; Received in revised form 1 September 2022; Accepted 6 September 2022 Available online 8 September 2022 0300-5712/© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

^{*} Corresponding author at: Restorative Dentistry Department, University Dental School & Hospital, University College Cork, Wilton, Co. Cork, T12 E8YV, Ireland. *E-mail address:* k.alkhalaf@umail.ucc.ie (K. Al Khalaf).

computer technology, internet access, statistics and editorial policies have changed the practice of data collection.

Nowadays, electronic survey has become widely used among researchers. It minimizes the processing costs because it automates the transformation of raw data into electronic form, and combines the advantages of interviews (e.g., complex branching) with those of paperform surveys (e.g., standardization, anonymity) [8]. In 1984, Kraut used a computer network in an international corporation to conduct research on work behaviours and attitudes [9]. Then in 1986, a study aimed to conduct an experimental study by using both electronic and paper mail sample survey. The authors found that more respondents returned the paper survey (75%) than the electronic survey (67%). However, they reported less completion mistakes and faster returned response for the electronic surveys [8]. In literature, there is a conflict on whether email or paper-form questionnaires result in higher response rates. Some papers reported higher response rate for electronic questionnaires [10], others reported higher response rates for mail questionnaires [8,11], while others reported comparable response rates [12, 13].

Furthermore, email propagation or social media can be utilized to ensure reaching the most significant number of the targeted population. Hence, a higher response rate for the questionnaire-based study and a minimum non-responder bias might be obtained. However, participant recruitment through social media platforms or email propagation might be a challenging process, and the response rate cannot always be measured. A recent descriptive WhatsApp messenger-based crosssectional survey study found that social media platforms are reliable and could be used for disseminating information as well as a research tool among medical students and healthcare professionals. The study also found that more than two-thirds of medical students and healthcare professionals routinely use social media, although a 21% response rate was reported [14].

It has been claimed that using a mixed-mode approach will enhance the survey response rates or if mailed surveys are combined with e-mail follow-up [15]. In a study comparing different methods of administration, response rates close to 60% were achieved by mixed-mode contacts [12]. This approach, combining both mailed and e-mailed survey instruments with an Internet-based response mechanism, also is an approach to help reduce the problem of coverage error in the administration of surveys [12]. In comparison, Dillman et al. suggested that mixed-mode distribution of questionnaires may increase response rates with respondents choosing their preferred method of response [16]. Similarly, it was reported that 41% of electronic questionnaire respondents would not complete a telephone interview on the survey, confirming the potential of mixed-mode distribution to reduce non-response bias [17]. Moreover, a response rate of 72% was reported when a mixed-mode method was used, noticing that this technique "improved representativeness of the sample without biasing other results" [18].

Asch *et al.* reported that questionnaires distributed to medical professionals yield low response rates (mean response rate among medical doctors was 54% and it is $\approx 60\%$ among mail surveys published in medical journals) [19]. This is confirmed in a systematic review by Cook *et al.* in 2009 (median response rate of 60%) on healthcare professionals' response rates which was significantly lower than the estimate for the prior 10 year period. Authors also highlight the importance of non-response analysis and indicated that sending reminders and conducting studies on less than 1000 population will increase the response rate [20].

The research question for this study is to determine "what is considered to be a 'reasonable' response rate to merit publication of a questionnaire-based study in the contemporary dental literature". Hence, we conducted a systematic review to investigate what is a reasonable response rate for dental questionnaire-based studies in recent literature and to assess the factors that affect the response rates.

2. Materials and methods

The Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guidelines were used for all procedures and reporting [21].

2.1. Information sources and search strategy

- i) <u>Electronic database:</u> In February 2022, an electronic search of the online MEDLINE/PubMed 2018 – present database using Ovid-dc2 interface was performed using the MeSH terms related to questionnaire and dental literature.
- ii) <u>Manual searching</u>: To ensure a manageable number of articles, the search was restricted to articles published in the January-October 2021 period inclusive. At this stage, of the 782 citations, 556 articles were excluded after screening the title/abstract because they were not questionnaire-based studies or the date of publication was not within our study frame-time. When publication dates were unclear (e.g., Winter, Spring, etc.), journal websites were reviewed. Then, the remaining 226 articles were screened by 2 independent reviewers (K.A.) and (S.O'D.K.) to determine whether eligibility criteria were met. Following this, 54 articles were excluded and a total of 172 studies were eligible for inclusion in this review, of these 39 studies did not report information regarding to the response rate, therefore these studies were excluded (Fig. 1).

2.2. Selection process and eligibility criteria

Inclusion and exclusion criteria were set prior to the selection of articles for this systematic review. Inclusion criteria included: papers published between January and October 2021 (even if available online earlier), methodology sections reporting questionnaires as tools of investigation and subsequently reported on in the results section. In addition, the questionnaire in the study must be directed to the dental schools' faculties/academics, dental health practitioners (dentists, hygienist, therapists, etc.) or dental students. Studies were excluded if: the reported questionnaire was used to collect participant baseline characteristics as part of a larger study (e.g., randomised controlled trial) or if the questionnaire was not subsequently reported on in the article.

2.3. Data collection process and quality assessment

An electronic standardized data extraction form was used to extract the data from the eligible studies. Two investigators (K.A.) and (S.O'D. K.) extracted data for all included studies. We extracted data on the article title, reference details (ISSN/Doi), study ID, country(ies), discipline, reported response rate, sample size, area of distribution (e.g., national or international), actual number of questionnaires distributed, participant population and sampling frame, method of distribution (electronic or paper questionnaire), provision of stamped-addressed envelope for return of postal questionnaires, ethical approval, questionnaire piloting, follow-up, incentives, questionnaire reproduced in published article, number of questions in questionnaire, reference to response rate in discussion section, validity testing of questionnaire and the impact factor. In addition, a third investigator (C.D.L.) resolved any inconsistencies between the two investigators about included articles and data extraction.

2.4. Statistical analysis

While this project was mainly exploratory in nature, some potential associations of interest were investigated for statistical significance. Data distribution was considered to facilitate appropriate use of parametric/ non-parametric tests. We assessed the distribution of response rate and as it was not normally distributed, we used non-parametric tests for almost all reported results in this review. A Spearman's correlation



Fig. 1. PRISMA flow diagram of the identified studies.

was performed to assess the relationship between the response rates and the following numerical variables (actual number of distributed questionnaires, number of questions in questionnaire and journals impact factor). Furthermore, Kruskal-Wallis test was used to assess the pattern of response rate across the following categorical variables (area of distribution, geographical distribution, method of distribution, piloting, and follow up). A statistically significant P value was based on a threshold of <0.05 and all analyses were using Stata/MP software (version 16).

3. Results

A total of 172 articles were eligible to be included in this review. Of these, 14 articles had more than one population (e.g., dentists and dental assistants) with different response rate distribution. Thirty-nine articles did not report information regarding to the response rate; therefore, these studies were excluded. Eventually, a total of 133 articles with 149 reported response rates were included. The characteristics of included studies are presented in Table 1 according to the response rate.

3.1. Reported response rate

Table 2 shows the distribution of response rates in the sample. Ninety-two of the reported response rates (61.7%) were greater than 70%. Thirty-seven reported response rates (24.8%) were of 40–69%, and 20 reported response rates (13.4%) were less than 40%.

3.2. Actual number of questionnaires distributed

Of the 149 reported response rates, 140 reported information about the number of questionnaires distributed to participants (93.9%). Nineteen studies reported questionnaires distributed to \leq 50 participants (13.5), 24 (17.1%) reported questionnaires distributed to participants between 51 and 100, 40 (28.5%) reported questionnaires distributed to 101–300 participants, 37 (26.4%) reported questionnaires distributed to 301–1000 participants and 20 (14.2%) reported questionnaires distributed to >1000 participants (Table 1). We also found a statistically significant moderate negative correlation between the number of questionnaires distributed and response rate (Spearman's Correlation coefficient = -0.4127; P < 0.001) suggesting that when the questionnaire was distributed to a smaller sample population, a higher response rate was achieved.

3.3. Area of distribution

A total of 55 (36.9%) studies were conducted within the same

Table 1

Study characteristics based on response rate.

Response rateVariables	<50%	50-<70%	≥70%		
Actual number of questionnaires distributed n (%)					
50 or less	0	3 (12)	16 (18.6)		
51–100	5 (17.2)	3 (12)	16 (18.6)		
101–300	4 (13.8)	6 (24)	30 (34.9)		
301-1000	11 (37.9)	8 (32)	18 (20.9)		
>1000	9 (31)	5 (20)	6 (6.9)		
Geographical area n(%)					
Africa	2 (6.25)	1 (4)	2 (2.17)		
Asia	8 (25)	15 (60)	48 (52.2)		
Europe	14 (43.8)	8 (32)	28 (30.4)		
North America	4 (12.5)	0	5 (5.43)		
South America	3 (9.38)	0	5 (5.4)		
Other*	1 (3.13)	1 (4)	4 (4.35)		
Method of distribution					
Paper	8 (25.8)	5 (25)	32 (41)		
Electronic	23 (74.2)	14 (70)	45 (<i>57.7</i>)		
Both	0	1 (5)	1 (1.28)		
Number of questions n (%)					
< 50 questions	20 (100)	10 (83.3)	57 (86.4)		
\geq 50 questions	0	2 (16.7)	9 (13.6)		
Follow up reminders n (%)					
Yes	11 (34.4)	9 (36)	18 (19.6)		
Not reported	20 (62.5)	15 (60)	60 (65.2)		
Not applicable	1 (3.13)	1 (4)	14 (15.2)		
Pilot study					
Yes	16 (50)	13 (52)	35 (38)		
Not reported	16 (50)	12 (48)	57 (62)		
Ethical Approval obtained, n (%)					
Yes	26 (81.3)	25 (100)	76 (82.6)		
No or not reported	6 (18.8)	0	16 (17.4)		

Journal of Dentistry 126 (2022) 104284

Table 2

Reported	response rate	within the sa	ample included	l in t	he review.
----------	---------------	---------------	----------------	--------	------------

Response rate (%)	Articles			
	n	%		
<10	2	1.34		
10–19	3	2.01		
20-29	7	4.69		
30–39	8	5.36		
40–49	12	8.05		
50–59	12	8.05		
60–69	13	8.72		
70–79	20	13.42		
80-89	36	24.16		
≥ 90	36	24.16		
Total	149	100		
The median response rate was 77% (min=7%, max=100%) and the mean was 70.8%				

institution as the study investigators, 81 studies (54.3%) were conducted nationally, and 13 (8.7%) were international studies. Table 3 shows the pattern of response rate according to the area of distribution. Significant differences in response rates based on the area of distribution were found, with higher response rates among studies that conducted within the institute (Kruskal-Wallis test P value = 0.0012). However, it should be noted that few international studies were included in our review (n = 13).

3.4. Geographical distribution

The 149 reported response rate were categorised into six geographical areas namely, Africa, Asia, Europe, North America, South America and other. Table 4 illustrates the pattern of response rate in regard to the geographical distribution. There were significant differences in response rate across the geographical distribution (Kruskal-Wallis test P value = 0.067). Approximately, two-third of studies in Asia had response rates \geq 70%, whereas about half of European and north American studies had response rates \geq 70%.

3.5. Method of distribution

The majority of the included studies reported electronic/internet distribution of the questionnaires (82, 55%). Forty-five studies (30%) reported paper distribution whereas two studies (1.5%) reported mixed-mode distribution. No information was reported in the remaining 20 studies (13%). Table 4 shows the pattern of response rate according to the method of distribution. There is weak evidence of differences between the method of distribution and the response rate (Kruskal-Wallis test P value = 0.0489).

Table 3Response rates according to the area of distribution.

Response rate (%)	Area of distribution Within institute (%)	National (%)	International (%)
<10	0 (0)	2 (2.47)	0 (0)
10–19	0 (0)	3 (3.70)	0 (0)
20-29	0 (0)	6 (7.41)	1 (7.69)
30–39	1 (1.82)	7 (8.64)	0 (0)
40–49	4 (7.27)	7 (8.64)	1 (7.69)
50–59	2 (3.64)	9 (11.11)	1 (7.69)
60–69	6 (10.91)	5 (6.17)	2 (15.38)
70–79	12 (21.82)	8 (9.88)	0 (0)
80-89	10 (18.18)	22 (27.16)	4 (30.77)
≥ 90	20 (36.36)	12 (14.81)	4 (30.77)
Total	55	81	13
Minimum	36.1	7	20.4
Median	84	73	84
Maximum	100	100	99.4

Table 4

Response rates according to the geographical distribution and method of distribution.

Variable	Number of studies (%)	Min. response rate (%)	Median (%)	Max. response rate (%)	$\frac{\text{Mean}}{\text{SD}} \pm$
Geographica	al distribution				
Africa	5 (3.36)	25	60.8	91	$60\pm\ 25.6$
Asia	71 (47.65)	21.7	81.5	100	$\textbf{77} \pm \textbf{19.4}$
Europe	50 (33.56)	20.4	73.5	100	65 ± 22.7
North	9 (6.04)	9.4	80	100	56 ± 38.3
America					
South	8 (5.37)	16.8	73.7	97.4	65 ± 31.6
America					
Other	6 (4.03)	7	88.9	100	71 ± 36.1
Method of distribution					
Electronic	82 (63.5)	7	72.8	100	65 ± 25.6
Paper	45 (34.8)	20.6	84.3	100	75 ± 21.6
Mixed-	2 (1.5)	52.6	68.1	83.7	68 ± 21.9
mode					

3.6. Provision of a stamped-addressed envelope for return of postal questionnaires

Only three studies out of the 45 (6.6%) that used paper distribution of the questionnaire reported provision of a stamped-addressed envelope for return.

3.7. Ethical approval

Of the 149 reported response rate, 127 (85.2%) reported ethical approval being sought, 20 studies (13.4%) did not report seeking ethical approval and two studies (1.3%) reported that ethical approval was not sought or being advised that ethical approval was not required.

3.8. Piloting

Of the 149 reported response rates, 64 (43%) had been piloted, while no information on piloting was reported in the remaining studies (85, 57%) (Table 5). There was no statistical significance differences in the response rates across these groups (Kruskal-Wallis test P value = 0.0877).

3.9. Follow-up

Of the 149 reported response rate, 38 studies (25.5%) reported follow-up was performed. Sixteen studies (10.7%) did not allow for follow-up and the remaining 95 studies (63.7%) did not report any follow-up (Table 5). Comparison of response rates across these groups was statistical significance (Kruskal-Wallis test P value = 0.0047).

3.10. Incentives

One-hundred and forty-two studies (95.3%) did not report any incentives. Five studies (3.3%) reported that no incentives had been offered. Two studies (1.3%) reported incentives had been offered.

Table 5

Response rates according to piloting and follow-up.

Variable	Median (min, max)%
Piloting or not (n,%)	
Piloting (64, 43)	79.5% (7%, 100%)
No piloting (85, 57)	74.8% (13.5%, 100%)
Follow-up or not (n,%)	
Follow-up (38, 25.5)	67.3% (7%, 100%)
No follow-up (16, 10.7)	84.9% (42.6%, 100%)
Did not refer to follow-up (95, 63.7)	77% (9.4%, 100%)

3.11. Questionnaire reproduced

Sixty-four out of the 133 included articles (48%) reproduced the original questionnaire within the published article. The remaining 69 articles (52%) did not reproduce their questionnaire.

3.12. Number of questions in the questionnaire

Table 6 shows the number of questions included in the questionnaires and Fig. 2 illustrates the association between number of questions and the response rate. It is clear from the scatter plot that there is no correlation between number of questions and response rate (Spearman's correlation coefficient: 0.1148; P = 0.2604).

3.13. Reference to response rate in discussion section

Of the 149 reported response rate, 53 studies (35.5%) commented on the response rate achieved in their discussion or conclusion section. Ninety-six studies (64.4%) did not comment on the response rate.

3.14. Validity testing of questionnaire

Fifty-nine studies (39%) reported that the questionnaire had been validated or was based on a previously-validated questionnaire. Three studies (2%) did not validate the used questionnaire and no information on validation was included in the remaining 87 studies (58%) out of the 149.

3.15. Impact factor

The range of journal impact factors was from 0.406 to 6.99. The scatter plot (Fig. 3) shows no association between impact factor and response rate (Spearman's correlation coefficient: -0.0939; P = 0.2548).

4. Discussion

This systematic review included 133 questionnaire-based articles with a total of 149 reported response rates. The median of response rate across the dental literature was 77% (mean=70.8%). In 1997, Tan and Burke reviewed 77 articles that were published between 1988 and 1992, the response rates to "questionnaires mailed to dentists" had an average of 64% [7]. Thus, based on the information included in this systematic review, the response rate of the questionnaire-based dental studies is higher than what it was reported 25 years ago.

Furthermore, this review revealed that studies conducted within the institute (e.g., university) had higher response rates than the studies that conducted nationally or internationally. Moreover, we found that about two-third of studies in Asia had response rates \geq 70%, whereas about half of the European and north American studies had response rates \geq 70%. However, we found weak evidence that the method of distribution (e.g., electronic) could affect the response rate, and comparable response rates could be achieved with any method of distribution as previously reported [12,13]. Thus, this finding does not support using of electronic questionnaire [10] or the mixed-mode method [15,16,18] to

Table 6

Number of questions used within the questionnaires.

Number of questions	Number of studies (%)	Median (min, max)%
≤ 10	13 (8.7)	68 (13.5, 100)
11-20	33 (22.1)	83 (7, 100)
21-30	27 (18.1)	75.79 (15.9, 100)
31–40	9 (6)	81 (68.3, 89.3)
>40	11 (7.3)	88 (53.4, 100)
No information	56 (37.5)	72.45 (9.4, 100)
Total	149 (<i>100</i>)	P value = 0.2604



Fig. 2. Association between number of questions and response rate.



Fig. 3. Association between impact factor and response rate.

increase the response rate.

Beside the many advantages of electronic distribution of the questionnaire-base studies (e.g., automates the transformation of raw data into electronic form), concern has been expressed in relation to potential reduced response rates associated with this method. Yun *et al.* stated that "*e-mail research raises many ethical concerns because unsolicited e-mail invades a person's private space*" [18]. Likewise, Dillman *et al.* also noted that junk email risks reducing response rates [16]. Comparable to older studies [22,23], this systematic review has a median response rate for electronic distribution (72.8%) which was lower than that of paper

distribution (84.3%).

More than two-third of the studies that reported \geq 70% response rate had distributed their questionnaire to participants of 300 or less. This indicates that a higher response rate could be achieved when the questionnaire is distributed to a smaller population. Similarly, Cook *et al.* suggested that conducting studies on a population less than 1000 participants would increase the response rate [20]. Cook *et al.* also suggested that sending a reminder would increase the response rate [20] which is in keeping with the original recommendations of Dillman *et al.* [16]. However, we found that studies that reported sending a follow-up reminder had lower response rates than the studies that did not report any information on sending reminders. Therefore, our results could not support the importance of sending reminder to increase the response rate as 63.7% of articles did not report any information on follow-up.

While piloting is deemed essential and a recognised means of increasing response rate [16], the difference in response rates did not reach a significant level between the included studies reporting piloting (43%) and those which did not report any information (57%). A wide variation was also noted in the quality of information reported within the systematic review sample of publications. Thirteen percent of articles did not report seeking ethical approval for their questionnaire, 58% did not report whether or not their questionnaire had been validated, and 56% did not report whether their questionnaire was reproducible or not (i.e., can be used again). Additionally, we found no relationship between the number of questions in questionnaire and the response rate. In addition, 96 articles (64.4%) did not comment or mention the achieved response rate in the discussion section of their study and 142 studies (95.3%) did not report offering any incentives.

Ouestionnaire-based studies and the response rate achieved are controversial. There is no evidence base for specific thresholds for response rates among these studies [6]. However, conducting good research practice by following the core principle based on Dillman's work [16] which have achieved high levels of acceptance within the scientific community and considered as fundamental to high quality surveys [6]. Briefly, Dillman et al. recommended that: (1) more work and scientific rigor goes into the planning and execution of a questionnaire survey, the more likely the results are to be valid, (2) the quality and clarity of the survey's covering letter will improve response rates, (3) the questions set should be simple, short, specific without being too specific, unambiguous and should avoid bias, (4) the questionnaire should be piloted amongst colleagues, potential users of the information and also amongst the population to be surveyed, (5) the questionnaire recipients should be randomly selected, representative of the population as a whole (to avoid selection bias and ensure that results are 'generalisable') and of sufficient size, and (6) if one mailing achieves a poor response rate, then repeat the mailing twice and try to identify how non-responders differ (demographically) from responders [16].

Systematic reviews offer advantages such as efficiency, integration of information from a diverse range of related articles, while reducing bias and increasing reliability and accuracy of recommendations because of its formalised and thorough method of investigation [24]. We conducted a comprehensive search strategy and the investigation was exploratory in nature; some potential associations of interest were investigated for statistical significance. Due to the non-normal distribution of data, non-parametric tests were selected. These tests are valuable; however, they are less sensitive than parametric tests [25]. In addition, our review was limited to the English language studies that published between January and October 2021 which could be considered as limitations of this systematic review.

5. Conclusion

The findings of this systematic review confirm that high response rate is associated with smaller sample size and conducting the study within the same institute. However, we noted considerable variation exists amongst response rates and the reporting of other information. Questionnaire-based publications can effectively contribute to dental research; thus, dental journals should consider development of a minimum set of guidelines in the reporting of questionnaire-based manuscripts.

CRediT authorship contribution statement

Khaleel Al Khalaf: Conceptualization, Electronic database search,

Screening, Data extraction, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Validation, Visualization, Writing - original draft, Writing - review & editing. Shane O'Dowling Keane: Screening, Data extraction & Validation. Cristiane da Mata: Conceptualization, Project administration, Supervision, Validation, Review & editing. Catherine T McGillycuddy: Conceptualization, Validation, Review & editing. Barbara L Chadwick: Conceptualization, Validation, Review & editing. Christopher D. Lynch: Conceptualization, Investigation, Methodology, Project administration, Supervision, Validation, Writing - review & editing.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- A.N. Oppenheim, Questionnaire Design, Interviewing and Attitude Measurement, Bloomsbury Publishing, 2000.
- [2] P.M. Boynton, T. Greenhalgh, Selecting, designing, and developing your questionnaire, BMJ 328 (7451) (2004) 1312–1315.
- [3] J. Bell, S. Waters, Ebook: Doing Your Research Project: a Guide For First-Time Researchers, Mcgraw-Hill Education, 2018 (uk).
- [4] S. Kugananthan, et al., Knowledge, experiences and attitudes of dental health professionals towards reporting child abuse in Western Australia, Aust. Dent. J. 66 (2) (2021) 194–200.
- [5] H.N. Yilmaz, E.O. Ozbilen, G. Karabiber, Assessment of the diagnostic skills of general dentists in different types of orthodontic malocclusions, Turk. J.Orthod. 34 (3) (2021) 189–199.
- [6] I.L. Chapple, Questionnaire research: an easy option? Br. Dent. J. 195 (7) (2003), 359-359.
- [7] R. Tan, F. Burke, Response rates to questionnaires mailed to dentists. A review of 77 publications, Int. Dent. J. 47 (6) (1997) 349–354.
- [8] S. Kiesler, L.S. Sproull, Response effects in the electronic survey, Public Opin. Q. 50 (3) (1986) 402–413.
- [9] R. Kraut, Telecommuting: Cautious pessimism, Bell Communications Research, Murray Hill, NJ, 1984.
- [10] L. Parker, Collecting data the e-mail way, Train. Dev. 46 (7) (1992) 52–55.
- [11] D. Bachmann, J. Elfrink, G. Vazzana, Tracking the progress of e-mail vs. snail-mail, Market. Res. 8 (2) (1996) 30.
- [12] D.R. Schaefer, D.A. Dillman, Development of a standard e-mail methodology: results of an experiment, Public Opin. Q. (1998) 378–397.
- [13] M.D. Kaplowitz, T.D. Hadlock, R. Levine, A comparison of web and mail survey response rates, Public Opin. Q. 68 (1) (2004) 94–101.
- [14] R. Olum, F. Bongomin, Social media platforms for health communication and research in the face of COVID-19 pandemic: a cross sectional survey in Uganda, MedRxiv (2020).
- [15] P.D. Converse, et al., Response rates for mixed-mode surveys using mail and e-mail/web, Am. J. Eval. 29 (1) (2008) 99–107.
- [16] D.A. Dillman, J.D. Smyth, L.M. Christian, Internet, Phone, Mail, and Mixed-Mode Surveys: The Tailored Design Method, John Wiley & Sons, 2014.
- [17] W.J. Swoboda, et al., Internet surveys by direct mailing: an innovative way of collecting data, Soc. Sci. Comput. Rev. 15 (3) (1997) 242–255.
- [18] G.W. Yun, C.W. Trumbo, Comparative response to a survey executed by post, email, & web form, J. Comput.-Mediat. Commun. 6 (1) (2000) JCMC613.
- [19] D.A. Asch, M.K. Jedrziewski, N.A. Christakis, Response rates to mail surveys published in medical journals, J. Clin. Epidemiol. 50 (10) (1997) 1129–1136.
- [20] J.V. Cook, H.O. Dickinson, M.P. Eccles, Response rates in postal surveys of healthcare professionals between 1996 and 2005: an observational study, BMC Health Serv. Res. 9 (1) (2009) 1–8.
- [21] D. Moher, et al., Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement, Syst. Rev. 4 (1) (2015) 1–9.
- [22] S.E. Anderson, B.M. Gansneder, Using electronic mail surveys and computermonitored data for studying computer-mediated communication systems, Soc. Sci. Comput. Rev. 13 (1) (1995) 33–46.
- [23] M.J. Kittleson, Response rate via the postal service and e-mail, Health Values 18 (2) (1995) 27–29.
- [24] A. Petrie, J. Bulman, J. Osborn, Further statistics in dentistry Part 8: systematic reviews and meta-analyses, Br. Dent. J. 194 (2) (2003) 73–78.
- [25] R.J. Dailey, J.S. Kim, Biostatistics For Oral Healthcare, John Wiley & Sons, 2008.