

Title	Health-related quality of life in women after colposcopy: results from a longitudinal patient survey
Authors	O'Connor, Mairead; Ó Céilleachair, A.; O'Brien, K.; O'Leary, J.; Martin, C.; D'Arcy, T.; Flannelly, G.; McRae, J.; Prendiville, W.; Ruttle, C.; White, C.; Pilkington, L.; Sharp, L.
Publication date	2021-04-01
Original Citation	O'Connor, M., Ó Céilleachair, A., O'Brien, K., O'Leary, J., Martin, C., D'Arcy, T., Flannelly, G., McRae, J., Prendiville, W., Ruttle, C., White, C., Pilkington, L. and Sharp, L. (2021) 'Health-related quality of life in women after colposcopy: results from a longitudinal patient survey', Quality of Life Research, 30(9), pp.2509-2520. https://doi.org/10.1007/s11136-021-02831-3
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1007/s11136-021-02831-3
Rights	© 2021, the Authors, under exclusive licence to Springer Nature Switzerland AG, part of Springer Nature. This is a post-peer-review, pre-copyedit version of an article published in Quality of Life Research. The final authenticated version is available online at: https://doi.org/10.1007/s11136-021-02831-3
Download date	2026-08-30 07:55:45
Item downloaded from	https://hdl.handle.net/10468/11244

Health-related quality of life in women after colposcopy: Results from a longitudinal patient survey

O'Connor M^{1*}¶, Ó Céilleachair A^{2¶}, O'Brien K², O'Leary J³, Martin C³, D'Arcy T³, Flannelly G⁴, McRae J², Prendiville W³, Ruttle C³, White C³, Pilkington L³, Sharp L⁵

¹ *School of Public Health, University College Cork, Ireland*

² *National Cancer Registry Ireland, Cork, Ireland*

³ *Coombe Women and Infants University Hospital, Dublin, Ireland*

⁴ *National Maternity Hospital, Dublin, Ireland*

⁵ *Population Health Sciences Institute, Newcastle University Centre for Cancer, Newcastle University, England*

Running title: Health-related quality of life in women after colposcopy

***Corresponding author:** Dr. M O'Connor, School of Public Health, 4th Floor, Western Gateway Building, Western Road, University College Cork, Cork, Ireland. email: maireadoconnor@ucc.ie

¶Joint first co-authors

Word count: 3,737

Abstract

Background: Information concerning the health-related quality of life (HRQoL) consequences of colposcopy is limited, particularly over time. In a longitudinal study, we investigated women's HRQoL at 4, 8 and 12 months post-colposcopy and the factors associated with this.

Methods: Women attending colposcopy at two large hospitals affiliated with the national screening programme in Ireland were invited to complete questionnaires at 4, 8 and 12 months post-colposcopy. HRQoL was measured using the EQ-5D-3L and compared across a range of socio-demographic, clinical and attitudinal variables. A mixed-effects logistic multivariable model was employed to investigate associations between these variables and low HRQoL.

Results: Of 584 women initially recruited, 429, 343 and 303 completed questionnaires at 4, 8 and 12 months, respectively. The mean overall HRQoL score for the sample across all time points was 0.90 (sd=0.16). Approximately 18% of women experienced low HRQoL at each of the three time points. In multivariable testing, over the entire 12-month follow-up period, non-Irish nationals (OR=8.99, 95% CI 2.35-34.43) and women with high grade referral cytology (OR=2.78, 95% CI 1.08-7.13) were at higher odds of low HRQoL. Women who were past (OR=0.20, 95% CI 0.07-0.58) or never (OR=0.42, 95% CI 0.16-1.12) smokers were at lower odds of low HRQoL than current smokers. As women's satisfaction with their healthcare increased their odds of experiencing low HRQoL fell (OR per unit increase=0.51, 95% CI 0.34-0.75).

Conclusions: Women's HRQoL did not change over the 12 months post-colposcopy, but some subgroups of women were at higher risk of experiencing low HRQoL. These subgroups may benefit from additional support.

Keywords: Health-related quality of life, colposcopy, cervical screening, longitudinal study design

Introduction

Cervical cancer screening is effective in reducing incidence and mortality from cervical cancer among target populations [1]. To ensure effectiveness of screening, women who receive positive cytology test results require follow-up investigation, usually a hospital-based colposcopy. Colposcopy is a very common procedure with 171,874 women undergoing colposcopy for the first time in England in 2018[2].

Colposcopy and its related interventions (e.g. biopsies of any areas of the cervical transformation zone that appear abnormal, or treatment to remove abnormal cervical cells) are associated with distress and anxiety in women before and during the procedure(s) [3]. Concerns around cancer, sex and fertility can also persist after colposcopy and related interventions [4-8]. While an extensive body of literature exists on these dimensions of quality of life, there is considerably less evidence on validated health-related quality-of-life (HRQoL) measures, particularly those that are typically used to impute quality-adjusted life years (QALYS) in health economic evaluations, such as the widely-used EuroQol EQ-5D instrument [9]. Information is also limited on factors associated with HRQoL in women following colposcopy and related procedures and how colposcopy related HRQoL varies over time [10]. What evidence does exist, particularly from longitudinal studies, is often based on small samples [6, 11, 12] or includes only women with low-grade abnormal cytology [13]. While it could be hypothesised, given the relationship between anxiety and undergoing colposcopy, that the procedure would negatively impact HRQoL, it is difficult to draw such conclusions from the existing literature with some studies finding little to no relationship between receipt of colposcopy and HRQoL [6, 11], others a modest positive relationship [13] and still others a negative relationship, albeit short-term [12].

The discovery of the causal relationship between infection with certain high-risk strains of human papillomavirus (HPV) and cervical neoplasia [14] has transformed cervical cancer prevention. Many screening programmes have introduced (or intend to introduce) HPV testing as their primary screening tool instead of cytology-based testing and many countries offer vaccination against certain HPV types to young adolescents [15]. To determine how best to implement cervical cancer prevention in this changing context, the cost-effectiveness of these newer strategies are being assessed [16-18]. Consequently, new estimates of HRQoL for use in the calculation of QALYs would appear timely. Although primary HPV screening has been adopted by programmes, colposcopy is likely to remain a central pillar of follow-up of women with abnormal screening tests into the future.

The current study aimed to: (1) assess how women's HRQoL changes over the 12 months following colposcopy; (2) provide mean utility scores for different clinical and socio-demographic subgroups for use in (future) cost-effectiveness modelling; and (3) investigate the factors associated with low HRQoL in the year after colposcopy, as this may identify subgroups of women who could benefit from additional interventions or supports.

98 **Methods**

99 *Setting*

100 The study setting was Ireland, which has a mixed public–private healthcare system. The
101 national cervical screening programme (CervicalCheck) was introduced in 2008, offering free
102 cervical cytology tests and follow-up, if needed, to women aged 25–60 years. Women with
103 two or more low-grade abnormal cervical cytology test results, or one high-grade result, were
104 referred for colposcopy in a clinic affiliated with the screening programme located in one of
105 15 maternity hospitals throughout Ireland.

106 *Participants and procedures*

107 Women attending two large colposcopy clinics in Dublin, Ireland (The Coombe Women and
108 Infants University Hospital and the National Maternity Hospital) were recruited to the study
109 between September 2010–July 2011. Women were eligible if they were referred to
110 colposcopy because of an abnormal cytology test result in the context of routine screening.
111 They were eligible irrespective of the management they received at their initial clinic
112 appointment (i.e. colposcopy only, loop excision, or another form of intervention or
113 treatment) or subsequent follow-up. The exclusion criteria for the study were pregnancy at
114 recruitment or treatment for previous cervical abnormalities. A member of the research team
115 (CW) approached women at the clinics and invited them to participate in the study. For
116 reasons relating to the clinical and operational circumstances prevailing in the two
117 recruitment centres, it was not possible to consecutively approach all women attending
118 clinics. The women were given a study information sheet with more details about the
119 research. Interested women signed a consent form and returned it to the research team and
120 were mailed questionnaire packs to complete at 4, 8 and 12 months after their clinic visit.

121

122 The women received no reward or payment for being recruited to the study, however, women
123 who took part in the study and completed at least one of the questionnaires were offered the
124 chance to enter into a prize draw to win one of two €150 high street shopping vouchers.
125 Offering modest incentives such as these has been shown to be the most effective way of
126 maximizing response rates [19]. Women were not be told about this incentive until they
127 received their first (~4 month) questionnaire so that it did not affect their willingness to
128 consent to the study. Ethical approval was obtained from the ethics committees of the
129 Coombe Women and Infants University Hospital (ref:21-2006) and the National Maternity
130 Hospital, Dublin (no reference number given).

131 *Assessment of health-related quality of life*

132 HRQoL was measured using the EuroQol EQ-5D 3L [9] included as part of the psychosocial
133 questionnaire administered at 4,8 and 12 months after the women's initial colposcopy
134 appointment. The EQ-5D is a widely used instrument for measuring HRQoL, particularly for
135 health technology assessments (HTAs) and other economic evaluations. A broader
136 description of the instrument can be found in the EuroQol Research Foundation's User Guide
137 [20]. The EQ-5D consists of five dimensions: mobility; self-care; usual activities;
138 pain/discomfort; and anxiety/depression. Each of these dimensions has three response levels
139 consisting of 1 'no problems', 2 'some problems' and 3 'extreme problems'. These responses
140 can be linked to utility values ranging from 0-1 (death-full health). No Irish value set is
141 available, consequently health states were valued using value sets for the UK [21].

The distribution of the HRQoL index values was discrete and non-normal. For modelling purposes, therefore, we defined low HRQoL as a response of “3” in at least one dimension or an answer of “2” in at least two dimensions.

Assessment of potential predictors of health-related quality of life

Information on potential predictors of low HRQoL was obtained from the four-month questionnaire, which contained questions on a range of socio-demographic and lifestyle factors [4, 7]. Women’s satisfaction with life and general healthcare were assessed by means of items adapted from McCaffery et al [22]. Colposcopy clinic records were used to provide information on colposcopy referral cytology, initial colposcopic impression, initial management and histology result. Clinical data were extracted from women’s records at the four month timepoint.

Statistical Analyses

Analyses were conducted in Stata 12. Mean HRQoL scores at each follow-up time point (i.e. 4, 8 and 12 months were computed overall and for a range of key socio-demographic and clinical variables.

Because we had longitudinal data, a mixed-effects logistic multivariable regression model was used to explore associations between socio-demographic, clinical and attitudinal variables and low HRQoL. This allowed inclusion of all observations across all points [23]. Variables that were significant at the 5% level in univariate logistic analysis were included in initial multivariable analysis and Wald tests used to determine which remained significant

once other independent variables were included. The final model only included variables that remained statistically significant. Time-point was included in all models, in order to test for a temporal trend in low HRQoL.

As sensitivity analyses, we re-ran the final model with utility scores as a continuous dependent variable in a mixed-effects linear model. We also changed the specification of “low” HRQoL so that it included women in the bottom three quartiles and re-ran the final model. These alternative analyses did not change the variables that attained statistical significance, nor the direction of their relationship with “low” HRQoL (data not shown).

Our study conformed with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement [24]. A copy of the completed STROBE checklist for this study is available in supplementary file [Online Resource 1].

Results

Characteristics of participants

Of 584 women who consented to participate in the study: 429 (73%); 343 (59%) and 303 (52%) completed questionnaires at 4, 8 and 12 months post-colposcopy respectively.

Women’s socio-demographic, clinical lifestyle and attitudinal characteristics at four months are summarised in Table 1.

Temporal trends in HRQoL post-colposcopy

Mean overall HRQoL was static at 0.90 across all three time points (standard deviation (SD)=0.16 at 4 and 8 months and 0.14 at 12 months). Table 2 details the mean HRQoL scores

over time by several key clinical and socio-demographic characteristics. While broadly HRQoL appeared similar across groups, those without tertiary education had lower mean HRQoL than those who had. Those who had undergone large loop excision of the transformation zone (LLETZ) had lower HRQoL than those who had not and those with a medical card (which is available to those with limited means and provides free healthcare in the public system) had lower HRQoL than those who did not.

75 (18%) women were classified as having “low” HRQoL at four months post colposcopy. This percentage was unchanged at 8 months (n=57;18%) and 12 months (n=50;18%) post-colposcopy.

Factors that influence HRQoL over 12 months post colposcopy

The results of univariate analysis of associations between socio-demographic, clinical and attitudinal variables and low HRQoL testing are presented in Table 3. In the multivariable model (Table 4), nationality, the woman’s referral cytology, smoking status and satisfaction with healthcare were significantly related to low HRQoL. The odds of having low HRQoL in non-Irish women were almost nine times higher than for Irish women (adjusted OR=8.99, 95% CI 2.35-34.43). Odds of low HRQoL was also more than two and half times higher for women referred for colposcopy on the basis of a high-grade cytology result (OR=2.78, 95% CI 1.08-7.13). Conversely, women who were either past smokers or who never smoked were at significantly lower odds of experiencing low HRQoL compared to current smokers (past vs current smoker: OR=0.20, 95% CI 0.07-0.58; never vs current smoker: OR=0.42, 95% CI 0.16-1.12). Higher levels of satisfaction with general healthcare received were associated with a lower odds of low HRQoL; with every unit increase in satisfaction on a Likert scale of 1-10 (where 10 indicated complete satisfaction), the odds of experiencing low HRQoL were reduced by approximately one half (OR=0.51, 95% CI 0.34-0.75).

Discussion

In this longitudinal study we found no discernible change in women's HRQoL over the 12 months post colposcopy; mean HRQoL values were identical at each time point as was the proportion of women with low HRQoL. In multivariable analysis, we identified two subgroups of women at higher odds of experiencing low HRQoL: (1) non-Irish women and (2) women with high-grade abnormal referral cytology. We also found that women who were not current smokers were less likely to experience low HRQoL and that as women's satisfaction with their healthcare increased, their odds of experiencing low HRQoL decreased.

Temporal trends in HRQoL post-colposcopy

Over the course of the three follow-up time points, mean HRQoL remained stable and high at 0.90. It is likely that the high index score reflects the relatively young age distribution of our study population; EQ-5D Index scores have been shown to decrease with age and younger respondents report better health status [25]. The pattern of little change over time is in keeping with most of the (limited) literature in this area. Korfage et al [12] found that both EQ-5D and EQ-VAS (a visual analogue scale running from 1-100) scores were similar at 1, 3 and 6 months post-colposcopy. Similarly, while Whynes et al [13] found some differences across time, none of these were greater than the minimum important difference for the EQ-VAS [26]. Heinonen et al [11] found that HRQoL was worse at 12 months than at baseline, but this may, in part, be because they used a more sensitive instrument to assess outcomes [27-29]. The issue of (lack of) sensitivity with respect to the EQ-5D-3L, has been documented before [28, 29]; this may also be an issue in the context of investigation and treatment of abnormal cervical cytology [30].

233 *Factors that influence HRQoL over 12 months post colposcopy*

234 Women whose referral cytology indicated moderate or severe dyskaryosis (i.e. “high-grade”)
235 were more likely to be report responses on the EQ-5D consistent with low HRQoL than
236 women referred to colposcopy with (multiple) low-grade results. This finding appears new to
237 the literature. Neither Heinonen et al [11] nor Korfage et al [12] found evidence of a
238 relationship between HRQoL and women’s original cytological abnormality. However, in
239 both of these studies the numbers of women in the groups with different referral cytology
240 were substantially smaller than those in our study. In a study of women with low-grade
241 referral cytology, Whynes et al [13] found a relationship between colposcopy plus LLETZ
242 management (compared to colposcopy only and colposcopy plus punch biopsies) and lower
243 HRQoL at some follow-up time points. This association suggests that it is possible our
244 finding reflects women with high-grade cytology being more likely, than those with low-
245 grade cytology, to have underlying cervical intraepithelial neoplasia (CIN) grade 2/3 and
246 undergo LLETZ. Women with high-grade cytology are also more likely to have undergone
247 additional follow-up procedures. We did not have information on any interventions or
248 procedures after the initial colposcopy appointment, and this is a limitation of the study.

249 We and others have shown that colposcopy itself, and attending for follow-up, can have an
250 adverse psychological impact on women with high grade findings, causing significant
251 anxiety, distress and worries which may be long lasting [31-33]. It is possible that these
252 adverse psychological effects translate into decrements in HRQoL. For example, Whynes et
253 al. [34] found the EQ-5D to be responsive to differing degrees of HADS-assessed anxiety and
254 depression among a large sample (n=3119) of UK women with low grade cervical
255 abnormalities detected through screening. In that study, a higher likelihood of HADS-
256 identified anxiety and/or depression was associated with significantly lower EQ-5D index.

This suggests that our observed association between high grade dyskaryosis and low HRQoL may reflect more worries, distress or anxiety in that group.

In a previous study analysing this population, non-Irish women were found to be at increased risk of anxiety after colposcopy [4] and our analysis here finds the same pattern with HRQoL (i.e. non-Irish women were significantly more likely to be considered to have low HRQoL in the 12 months after colposcopy than Irish women). To the best of our knowledge, our study is the first to show this relationship with HRQoL after colposcopy. Disparities in cervical screening participation among immigrants compared with general population women (e.g. in Finland and England) have been well documented [35-37], but less is known about the impacts of screening participation in migrants compared to native populations. The previous cervical screening history of the women included in our study (both Irish and non-Irish) was not available. However, it seems unlikely that lack of previous access to screening among non-Irish women would explain our findings, as all but seven of the thirty-nine non-Irish women were from another OECD country, and would likely have had some access to smear tests. Referral cytology was not associated with nationality ($p=0.581$), and the model was adjusted for this, so this cannot explain the finding either. The number of non-Irish women in our study population was broadly in line with the number expected based on the nationality of people resident in Ireland [38]. Historically, the largest group of non-Irish nationals were UK citizens, but more diverse migration patterns mean that increasingly, non-Irish women may not speak good English. This may impede them in navigating the healthcare system and limit their understanding of information about their colposcopy. The Irish cervical screening programme has produced information leaflets in 11 languages, but these do not include information about colposcopy and follow-up, nor is information in a suitable language likely to be the only thing needed to enable a migrant to understand and navigate, and feel confident within, an unfamiliar healthcare system. More culturally appropriate screening and follow-up

literature for non-Irish women is needed for women who have limited English. In addition, more support, and assistance with patient navigation, for non-Irish women, could help reduce the risk of low HRQoL in such women following colposcopy.

Satisfaction with healthcare also determined whether women reported EQ-5D responses considered to indicate low HRQoL in the 12 months post colposcopy. While evidence on the relationship between satisfaction with healthcare and HRQoL is comparatively sparse, a review by Bamm et al. [39] and colleagues of people living with chronic conditions found a generally positive relationship between HRQoL and satisfaction with care, consistent with our findings.

In terms of our final predictor of HRQoL, the observed association between current smoking and low HRQoL measured by the EQ-5D is consistent with relationships in the general population [40-42], though it is possible that this finding reflects uncontrolled confounding by comorbidities; smokers have higher prevalence of comorbidities than non-smokers [43]. However, given the young age profile of participants (approximately 70% of participants were aged forty or younger) and the exclusion of women managed for previous cervical abnormalities, it is likely that the number of women with significant comorbidities would be small. There was a greater proportion of current smokers among women with high-grade disease compared to those with low-grade disease in our study (44% vs 29%) and smoking status and referral cytology result were significantly associated ($p=0.018$). Previous studies have observed an association between being a current smoker and low HRQoL [41,42]. However, in our model the effect of smoking was adjusted for referral cytology, so this cannot account for the observed association with smoking. The likelihood of regression of cervical precancerous lesions is significantly lower among women who smoke than in never-smokers [44] and women who smoke are more likely to require follow-up visits to

colposcopy clinics and require further treatments to remove cervical abnormalities compared to non-smokers [45]. This may explain our findings.

HRQoL, colposcopy and the implications for cost-effectiveness modelling

Many healthcare systems are re-evaluating the cost-effectiveness of cervical cancer screening protocols. Accurate estimates of HRQoL are of great importance to this process. The numbers of women involved in screening and follow-up are large, meaning that it is arguably more important to have robust HRQoL data for these health states than for states for which HRQoL might be lower, but which affect fewer women (e.g. cancer). To the best of our knowledge, our study represents one of the very few longitudinal studies on HRQoL of follow-up and which provides QALY weights for use in economic analyses. We have reported weights both by treatment received and also by histological finding. Ours is one of the few studies to do both. With respect to treatment received results are largely in keeping with the available literature. Whynes et al [13], for example, found mean QALY weights for women who had received a colposcopy (0.906), a colposcopy with biopsy (0.912) and LLETZ (0.919). These values are close to our own (0.92-0.89, 0.91 and 0.86-0.89 respectively). Our QALY weights by histology are also similar to those found in other studies [12, 46]. While there were not a sufficient number of women in our sample who had developed a cancer to report reliable weights for cervical cancer, we feel that these weights will be of use to cost-effectiveness modellers.

329 *Strengths and limitations*

330 Our study is one of only a handful to examine the predictors of low HRQoL in the year after
331 colposcopy employing a longitudinal study design. With the exception of one UK trial [47],
332 which only included women with low-grade cytology, our study is also the largest to have
333 investigated HRQoL after-effects of colposcopy and related interventions. As a generic
334 questionnaire, the comparative lack of sensitivity of the EQ-5D-3L (and potential
335 ceiling/floor effects) [48] might mean some factors that concern women following
336 colposcopy, such as worries about fertility, sex and cancer [4, 49, 50], are not adequately
337 captured by the instrument. In terms of the explanatory variables considered in the study,
338 women's clinical data (e.g. initial management received) was extracted from colposcopy
339 clinic records at 4 months and we lacked information on any procedures that happened
340 between then and 12 months. The other variables explored for associations with HRQoL also
341 related to the 4-month time point; if these changed during follow-up the results may be
342 impacted by some non-differential misclassification. Related to this, although life events have
343 been shown to have a modest impact on HRQoL [51], we did not collect information on these
344 or some other variables that influence HRQoL (such as social support) [52], meaning there is
345 a possibility of some uncontrolled confounding.

346 Women were recruited from only two colposcopy clinics, both were part of the
347 CervicalCheck screening programme. However, the hospitals in which they were located
348 serve large numbers of patients from diverse backgrounds. For logistical reasons related to
349 running a research study within busy hospital clinics, we were not able to systematically
350 approach every woman in every clinic. It is possible that some particular groups of women
351 were not approached (e.g. those visibly upset). While the way the women who took part in
352 the study were recruited may affect overall estimates of HRQoL (i.e. we may have
353 overestimated this), internal comparisons within the study population will not necessarily be

354 impacted. Women who completed and returned at least one of the three questionnaires were
355 offered the chance to enter into a prize draw to win one of two high street shopping vouchers.
356 While modest incentives such as these are effective in improving questionnaire response rates
357 [19], we cannot entirely preclude the possibility that they influenced (directly or indirectly)
358 HRQoL as measured in the study participants. However, the authors are unaware of any
359 empirical evidence to support this; indeed most studies show that incentives have no effect on
360 quality of response, sample composition or response distributions [53].

362 **Conclusion**

363 While mean HRQoL the proportion of the sample considered to have low HRQoL, remained
364 static over 12 months following colposcopy, some subgroups of women were at greater risk
365 of low HRQoL. In particular non-Irish women and women whose referral cytology was high-
366 grade had a significantly greater chance of low HRQoL. As well as providing information to
367 inform cost-effectiveness analyses, our findings may aid policymakers, service providers and
368 patient advocacy groups to better target supports and advice for women who have received a
369 colposcopy thereby minimising the risk of them experiencing low health-related quality of
370 life.

Declarations

Acknowledgements

This study was undertaken as part of the CERVIVA research consortium (www.cerviva.ie).

We thank the women who completed questionnaires. We are grateful to the clinicians and nurses and staff at the two colposcopy centres for facilitating this research.

Author contributions

AÓC analysed the data and wrote the paper. MOC conducted the survey, analysed the data and wrote the paper. KOB analysed the data and wrote the paper. JOL, CM, TD and GF conceived and designed the experiments and wrote the paper. JM conducted the survey and wrote the paper. WP and CR conceived and designed the experiments and wrote the paper. CW conducted the survey and wrote the paper. LP conceived and designed the experiments and wrote the paper. LS conceived and designed the experiments, analysed the data and wrote the paper. All authors read and approved the manuscript

Funding

The data collection was funded by the Health Research Board, Ireland (HS-05-09). MOC (ICE 2011/2 and ICE 2015-1037, AÓC (HRA-HSR/2012/30) and CW (HRA-HSR/2012/30)) were supported by grants from the Health Research Board.

Compliance with ethical standards

Conflict of interest

The authors have no conflicts of interest relevant to this manuscript.

Ethical approval

Ethical approval was obtained from the ethics committees of the Coombe Women and Infants University Hospital (ref:21-2006; approval granted 27 April 2010) and the National Maternity Hospital, Dublin (no ref. number given; approval granted 26th October 2010).

Informed consent

Written informed consent was obtained from women interested in partaking in the study.

References

1. IARC (2005). IARC Handbooks of Cancer Prevention Volume 10 Cervix Cancer Screening. <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Handbooks-Of-Cancer-Prevention/Cervix-Cancer-Screening-2005>. Accessed 25 August 2020.
2. PHE Screening (2019). Cervical Screening Programme England, 2018-19 – Report. <https://digital.nhs.uk/data-and-information/publications/statistical/cervical-screening-annual/england--2018-19>. Accessed 13 August 2020.
3. Galaal K, Bryant A, Deane KH, Al-Khaduri M, Lopes AD., (2011). Interventions for reducing anxiety in women undergoing colposcopy. *Cochrane Database Syst Rev Online* 12, CD006013.
4. O'Connor, M., O'Leary, E., Waller, J., Gallagher, P., D'arcy, T., Flannelly, G., et al. (2016). Trends in, and predictors of, anxiety and specific worries following colposcopy: a 12-month longitudinal study. *Psycho-oncology*, 25(5), 597–604. <https://doi.org/10.1002/pon.3980>
5. Sparić, R., Papoutsis, D., Kadija, S., Stefanović, R., Antonakou, A., Nejković, L., & Kesić, V.(2019). Psychosexual outcomes in women of reproductive age at more than two-years from excisional cervical treatment - a cross-sectional study. *Journal of psychosomatic obstetrics and gynaecology*, 40(2), 128–137.
6. Hellsten, C., Sjöström, K., & Lindqvist, P. G. (2009). A longitudinal 2-year follow-up of quality of life in women referred for colposcopy after an abnormal cervical smear. *European journal of obstetrics, gynecology, and reproductive biology*, 147(2), 221–225. <https://doi.org/10.1016/j.ejogrb.2009.09.002>.
7. O'Connor, M., O'Brien, K., Waller, J., Gallagher, P., D'Arcy, T., Flannelly, G., et al. (2017). Physical after-effects of colposcopy and related procedures, and their inter-relationship with psychological distress: a longitudinal survey. *BJOG : an international journal of obstetrics and gynaecology*, 124(9), 1402–1410. <https://doi.org/10.1111/1471-0528.14671>.
8. Swancutt, D. R., Greenfield, S. M., Luesley, D. M., & Wilson, S. (2011). Women's experience of colposcopy: a qualitative investigation. *BMC women's health*, 11, 11. <https://doi.org/10.1186/1472-6874-11-11>.
9. Rabin, R., & de Charro, F. (2001). EQ-5D: a measure of health status from the EuroQol Group. *Annals of medicine*, 33(5), 337–343. <https://doi.org/10.3109/07853890109002087>.
10. Cullen, J., Schwartz, M. D., Lawrence, W. F., Selby, J. V., & Mandelblatt, J. S. (2004). Short-term impact of cancer prevention and screening activities on quality of life. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, 22(5), 943–952. <https://doi.org/10.1200/JCO.2004.05.191>.
11. Heinonen, A., Tapper, A. M., Leminen, A., Sintonen, H., & Roine, R. P. (2013). Health-related quality of life and perception of anxiety in women with abnormal cervical cytology referred for colposcopy: an observational study. *European journal of obstetrics, gynecology, and reproductive biology*, 169(2), 387–391. <https://doi.org/10.1016/j.ejogrb.2013.03.033>.
12. Korfage, I. J., Essink-Bot, M. L., Westenberg, S. M., Helmerhorst, T., Habbema, J. D., & van Ballegooijen, M. (2014). How distressing is referral to colposcopy in cervical cancer screening?: a prospective quality of life study. *Gynecologic oncology*, 132(1), 142–148. <https://doi.org/10.1016/j.ygyno.2013.11.001>.
13. Whynes, D. K., Sharp, L., Cotton, S. C., Cruickshank, M., Gray, N. M., Little, J., & Waugh, N. (2013). Impact of alternative management policies on health-related quality of life in women with low-grade abnormal cervical cytology referred for colposcopy. *European journal of obstetrics, gynecology, and reproductive biology*, 169(1), 88–92. <https://doi.org/10.1016/j.ejogrb.2013.02.006>.

14. Bosch, F. X., Burchell, A. N., Schiffman, M., Giuliano, A. R., de Sanjose, S., Bruni, L., Tortolero-Luna, G., Kjaer, S. K., & Muñoz, N. (2008). Epidemiology and natural history of human papillomavirus infections and type-specific implications in cervical neoplasia. *Vaccine*, 26 Suppl 10, K1–K16. <https://doi.org/10.1016/j.vaccine.2008.05.064>
15. Nguyen-Huu, N. H., Thilly, N., Derrough, T., Sdona, E., Claudot, F., Pulcini, C., Agrinier, N., & HPV Policy working group (2020). Human papillomavirus vaccination coverage, policies, and practical implementation across Europe. *Vaccine*, 38(6), 1315–1331.
16. Lew, J. B., Simms, K. T., Smith, M. A., Hall, M., Kang, Y. J., Xu, X. M., Caruana, M., Velentzis, L. S., Bessell, T., Saville, M., Hammond, I., & Canfell, K. (2017). Primary HPV testing versus cytology-based cervical screening in women in Australia vaccinated for HPV and unvaccinated: effectiveness and economic assessment for the National Cervical Screening Program. *The Lancet. Public health*, 2(2), e96–e107. [https://doi.org/10.1016/S2468-2667\(17\)30007-5](https://doi.org/10.1016/S2468-2667(17)30007-5).
17. Barré, S., Massetti, M., Leleu, H., & De Bels, F. (2017). Organised screening for cervical cancer in France: a cost-effectiveness assessment. *BMJ open*, 7(10), e014626. <https://doi.org/10.1136/bmjopen-2016-014626>.
18. Termrungruanglert, W., Khemapech, N., Tantitamit, T., & Havanond, P. (2019). Cost effectiveness analysis of HPV primary screening and dual stain cytology triage compared with cervical cytology. *Journal of gynecologic oncology*, 30(2), e17. <https://doi.org/10.3802/jgo.2019.30.e17>.
19. Edwards, P. J., Roberts, I., Clarke, M. J., Diguiseppi, C., Wentz, R., Kwan, I., Cooper, R., Felix, L. M., & Prata, S. (2009). Methods to increase response to postal and electronic questionnaires. *The Cochrane database of systematic reviews*, 3, MR000008. <https://doi.org/10.1002/14651858.MR000008.pub4>
20. EuroQol Research Foundation. (2018). EQ-5D-3L User Guide. EuroQol Research Foundation. https://euroqol.org/wp-content/uploads/2019/10/EQ-5D-3L-User-Guide_version-6.0.pdf. Accessed 13 August 2020.
21. Szende A, Oppe M, Devlin NJ, EuroQol Group, editors. EQ-5D value sets: inventory, comparative review, and user guide. Dordrecht: Springer; 2007.
22. McCaffery, K. J., Irwig, L., Turner, R., Chan, S. F., Macaskill, P., Lewicka, M., Clarke, J., Weisberg, E., & Barratt, A. (2010). Psychosocial outcomes of three triage methods for the management of borderline abnormal cervical smears: an open randomised trial. *BMJ (Clinical research ed.)*, 340, b4491. <https://doi.org/10.1136/bmj.b4491>.
23. Brown H, Prescott R., Applied Mixed Models in Medicine, in Statistics in Practice. 2006, John Wiley & Sons: West Sussex UK
24. von Elm, E., Altman, D. G., Egger, M., Pocock, S. J., Gøtzsche, P. C., Vandenbroucke, J. P., & STROBE Initiative (2007). The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet (London, England)*, 370(9596), 1453–1457. [https://doi.org/10.1016/S0140-6736\(07\)61602-X](https://doi.org/10.1016/S0140-6736(07)61602-X).
25. Feng, Y., Devlin, N., & Herdman, M. (2015). Assessing the health of the general population in England: how do the three- and five-level versions of EQ-5D compare?. *Health and quality of life outcomes*, 13, 171. <https://doi.org/10.1186/s12955-015-0356-8>.
26. Pickard, A. S., Neary, M. P., & Cella, D. (2007). Estimation of minimally important differences in EQ-5D utility and VAS scores in cancer. *Health and quality of life outcomes*, 5, 70. <https://doi.org/10.1186/1477-7525-5-70>.

27. Heiskanen, J., Tolppanen, A. M., Roine, R. P., Hartikainen, J., Hippeläinen, M., Miettinen, H., & Martikainen, J. (2016). Comparison of EQ-5D and 15D instruments for assessing the health-related quality of life in cardiac surgery patients. *European heart journal. Quality of care & clinical outcomes*, 2(3), 193–200. <https://doi.org/10.1093/ehjqcco/qcw002>.
28. Vartiainen, P., Mäntyselkä, P., Heiskanen, T., Hagelberg, N., Mustola, S., Forssell, H., Kautiainen, H., & Kalso, E. (2017). Validation of EQ-5D and 15D in the assessment of health-related quality of life in chronic pain. *Pain*, 158(8), 1577–1585. <https://doi.org/10.1097/j.pain.0000000000000954>.
29. Sintonen H. (2001). The 15D instrument of health-related quality of life: properties and applications. *Annals of medicine*, 33(5), 328–336. <https://doi.org/10.3109/07853890109002086>.
30. Payakachat, N., Ali, M. M., & Tilford, J. M. (2015). Can The EQ-5D Detect Meaningful Change? A Systematic Review. *Pharmacoeconomics*, 33(11), 1137–1154. <https://doi.org/10.1007/s40273-015-0295-6>.
31. O'Connor, M., Waller, J., Gallagher, P., Martin, C. M., O'Leary, J. J., D'Arcy, T., Prendiville, W., Flannelly, G., Sharp, L., & Irish Screening Research Consortium (CERVIVA) (2015). Understanding Women's Differing Experiences of Distress after Colposcopy: A Qualitative Interview Study. *Women's health issues : official publication of the Jacobs Institute of Women's Health*, 25(5), 528–534. <https://doi.org/10.1016/j.whi.2015.05.009>.
32. O'Connor, M., Gallagher, P., Waller, J., Martin, C. M., O'Leary, J. J., Sharp, L., & Irish Cervical Screening Research Consortium (CERVIVA) (2016). Adverse psychological outcomes following colposcopy and related procedures: a systematic review. *BJOG : an international journal of obstetrics and gynaecology*, 123(1), 24–38. <https://doi.org/10.1111/1471-0528.13462>.
33. Sharp, L., Cotton, S., Cruickshank, M., Gray, N., Smart, L., Whynes, D., Little, J., & TOMBOLA Group (2016). Impact of post-colposcopy management on women's long-term worries: results from the UK population-based TOMBOLA trial. *The journal of family planning and reproductive health care*, 42(1), 43–51. <https://doi.org/10.1136/jfprhc-2015-101170>.
34. Whynes, D. K., & TOMBOLA Group (2009). Responsiveness of the EQ-5D to HADS-identified anxiety and depression. *Journal of evaluation in clinical practice*, 15(5), 820–825. <https://doi.org/10.1111/j.1365-2753.2008.01102.x>.
35. Idehen, E. E., Koponen, P., Härkänen, T., Kangasniemi, M., Pietilä, A. M., & Korhonen, T. (2018). Disparities in cervical screening participation: a comparison of Russian, Somali and Kurdish immigrants with the general Finnish population. *International journal for equity in health*, 17(1), 56. <https://doi.org/10.1186/s12939-018-0768-2>.
36. Moser, K., Patnick, J., & Beral, V. (2009). Inequalities in reported use of breast and cervical screening in Great Britain: analysis of cross sectional survey data. *BMJ (Clinical research ed.)*, 338, b2025. <https://doi.org/10.1136/bmj.b2025>.
37. Vahabi, M., Lofters, A. K., Kopp, A., & Glazier, R. H. (2021). Correlates of non-adherence to breast, cervical, and colorectal cancer screening among screen-eligible women: a population-based cohort study in Ontario, Canada. *Cancer causes & control : CCC*, 32(2), 147–155. <https://doi.org/10.1007/s10552-020-01369-y>.
38. Central Statistics Office. Census 2016 - Non-Irish nationals in Ireland. Cork; 2018. <https://www.cso.ie/en/releasesandpublications/ep/p-cpnin/cpnin/>. Accessed 01 September 2020
39. Bamm, E. L., Rosenbaum, P., & Wilkins, S. (2013). Is Health Related Quality Of Life of people living with chronic conditions related to patient satisfaction with care?. *Disability and rehabilitation*, 35(9), 766–774. <https://doi.org/10.3109/09638288.2012.707746>.

40. Vogl, M., Wenig, C. M., Leidl, R., & Pokhrel, S. (2012). Smoking and health-related quality of life in English general population: implications for economic evaluations. *BMC public health*, 12, 203. <https://doi.org/10.1186/1471-2458-12-203>.
41. Maheswaran, H., Petrou, S., Rees, K., & Stranges, S. (2013). Estimating EQ-5D utility values for major health behavioural risk factors in England. *Journal of epidemiology and community health*, 67(2), 172–180. <https://doi.org/10.1136/jech-2012-201019>.
42. Orji, C. C., Ghosh, S., Nwaobia, O. I., Ibrahim, K. R., Ibiloye, E. A., & Brown, C. M. (2021). Health Behaviors and Health-Related Quality of Life Among U.S. Adults Aged 18-64 Years. *American journal of preventive medicine*, S0749-3797(20)30516-X. Advance online publication. <https://doi.org/10.1016/j.amepre.2020.10.020>.
43. U.S. Department of Health and Human Services. (2014). The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General. Atlanta. https://www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf_NBK179276.pdf. Accessed 20 August 2020.
44. Matsumoto, K., Oki, A., Furuta, R., Maeda, H., Yasugi, T., Takatsuka, N., Hirai, Y., Mitsuhashi, A., Fujii, T., Iwasaka, T., Yaegashi, N., Watanabe, Y., Nagai, Y., Kitagawa, T., Yoshikawa, H., & Japan HPV And Cervical Cancer (JHACC) Study Group (2010). Tobacco smoking and regression of low-grade cervical abnormalities. *Cancer science*, 101(9), 2065–2073. <https://doi.org/10.1111/j.1349-7006.2010.01642.x>.
45. Lamb, J., Dawson, S. I., Gagan, M. J., & Peddie, D. (2013). Cigarette smoking and the frequency of colposcopy visits, treatments and re-referral. *Nursing praxis in New Zealand inc*, 29(1), 24–33.
46. Marcellusi, A., Capone, A., Favato, G., Mennini, F. S., Baio, G., Haeussler, K., Bononi, M., & HPV Italian Collaborative Study Group (2015). Health utilities lost and risk factors associated with HPV-induced diseases in men and women: the HPV Italian collaborative study group. *Clinical therapeutics*, 37(1), 156–167.e4. <https://doi.org/10.1016/j.clinthera.2014.11.002>
47. Cotton, S. C., Sharp, L., Little, J., Duncan, I., Alexander, L., Cruickshank, M. E., Gray, N. M., Jenkins, D., Philips, Z., Robertson, A., Seth, R., & TOMBOLA group (2006). Trial of management of borderline and other low-grade abnormal smears (TOMBOLA): *Trial design. Contemporary clinical trials*, 27(5), 449–471. <https://doi.org/10.1016/j.cct.2006.04.001>.
48. Brooks RG. The EuroQol Group after 25 years. Dordrecht; New York: Springer; 2013. <http://public.eblib.com/choice/publicfullrecord.aspx?p=1030747>. Accessed 20 August 2020.
49. Gray, N. M., Sharp, L., Cotton, S. C., Masson, L. F., Little, J., Walker, L. G., Avis, M., Philips, Z., Russell, I., Whynes, D., Cruickshank, M., Woolley, C. M., & TOMBOLA group (2006). Psychological effects of a low-grade abnormal cervical smear test result: anxiety and associated factors. *British journal of cancer*, 94(9), 1253–1262. <https://doi.org/10.1038/sj.bjc.6603086>.
50. Sharp, L., Cotton, S., Cruickshank, M., Gray, N. M., Harrild, K., Smart, L., Walker, L. G., Little, J., & TOMBOLA Group (2014). The unintended consequences of cervical screening: distress in women undergoing cytologic surveillance. *Journal of lower genital tract disease*, 18(2), 142–150.
51. Pocnet, C., Antonietti, J. P., Strippoli, M. F., Glaus, J., Preisig, M., & Rossier, J. (2016). Individuals' quality of life linked to major life events, perceived social support, and personality traits. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*, 25(11), 2897–2908. <https://doi.org/10.1007/s11136-016-1296-4>
52. Staniute, M., Brozaitiene, J., & Bunevicius, R. (2013). Effects of social support and stressful life events on health-related quality of life in coronary artery disease patients. *The Journal of cardiovascular nursing*, 28(1), 83–89. <https://doi.org/10.1097/JCN.0b013e318233e69d>

53. Singer, E., & Ye, C. (2013). The Use and Effects of Incentives in Surveys. *The Annals of the American Academy of Political and Social Science*, 645, 112-141. doi:10.2307/23479084