

Title	Topology lecture notes
Authors	McKay, Benjamin
Publication date	2021-11-29
Original Citation	McKay, B. (2021) Topology Lecture Notes, Cork.
Type of publication	Book
Rights	© Benjamin McKay. This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 Unported License - https://creativecommons.org/licenses/by-sa/4.0/
Download date	2024-04-20 00:15:05
Item downloaded from	https://hdl.handle.net/10468/13431

Benjamin McKay

Topology Lecture Notes

November 29, 2021

Preface

There are many excellent books of topology; from the bibliography at the end of this book, the reader might look at [1, 2, 3, 5, 6, 7, 8, 9, 11, 10, 12, 13, 14, 15]. So it might have been wise to avoid writing these notes. They explain what I tried to cover in my 2017 lectures on topology for undergraduate students at University College Cork. I assume the reader is familiar with elementary theory of metric spaces. I will also give a few examples from differential geometry, which the reader can ignore, and which assume familiarity with manifolds and diffeomorphisms.

Contents

1	Topology	1
2	Continuity	17
3	Fundamental Groups	29
4	Covering Spaces	39
	Hints	65
	Bibliography	73
	Index	75

Chapter 1

■ Topology

We define topological spaces and their essential properties.

Motivation

When we study polynomial functions on the plane, there is a natural notion of “open set” different from the usual one: a *Zariski open set* is the set of points on which some polynomial function p is not zero. For example,

$$U = \{ (x, y) \in \mathbb{R}^2 \mid x^2 + y \neq 0 \}$$

is a Zariski open set. Of course, U is also an open set in the usual sense, but

$$W = \{ (x, y) \in \mathbb{R}^2 \mid x > 0 \}$$

is an open set in the usual sense, but not a Zariski open set.

1.1 Prove that W is not Zariski open.

So we can have different concepts of open set playing different but no less useful roles in the same space \mathbb{R}^2 . Intuitively, an open set is like a little “fat blob”. If you want to study polynomial functions, then sets like W above are not as fat as they “should” be, because any polynomial which doesn’t vanish on W *also* doesn’t vanish on some much larger set.

Definition

A *topology* on a set X is a collection of subsets of X , called the *open sets* of the topology, so that

- a. the union of any collection of open sets is an open set,
- b. the intersection of any finite collection of open sets is an open set,
- c. the empty set and the whole of X are open sets.

A *topological space* is a set X equipped with a topology; we usually leave the topology as implicitly understood somehow. The elements of the set X are called *points*.

As in our previous experience, every subset $X \subseteq \mathbb{R}^n$ is a topological space, with open sets being the intersections $X \cap U$ where $U \subseteq \mathbb{R}^n$ is the union of a collection of open balls, the *Euclidean topology*. If no topology is otherwise specified, we always mean the Euclidean topology.

If X is a metric space, then the usual open sets (unions of open balls) form a topology, making every metric space into a topological space, its *metric topology*. If no topology is otherwise specified, we always mean the metric topology.

Take any set X and let the open sets be all of the subsets of X , the *discrete topology*.

Take any set X and let the only open sets be the empty set and X , the *indiscrete topology*.

Take $X = \mathbb{R}^n$ and take as open sets the sets

$$\{x \in \mathbb{R}^n \mid p(x) \neq 0\}$$

for some polynomial function p on \mathbb{R}^n : the *Zariski topology*.

The *periodic topology* on $X = \mathbb{R}$ is the topology whose open sets are just those from among the usual (Euclidean) topology which happen to be 2π periodic.



Let X be the set of nonnegative real numbers, and take as open sets all sets of the form

$$\{x \mid x_0 < x\},$$

for any real number $x_0 \geq 0$. This is yet another topology.

If $S \subseteq X$ is any subset of a topological space, the *subspace topology* on S has open sets $S \cap U \subseteq S$ for $U \subseteq X$ any open set.

If X is any set, the *cofinite topology* has open sets just precisely (i) the empty set and (ii) the sets $X - F$ where F is any finite set.

1.2 Check that each of these examples correctly defines a topology.

1.3 What are all topologies on the empty set? On a set with one element $X = \{0\}$? On a set with two elements $X = \{0, 1\}$? On a set with three elements $X = \{0, 1, 2\}$? On a set with 4 elements $X = \{0, 1, 2, 3\}$?

Take distinct points p, q , let r be the distance between them, and look at the balls of radius $r/2$ around those points. Those balls are not empty, but don't intersect. Hence any metric space with two or more points contains nonintersecting nonempty open sets.

If we take nonempty Zariski open sets $U, W \subseteq \mathbb{R}^n$, we claim they intersect. Write

$$U = \{x \in \mathbb{R}^n \mid p(x) \neq 0\}$$

and

$$W = \{x \in \mathbb{R}^n \mid q(x) \neq 0\}.$$

We need to find a point in $U \cap W$, i.e. a point where neither of these polynomials vanish. Since U and W are not empty, we can take points x in U and y in W . Take the line between them. It is enough to find a point of that line on which neither of those polynomials vanish. We can rotate and translate to get that line to the x_1 -axis. Set all of the other variables except x_1 to zero. So it is enough to assume that both of our polynomials depend on only one variable. Each polynomial, being not everywhere zero, vanishes on a finite set of points. Remove those points, and neither of the polynomials vanish on any of the remaining points: U and W contain a point in common. Therefore the Zariski topology is *not* a metric topology of *any* metric (including, in particular, the usual metric).

Closed sets

If A is a subset of a set B , we write $B - A$ to mean the set of points of B not lying in A , the *complement* of A in B . A subset $C \subseteq X$ of a topological space X is *closed* if its complement $X - C \subseteq X$ is open.

1.4 Prove that the intersection of any closed sets is closed.

1.5 Prove that union of finitely many closed sets is closed.

1.6 Prove that the empty set and X are closed subsets of any topological space X . In particular, sets can be *both* open and closed (sets are *not* doors).

The *closure* \bar{A} of a set $A \subseteq X$ in a topological space X is the intersection of all closed sets containing A .

1.7 Find the closure of the rational numbers in the real numbers.

1.8 Find the closure of the open unit ball in \mathbb{R}^n .

1.9 Find the closure of the open unit ball in \mathbb{R}^n with the Zariski topology.

1.10 Prove that

- a. the closure \bar{A} of any subset $A \subseteq X$ of any topological space X is closed
- b. A is closed just when $A = \bar{A}$
- c. $A \subseteq \bar{A}$
- d. \bar{A} lies inside any closed set containing A .

Bases

A *neighborhood* of a point $x \in X$ in a topological space is a subset $S \subseteq X$ containing x so that there is an open subset $U \subseteq X$ lying inside S and containing x . A *basis* is a collection of open sets so that every open set is a union of open sets from the basis.

The open balls of a metric space form a basis for the metric topology.

The only basis of the cofinite topology on any infinite set is the entire cofinite topology.

All open sets taken together form a basis (there is no reasonable notion here of “independence” like in linear algebra).

1.11 Prove that every basis S of \mathbb{R}^n contains an infinite set $T \subset S$ so that $S - T$ is also a basis.

1.12 A collection of open sets of a topological space X forms a basis just when, for any point $x \in X$ and neighborhood $N \subseteq X$ of x , there is an element U in that collection so that $x \in U \subseteq N$.

1.13 Prove that the topology of \mathbb{R}^n has a countable basis.

1.14 Prove that the Zariski topology on \mathbb{R}^n does not have a countable basis. Hint: first try $n = 1$.

1.15* Give an example of an open set $U \subseteq \mathbb{R}$ containing the rational numbers so that $\mathbb{R} - U$ is uncountable.

A *basis* S on a set X is a collection of subsets of X so that any finite intersection of those subsets is also expressible as a union of some of those subsets. Clearly S is then a basis for a unique topology: the one whose open sets are unions of any collections of those subsets, called the topology by or by S .

Boundaries

Given a subset $A \subseteq X$ of a topological space, a point $x \in X$ is an *interior point* of A if A is a neighborhood of x , an *exterior point* of A if $X - A$ is a neighborhood of x , and a *boundary point* otherwise, i.e. if every neighborhood of x contains points both inside A and outside A . The *exterior* is the set of exterior points, and so on.

1.16 Find the interior, exterior and boundary of

- the set $A \subset X$ of rational numbers inside the set X of real numbers.
- the set $A \subset X$ of irrational numbers inside the set X of real numbers.
- the set $A \subset X$ of positive numbers inside the set X of real numbers.
- the set $A \subset X$ of positive numbers inside the set X of rational numbers.
- the unit interval $A \subset X$ inside the set of $X = \mathbb{R}$ in the Zariski topology.
- the rational numbers $A \subset X$ inside the set of $X = \mathbb{R}$ in the Zariski topology.

1.17 Find a subset $A \subset \mathbb{R}$ whose boundary has nonempty interior.

1.18 Take any set X with the discrete topology and any subset $A \subseteq X$. Find the interior, exterior and boundary of A .

1.19 Take any set X with the indiscrete topology and any subset $A \subseteq X$. Find the interior, exterior and boundary of A .

1.20 Find 3 different open subsets of the real number line that have the same boundary.

1.21 Suppose that U is an open subset of the plane and that the boundary of U is a finite set of points, say $\{p_1, p_2, \dots, p_k\}$. Prove that $U = \mathbb{R}^2 - \{p_1, p_2, \dots, p_k\}$.

Density

A subset $A \subseteq X$ of a topological space is *dense* in a subset $B \subset X$ if $B \subset \bar{A}$, and *everywhere dense* if $\bar{A} = X$.

1.22 Prove that every nonempty open set in \mathbb{R}^n in the Zariski topology is everywhere dense.

1.23 Prove that the rational numbers are dense in the real numbers.

In the indiscrete topology, every nonempty subset is everywhere dense.

In the discrete topology on a set X , only X is everywhere dense.

In \mathbb{R}^n , the points with rational coordinates are everywhere dense, as are the points with irrational coordinates.

1.24 If $A \subset X$ is dense and $U \subset X$ is open, prove that $A \cap U$ is dense in U .

Separability

A topological space X is *separable* if it contains a dense sequence of points, i.e. a sequence which enters every open set.

In the indiscrete topology, every sequence is dense, so the space is separable.

In the discrete topology on a set X , only X is everywhere dense, so if the points of X do not all lie on a single sequence, then X is not separable.

The real numbers are separable: put the rational numbers into a sequence.

Euclidean space is separable: put the points with rational coordinates into a sequence.


1.25 Which of the following topologies on \mathbb{R}^n make \mathbb{R}^n separable?

- a. The Euclidean topology.
- b. The discrete topology.
- c. The cofinite topology.
- d. The Zariski topology.


1.26* The *half-open topology* on \mathbb{R} is the topology generated by the half intervals $a \leq x < b$ for numbers $a < b \in \mathbb{R}$. Prove that this topology has a countable basis, is separable, and is *not* the Euclidean topology.

Products

Recall that the plane \mathbb{R}^2 is the product $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$. But the open sets of the plane are not all products; there are disks, interiors of triangles, and so on, as well.



The product open sets in the plane are only the products of intervals, i.e. the open boxes:



If X and Y are topological spaces, the *product topology* on $X \times Y$ has as open sets precisely those sets whose every point lies in a product $U_X \times U_Y$ of an open set $U_X \subset X$ and an open set $U_Y \subset Y$.

1.27 Take any basis for the topology of X , and any basis for the topology of Y . Take the open sets of the form of a product $U_X \times U_Y$ of a basis element $U_X \subset X$ and a basis element $U_Y \subset Y$. Prove that these form a basis for the product topology.

1.28 Suppose that $A \subset X$ and $B \subset Y$ are closed sets in topological spaces. Prove that $A \times B$ is a closed set in $X \times Y$ in the product topology.

1.29 Suppose that $A \subset X$ and $B \subset Y$ are sets in topological spaces and write A° for the interior of a set and \bar{A} for its closure. Prove that $(A \times B)^\circ = A^\circ \times B^\circ$ and that $\bar{A} \times \bar{B} = \overline{A \times B}$ in the product topology.

1.30 If $X = \mathbb{R}^p$ and $Y = \mathbb{R}^q$ with the Zariski topology, is the product topology on $X \times Y = \mathbb{R}^{p+q}$ equal to the Zariski topology on \mathbb{R}^{p+q} ?

Subspaces

If $A \subset X$ is a subset of a topological space X , the *subspace topology* has as open sets the sets $A \cap U$ for any open set $U \subset X$.

1.31 Prove that the subspace topology is a topology.

1.32* If a topological space is separable, is every subset separable in its subspace topology?

1.33 Suppose that $A \subset B \subset X$ are subsets of a topological space. Then A has a subspace topology as a subset of X , as does B . But then A has another subspace topology, as a subspace of B where B has its subspace topology as a subspace of X . Prove that these two topologies on A are the same topology.

If $X = \mathbb{R}$ and $A = \mathbb{R}^{>0}$, then the closure of A as a subset of A is A , but as a subset of X the closure of A inside X is $\mathbb{R}^{\geq 0}$.

Lemma 1.1. *If $A \subset X$ is a subset of a topological space X , then A has as closed sets just exactly the sets $A \cap C$ for any closed set $C \subset X$. Moreover, the closure of a subset $S \subset A$ inside A is the intersection of A with the closure of S in X .*

Proof. Take a subset $S \subset A$ and let $C_X \subset X$ be its closure in X and $C_A \subset A$ be its closure in A . Let $U_A = A - C_A$ and $U_X = X - C_X$. Then

$$C_A = \bigcap_{S \subset C} C$$

where the intersection is over the A -closed subsets of A containing S . So

$$\begin{aligned} U_A &= A - C_A, \\ &= \bigcap_{S \cap U \text{ empty}} U, \end{aligned}$$

where the intersection is over the A -open sets $U \subset A$ not intersecting S ; write those open sets as $U \cap A$, for X -open sets $U \subset X$:

$$= \bigcap_{S \cap U \cap A \text{ empty}} U \cap A,$$

but then $S \cap U \cap A = S \cap U$ since $S \subset A$, so

$$= \bigcap_{S \cap U \text{ empty}} U \cap A,$$

where the intersection is over the X -open sets $U \subset X$ not intersecting S ,

$$= A \cap \bigcap_{S \cap U \text{ empty}} U,$$

where the intersection is over the X -open sets $U \subset X$ not intersecting S ,

$$= A \cap U_X.$$

Therefore $C_A = A \cap C_X$. \square

1.34 A subset $A \subset X$ of a topological space is *locally closed* if each point a of A lies in an open subset $U \subset X$ of X so that $A \cap U \subset U$ is a closed subset of U . Given an example of a locally closed subset of $X = \mathbb{R}$ which is not closed. Given an example of a locally closed subset of $X = \mathbb{R}^3$ which is not closed.

The *disjoint union* of two topological spaces X, Y is the set $X \sqcup Y$ of all points of the form $(x, 1)$ for $x \in X$ or $(y, 2)$ for $y \in Y$. We give it the topology for which a basis of open sets consists of sets of the form $U \sqcup W$ for $U \subset X$ open and $W \subset Y$ open. Intuitively, $X \sqcup Y$ means X and Y “drawn separated from one another”. Typically, we will denote points of $X \sqcup Y$ as $x \in X$ or $y \in Y$ rather than as pairs $(x, 1)$ or $(y, 2)$, as long as this doesn’t confuse matters.

Hausdorff

Two point $x, y \in X$ of a topological space X are *housed off* from one another if there are open sets $U, V \subset X$ with $x \in U$, $y \in V$ and $U \cap V$ is empty.



If any two distinct points of X are housed off, then X is *Hausdorff*.

1.35 If X is any set with the cofinite topology, prove that X is Hausdorff just when X is finite.

1.36 Prove that every metric space is Hausdorff.

Lemma 1.2. *The Zariski topology is not Hausdorff.*

Proof. Take two points $x, y \in \mathbb{R}^n$ and two Zariski open sets U, V with $x \in U$ and $y \in V$. We write U as the set of points at which some polynomial p doesn't vanish, and V as the set of points at which some polynomial q doesn't vanish. Draw the line from x to y . Restrict the polynomials to that line. None of our polynomials vanishes everywhere on that line, since each is nonzero at x or at y . So each vanishes at only a finite number of points. Remove those points to see that U and V intersection. \square

1.37 Prove that in any Hausdorff space X , for any point x of X , the set $\{x\}$ is closed: “points are closed”.

1.38 Prove that, in the Zariski topology, “points are closed”, even though the Zariski topology is not Hausdorff.

1.39 Prove that in any topological space, “points are closed” if and only if finite sets are closed.

If X is a topological space, the *diagonal* $\Delta_X \subset X \times X$ be, i.e. the set of points (x, x) for x in X .

1.40* A topological space is Hausdorff just when its diagonal is closed.

1.41 If $X = \mathbb{R}^n$ with the Zariski topology, prove that the diagonal in $X \times X$ is dense in the product topology, but not in the Zariski topology on \mathbb{R}^{2n} .

1.42 The product $X \times Y$ of two Hausdorff spaces X, Y is also a Hausdorff space.

1.43 Prove that, for any subset $A \subset X$ of a Hausdorff space, A is Hausdorff in the subspace topology on A .

Compactness

Recall that a subset $A \subset \mathbb{R}^n$ is compact just when every infinite sequence of points has a convergent subsequence. We will try (and fail) to imitate this in topological spaces. A sequence x_1, x_2, \dots of points of a topological space X *converges* to a point $x \in X$ if every open set containing x contains all but finitely many points of that sequence.

Take the sequence of points $x_j = (j, 2^j) \in \mathbb{R}^2$ and let $X = \mathbb{R}^2$ with the Zariski topology. Take any open set U , not empty. Then U has the form $U = \mathbb{R}^2 - C$ where $C \subset \mathbb{R}^2$ is an algebraic curve in the plane (or C is the empty set). Only finitely many points of our sequence can lie inside C , because the points lie on the graph of $y = 2^x$, which is not an algebraic function. (The reader should explain why that is the case.) So, for any nonempty open set, all but finitely many points of our sequence lie in that open set. In particular, this sequence converges to every point in the plane, in the Zariski topology.

Sequences turn out to be the wrong objects to work with in topology. We work with open sets.

Recall another defining property of compact sets in \mathbb{R}^n : every open cover has a finite subcover. An *open cover* of a set $A \subset X$ in a topological space X is a collection of open sets whose union contains A . A *subcover* is a smaller collection of open sets, each one belonging to the original collection.

The set of open balls of radius $1/2$ is an open cover of the plane \mathbb{R}^2 . The subset of open balls of radius $1/2$ with rational centre point is a subcover. The set of open balls of radius $1/3$ is *not* a subcover: these balls are not drawn from our original set. The set of open balls of radius $1/2$ around integer points is not a subcover: the point $(1/2, 1/2)$ is more than $1/2$ unit from any integer point.

A topological space X is *compact* if every open cover has a finite subcover.

1.44 Prove that every finite subset $A \subset X$, of any topological space X , is compact.

Lemma 1.3. *Every locally bounded function $f: X \rightarrow \mathbb{R}$ on a compact topological space X is bounded.*

Proof. Each point of X lies in an open set in which f is bounded. These open sets cover X . Take a finite subcover. So we have finitely many open sets, and a bound of f on each, from above and from below. Take the maximum of the upper bounds, and minimum of the lower bounds. \square

1.45 Suppose that $A \subset X$ is a subset of a compact topological space. Prove that if A is closed then A is compact. If X is also Hausdorff, prove that A is closed just when A is compact.

1.46 Give an example of a compact space X which is not Hausdorff and a compact set $A \subset X$ which is not closed.

1.47 Prove that, in any topological space, the union of finitely many compact sets is compact. Give an example to prove that the union of infinitely many compact sets need not be compact.

Lemma 1.4. *Topological spaces X and Y are both compact just when their product is compact.*

Proof. Suppose that $X \times Y$ is compact. Take an open cover X_a of X . Then $X_a \times Y$ is an open cover of $X \times Y$. Take a finite subcover, say $X_i \times Y$, $i = 1, 2, \dots, n$. Then X_1, X_2, \dots, X_n is a finite subcover of the collection of X_a . The same trick for Y in place of X .

Suppose that X and Y are compact. Take an open cover by open sets $U_a \subset X \times Y$. Each point $(x, y) \in X \times Y$ lies in one of these open sets, call it

$U_{x,y}$. The product open sets $X_b \times Y_c \subset X \times Y$ form a basis. So there is such a product, call it $X_{x,y} \times Y_{x,y}$, inside $U_{x,y}$.

For each point $x \in X$, the various $Y_{x,y}$ cover Y . Take a finite subcover, say

$$Y_{x,y_1(x)}, \dots, Y_{x,y_{n_x}(x)}.$$

Let

$$X_x = X_{x,y_1(x)} \cap \dots \cap X_{x,y_{n_x}(x)}.$$

Since $x \in X_x$, these various open sets $X_x \subset X$ form an open cover of X . Take a finite subcover X_1, X_2, \dots, X_n , say containing points $x_1 \in X_1, \dots, x_n \in X_n$. Then $X \times Y$ is covered by the finitely many sets $U_{x_i,y_j(x_i)}$ for all possible values of i and j for which this is defined. \square

Lemma 1.5. *In any Hausdorff space, every compact set is closed.*

Proof. Suppose that X is Hausdorff. Since X is Hausdorff, any two points $x, y \in X$ have disjoint “houses”. Suppose that x is outside a compact set $K \subseteq X$. Fixing x and letting y vary over K , finitely many of our “houses” cover those points of K . The finite intersection of the corresponding “houses” around x give an open set around x not intersecting K . So every point x outside K lies in an open set outside K , i.e. the complement of K is open. \square

1.48 Give an example of a compact but not closed subset of some non-Hausdorff space.

1.49 Prove that, in any Hausdorff space, for any collection of compact sets, the intersection of all of those sets is compact. If the intersection of any finite number of those compact sets is not empty, prove that the intersection of them all is not empty.

1.50 Give an example of a space, not Hausdorff, and two compact subsets of that space, whose intersection is not compact.

1.51 Take compact sets $K \subset X$, $L \subset Y$ of topological spaces. Suppose that $W \subset X \times Y$ is an open set containing $K \times L$. Prove that W contains an open set of the form $U \times V$ so that $U \subset X$ and $V \subset Y$.

Connectivity

A topological space X is *connected* if it is not expressible as a disjoint union $X = U \cup V$ of two nonempty open sets $U, V \subset X$.

1.52* Prove that any interval of the real number line is connected.

1.53* If a subset $S \subseteq X$ of a topological space is connected, prove that its closure is connected.

1.54* If a subset $S \subseteq X$ of a topological space is connected, and $f: X \rightarrow Y$ is a continuous map, prove that $f(S) \subset Y$ is connected.

1.55* A *coprime arithmetic progression* is a set of integers of the form

$$\dots, a - 3b, a - 2b, a - b, a, a + b, a + 2b, a + 3b, \dots$$

where a and b are integers with no common prime factor. The *Golomb topology* on the integers is the topology for which the coprime arithmetic progressions form a basis for the open sets. Prove that the set of integers, in the Golomb topology, is Hausdorff, connected and not compact.

1.56 Prove that every topological space is uniquely expressed as a union of its maximal connected subsets, called its *connected components*.

1.57 Suppose that $f: X \rightarrow Y$ is a continuous open map of topological spaces and that all fibers $f^{-1}\{y_0\}$ are connected, $y_0 \in Y$. Prove that f bijectively identifies the connected components of X with those of Y .

Path connectivity

A topological space X is *path connected* if any two points lie on the image of a path, i.e. a continuous map from an interval of the real number line.

1.58 Prove that any path connected space is connected.

1.59 Prove that every topological space is uniquely expressed as a union of its maximal path connected subsets, called its *path components*.

1.60 Suppose that $f: X \rightarrow Y$ is a continuous open map of topological spaces and that all fibers $f^{-1}\{y_0\}$ are path connected, $y_0 \in Y$. Prove that f bijectively identifies the path components of X with those of Y .

The union of the graph of $y = \sin(1/x)$ with the line $x = 0$ is connected, but not path connected: the graph is path connected, as is the line, so each lies in a single path component, so in a single component. Any open set around the line $x = 0$ intersects the graph, so intersects any open set containing the graph: there is only one component. But there are two path components, since a continuous path along the graph hits the infinitely many peaks and troughs, with infinitely many points where $y = 1$, and where $y = -1$, so no limit as it approaches the line.

1.61* Give an example of a connected space with infinitely many path components.

1.62 If a subset $S \subseteq X$ of a topological space is path connected, and $f: X \rightarrow Y$ is a continuous map, prove that $f(S) \subset Y$ is path connected.

A topological space X is *locally path connected* if, for any point $x_0 \in X$ and open set $U \subset X$ containing x_0 , there is a path connected open set $W \subset U$ containing x_0 with $W \subset U$.

1.63 If a topological space is locally path connected, prove that its path components are its components.

The Baire category theorem

A subset $A \subset X$ of a topological space is *everywhere dense* if A is dense in X .

For each rational number $q \in \mathbb{Q}$, take the set $U_q := \{x \in \mathbb{R} \mid x \neq q\}$. So the various $U_q \subset \mathbb{R}$ are dense open sets. Their intersection

$$\bigcap_{q \in \mathbb{Q}} U_q \subset \mathbb{R}$$

is precisely the set of irrational numbers, not open, but still dense. Roughly speaking, if we only pull out a single rational at each step, we still have a lot left over after an infinite sequence of steps.

A set $B \subset X$ of a topological space is *nowhere dense* if $B \cap U$ is not dense in any open set $U \subset X$. A *meager set* is a subset $S \subset X$ of a topological space, which can somehow be written as

$$S = C_1 \cup C_2 \cup \dots$$

as the union of a sequence of nowhere dense closed sets. A *comeager set* is a subset $A \subset X$ of a topological space, which can somehow be written as

$$A = U_1 \cap U_2 \cap \dots$$

as the intersection of a sequence of dense open sets.

1.64 Prove that a subset of a topological space is meager just when its complement is comeager.

A topological space X is *Baire* if every meager set has dense complement, or equivalently if every comeager set is dense.

1.65 Prove that every complete metric space is a Baire space, in its metric topology.

1.66 Prove that every open subset of any Baire space is Baire. Use this to prove that every meager subset of an open set in a complete metric space is nowhere dense.

A topological space X is *locally compact* if every point of X lies in the interior of a compact set.

1.67 Suppose that $K \subset X$ is a compact subset of a locally compact topological space. Prove that there is an open set $V \subset X$ containing K so that $\bar{V} \subset X$ is compact.

Theorem 1.6 (Baire category theorem). *In any Hausdorff locally compact space, every meager set has dense complement, i.e. every Hausdorff locally compact space is a Baire space.*

Proof. Take a comeager set $A = U_1 \cap U_2 \cap \dots$, with each U_i open and dense. Since U_1 is dense, it is not empty; take a point $x_1 \in U_1$. Take a compact set K_1 with x_1 in the interior of K_1 . Since U_2 is open and dense, it intersects the interior of K_1 ; pick a point x_2 in that intersection. Take a compact set K_2 with x_2 in the interior of K_2 . We can replace K_2 by $K_2 \cap K_1$, so assume that $K_2 \subset K_1$. By induction, generate nested compact sets $\dots \subset K_3 \subset K_2 \subset K_1$. The intersection is not empty, by the solution to problem 1.49 on page 12. \square

Given a topological space X , we say that “the generic element of X has the property ...” to mean that the set of elements of X which do *not* have the property ... is meager.

The generic real number is irrational (has the property of being irrational), i.e. the rationals form a countable union of closed, nowhere dense sets in the set X of real numbers.

The Baire category theorem says that, in any Baire space, any generic property occurs on a dense set.

The generic real number is irrational, and the real numbers form a complete metric space (so Baire), so the irrational numbers are dense in the real numbers.

The generic point of the unit ball in Euclidean space \mathbb{R}^n is irrational (i.e. has all coordinates irrational), and the ball is an open subset of a complete metric space (so Baire), so the irrational points are dense in the ball.

Take a sequence of polynomial functions $p_1(x, y), p_2(x, y), \dots$ in two variables x, y , each function nonzero somewhere. Associate to each polynomial $p_j(x, y)$ the set of its zeroes: $C_j := \{(x, y) \mid p_j(x, y) = 0\}$. The plane is Baire and contains the nowhere dense closed sets C_j . So the union has dense complement: you can avoid satisfying *all* of the polynomial equations $p_j(x, y) = 0$ by slight perturbation of any point (x, y) .

A *transcendental point* of \mathbb{R}^n is a point not satisfying any nonconstant polynomial equation $0 = p(x)$ with rational coefficients. Transcendental points of \mathbb{R}^n are generic, so dense, as there are countably many such polynomials, each with a closed, nowhere dense, zero set.

Local compactness

In any topological space, a subset with compact closure is *precompact*.

1.68 For any Hausdorff space X , prove that the following are equivalent:

- a.* X is locally compact.
- b.* Every point of X lies in a precompact open set.
- c.* X has a basis of precompact open sets.

So if X is locally compact Hausdorff, every open set around any point contains a precompact open set around that same point.

1.69 Prove that every closed or open set in any locally compact Hausdorff space is locally compact Hausdorff.

1.70 Prove that the product of two locally compact Hausdorff spaces is locally compact Hausdorff.

Chapter 2

Continuity

If $f: X \rightarrow Y$ is a map, the *preimage* $f^{-1}S$ of a subset $S \subseteq Y$ is the set

$$f^{-1}S := \{x \in X \mid f(x) \in S\} \subseteq X.$$

A *continuous map* $f: X \rightarrow Y$ is a map so that the preimage of any open set in Y is an open set in X .

2.1 If X and Y are subsets of Euclidean space, prove that this agrees with the usual definition.

Another way to say this: for any point $x_0 \in X$ and associated point $y_0 = f(x_0)$, if you want $y = f(x)$ to stay in some neighborhood of the point y_0 , you only have to keep x in a suitable neighborhood of x_0 . Continuity demands precisely that the preimage of a neighborhood is also a neighborhood.

2.2 Suppose that X and Y are sets equipped with the cofinite topology. Prove that a map $f: X \rightarrow Y$ is continuous just when each point has finite preimage.

2.3 Prove that a map $f: X \rightarrow Y$ is continuous just when the preimage of any closed set is a closed set.

2.4 We don't need to check *all* open sets: prove that, for any basis $Y_a \subset Y$ of open sets of a topological space Y , a map $f: X \rightarrow Y$ is continuous just when the preimage of any Y_a is open.

2.5 Prove that a map $f: X \rightarrow Y$ is continuous just when preimage commutes with closure, i.e. for any subset $A_Y \subset Y$, with closure $\bar{A}_Y \subset Y$, and with preimage denoted $A_X = f^{-1}A_Y$, we have $\bar{A}_X = f^{-1}\bar{A}_Y$.

2.6* If X and Y are Hausdorff spaces and $f: X \rightarrow Y$ is a continuous map, prove that the graph of f is a closed subset of $X \times Y$.

2.7 Prove that every polynomial function $f: \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous in the Zariski topologies of \mathbb{R}^n and \mathbb{R} .

2.8 Prove that the function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = \sin x$ is discontinuous in the Zariski topology of \mathbb{R} .

2.9 Prove that the function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = e^x$ is continuous in the Zariski topology of \mathbb{R} .

2.10 Prove that the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by $f(x, y) = y + e^x$ is discontinuous in the Zariski topology of \mathbb{R} .

2.11 Prove that constant maps are continuous.

2.12 Suppose that X is a set with the discrete topology, and Y is any topological space. What are all continuous maps $f: X \rightarrow Y$?

2.13 Suppose that X is the real number line, and Y is a set with the discrete topology. What are all continuous maps $f: X \rightarrow Y$?

Lemma 2.1. *The composition $g \circ f$ of any continuous maps $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ is continuous.*

Proof. For any open set $U_Z \subset Z$, note that the set

$$U_X = (g \circ f)^{-1} U_Z$$

is just

$$U_Z = g^{-1} U_Y$$

where

$$U_Y = f^{-1} U_X.$$

□

If $A \subset X$ is a subset and $f: X \rightarrow Y$ is a map, define

$$f|_A: A \rightarrow Y$$

by

$$f|_A(a) = f(a)$$

for every a in A ; the map $f|_A$ is the *restriction* of f to A .

2.14 Prove that the restriction of a continuous map to any set is continuous.

2.15 Continuity is “local”: take some open sets $X_a \subset X$ whose union is X . Prove that any map $f: X \rightarrow Y$ is continuous just when all of its restrictions $f|_{X_a}: X_a \rightarrow Y$ are continuous.

2.16 Continuity is “very local”. A map $f: X \rightarrow Y$ is *continuous at a point* x in X if, for any open set $U_Y \subset Y$ containing $f(x)$, there is an open set $U_X \subset X$ containing x so that $fU_X \subset U_Y$. Prove that any map is continuous just when it is continuous at every point.

2.17 Take topological spaces X, Y and let $p: X \times Y \rightarrow X$ be the map $p(x, y) = x$, the *projection map*. Prove that the projection map is continuous.

2.18 Take topological spaces X, Y, Z and continuous maps $f: X \rightarrow Y$ and $g: X \rightarrow Z$ and let $(f, g): X \rightarrow Y \times Z$ be $(f, g)(x) = (f(x), g(x))$. Prove that (f, g) is continuous.

2.19 Take topological spaces X, Y and let $f: X \times Y \rightarrow Y \times X$ be the map $f(x, y) = (y, x)$. Prove that f is continuous.

Lemma 2.2. Suppose that $f, g: X \rightarrow \mathbb{R}^n$ are continuous. Then $f+g: X \rightarrow \mathbb{R}^n$ is continuous.

Proof. Compose $(f, g): X \rightarrow \mathbb{R}^n \times \mathbb{R}^n$ with the map $+: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, continuous maps. \square

Similarly, multiplication and division are continuous, etc.

2.20* Any continuous map $f: X \rightarrow Y$ takes any compact subset $K \subset X$ to a compact set $f(K) \subset Y$.

Density

The zeroes of a continuous function on \mathbb{R}^n form a closed set. The same is true in much more generality:

Lemma 2.3. Take two continuous maps $f, g: X \rightarrow Y$. Let $A \subset X$ be the set of points at which $f = g$. If Y is Hausdorff, then A is closed.

Proof. Consider the map $F: X \rightarrow Y \times Y$ given by $F(x) = (f(x), g(x))$. Let $\Delta_Y \subset Y \times Y$ be the diagonal, i.e. the set of points of the form $(y, y) \in Y \times Y$ for any $y \in Y$. Then clearly $A = F^{-1}\Delta_Y$. By problem 1.40 on page 10, $\Delta_Y \subset Y \times Y$ is closed just when Y is Hausdorff. So then A is closed. \square

Lemma 2.4. If two continuous maps $f, g: X \rightarrow Y$ agree on a dense subset of a Hausdorff space X , then they agree everywhere.

Proof. By lemma 2.3, the set of points where $f = g$ is closed, so if dense, is X . \square

Homeomorphisms

A *homeomorphism* is a continuous map $f: X \rightarrow Y$ with a continuous inverse $f^{-1}: Y \rightarrow X$. If a homeomorphism $f: X \rightarrow Y$ exists, X and Y are *homeomorphic*; for purposes of topology, they are essentially identical.

The map $f: x \in \mathbb{R} \mapsto \arctan x \in (-1, 1)$ is a homeomorphism: the real number line is homeomorphic to any open interval of the real number line.

Let X be the interval $[0, 2\pi) \subset \mathbb{R}$. Let Y be the unit circle in the plane $Y \subset \mathbb{R}^2$. The map $f: \theta \in X \mapsto (\cos \theta, \sin \theta) \in Y$ is continuous, and has an inverse $f^{-1}(\cos \theta, \sin \theta) = \theta$. But f is *not* a homeomorphism. The map f^{-1} is discontinuous: for points just above the positive horizontal

axis, and points just below it, f^{-1} takes these close points far apart. To make a proof out of this idea: the f^{-1} preimage (i.e. the f image) of the open set $[0, \varepsilon)$ is not open in the circle for $\varepsilon < 2\pi$.

2.21 Prove that the plane is homeomorphic to any open ball in the plane.

Theorem 2.5. *A continuous bijection $f: X \rightarrow Y$ from a compact space X to a Hausdorff space Y is a homeomorphism.*

Proof. By continuity, open sets have open preimages; equivalently, closed sets have closed preimages. Closed sets of X are compact, so their images are compact, so closed: closed sets have closed images. Equivalently, open sets have open images. Hence the inverse is continuous. \square

2.22 Prove that the open unit disk in the plane is not homeomorphic to the real number line.

Proper maps

A *proper map* $f: X \rightarrow Y$ is a continuous map so that the preimage of any compact set is compact.

On $X = \mathbb{R}$, every nonconstant polynomial is proper.

On $X = \mathbb{R}^2$, the polynomial $x^2 + y^2$ is proper, while $x, x^3, x^2y^2, x^2 + y^3$ are not proper.

2.23 Which trigonometric and inverse trigonometric functions are proper maps?


2.24 The *one point compactification* of a topological space X is the topological space \bar{X} whose points are $X \cup \{\infty\}$ for some element ∞ not in X , and whose open sets are (a) the open sets of X and (b) the sets $\{\infty\} \cup (X - C)$ for $C \subset X$ compact. Prove that these sets form the open sets of a topology. Prove that a continuous map $f: X \rightarrow Y$ of Hausdorff spaces is proper if and only if it extends to a continuous map $\bar{f}: \bar{X} \rightarrow \bar{Y}$ with $\bar{f}(\infty) = \infty$.

2.25 For a topological space X , prove that X is a locally compact Hausdorff space if and only if there exists a topological space X' so that

- X is embedded into X' and
- $X' - X$ is a single point (to be denoted ∞) and
- X' is a compact Hausdorff space.

Moreover X' is then homeomorphic to the one point compactification of X by the unique homeomorphism which is the identity on X .

2.26 Prove that the one point compactification of Euclidean space \mathbb{R}^n is homeomorphic to the sphere S^n .



A map $f: X \rightarrow Y$ is *open* if the image of any open set is open, *closed* if the image of any closed set is closed.

2.27 Prove that any closed injection $f: X \rightarrow Y$ is a homeomorphism to its image.

2.28* A test to decide if a map is proper: prove that a continuous map $f: X \rightarrow Y$ is proper if and only if it satisfies the two conditions (a) preimages of points are compact and (b) f is closed.

2.29 Prove that every continuous map $f: X \rightarrow Y$ from a compact space to a Hausdorff space is proper.

Lemma 2.6. *Every proper map $f: X \rightarrow Y$ to a locally compact Hausdorff space Y is closed.*

Proof. Every open set $U \subseteq Y$ lying in some compact set has compact closure, i.e. is precompact, so $f^{-1}\bar{U}$ is compact.

Take $A \subseteq X$ closed. Pick a point $y \in Y - f(A)$. We need only prove there is an open set U in Y containing y avoiding $f(A)$. Take any precompact open set $U \subset Y$ containing y . Write A as a union of the compact set $A' = A \cap f^{-1}\bar{U}$ and the subset $A'' = A \cap f^{-1}(X - \bar{U})$. Since $f(A'')$ avoids U , we need only arrange that $f(A')$ avoids some perhaps smaller open set W around y , and then $f(A)$ will avoid $U \cap W$. So we can replace A by A' , X by $f^{-1}\bar{U}$, Y by \bar{U} , so assume both X and Y are compact. By problem 1.45 on page 11, $A \subseteq X$ is compact. By problem 2.20 on page 19, $f(A)$ is compact, and so closed. \square

A test for homeomorphism

Corollary 2.7. *Take a locally compact Hausdorff space Y . Any closed injection $f: X \rightarrow Y$ is a homeomorphism to its image. In particular, any proper injection $f: X \rightarrow Y$ is a homeomorphism to its image.*

Proof. One proof: by lemma 2.6 on the previous page, f is closed; apply problem 2.27 on the preceding page.

Another proof, which might give more intuition: write Y as a union of precompact open sets $Y_a \subset Y$. Let $\bar{X}_a := f^{-1}\bar{Y}_a$. Then

$$f|_{\bar{X}_a} : \bar{X}_a \rightarrow \bar{Y}_a$$

is a homeomorphism to its image by theorem 2.5 on page 20. In particular, if we let $X_a := f^{-1}Y_a$, then

$$f|_{X_a} : X_a \rightarrow Y_a$$

is also a homeomorphism to its image. Take any open set $U \subset X$ and let $U_a := X_a \cap U$. Then

$$f(U) = f\left(\bigcup_a U_a\right) = \bigcup_a f(U_a)$$

is open: open sets have open images. Hence the inverse is continuous. \square

Quotient topologies

Let's glue things together.

The *Möbius strip* is given by gluing two sides of a square together, in opposite directions. Imagine X is the square, and Y is the square after we identify those points together; define the map $X \rightarrow Y$ which takes each unglued point to the corresponding point after gluing. But what is the topology on Y ?



Suppose that we have a map $f: X \rightarrow Y$ between two sets X and Y . If X has a topology given, but Y doesn't, the *quotient topology* on Y has an open sets precisely those subsets of Y whose preimages in X happen to be open. Clearly for any map $f: X \rightarrow Y$, the quotient topology is the "simplest" (has the fewest open sets) so that f becomes continuous. In a picture, if we draw X as a looking like a box, and Y as its shadow on the plane:



and our map takes each point



to its shadow:



then open sets in Y :



are sets whose preimage is open in X :



A *pointed space* or *space with marked point* is a topological space X together with a point x_0 of X , often denoted (X, x_0) . If (X, x_0) and (Y, y_0) are spaces with marked points, the *join* of those spaces is the quotient space $(X \sqcup_{x_0=y_0} Y, x_0)$ defined as follows: let $X \sqcup_{x_0=y_0} Y$ be $X \sqcup Y$ but removing the point y_0 . Map $f: X \sqcup Y \rightarrow X \sqcup_{x_0=y_0} Y$ by $f(x) = x$ but

$$f(y) = \begin{cases} y & \text{if } y \neq y_0, \\ x_0 & \text{if } y = y_0. \end{cases}$$

Give $X \sqcup_{x_0=y_0} Y$ the marked point x_0 and the quotient topology from this map f .

Take n circles, each with a marked point, and join to get the *bouquet of circles*:



2.30 Prove that if $f: X \rightarrow Y$ is a surjective map and X is a complete metric space and every fiber $f^{-1}\{y\} \subset X$ of f is a closed subset, and any two distinct fibers $f^{-1}\{y_1\}, f^{-1}\{y_2\}$ stay a positive distance apart, then the quotient topology on Y arises from a metric on Y .

Another language we use to talk about surjective maps $f: X \rightarrow Y$ is the language of equivalence relations. We can say that two points of X are equivalent if they have the same value under f . On the other hand, given an equivalence relation \sim on X , define the *quotient* X/\sim , to be the set of all equivalence classes, and the *quotient map* $X \rightarrow X/\sim$ by sending each point $x \in X$ to its equivalence class $[x]$. If we write X/\sim as Y and the quotient map as f (to avoid the intimidating notation), we define the *quotient topology* by an equivalence relation to be the quotient topology via the quotient map. It is just a change of notation and language, but we get just the same quotient spaces if we think about surjective maps $f: X \rightarrow Y$ or equivalence relations \sim on a topological space X .

Theorem 2.8. Take a topological space X and an equivalence relation \sim on X with quotient map $X \rightarrow X/\sim$. Any continuous map $X/\sim \rightarrow Y$ induces a continuous map $X \rightarrow Y$ by composing with the quotient map: $X \rightarrow X/\sim \rightarrow Y$.

Conversely, a continuous map $X \rightarrow Y$ arises in this way, i.e. has the form of a composition $X \rightarrow X/\sim \rightarrow Y$, for a unique continuous map $X/\sim \rightarrow Y$ just when it is constant on equivalence classes, i.e. equivalent elements go to the same point of Y . We then say that that our map $X \rightarrow Y$ descends to the quotient.

Proof. Suppose that $f: X \rightarrow Y$ is a continuous map taking equal values in Y for any equivalent points of X . We define a map, perhaps not continuous, by $\tilde{f}([x]) = f(x)$ on each equivalence class $[x] \in X/\sim$. Clearly this is well

defined, but we need to prove that it is continuous. Let $q: X \rightarrow X/\sim$ be the quotient map taking each point $x \in X$ to its equivalence class $[x] \in X/\sim$. Note also that by definition $f(x) = \bar{f}([x])$, i.e. $f = \bar{f} \circ q$. Take an open set $U_Y \subset Y$. Take its preimage $U_{X/\sim} \subset X/\sim$. We need to prove that $U_{X/\sim} \subset X/\sim$ is open. By definition of the quotient topology, this is precisely proving that $U_X := q^{-1}U_{X/\sim}$ is open. But this is just

$$\begin{aligned} U_X &= q^{-1}\bar{f}^{-1}U_Y, \\ &= (\bar{f} \circ q)^{-1}U_Y, \\ &= f^{-1}U_Y, \end{aligned}$$

open in X . □

2.31 Suppose that $f: X \rightarrow Y$ is a surjective map from a topological space X . Suppose that X contains a compact set K for which $f(K) = Y$. Prove that Y is compact in the quotient topology.

2.32 A map $f: X \rightarrow Y$ is *strict* if the induced topology on $f(X)$ agrees with the quotient topology. Prove that every surjective open map is strict, and that every surjective closed map is strict.

Often we can guess what the quotient space by a map should be, and we need the following theorem to check our guess.

Theorem 2.9. *Suppose that $f: X \rightarrow Y$ is a continuous surjective map and Y is Hausdorff. Suppose that every point of Y lies in the interior of the image of a compact subset of X . Define an equivalence relation: $x_1 \sim x_2$ just when $f(x_1) = f(x_2)$ for any x_1, x_2 in X . Then $f: X \rightarrow Y$ descends to the quotient to a homeomorphism $X/\sim \rightarrow Y$.*

Proof. Clearly the quotient map $X/\sim \rightarrow Y$ is a bijection and constant on equivalence classes, so continuous. Pick an open subset $U_Y \subset Y$, which lies in the image of some compact set $K_X \subset X$. We can replace K_X by its intersection with the preimage of \bar{U}_Y , so that $f(K_X) = \bar{U}_Y$. By problem 2.20 on page 19, the image of K_X in $K_{X/\sim} \subset X/\sim$ is compact. By theorem 2.5 on page 20, $K_{X/\sim} \cong \bar{U}_Y$ is a homeomorphism. So near each point of Y , the quotient map is a homeomorphism. □

2.33* Suppose that f is a continuous function

$$f(v_1, v_2, \dots, v_n)$$

of vector variables $v_j \in \mathbb{R}^n$. Suppose that f is invariant under simultaneous orthogonal transformation of all of the vectors:

$$f(Uv_1, Uv_2, \dots, Uv_n) = f(v_1, v_2, \dots, v_n),$$

for any orthogonal $n \times n$ matrix U . Let Y be the set of all positive semidefinite symmetric matrices, i.e. symmetric matrices A so that $\langle Av, v \rangle \geq 0$ for any

vector $v \in \mathbb{R}^n$. Prove that there is a continuous function $g: Y \rightarrow \mathbb{R}$ so that f can be written as

$$f(v_1, v_2, \dots, v_n) = g(A)$$

where A is the matrix with entries $A_{ij} = \langle v_i, v_j \rangle$.

2.34 Suppose that \mathcal{B} is a collection of open subsets of a topological space X . Suppose that, for any $U, V \in \mathcal{B}$, and point $x \in U \cap V$, there is an open set $W \in \mathcal{B}$ so that $x \in W \subseteq U \cap V$. Prove that there is a topological space Y and an onto continuous map $f: X \rightarrow Y$ so that f takes every set $U \in \mathcal{B}$ to an open set $f(U)$, and $f^{-1}f(U) = U$, and these sets $f(U)$ form a basis of open sets of Y .

If $A \subset X$ is a subset of a topological space, X/A means the quotient where we make any two points of A equivalent, and no other points equivalent to one another. If X is a metric space and A is closed then X/A is also a metric space, so we have an enormous collection of examples of metric spaces.

2.35 Let X be the closed unit interval $[0, 1] \subset \mathbb{R}$ and Y the unit circle in the plane. Let $f: X \rightarrow Y$ by $f(x) = (\cos 2\pi x, \sin 2\pi x)$. Prove that $X/\{0, 1\}$ is homeomorphic to Y .

2.36* Let X be the closed unit ball $X = \bar{B} \subset \mathbb{R}^n$ and $Y \subset \mathbb{R}^{n+1}$ the unit sphere. Let $A \subset X$ be the unit sphere. Write points of \mathbb{R}^{n+1} as (t, x) for $t \in \mathbb{R}$ and $x \in \mathbb{R}^n$. Consider the map $f: X \rightarrow Y$ given by

$$f(x) = \begin{cases} \left(\cos(\pi|x|), \sin(\pi|x|)\frac{x}{|x|} \right), & \text{if } |x| \neq 0, \\ (1, 0), & \text{if } |x| = 0. \end{cases}$$

Prove that f is continuous. Prove that f is injective from the interior of X to $Y - \{(-1, 0)\}$ and that f takes every point of A to the point $(-1, 0) \in Y$. Prove that X/A is homeomorphic to Y .

The *Möbius strip* is the quotient of the closed unit square $X = [0, 1] \times [0, 1]$ by the equivalence relation $(0, y) \sim (1, 1 - y)$:



The *Klein bottle* is the quotient of the closed unit square $X = [0, 1] \times [0, 1]$ by the equivalence relation $(x, 0) \sim (x, 1)$ and $(0, y) \sim (1, 1 - y)$:



The *torus* is the quotient of the closed unit square $X = [0, 1] \times [0, 1]$ by the equivalence relation $(x, 0) \sim (x, 1)$ and $(0, y) \sim (1, y)$.



The *real projective space* is the quotient $\mathbb{RP}^n = S^n / (x \sim -x)$. Every line through the origin strikes the unit sphere at two antipodal points. In other words, a point of real projective space corresponds to a line through the origin in \mathbb{R}^{n+1} .

The *complex projective space* is the quotient $\mathbb{CP}^n = \mathbb{RP}^{2n+1} / \sim$ where a unit length vector $z \in \mathbb{C}^{n+1}$ is equivalent to $e^{i\theta}z$ for any real number θ . In other words, a point of complex projective space corresponds to a complex line through the origin in \mathbb{C}^{n+1} .

The *quaternionic projective space* is the quotient $\mathbb{HP}^n = \mathbb{CP}^{2n+1} / \sim$ where a unit length vector $z \in \mathbb{H}^{n+1}$ is equivalent to λz for any unit length quaternion λ . In other words, a point of quaternionic projective space corresponds to a quaternionic line through the origin in \mathbb{H}^{n+1} .

Let \bar{D} be the closed unit disk, i.e. the set of points (x, y) so that $x^2 + y^2 \leq 1$. The map

$$(x, y) \in \bar{D} \mapsto (x, y, z(x, y))$$

where


$$z(x, y) = \sqrt{1 - x^2 - y^2}$$

parameterizes precisely the top half of the unit sphere, the upper hemisphere.

Let $\pi: S^2 \rightarrow \mathbb{RP}^2$ be the quotient map, taking each unit vector (x, y, z) to the line through that vector and the origin, i.e. quotienting by $(x, y, z) \sim -(x, y, z)$. Consider the map

$$f: (x, y) \in \bar{D} \mapsto \pi(x, y, z(x, y)).$$

This map is onto, as every line through the origin strikes a point of the upper hemisphere. But horizontal lines strike the equator twice at opposite points. So f is not injective; indeed f is injective precisely in the interior of the disk, and 2 to 1 along the boundary of the disk, taking (x, y) and $(-x, -y)$ on the unit circle to the same line: the one spanned by $(x, y, 0)$. Define an equivalence relation \sim : two points are equivalent if their images are the same under f . The map f descends to a homeomorphism $\bar{D}/\sim \cong \mathbb{RP}^2$, since \bar{D} is compact, and f is surjective, every point of \mathbb{RP}^2 lies in the interior of the image of \bar{D} .



Chapter 3

Fundamental Groups

Homotopy

Continuous maps $f, g: X \rightarrow Y$ between topological spaces X, Y are *homotopic* if there is a *homotopy* between them, i.e. a map $F: [0, 1] \times X \rightarrow Y$, denoted by $F_s(x)$ instead of $F(s, x)$, so that $F_0 = f$ and $F_1 = g$. If there is a subset $X_0 \subset X$ on which $f = g$, f is *homotopic to g relative to X_0* if there is a homotopy F between f and g so that $F_s(x) = f(x)$ for all $x \in X_0$. A map is *null homotopic* if it is homotopic to a constant map, i.e. a map whose image is a single point.

3.1 Prove that the relation of being homotopic relative to some set is an equivalence relation.

Given points $x_0, x_1 \in X$, a *path* from x_0 to x_1 is a continuous map $x: [0, 1] \rightarrow X$ so that $x(0) = x_0$ and $x(1) = x_1$, i.e. a map $x: ([0, 1], 0, 1) \rightarrow (X, x_0, x_1)$. We often omit to write the expression “relative to $\{0, 1\}$ ” when discussing homotopies of paths, and the reader will have to decide when we should have written that expression.

Intuitively, a homotopy between paths looks something like



3.2 Take two paths $x, y: [0, 1] \rightarrow X$ for one is a *reparameterisation*, of the other, i.e. there is a continuous map $\tau: [0, 1] \rightarrow [0, 1]$ so that $y \circ \tau = x$ with $\tau(0) = 0$ and $\tau(1) = 1$. Prove that x is homotopic to y relative to $\{0, 1\}$.

Lemma 3.1. *Cover a topological space X in open sets $X_a \subset X$. Take a path $x: [0, 1] \rightarrow X$. Then there are real numbers $0 = t_0 < t_1 < \dots < t_n = 1$ so that $x(t)$ stays in one open set X_{a_i} for $t_i \leq t \leq t_{i+1}$.*

Proof. Each point of $[0, 1]$ lies in an open interval lying entirely inside one $x^{-1}X_a$. Replace that open interval with a smaller open interval: each point of $[0, 1]$ lies in an open interval whose closure lies entirely inside one $x^{-1}X_a$.

Since $[0, 1]$ is compact, finitely many such open intervals cover $[0, 1]$. Take the endpoints of those intervals: $0 = t_0 < t_1 < \cdots < t_n = 1$. \square

Some inessential differential geometry

Think of Euclidean space of dimension n as having n coordinates:

$$x = (x_1, \dots, x_n).$$

An n -dimensional *manifold* is a subset of Euclidean space \mathbb{R}^{n+k} , locally expressed as the graph of k of the coordinates as smooth functions of the other n coordinates. An n -dimensional *manifold with corners* is a subset of Euclidean space \mathbb{R}^{n+k} , locally expressed as the graph of k of the coordinates as smooth functions of the other n coordinates, constrained to be inside an n -dimensional box. A *surface* is a 2-dimensional manifold (perhaps with corners). A continuous map of manifolds is *smooth* if it is smooth as a map of those coordinates. We won't prove:

Theorem 3.2. *Every continuous map of manifolds is homotopic to a smooth map.*

Lemma 3.3. *Any path $x: [0, 1] \rightarrow M$ in a manifold (perhaps with corners) M is homotopic relative to $\{0, 1\}$ to a smooth path.*

Proof. If $M = \mathbb{R}^n$, and $x: [0, 1] \rightarrow M$ is a path then let $y(t) = (1-t)x(0) + tx(1)$ and let $F(s, t) = (1-s)x(t) + sy(t)$. More generally, the same trick works for M any convex domain in \mathbb{R}^n , and in particular for M a box.

Suppose next that we have a path $x(t)$ in a box M , and we want to smooth out that path only inside some interval $a < t < b$. Take a smooth increasing function $h(t)$ equal to 0 in a small neighborhood of a , and equal to 1 in a small neighborhood of b . Let

$$y(t) = \begin{cases} x(t), & \text{if } 0 \leq t \leq a, \\ (1-h(t))x(a) + h(t)x(b), & \text{if } a \leq t \leq b, \\ x(t), & \text{if } b \leq t \leq 1. \end{cases}$$

and let

$$F(s, t) = (1-s)x(t) + sy(t).$$

Cover M in open sets, each a graph over a convex open set in a box in \mathbb{R}^n . Apply lemma 3.1 on the preceding page to split $x(t)$ into intervals on which it stays in these open sets. On each interval, we use the first trick to smooth x . This done, x is now piecewise smooth. We use the second trick near each corner to smooth out corners. \square


Loops

A *loop* is a path $x: [0, 1] \rightarrow X$ with $x(0) = x(1)$. Given a path x from x_0 to x_1 and a path y from x_1 to x_2 , define a path $x * y$ from x_0 to x_2 by

$$(x * y)(t) = \begin{cases} x(2t), & \text{if } 0 \leq t \leq \frac{1}{2}, \\ y(2t - 1), & \text{if } \frac{1}{2} \leq t \leq 1. \end{cases}$$

and define $\bar{x}(t) = x(1 - t)$. Also, for any point x_0 , define the null path $x(t) = x_0$ for $0 \leq t \leq 1$ which we just denote by x_0 .

If you have two paths and I have two paths, with homotopies fixing endpoints between your first and my first, and between your second and my second, then we can make a homotopy between paths glued together:



Hence gluing paths together commutes with homotopy relative to $\{0, 1\}$.

3.3 For any three paths x, y, z , if $(x * y) * z$ is defined then so is $x * (y * z)$, and vice versa, and they are homotopic relative to $\{0, 1\}$.

3.4 For any path x , the path $x * \bar{x}$ is homotopic to $x(0)$ relative to $\{0, 1\}$ while $\bar{x} * x$ is homotopic to $x(1)$ relative to $\{0, 1\}$.

Therefore, for any topological space X and point $x_0 \in X$, the homotopy classes of loops relative to $\{0, 1\}$ form a group, called the *fundamental group* of X and denoted $\pi_1(X, x_0)$.

3.5 If $x_0, x_1 \in X$ are two points connected by a path x , then any loop y at x_0 has an associated loop $\bar{x} * (y * x)$ at x_1 . Prove that the homotopy class of $\bar{x} * (y * x)$ in $\pi_1(X, x_0)$ depends only on the homotopy class of y in $\pi_1(X, x_1)$, and that this gives an isomorphism of groups

$$\pi_1(X, x_0) \rightarrow \pi_1(X, x_1).$$

A topological space X is *path connected* if any two points of X are the endpoints of a path in X . If X is path connected, then we usually ignore the point x_0 and write $\pi_1(X)$ instead of $\pi_1(X, x_0)$, even though this is not strictly speaking well defined. A path connected topological space X is *simply connected* if $\pi_1(X) = \{1\}$.

A *star shaped set* is a set $X \subset \mathbb{R}^n$ so that there is some point $x_0 \in X$ so that for every point $x_1 \in X$, the line segment from x_0 to x_1 lies entirely in X .



In other words, X admits the homotopy $F: (s, x) \in [0, 1] \times X \mapsto x_0 + s(x - x_0) \in X$. Clearly $\pi_1(X) = \{1\}$ is the trivial group, where 1 is the constant path x_0 . In particular, $\pi_1(\mathbb{R}^n) = \{1\}$.

For the circle S^1 , take any path $z(t) = x(t) + iy(t)$ on S^1 and write it as $z(t) = e^{i\theta(t)}$ using trigonometry to prove that one can continuously pick out a value of $\theta(t)$ for any continuous $z(t)$. The number

$$n(z) := \frac{1}{2\pi} (\theta(1) - \theta(0))$$

is an integer, since z is periodic. Suppose that two loops $z_0(t)$ and $z_1(t)$ have the same values of $n(z_0) = n(z_1)$ and start (and thus end) at the same point of S^1 , say at $e^{i\alpha_0}$. Pick continuous angles $\theta_0(t), \theta_1(t)$ with $\theta_0(0) = \theta_1(0) = \alpha_0$ and

$$z_0(t) = e^{i\theta_0(t)}, z_1(t) = e^{i\theta_1(t)}.$$

Clearly

$$\theta_0(1) = \theta_1(1).$$

Let

$$\theta_s(t) = (1 - s)\theta_0(t) + s\theta_1(t)$$

and

$$z_s(t) = e^{i\theta_s(t)},$$

a homotopy. Therefore $\pi_1(S^1) = \mathbb{Z}$, identifying the homotopy class of a loop z with the number $n(z)$.

For an annulus

$$A := \{ x \in \mathbb{R}^n \mid r_0 < \|x\| < r_1 \},$$

the map

$$F_s(x) = \frac{x}{(1-s) + s\|x\|}$$


retracts the annulus to the unit sphere, and retracts paths on the annulus to those on the sphere, homotopies on the annulus to homotopies on the sphere, so that $\pi_1(A) = \pi_1(S^{n-1})$.

It turns out to be more difficult to find the fundamental group of the plane punctured at two points, or at three points, etc.

The spaces \mathbb{R} and \mathbb{R}^2 are not homeomorphic, as \mathbb{R} becomes path disconnected when we remove any point, while \mathbb{R}^2 does not.

The spaces \mathbb{R}^2 and \mathbb{R}^3 are not homeomorphic, as \mathbb{R}^2 punctured at any point has fundamental group \mathbb{Z} , while \mathbb{R}^3 punctured at any point remains simply connected.

The Hawaiian earring has a very complicated and uncountable fundamental group.



The union of all circles with irrational radius passing through the origin and tangent to the vertical axis is even more complicated than the Hawaiian earring.

Lemma 3.4. *Any connected finite graph has finitely generated fundamental group.*

Proof. Take a connected finite graph X and a vertex $x_0 \in X$. A *tree* is a simply connected graph. A *maximal subtree* is a maximal simply connected subgraph; take one, say $T \subset X$. Then T contains all vertices, since otherwise we could add an edge that attaches one more vertex, without creating a loop (any loop would have two edges reaching that vertex). Each path in X is determined up to homotopy by listing the edges it passes through. Each path in T is uniquely

determined, up to homotopy, by its end points, since there is then a unique list of edges giving a path between the end points. For each edge e_a of X not in T , pick a loop $[x_a]$ in X starting at x_0 , and passing once along e_a . Every loop $[x]$ in X , starting and ending at x_0 , passes finitely many times through each e_a , either in the same direction or the opposite direction to $[x_a]$. Picture the last edge e_a which $[x]$ passes along, and picture direction in which it passes. We can suppose for simplicity that $[x_a]$ traverses e_a in the opposite direction. Consider the loop $[y] = [x_a] * [x]$. It goes through e_a , say through the two vertices x_b, x_c of e_a , and then passes through T to x_0 , and then back again to x_b and then x_c . By uniqueness of paths in T , up to homotopy, with given end points, we can arrange that $[y]$ takes the same route from x_c to x_0 and then back again. So $[y]$ is homotopic to cutting that part of $[y]$ out, i.e. up to homotopy $[x_a][x]$ has one fewer pass through e_a than $[x]$ did. By induction, we arrange that $[x]$ is a product of these various $[x_a]$. \square

Lemma 3.5. *Any graph with countably many vertices and edges has countable fundamental group.*

Proof. By compactness of $[0, 1]$, each loop in the graph can only hit finitely many vertices and so finitely many edges. Listing the vertices and edges of the loop in order gives the loop up to homotopy. \square

More inessential remarks on differential geometry

A *diffeomorphism* is a smooth of manifolds with smooth inverse. We will not prove:

Theorem 3.6 (Sard). *Take a smooth map $\varphi: P \rightarrow Q$ of manifolds. If P has smaller dimension than Q , then the image of φ is nowhere dense. If P and Q have equal dimension, there is a dense set of points $q_0 \in Q$ whose preimage $\varphi^{-1}\{q_0\}$ consists entirely of points $p_0 \in P$ near which φ is a local diffeomorphism.*

Careful: this point q_0 could have empty preimage, for example if φ maps all of P to a single point of Q .

For the spheres S^2, S^3, \dots , any path $x: [0, 1] \rightarrow S^n$ is smoothly approximated by some smooth path $y: [0, 1] \rightarrow \mathbb{R}^{n+1}$ homotopic to x , which we then divide by $\|y\|$ to get a smooth path homotopic to x lying in S^n . By Sard's theorem, y misses some point of the sphere. Recall that stereographic projection from that point identifies the rest of the sphere with \mathbb{R}^n , where we can use our previous result to homotope to a constant map: $\pi_1(S^n) = \{1\}$. Note that this doesn't work for the circle S^1 , where the application of Sard's theorem doesn't tell us anything.

A more difficult theorem, which we won't prove, but which may provide some comfort:

Theorem 3.7 (Whitney [4] p. 49 Theorem 2.6). *The smooth maps between any two manifolds are dense in the continuous maps, in the topology of uniform convergence on compact sets, and every continuous map is homotopic to a smooth map.*

Maps and fundamental groups

A *diagram* is a collection of maps between sets, drawn as a graph like:

$$\begin{array}{ccc} X & \xrightarrow{f} & Z \\ & \searrow g & \swarrow h \\ & Y & \end{array}$$

Start at one of the sets, and follow a path along the maps, in the direction of their arrows: compose those maps. The diagram *commutes* if any two paths with the same starting and ending points give the same composition. In our example, this means that $g = h \circ f$.

Lemma 3.8. *A continuous map $f: X \rightarrow Y$ between topological spaces yields a group morphism*

$$f_*: \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$$

where $y_0 = f(x_0)$, given by

$$f_*[x] = [f \circ x]$$

for any path $x: [0, 1] \rightarrow X$. The group morphism doesn't change if f varies through a homotopy of maps taking x_0 to y_0 . More generally, if we take homotopy f_t of f through a family of maps taking, say $x(t)$ to $y(t)$, for some paths $x(t), y(t) = f_t(x(t))$, then there is a commutative diagram between the maps f_{0*} and f_{1*} :

$$\begin{array}{ccc} \pi_1(X, x_0) & \xrightarrow{f_0} & \pi_1(Y, y_0) \\ \downarrow x_* & & \downarrow y_* \\ \pi_1(X, x_1) & \xrightarrow{f_1} & \pi_1(Y, y_1) \end{array}$$

Under composition of continuous maps

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

the group morphisms

$$\pi_1(X, x_0) \xrightarrow{f} \pi_1(Y, y_0) \xrightarrow{g} \pi_1(Z, z_0)$$

compose: $(g \circ f)_* = g_* \circ f_*$.

A *homotopy equivalence* is a continuous map $f: X \rightarrow Y$ between topological spaces so that there is a continuous map $g: Y \rightarrow X$ for which $f \circ g$ and $g \circ f$ are both homotopic to identity maps. A homotopy equivalence identifies fundamental groups.

The annulus in \mathbb{R}^n is homotopy equivalent to the unit sphere, by the map including the sphere into the annulus.

The sphere in \mathbb{R}^n punctured at one point is homotopy equivalent to \mathbb{R}^{n-1} by Ptolemaic projection, and so homotopy equivalent to a point.

The sphere in \mathbb{R}^n punctured at two points is homotopy equivalent to the annulus in \mathbb{R}^{n-1} by Ptolemaic projection, and so to the sphere in \mathbb{R}^{n-1} .

If $L \subset \mathbb{R}^3$ is a line, the topological space $X = \mathbb{R}^3 - L$ is homotopy equivalent to S^1 , by projecting to a plane perpendicular to L , punctured where the plane strikes L , and then taking a homotopy to a circle.

Theorem 3.9 (Fundamental theorem of algebra). *Every nonconstant polynomial function of one complex variable has a complex root.*

Proof. Take a nonconstant polynomial function

$$p(z) = a_0 + a_1z + \cdots + a_nz^n,$$

with $a_n \neq 0$. Clearly $p(z)$ has a root just where $p(z)/a_n$ has a root, so we can assume that $a_n = 1$. Let α be the maximum of $|a_0|, |a_1|, \dots, |a_{n-1}|$. If we pick any z with $|z| > (n-1)\alpha$ then clearly the leading term of $p(z)$ is larger than all other terms added together, so $p(z) \neq 0$. Rescale the z variable if needed to ensure that $(n-1)\alpha < 1$. So if $|z| \geq 1$ then $p(z) \neq 0$. By the same reasoning, none of the functions

$$p_t(z) = z^n + (1-t)(a_0 + a_1z + \cdots + a_{n-1}z^{n-1}),$$

vanishes as long as $|z| \geq 1$, a homotopy between $p_0(z) = p(z)$ and $p_1(z) = z^n$. Let

$$g_t(e^{i\theta}) = \frac{p_t(z)}{p_t(1)},$$

where $z = e^{i\theta}$, for $0 \leq t \leq 1$. This map is a homotopy between the loop

$$g_0(z) = \frac{p(z)}{p(1)}$$

with $z = e^{i\theta}$ and the loop

$$g_1(z) = z^n.$$

Note that $g_t(1)$ is fixed during the homotopy.

Let

$$f_t(z) = \frac{p(tz)}{p(t)},$$

where $z = e^{i\theta}$ and $0 \leq t \leq 1$. The map f is a homotopy from the trivial loop at $t = 0$ to a loop at $t = 1$:

$$f_1(z) = \frac{p(z)}{p(1)} = g_0(z),$$

for $z = e^{i\theta}$. So, inside the plane punctured at the origin, the trivial loop is homotopic to the loop winding n times and so $n = 0$. \square


3.6 Prove that the fundamental group of a product is the product of the fundamental groups:

$$\pi_1(X \times Y, (x_0, y_0)) \cong \pi_1(X, x_0) \times \pi_1(Y, y_0).$$

The torus $T^n = S^1 \times S^1 \times \cdots \times S^1$ has fundamental group

$$\pi_1(T) = \mathbb{Z}^n.$$

3.7 Give some explanation (not a rigorous proof) why the surface called *Alexander's horned sphere* has uncountable fundamental group, while the open subset of \mathbb{R}^3 inside (called *Alexander's horned ball*) is simply connected.



Alexander's horned sphere, Krzysztof Rykaczewski

Corollary 3.10. *If a path connected topological space X admits a countable basis of simply connected open sets, then its fundamental group is countable.*

Proof. The homotopy class of any path is determined by listing off finitely many simply connected open sets that cover it, in the order that it enters them, as in lemma 3.1 on page 29. \square

Corollary 3.11. *Every compact and locally simply connected topological space X has finitely generated fundamental group.*

Proof. By compactness, we can find a finite covering by simply connected open sets X_a and cover the overlaps $X_a \cap X_b$ by finitely many path connected open sets X'_c . The homotopy class of any path is determined by listing off finitely many simply connected open sets that cover it, in the order that it enters them, as in lemma 3.1 on page 29. The homotopy class of any path is determined by listing off finitely many X_a and X'_c that cover it, in the order it enters them, as in lemma 3.1 on page 29. \square


Chapter 4

Covering Spaces

Covering maps

A continuous map $f: X \rightarrow Y$ of topological spaces *evenly covers* an open set $U_Y \subset Y$ if $f^{-1}U_Y \subset X$ is a disjoint union of open sets mapped homeomorphically to U_Y by f , called *sheets*. The map $f: X \rightarrow Y$ is a *covering map* if Y has an open cover by evenly covered open sets. A *covering space* of a topological space Y is a topological space X equipped with a covering map $f: X \rightarrow Y$. Covering spaces are the main tool to calculate fundamental groups.

The map $\theta \in \mathbb{R} \mapsto (\cos \theta, \sin \theta) \in S^1$ is a covering map. Every open set of S^1 which is not all of S^1 is evenly covered: use trigonometry to write out an angle θ for each point of your open set, well defined and unique up to adding 2π multiples. The sheets correspond to the 2π multiples.



The map $e^{i\theta} \in S^1 \mapsto e^{in\theta} \in S^1$ is an n -sheeted covering.

The map $x \in \mathbb{R}^n \mapsto (e^{ix_1}, e^{ix_2}, \dots, e^{ix_n})$ is a covering map of the n -dimensional torus $T^n := S^1 \times S^1 \times \dots \times S^1$.


The map $x \in S^2 \mapsto [x] \in \mathbb{RP}^2$ taking a point to the line through that point is a 2-sheeted covering of the projective plane.

Take a manifold M . Let \hat{M} be the set of pairs (m, o) where $m \in M$ and o is an orientation of $T_m M$, i.e. a choice of oriented basis up to equivalence, with two bases equivalent if the change of basis matrix between them has positive determinant. Then $\hat{M} \rightarrow M$ is a 2-1 covering

map. If M is orientable, then \hat{M} is a disjoint union of two copies of M and the covering map is a homeomorphism on each copy. If M is not orientable, then \hat{M} is connected.

4.1 Let X be the open interval $(0,3) \subset \mathbb{R}$. Let Y be the unit circle in the complex plane. Prove that the map $f: X \rightarrow Y$ given by $f(x) = e^{2\pi i x}$ is not a covering map.

In the complex plane \mathbb{C} , the map $z \in \mathbb{C} - \{0\} \mapsto z^3 \in \mathbb{C} - \{0\}$ is a covering map. We like to pretend that we can draw every covering map as a picture of “sheets”, but in this case the best picture we get is an immersed surface.



4.2 Prove that the number n of sheets (which might be ∞) above an evenly covered open set is constant along any path in Y . In particular, if Y is path connected, this number n is constant, and we say that the covering map is n to 1.

The *fiber* of a map $f: X \rightarrow Y$ over a point $y_0 \in Y$ is the set $f^{-1}\{y_0\}$, usually denoted X_{y_0} .

4.3 Prove that $X_{y_0} \subset X$ is discrete, i.e. has the discrete topology as a subset of X .

4.4 Suppose that $f: X \rightarrow Y$ is a covering map. Prove X is Hausdorff if and only if Y is Hausdorff.

4.5 Prove that every proper local diffeomorphism $f: P \rightarrow Q$ between manifolds without boundary, with Q connected, is a covering map.

Theorem 4.1 (Fundamental theorem of algebra). *Suppose that k is a field containing \mathbb{R} and of finite dimension as a real vector space. Then $k = \mathbb{R}$ or $k = \mathbb{C}$, up to isomorphism. In particular, the splitting field of any real or complex polynomial is \mathbb{R} or \mathbb{C} , i.e. every complex polynomial in one variable splits into a product of linear factors over \mathbb{C} .*

Proof. Pick any inner product on the real vector space k . Take the unit sphere $S \subset k$. The map $f: x \in S \mapsto x^2/|x^2| \in S$ is smooth. Compute that

$$f'(x)y = \frac{2x}{|x^2|} (y - \langle f(x), xy \rangle x).$$

In particular, if $f'(x)y = 0$ then y is a scalar multiple of x . So for y tangent to S , $f'(x)y = 0$ just when $y = 0$, i.e. f is a local diffeomorphism. The sphere is compact, so $f: S \rightarrow S$ is a covering map. If $f(x) = f(y)$, pick a real number $\lambda > 0$ so that $\lambda^2 = |x^2|/|y^2|$. Check that $(x - \lambda y)(x + \lambda y) = 0$, so $x = \pm \lambda y$, i.e. up to scaling $x = \pm y$, so $f: S \rightarrow S$ is 2-1. But the sphere of dimension 2 or more is simply connected, so does not admit a covering map of degree 2 from a connected space. So S is the 0-dimensional or 1-dimensional sphere, i.e. $k = \mathbb{R}$ or k is a 2-dimensional real algebra. Suppose that k is 2-dimensional. Pick some $y \in k$ nonzero. Then $y/|y|$ lies in the image of f , i.e., for some x with $|x| = 1$,

$$\frac{x^2}{|x^2|} = \frac{y}{|y|}.$$

So if we let

$$\lambda = \sqrt{\frac{|y|}{|x^2|}},$$

and replace x by λx , we find $x^2 = y$. So every nonzero element of k has a square root, and so in particular, $-1 \in \mathbb{R}$ has a square root in k , so k contains \mathbb{C} , and hence $k = \mathbb{C}$. Note that we could pick k to be the splitting field of any real or complex polynomial. \square

Quotients by group actions

An *action* of a group Γ on a topological space X is a map associating to each $g \in \Gamma$ a continuous map $X \rightarrow X$ denoted $x \mapsto gx$ or sometimes denoted $x \mapsto g \cdot x$, so that $g(hx) = (gh)x$ for any $g, h \in \Gamma$ and $x \in X$ and so that $1x = x$ for any $x \in X$. The action is *free* if for any $x \in X$, the only $g \in \Gamma$ for which $gx = x$ is $g = 1$. The action is a *covering action* if any $x \in X$ lies in an open set $U \subset X$ so that the only $g \in \Gamma$ for which gU intersects U is $g = 1$. A group action is *proper* when any points $x, y \in X$ lie in open sets U_x, U_y so that gU_x intersects U_y for only finitely many $g \in \Gamma$. The *orbit* of a point $x \in X$ is the set $\Gamma x := \{gx \mid g \in \Gamma\}$. The *quotient space* X/Γ of the group action is the set of orbits, with the *quotient map* $x \in X \mapsto \Gamma x \in X/\Gamma$.

4.6 Suppose that X is a metric space and that Γ acts on X by isometries. Prove that Γ acts on X as a covering action if and only if the action is free with discrete orbits.

4.7 Take an invertible matrix A with at least one eigenvalue λ satisfying $\lambda > 1$ and at least one eigenvalue μ satisfying $0 < \mu < 1$. Let $M := \mathbb{R}^n - \{0\}$ and let $\Gamma := \{A^n \mid n \in \mathbb{Z}\}$. Show that the action of Γ on M has discrete orbits, but the quotient space is not Hausdorff.

4.8 If a group Γ acts on a topological space X and X contains a compact set intersecting every Γ -orbit, then \bar{X} is also compact.

Theorem 4.2. *Take a group Γ acting on a topological space X . The quotient map $X \rightarrow \bar{X} = X/\Gamma$ is a covering map just when the action is a covering action.*

Proof. For each point $x \in X$ pick some open set $U_x \subset X$ containing x so that gU_x doesn't intersect U_x for any $g \in \Gamma$ unless $g = 1$. The image $\bar{U}_x \subset \bar{X}$ has preimage in X precisely the union of the nonoverlapping translates gU_x for $g \in \Gamma$. The open sets in \bar{U}_x have preimages precisely the Γ -invariant open sets in the translates of U_x , precisely one of which lies in U_x . Hence the quotient map restricted to U_x is a homeomorphism $U_x \rightarrow \bar{U}_x$. Each element of Γ interchanges the sheets. Conversely, if the quotient map is a covering map, then evenly covered open sets have preimages precisely open sets U not intersecting their translates gU . \square

Theorem 4.3. *Take an action of a group Γ on a Hausdorff space X . The quotient space is Hausdorff just when any two points of X lie in disjoint Γ -invariant open sets.*

Proof. Take two points $\bar{x} \neq \bar{y} \in \bar{X}$ in the quotient space $\bar{X} := X/\Gamma$. Take points $x, y \in X$ mapping to them. If \bar{x}, \bar{y} lie in disjoint open sets $\bar{U}_{\bar{x}}, \bar{U}_{\bar{y}}$, then the preimages of these open sets are disjoint Γ -invariant open sets around x and y .

Conversely, take any disjoint Γ -invariant open sets around x and y , U_x, U_y . Let \bar{U}_x, \bar{U}_y be their images in \bar{X} . By Γ -invariance, the preimage of \bar{U}_x is U_x so \bar{U}_x is open and $\bar{U}_x \cap \bar{U}_y$ has preimage $U_x \cap U_y$ empty so is empty. \square

Theorem 4.4. *Any proper group action on any Hausdorff space has Hausdorff quotient space.*

Proof. Surround points $x, y \in X$ with open sets U_x, U_y so that gU_x is disjoint from U_y except for finitely many $g \in \Gamma$, say g_1, g_2, \dots, g_n . Because X is Hausdorff, for each j we can take “houses”: disjoint open sets V_j, W_j so that $V_j \subset g_j U_x$ and $W_j \subset U_y$ and $g_j x_j \in U_j$ and $y \in W_j$. Replace U_x with

$$\bigcap_j g_j^{-1} U_j$$

and U_y with

$$\bigcap_j W_j.$$

The resulting sets U_x and U_y have all translates disjoint, i.e. x and y lie in disjoint invariant open sets. \square

A group Γ acting on a metric space X acts *by isometries* if $x \mapsto gx$ is an isometry for all $g \in \Gamma$, i.e. $d(gx, gy) = d(x, y)$ for any points x, y and $g \in \Gamma$.

Theorem 4.5. *Take a group action on a metric space X by a group of isometries Γ . The following are equivalent:*

- The orbits are closed.*
- Any two points of X lie in disjoint Γ -invariant open sets.*
- The quotient space is a metric space, under the quotient metric*

$$d(\bar{x}, \bar{y}) = \inf_{g \in \Gamma} d(gx, y),$$

so that the metric space topology agrees with the quotient topology.

Proof. The orbits are closed just when any point in any orbit lies at a positive minimum distance from some point in any other chosen orbit, which occurs just when the “quotient metric” expression d is a metric.

Suppose that the orbits are closed. Then the balls around distinct points $\bar{x}, \bar{y} \in \bar{X}$ have as preimages in X some disjoint Γ -invariant open sets.

Suppose that any two points $x \neq y \in X$ lie in disjoint Γ -invariant open sets U_x, U_y . It is clear that the expression d above is continuous on $\bar{X} \times \bar{X}$, since any open set of real numbers has open preimage inside $X \times X$. Next we want to prove that $d(\bar{x}, \bar{y}) = 0$ just when $\bar{x} = \bar{y}$. If $d(g_j x, y) \rightarrow 0$ for some $g_j \in \Gamma$, then $g_j x$ enters U_y for all but finitely many j , and so $U_x \cap U_y$ is not empty.

So d is a metric on \bar{X} . If $x \in X$ maps to $\bar{x} \in \bar{X}$ then the ball of radius r around each point \bar{x} has preimage precisely the set of points $y \in X$ so that $d(gy, x) < r$ for some $g \in \Gamma$, i.e. just exactly the union of Γ -translates of the ball of radius r around x . Take an open set $\bar{W} \subset \bar{X}$ around \bar{x} . Let $W \subset X$ be its preimage. Pick a radius r small enough that the ball of radius r around x lies inside W . Then so do all Γ -translates. So the ball of radius r around \bar{x} lies in \bar{W} . \square

A *locally isometric covering map* is a map of metric spaces $X \rightarrow Y$ so that every point of Y lies in an open set \bar{U} evenly covered, say by disjoint open sets $U_\alpha \subset X$ for which $U_\alpha \rightarrow \bar{U}$ is an isometry.

Theorem 4.6. *Take a free group action on a metric space X by a group of isometries Γ with discrete orbits. Then the quotient map $X \rightarrow X/\Gamma$ is a locally isometric covering map.*

Proof. The quotient is a metric space by theorem 4.5 on the preceding page. Take a point $x \in X$ mapping to a point $\bar{x} \in \bar{X}$ and a ball $B \subset \bar{X}$ about \bar{x} of radius r small enough that it is evenly covered by balls $B_\alpha \rightarrow B$ so that gB_α doesn't intersect B_α unless $g = 1$. Let $B' \subset B$ be the ball of radius $r/2$ about \bar{x} . Pick points $y, z \in X$ in the same sheet B'_α mapping to points \bar{y}, \bar{z} in B' . The orbits are closed, so we can pick a point gz to be the closest point to y in the orbit Γz . So $d(gz, y) \leq d(z, y) < r/2$ so $d(gz, x) < r$. But then gB_α intersects B_α . So $g = 1$. Hence z is the closest point to y mapping to \bar{z} . Therefore $d(z, y) = d(\bar{z}, \bar{y})$, for all $z, y \in B'_\alpha$. \square

The torus $T^n = \mathbb{R}^n / \mathbb{Z}^n$ is the quotient of the \mathbb{R}^n by the free action of \mathbb{Z}^n with discrete orbits. Hence the torus is a metric space in the quotient topology, and $\mathbb{R}^n \rightarrow T^n$ is a locally isometric covering map, as are all of the following examples.

The real projective plane \mathbb{RP}^2 is the quotient of the unit sphere $S^2 \subset \mathbb{R}^3$ by the covering action of $\Gamma = \{\pm 1\}$.

The *Möbius strip* is the quotient \mathbb{R}^2 / Γ by the covering action of rigid motions Γ of the plane generated by the transformation

$$(x, y) \mapsto (x + 1, -y).$$

It is covered by the cylinder, given as the quotient \mathbb{R}^2 / Γ_0 by the subgroup $\Gamma_0 \subset \Gamma$ generated by

$$(x, y) \mapsto (x + 2, y).$$




The *Klein bottle* is the quotient \mathbb{R}^2/Γ by the covering action of rigid motions Γ of the plane generated by the two transformations

$$(x, y) \mapsto (x, y + 1)$$

and

$$(x, y) \mapsto (x + 1, -y).$$

It is clearly covered by the Möbius strip, and by the cylinder.



Lifting

Lemma 4.7. *Take a covering map $f: X \rightarrow Y$ from a Hausdorff space X . Take a path connected space Z and continuous maps $g_1, g_2: Z \rightarrow X$ so that $f \circ g_1 = f \circ g_2$. If $g_1(z_0) = g_2(z_0)$ for some point $z_0 \in Z$ then $g_1 = g_2$.*

Proof. Suppose that $g_1(z) = g_2(z)$ at some point $z \in Z$. Let $x = g_1(z)$ and $y = f(z)$. Take an open set $U_Y \subset Y$ around y so that $f^{-1}U_Y$ is a disjoint union of sheets. Let U_X be the sheet containing x . Let

$$\tilde{f} := f|_{U_X}.$$

Let

$$U_Z := g_1^{-1}U_X \cap g_2^{-1}U_X.$$

Clearly $z \in U_Z$. Moreover

$$f \circ g_1 = f \circ g_2$$

implies that on U_Z ,

$$\tilde{f} \circ g_1 = \tilde{f} \circ g_2.$$

But \tilde{f} is a diffeomorphism, so on U_Z , $g_1 = g_2$.

The set E of points $z \in Z$ at which $g_1 = g_2$ is not empty because z_0 lies in E . But E is open, since it contains an open set U_Z around each of its points as above. But E is also closed, because X is Hausdorff. Any path in Z starting in E stays in E , because the set of points at which it lies in E is both an open and a closed subset of $[0, 1]$. Since Z is path connected, E is all of Z . \square

Proposition 4.8. *Take a covering map $f: X \rightarrow Y$ from a Hausdorff space X . Take a path $y: [0, 1] \rightarrow Y$, and a point $x_0 \in X$ so that $f(x_0) = y(0)$. There is a unique path $x: [0, 1] \rightarrow X$ so that $f \circ x = y$ and $x(0) = x_0$, the lift of the path y .*

Proof. Uniqueness follows from lemma 4.7 on the previous page. Near each point $y \in Y$ there is an evenly covered open set $U_Y \subset Y$, i.e. so that $f^{-1}U_Y$ is a disjoint union of open sets (the “sheets”), each mapped homeomorphically to U_Y by f . For example, picking y to be y_0 , we find x_0 on precisely one of these sheets, call it U_X , and we define

$$x(t) = f|_{U_X}^{-1}(y(t))$$

on the largest interval $0 \leq t < \tau$ for which $y(t)$ stays inside U_Y .

Cover $[0, 1]$ by open intervals I_a so that $y(t)$ stays inside an evenly covered set U_a on each open interval I_a . By compactness, extract a finite cover, and therefore a finite collection of intervals $0 \leq t \leq a_1, a_1 \leq t \leq a_2, \dots, a_{n-1} \leq t \leq 1$, so that on each one $y(t)$ stays inside an evenly covered open set U_1, U_2, \dots, U_n . Lift up $y(t)$ by inverting f over each U_i one at a time. \square

Proposition 4.9. *Suppose that $f: X \rightarrow Y$ is a covering map from a Hausdorff space, and $F: [0, 1] \times Z \rightarrow Y$ is a continuous map, written $F_s(z) := F(s, z)$, and that there is a continuous map $\hat{F}_0: Z \rightarrow X$ so that $f \circ \hat{F}_0 = F_0$. Then there is a unique map $\hat{F}: [0, 1] \times Z \rightarrow X$ so that $f \circ \hat{F} = F$ and $\hat{F}(0, z) = \hat{F}_0(z)$ for all $z \in Z$, the lift of the map F .*

Proof. For each $z_0 \in Z$, lift the path $F(s, z_0) \in Y$ to a path $\hat{F}(s, z_0)$, and this defines $\hat{F}(s, z_0)$. We need to prove that \hat{F} is continuous. This is a purely local problem, so it suffices to prove for a trivial covering map, a homeomorphism, for which it is obvious. \square

Lemma 4.10. *The morphism of fundamental groups $f_*: \pi_1(X) \rightarrow \pi_1(Y)$ of a covering map $f: X \rightarrow Y$ on a Hausdorff space X is injective. Its image is the set of loops in Y (modulo homotopy) which lift to loops in X .*

Proof. The kernel of f_* consists of the homotopy classes of the loops $x: [0, 1] \rightarrow X$ so that $f \circ x: [0, 1] \rightarrow Y$ is homotopic to the trivial loop. Lift that homotopy up to a homotopy of x to the trivial loop. Given a loop y with homotopy class in the image, x is its lift, a loop. \square

Take a covering $f: X \rightarrow Y$ with points $x_0 \in X$ and $y_0 := f(x_0) \in Y$. Denote the fiber as $X_{y_0} := f^{-1}\{y_0\}$. Given any loop ℓ based at y_0 , let $\hat{\ell}$ be its lift, a curve in X starting at x_0 . Suppose that we replace ℓ by a homotopy of loops ℓ_t based at y_0 . That homotopy lifts to a homotopy of curves $\hat{\ell}_t$, all starting at x_0 . Since ℓ_t ends at y_0 , all of those curves $\hat{\ell}_t$ end at points of X_{y_0} . Looking at our picture of sheets, we see that X_{y_0} is a discrete set, i.e. any continuous map to X_{y_0} is constant. By continuity, the map $t \mapsto \hat{\ell}_t(1)$ is constant. So the endpoint $\hat{\ell}(1)$ doesn't change if we move ℓ through a homotopy. Therefore we have a well defined map

$$\pi_1(Y, y_0) \mapsto X_{y_0}.$$

Take any loop x based at x_0 and consider the associated loop $y = f \circ x$ based at y_0 . The loop $y * \ell$ starts at y_0 and its lift is $x * \hat{\ell}$, with the same endpoints as $\hat{\ell}$. Therefore our map descends to a map

$$\pi_1(Y, y_0) / f_* \pi_1(X, x_0) \mapsto X_{y_0}.$$

Lemma 4.11. *If X and Y are path connected Hausdorff topological spaces and $f: X \rightarrow Y$ is a covering map, then the endpoint map*

$$\pi_1(Y, y_0) / f_* \pi_1(X, x_0) \mapsto X_{y_0}$$

is bijective.

Proof. Take any two points $x_0, x_1 \in X_{y_0}$ and connected a path x . Then the loop $y = f \circ x$ lifts to x , so the map takes y to x_1 . \square

Proposition 4.12. *Take a covering map $f: X \rightarrow Y$ from a Hausdorff space and a map $g: Z \rightarrow Y$ from a path connected and locally path connected topological space Z ,*

$$\begin{array}{ccc} & X & \\ & \downarrow & \\ Z & \longrightarrow & Y \end{array}$$

and points $z_0 \in Z$, $x_0 \in X$, $y_0 \in Y$ so that $y_0 = f(x_0) = g(z_0)$. Then there is a unique lift $\hat{g}: Z \rightarrow X$ so that

$$\begin{array}{ccc} & X & \\ & \downarrow & \\ Z & \xrightarrow{\hat{g}} & Y \end{array}$$

i.e. a map so that $f \circ \hat{g} = g$ and $\hat{g}(z_0) = x_0$, if and only if

$$g_* \pi_1(Z, z_0) \subset f_* \pi_1(X, x_0).$$

Proof. If \hat{g} exists then $g_* = f_* \circ \hat{g}_*$ so the image of g_* lies in the image of f_* . Suppose that the image of g_* lies in the image of f_* . Take a path z in Z starting at z_0 . Map it a path y in Y and lift to a path x in X . Then define $\hat{g}(z) = x(1)$. This is a map $z \mapsto \hat{g}(z)$ defined on paths. Lifting homotopy of paths, $\hat{g}(z)$ is clearly defined on homotopy classes of paths: $\hat{g}([z])$. Take two paths z, z' with the same endpoints in Z and let y, y' and x, x' be the corresponding paths in Y and lifts to X . Then $[z]^{-1}[z']$ is a loop in Z , mapping to the loop $[y]^{-1}[y']$ in Y , which lifts to a loop in X . By uniqueness of lifts, this loop is $[x]^{-1}[x']$. In other words, x and x' have the same endpoint, so $\hat{g}([z]) = \hat{g}([z'])$ is that endpoint. So $\hat{g}([z])$ depends only on the endpoint $z_1 = z(1)$ of a path z : $\hat{g}(z_1)$, i.e. $\hat{g}: Z \rightarrow X$. Invert the covering map f locally, in some evenly covered open set in Y . Pick a simply connected open set in Z mapping to that open set in Y . Then paths in that simply connected open set will stay in the domain of the local inverse of f . We see that \hat{g} and g are locally identified by that local inverse of f , so \hat{g} is continuous. \square

4.9 Suppose that $Z \subset \mathbb{C}$ is a domain in the complex plane and that $g: Z \rightarrow \mathbb{C}$ is a complex analytic function defined in Z . A *logarithm* for $g(z)$ is a complex analytic function $G: Z \rightarrow \mathbb{C}$ so that $g(z) = e^{G(z)}$. Prove that $g(z)$ has a logarithm $G(z)$ just when both of the following conditions are satisfied:

- a. $g(z) \neq 0$ for any $z \in Z$ and
- b. g takes every loop in Z to a null homotopic loop in $\mathbb{C} - \{0\}$.

Suppose that $f: X \rightarrow Y$ is a covering space over a path connected space Y . Take two points $y_0, y_1 \in Y$. Take a path $y(t) \in Y$ from y_0 to y_1 . For each point $x_0 \in X_{y_0}$, let $x_{x_0}(t)$ be the lift of $y(t)$ that satisfies $x_{x_0}(0) = x_0$. Let $h: [0, 1] \times X_{y_0} \rightarrow X$, $h_t(x_0) := x_{x_0}(t)$. Since the lift $x_{x_0}(t)$ is uniquely determined by the choice of x_0 , this map h is well defined. By problem 4.3 on page 41, every fiber X_{y_0} has the discrete topology as a subset of X . Hence h is continuous simply because each path $x_{x_0}(t)$ is continuous. The map $h_t: X_{y_0} \rightarrow X_{y_1}$ has inverse given by following the path $y(t)$ backwards and constructing the same sort of map as h . In particular, the fibers X_{y_0} and X_{y_1} are homeomorphic via $h_1: X_{y_0} \rightarrow X_{y_1}$, the *monodromy* of the path $y(t)$.

By the homotopy lifting lemma, the monodromy depends only on the homotopy class of $y(t)$, giving a group morphism from $\pi_1(Y, y_0)$ to the group of permutations of the elements of X_{y_0} , the *monodromy morphism*.

4.10 Let X be the unit sphere in \mathbb{R}^3 , let Y be the projective plane, and let $f: X \rightarrow Y$ be the usual covering map: $f(x)$ being the line through the origin passing through x . Take the path $x(t) = (\cos \pi t, \sin \pi t, 0) \in X$ and let $y(t) = f(x(t))$. Explain why $y(t)$ is a loop and calculate its monodromy morphism.

4.11 Take a path connected Hausdorff space X with covering action of a group Γ and let $\bar{X} = \Gamma \backslash X$. Explain how to find the monodromy of the covering map $x \in X \mapsto \Gamma x \in \bar{X}$ over any loop in \bar{X} .

4.12 Prove that every continuous map $X \rightarrow Y$ from the real projective plane X to the 2-dimensional torus Y is null homotopic.

The universal covering space

A *universal covering space* of a topological space Y is a simply connected covering space X of Y . The map $X \rightarrow Y$ is a *universal covering map*.

The map $\theta \in \mathbb{R} \mapsto e^{i\theta} \in S^1$ is a universal covering map.

The map $x \in S^2 \rightarrow [x] \in \mathbb{RP}^2$, taking a unit vector x to the line through 0 and x , is a universal covering map.

A *morphism* of covering spaces $X \rightarrow Y$ and $Z \rightarrow Y$ is a continuous map $X \rightarrow Z$ making a commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Z \\ & \searrow & \swarrow \\ & Y & \end{array}$$

4.13 Prove that $X \rightarrow Z$ is then also a covering map.

4.14 Suppose that Y is a Hausdorff topological space which admits a universal covering space. Prove that a covering map $(X, x_0) \rightarrow (Y, y_0)$ is universal just when every other covering map $(Z, z_0) \rightarrow (Y, y_0)$ has a unique morphism $(X, x_0) \rightarrow (Z, z_0)$.

An *isomorphism* is a morphism with an inverse morphism.

4.15 Suppose that $X_1 \rightarrow X$ and $X_2 \rightarrow X$ are universal covering spaces of a Hausdorff space X . Pick points $x_1 \in X_1$ and $x_2 \in X_2$ mapping to the same point $x_0 \in X$. Prove that there is a unique isomorphism taking x_1 to x_2 .

Theorem 4.13. *Every path connected and locally simply connected topological space X has a universal covering space $\tilde{X} \rightarrow X$.*

Proof. Pick a point $x_0 \in X$. Let \tilde{X} be the set of all paths starting at x_0 , modulo homotopy fixing endpoints. Let $\tilde{x}_0 = [1_{x_0}]$ be the homotopy class of the trivial loop. Map $p: [x] \in \tilde{X} \mapsto x(1) \in X$, a surjective map. The open sets $U \subset X$ which are simply connected form a basis of open sets on X . To such a set U and a path x from x_0 to a $x_1 \in U$, with homotopy class $[x]$ with fixed endpoints, we associate the set

$$U_{[x]} := \{ [x * y] \mid y \text{ is a path in } U \text{ and } y(0) = x(1) \} \subset \tilde{X}.$$

Note that $[x] \in U_{[x]}$, so these sets $U_{[x]}$ cover \tilde{X} . We take these sets $U_{[x]}$ be a basis of open sets in \tilde{X} . Note that

$$p([x * y]) = y(1) \in U,$$

so that

$$p(U_{[x]}) = U.$$

Since U is simply connected, any two paths y, z from $x(1)$ to $y(1) = z(1)$ are homotopic, so $[x * y] = [x * z]$, i.e. p is injective on $U_{[x]}$. So p is a bijection

$$p: U_{[x]} \rightarrow U.$$

If $[z] \in U_{[x]}$ then $[z] = [x * y]$ for some y . But then $[x] = [\bar{y} * z]$, so

$$[x] \in U_{[z]}.$$

Take any two such open sets $U_{[x]}, V_{[y]}$, containing some common point $[z]$, we see than that

$$U_{[x]} = U_{[z]}$$

and

$$V_{[y]} = V_{[z]}.$$

Take an open subset $W \subset U \cap V$, connected and simply connected. Then

$$W_{[z]} \subset U_{[x]} \cap V_{[y]}.$$

Therefore the sets $U_{[x]}$ form the basis of a topology: their unions are closed under finite intersections. Note that

$$p^{-1}U = \bigcup_{[x]} U_{[x]}$$

where the union is over all paths x from x_0 to a point of U . The map $p: \tilde{X} \rightarrow X$ is therefore continuous.

We want to show that the map

$$p|_{U_{[x]}}: U_{[x]} \rightarrow U$$

has a continuous inverse. This is a bijection, so has an inverse, call it q . We need only check that, for any open set of $U_{[x]}$, the inverse image via q is also open. In other words, we need only check that, for any open set of $U_{[x]}$, the image via p is also open. Since the various $\{W_{[y]}\}$ open sets form a basis for our topology, it is enough to check that, for any simply connected open set $W \subset U$, and path y from x_0 to a point of W , the image

$$pW_{[y]} \subset U$$

is open. But this is just exactly W , since W is path connected, so every point of W is the endpoint of a path in W .

Given any path $x(t)$ starting at x_0 , let

$$x_s(t) = \begin{cases} x(t), & \text{if } 0 \leq t \leq s, \\ x(s), & \text{if } s \leq t \leq 1. \end{cases}$$

The homotopy class $[x_t]$ is a path $\tilde{x}(t) = [x_t]$ in \tilde{X} from \tilde{x}_0 to $[x_1]$. So \tilde{X} is path connected.

Take a loop $x(t)$ in X starting and ending at x_0 . The loop lifts to a path $\tilde{x}(t)$ as above. By definition, the path $\tilde{x}(t)$ is a loop just when $\tilde{x}(1) = \tilde{x}(0) = [1_{x_0}]$. But $\tilde{x}(1) = [x_1] = [x]$ is the homotopy class of x , so the lift is a loop just when the original loop is null homotopic, in which case the lift is null homotopic. \square

4.16 Suppose that $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a continuously differentiable map, that $f'(x)$ is an invertible matrix for every $x \in \mathbb{R}^n$, and that, for any sequence x_1, x_2, \dots with $\|x_i\| \rightarrow \infty$, $\|f(x_i)\| \rightarrow \infty$. Prove that f is a diffeomorphism.

The fundamental group $\pi_1(X, x_0)$ acts on the universal covering space \tilde{X} by the action $[x][y] = [x * y]$, which is clearly continuous. The covering map $\tilde{X} \rightarrow X$ is invariant under the action.

Lemma 4.14. *Take a path connected and locally simply-connected topological space X . The action of the fundamental group on the universal covering space is a covering action.*

Proof. We use the notation of theorem 4.13 on page 49. Take two points $[x_1], [x_2] \in \tilde{X}$. If the paths $x_1(t), x_2(t)$ have distinct endpoints, say $x_1(1) \neq x_2(1)$, then pick disjoint open sets U, W around them, and then $U_{[x_1]}$ does not intersect $W_{[x_2]}$. If the paths have the same endpoint, take a simply connected open set U containing that endpoint: $x_1(1) = x_2(1) \in U$. If the open sets $U_{[x_1]}$ and $U_{[x_2]}$ are not disjoint, then they are equal (as we saw in the proof of theorem 4.13). The homeomorphism $p: U_{[x_1]} \rightarrow U$ over the evenly covered open sets U then identifies the points $[x_1], [x_2] \in \tilde{X}$, so the points of \tilde{X} are not distinct. \square

4.17 Prove that the only topological spaces with \mathbb{R} as a covering space are \mathbb{R} and S^1 .

Lemma 4.15. *Take a path connected and locally simply connected space X . Every subgroup $\Gamma \subset \pi_1(X)$ arises as the image of the fundamental group of a connected covering space $X_\Gamma \rightarrow X$. Any connected covering space $Z \rightarrow X$ whose fundamental group has image Γ is isomorphic to $X_\Gamma \rightarrow X$.*

Proof. Let $X_\Gamma := \tilde{X}/\Gamma$, with the topology generated by declaring open sets to be the preimages of the open sets of X under the quotient map. So $\tilde{X} \rightarrow X_\Gamma \rightarrow X$ are continuous maps, and indeed covering maps. Take a connected covering space $f: Z \rightarrow X$ for which $\pi_1(Z)$ maps to $f_*\pi_1(Z) = \Gamma$. By proposition 4.12 on page 47, the map $f: Z \rightarrow X$ lifts to a map $\hat{f}: Z \rightarrow X_\Gamma$, and the covering map $p: X_\Gamma \rightarrow X$ lifts to a map $\hat{p}: X_\Gamma \rightarrow Z$. We leave the reader to check that these maps are inverses of one another. \square

Corollary 4.16. *If a group Γ has a covering action on a simply connected and locally simply connected Hausdorff topological space X , then the quotient has fundamental group*

$$\pi_1(\Gamma \backslash X) = \Gamma$$

and the quotient map $X \rightarrow \Gamma \backslash X$ is the universal covering map.

The fundamental group of the Klein bottle is the nonabelian group generated by the two transformations

$$(x, y) \mapsto (x, y + 1)$$

and

$$(x, y) \mapsto (x + 1, -y)$$

of the plane. Picture how this acts on the 1×1 squares with integer corners. Since the (x, y) -plane is the universal covering space of the Klein bottle, the preimage of each point is an orbit of these transformations. Each element of the group is identified with such a point, so the group is identified with the integer points of the plane, but with a tricky group operation.

The fundamental group of the torus $\mathbb{R}^2/\mathbb{Z}^2$ is \mathbb{Z}^2 , abelian. Hence the Klein bottle is not homeomorphic to the torus.

The fundamental group of \mathbb{RP}^n is ± 1 , from the covering map $S^n \rightarrow \mathbb{RP}^n$ and the fact that S^n is simply connected.

Deck transformations

A *deck transformation* of a covering map $f: X \rightarrow Y$ is a homeomorphism $g: X \rightarrow X$ so that $f \circ g = f$.

For any integer n , the map $\theta \mapsto \theta + 2\pi n$ is a deck transformation of the universal covering space $\theta \in \mathbb{R} \mapsto e^{i\theta} \in S^1$.

The map $x \in S^2 \mapsto -x \in S^2$ is a deck transformation of the universal covering map $x \in S^2 \mapsto [x] \in \mathbb{RP}^2$.

Lemma 4.17. *Take a Hausdorff, path connected and locally simply connected space X . The group of deck transformations of the universal covering space $\tilde{X} \rightarrow X$ is precisely $\pi_1(X)$, acting by $[x][y] = [x * y]$.*

Proof. It is clear that the fundamental group acts in this manner by deck transformations. Write out universal covering as $p: \tilde{X} \rightarrow X$ and suppose that

$$p(\tilde{x}_0) = x_0.$$

Take a deck transformation $g: \tilde{X} \rightarrow \tilde{X}$, say taking \tilde{x}_0 to some point \tilde{x}_1 . Since \tilde{X} is simply connected, there is a unique path \tilde{x} from \tilde{x}_0 to \tilde{x}_1 , up to homotopy. Composing g with the map

$$[y] \mapsto [\tilde{x}^{-1} * y]$$

we can arrange that g fixes \tilde{x}_0 . Being a deck transformation, g is locally identified by p with the identity map $X \rightarrow X$, so if g fixes a point, then g fixes all nearby points, lifting the identity map to the unique lift up to \tilde{X} . Therefore g is the identity map. So every deck transformation of the universal covering space arises from a unique element of the fundamental group. \square

4.18 Let Y be the set of all quadratic polynomials of the form $z^2 + bz + c$, with two distinct roots and complex coefficients b, c , in a complex variable z . Let X be the set of all pairs (z_0, z_1) of distinct complex numbers $z_0 \neq z_1$. Map $f: X \rightarrow Y$ by taking $f(z_0, z_1)$ to be the quadratic polynomial in Y with roots $z = z_0$ and $z = z_1$. Explain why f is a covering map. Explain how to find the universal covering space $\tilde{Y} \rightarrow Y$. Explain how the fundamental group of Y acts on \tilde{Y} .

Classification of regular covering spaces

A covering space $X \rightarrow Y$ is *regular* if the deck transformations act transitively on the fibers, i.e. the quotient is Y . If Γ is the group of deck transformations, we also say that $Y \rightarrow X$ is a Γ -covering. It is convenient to restate lemma 4.15 on page 51 in terms of maps to groups, rather than in terms of subgroups, so that we can find all of the regular covering spaces. Take a path connected and locally simply connected space X , a group Γ , and a group morphism $\phi: \pi_1(X) \rightarrow \Gamma$. Write the universal covering space of X as $p: \tilde{X} \rightarrow X$. Give Γ and $\pi_1(X)$ the discrete topology. Let $\pi_1(X)$ acts on $\tilde{X} \times \Gamma$ by having any $h \in \pi_1(X)$ act on $(x, g) \in \tilde{X} \times \Gamma$ as

$$h(x, g) = (hx, g\phi(h)^{-1}).$$

Let

$$X_\phi := (\tilde{X} \times \Gamma) / \pi_1(X),$$

with quotient topology. The map $(x, g) \mapsto p(x)$ is $\pi_1(X)$ -invariant, so descends to a unique map $p_\phi: X_\phi \rightarrow X$. The group Γ acts on $\tilde{X} \times \Gamma$ by $g_0(x, g) = (x, g_0g)$. This action commutes with the $\pi_1(X)$ -action, so descends to an action on X_ϕ .

Theorem 4.18. *Take a path connected and locally simply connected space X , a group Γ , and a group morphism $\phi: \pi_1(X) \rightarrow \Gamma$. Then $p_\phi: X_\phi \rightarrow X$ is a Γ -covering. The space X_ϕ is path connected just when ϕ is surjective. Conversely, every regular covering space $Y \rightarrow X$ is isomorphic to X_ϕ , for a unique ϕ .*

Proof. The actions of $\pi_1(X)$ and of Γ on $\tilde{X} \times \Gamma$ are covering actions, so

$$\tilde{X} \times \Gamma \rightarrow X_\phi$$

is a covering map. But

$$\tilde{X} \times \Gamma \rightarrow X_\phi \rightarrow X$$

is also a covering map, so every point $x_0 \in X$ lies in an open set $U \subset X$ with preimage in \tilde{X} equivariantly homeomorphic to a product $U \times \pi_1(X)$. So the preimage of U in $\tilde{X} \times \Gamma$ is equivariantly $U \times \pi_1(X) \times \Gamma$. So X_ϕ is the quotient by the action, i.e. $U \times \Gamma$.

Suppose that Γ acts as deck transformations on some covering space $Y \rightarrow X$, with quotient X . So if we have points $x_0 \in X$ and $y_0 \in Y_{x_0}$ then Γ acts on Y_{x_0} transitively, and each element of Γ is uniquely determined by where it takes y_0 . Take a path on X , say $[x] \in \tilde{X}$, starting at x_0 . Lift to a path $[y]$ on Y , starting at y_0 , to get a map $\tilde{X} \rightarrow Y$. If $[x]$ is a loop, $[y]$ has end points above x_0 . Since Γ acts as deck transformations on Y with quotient X , $y(1) = g y_0$ for a unique deck transformation $g \in \Gamma$, say $g = \phi([x])$, so $\phi: \pi_1(X) \rightarrow \Gamma$ is a group morphism. So then $\tilde{X} \rightarrow Y$ is ϕ -equivariant, dropping to a morphism $X_\phi \rightarrow Y$, which is a bijection on fibers above x_0 , so an isomorphism. \square

van Kampen's theorem I

Theorem 4.19 (van Kampen I). *Suppose that X is a path connected and locally simply connected topological space, with a covering by path connected open sets $X_a \subset X$, all containing the same point $x_0 \in X$, and that every intersection $X_{ab} := X_a \cap X_b$ is also path connected. Let $\pi := \pi_1(X, x_0)$ and $\pi_a := \pi_1(X_a, x_0)$, and so on. Take a group Γ and group morphisms $\pi_a \rightarrow \Gamma$ which agree on every π_{ab} , i.e. a commutative diagram*

$$\begin{array}{ccc} \pi_{ab} & \longrightarrow & \pi_a \\ \downarrow & & \downarrow \\ \pi_b & \longrightarrow & \pi \end{array} \quad \begin{array}{c} \searrow \\ \Gamma \end{array}$$

for every a, b . Then there is a unique group morphism $\pi \rightarrow \Gamma$ which makes commutative all of these diagrams:

$$\begin{array}{ccc} \pi_{ab} & \longrightarrow & \pi_a \\ \downarrow & & \downarrow \\ \pi_b & \longrightarrow & \pi \end{array} \quad \begin{array}{c} \searrow \\ \Gamma \end{array}$$

Proof. Call the group morphisms $\phi_a: \pi_a \rightarrow \Gamma$. Returning to theorem 4.18 on page 53, consider the Γ -covering spaces $\check{X}_a := (X_a)_{\phi_a} \rightarrow X_a$. Each contains an open set $\check{X}_{ab} \subset \check{X}_a$: the preimage of $X_{ab} \subset X_a$. So $\check{X}_{ab} \rightarrow X_{ab}$ is the Γ -covering space $X_{\psi_{ab}}$ where ψ_{ab} is the composition $\pi_{ab} \rightarrow \pi_a \rightarrow \Gamma$. Since the two compositions $\pi_{ab} \rightarrow \pi_a \rightarrow \Gamma$ and $\pi_{ab} \rightarrow \pi_b \rightarrow \Gamma$ are equal by assumption, these Γ -covering spaces \check{X}_{ab} and \check{X}_{ba} are identical, so there is a Γ -equivariant homeomorphism $\check{X}_{ab} \rightarrow \check{X}_{ba}$. Let \check{X} be the join: quotient of the disjoint union of all \check{X}_a by the equivalence of these homeomorphisms. By equivariance, \check{X} has a Γ -action, free because it is free at each point of each \check{X}_a . The Γ -quotient is the join of all X_a along all X_{ab} , i.e. X , and $\check{X} \rightarrow X$ is a Γ -covering because it is locally. But then $\check{X} \rightarrow X$ arises from a unique group morphism $\phi: \pi \rightarrow \Gamma$, as $\check{X} = X_\phi$. This morphism then restricts to π_a to become the group morphism that builds \check{X}_a , i.e. ϕ_a . \square

Group presentations

Take an abstract set S . A *word* on the *alphabet* S is a finite sequence of choices of element of S and integer power. We write each word as a string of symbols

$$s_1^{a_1} s_2^{a_2} \dots s_k^{a_k}.$$

We allow the empty string, and write it as 1. *Reduce* a word by deleting any s^0 symbols and replacing any subword $s^p s^q$ by s^{p+q} . The *free group* $\langle S \rangle$ on an alphabet S is the collection of all reduced words on S . The multiplication operation: multiply two words by writing down one after the other, called *concatenation* and then reducing. The inverse operation: write down the word in reverse order, with opposite signs for the powers.

Take an abstract set S with associated free group $\langle S \rangle$. Take a subset $R \subset \langle S \rangle$. The group *generated* by S with *relations* R , denoted $\langle S | R \rangle$, is the quotient of $\langle S \rangle$ by the smallest normal subgroup containing R . The expression of a group in the form $\langle S | R \rangle$ is a *presentation* of the group with *generators* S and *relations* R . A group is *finitely presented* if it is isomorphic to a group $\langle S | R \rangle$ with finite S and R .

Lemma 4.20. *Every map of sets $f: S \rightarrow G$ to a group extends uniquely to a morphism of groups $f: \langle S \rangle \rightarrow G$. It therefore extends to an injective morphism of groups $\langle S | R \rangle \rightarrow G$ where R is the set of all words*

$$w = s_1^{a_1} s_2^{a_2} \dots s_k^{a_k}$$

on the alphabet S for which

$$f(s_1)^{a_1} \dots f(s_k)^{a_k} = 1.$$

In practice, we usually write the relations not as a set R , but as a collection of equations like

$$w_1 = w_2$$

between words. This equation means that we require $w_2^{-1} w_1 \in R$.

We have seen that the Klein bottle X has fundamental group generated by the two transformations

$$T: (x, y) \mapsto (x, y + 1)$$


and

$$F: (x, y) \mapsto (x + 1, -y)$$

of the plane, a translation and a flip. Check $TFT = F$. Let $\Gamma = \langle f, t \mid tft = f \rangle$; in other words Γ is the group generated by alphabet $\{t, f\}$ with relation $tft = f$, i.e. with $R = \{t^{-1}tft\}$. So the fundamental group $\pi = \pi_1(X)$ admits a surjective morphism of groups $\Gamma \rightarrow \pi$, mapping $t \mapsto T$ and $f \mapsto F$. Here t is a formal symbol, not an actual transformation of the plane, while T is the actual transformation. Take any word in Γ . Wherever we find tf we replace it by ft^{-1} . Wherever we find $t^{-1}f$ we replace it by ft . Wherever we find tf^{-1} we replace it by $f^{-1}t^{-1}$. Wherever we find $t^{-1}f^{-1}$ we replace it by $f^{-1}t$. The reader can check that these are all consequences of $tft = f$. Therefore any word in Γ can be written uniquely as $f^p t^q$. If the corresponding transformation $F^p T^q$ is the identity, then it fixes the origin, and so the translations of the x variable cancel each other out, i.e. $p = 0$. But then $T^q(x, y) = (x, y + q)$ fixes the origin just when $q = 0$. So the only element of Γ mapping to the trivial transformation of the plane is 1. Therefore $\Gamma = \pi$, i.e. the fundamental group of the Klein bottle X is

$$\pi_1(X) = \langle f, t \mid tft = f \rangle.$$

The image



is invariant under a vertical translation and under a horizontal translation with a flip (taking an upward pointing corner to a downward pointing one). The quotient by those two symmetries is thus the Klein bottle.

The fundamental group of the real projective space \mathbb{RP}^n is $\mathbb{Z}/2\mathbb{Z}$ (if $n \geq 2$), as we have seen, which clearly has the presentation

$$\pi_1(\mathbb{RP}^n) = \langle x | x^2 = 1 \rangle.$$

Amalgamations

Suppose that G and H are two groups. We would like to define a group $G * H$, which contains G and H , and is otherwise “as unconstrained as possible”. The product $G \times H$ is not “unconstrained”, because the elements of G commute with those of H inside $G \times H$.

First, note that any group G has an obvious group morphism $\langle G \rangle \rightarrow G$ given by $g \mapsto g$. It will help to write out concatenations using some symbol like

$$g_1 * g_2 * \cdots * g_n \in \langle G \rangle.$$

Then we can write our group morphism as

$$g_1 * g_2 * \cdots * g_n \in \langle G \rangle \mapsto g_1 g_2 \cdots g_n \in G.$$

This group morphism is clearly surjective, with kernel precisely the group $N_G \subset \langle G \rangle$ whose elements are the concatenations

$$g_1 * g_2 * \cdots * g_n$$

for which $g_1 g_2 \cdots g_n = 1$. So we can write

$$G = \langle G | N_G \rangle.$$

Think of N_G as encoding all of the equations satisfied inside the group G .

We define the *free product* $G * H$ to be the group $\langle G \sqcup H | N_G \sqcup N_H \rangle$ generated by the elements of G and H , subject to the relations consisting of all equations satisfied by elements of G together with all equations satisfied by elements of H .

Another way to look at this: a *word* in G, H is a finite sequence of elements of G and of H (perhaps empty), written beside one another with $*$ symbols inbetween, like

$$g_1 * g_2 * h_1 * g_3 * h_2 * h_3 * g_4,$$

et cetera. We denote the empty sequence as 1. We *reduce* a word by deleting any appearance of the identity element (of either group), and also by replacing any two neighboring elements from the same group by their product in that group:

$$g_1 * g_2 \mapsto g_1 g_2.$$

A word is *reduced* if we cannot further reduce it. The group $G * H$ is the set of reduced words, with multiplication being simply writing down one word after another and then reducing.

A further wrinkle: suppose that $K \subset G$ is a subgroup which also appears as a subgroup of H : $K \subset H$. The *amalgamation* of G and H over K , denoted $G *_K H$, is

$$G *_K H = \langle G \sqcup H \mid N_G \sqcup N_H \sqcup E \rangle$$

where E is the collection of equations $k_G = k_H$ where k_G is an element of K as a subgroup of G , and k_H is the associated element of $K \subset H$. Equivalently, we can think of allowing reduced words to be acted on by inserting into any reduced word an element of K right of an element of G , and left of the next element of H , and vice versa. There is no longer any straightforward description in terms of reduced words, but the trivial elements, once reduced, are products $k_1 * k_1^{-1} * \cdots * k_n * k_n^{-1}$.

Similarly, if we have a collection of groups G_a , for a in some set, the *free product* $*G_a$ is the quotient

$$\left\langle \bigsqcup_a G_a \mid \bigsqcup_a N_{G_a} \right\rangle,$$

and if we have some groups K_c and various morphisms $\phi_{ca}: K_c \rightarrow G_a$, the *amalgamation* $*_{K_c} G_a$ is

$$\left\langle \bigsqcup_a G_a \mid \bigsqcup_a N_{G_a} \sqcup E \right\rangle,$$

where E is the set of all equations $\phi_{ca}(k) = \phi_{cb}(k)$ where $k \in K_c$.

van Kampen's theorem II

Theorem 4.21 (van Kampen II). *Take a path connected and locally simply connected topological space X , and a cover by path connected open sets $X_a \subset X$, with path connected intersections $X_{ab} := X_a \cap X_b$, all containing some point $x_0 \in X$. Let $\pi := \pi_1(X, x_0)$, $\pi_a := \pi_1(X_a, x_0)$, and so on, with obvious commutative diagrams*

$$\begin{array}{ccc} \pi_{ab} & \longrightarrow & \pi_a \\ \downarrow & & \downarrow \\ \pi_b & \longrightarrow & \pi \end{array}$$

*Then $\pi = *_{\pi_{bc}} \pi_a$ is the amalgamation of all π_a over all π_{ab} .*

Proof. Let $\Gamma := *_{\pi_{bc}} \pi_a$. There are obvious morphisms of groups

$$\begin{array}{ccc} \pi_{ab} & \longrightarrow & \pi_a \\ \downarrow & & \downarrow \\ \pi_b & \longrightarrow & \pi \end{array} \quad \begin{array}{c} \searrow \\ \Gamma \end{array}$$

given by taking each word in π_a as giving a word in the larger alphabet of Γ . By theorem 4.19 on page 54, there is a unique group morphism $\pi \rightarrow \Gamma$ making all diagrams commutative

$$\begin{array}{ccc}
 \pi_{ab} & \xrightarrow{\quad} & \pi_a \\
 \downarrow & & \downarrow \\
 \pi_b & \xrightarrow{\quad} & \pi \\
 & \searrow & \searrow \\
 & & \Gamma
 \end{array}$$

This morphism of groups is surjective, because it includes the image of every π_a .

The group morphisms $\pi_a \rightarrow \pi$ determine a single group morphism $*\pi_a \rightarrow \pi$ of the free product. Take any loop in X ; lemma 3.1 on page 29 splits it into intervals, say $x_i(t) := x|_{[t_i, t_{i+1}]}$, remaining each in a single X_{a_i} . Adding a little on at the beginning and end of those intervals, because every X_a is path connected and contains x_0 , we can arrange that $x_i(t)$ is a loop starting and ending at x_0 . Hence $*\pi_a \rightarrow \pi$ is surjective.

The kernel of the composition $*\pi_a \rightarrow \pi \rightarrow \Gamma$ is precisely the subgroup generated by the various words $k_a * k_b^{-1}$ where k_a and k_b are the images in π_a and π_b of some $k \in \pi_{ab}$. The element k is a loop $[x]$ in X_{ab} , and so becomes that loop sitting in either X_a and X_b , and both of those become the same loop in X , so this word becomes trivial in π . So $\pi \rightarrow \Gamma$ is injective. \square

Suppose that X is the sphere of dimension $n \geq 2$. Let X_1 and X_2 be the sphere punctured at the south pole and the north pole. So X_{12} is the sphere with two punctures. These are all path connected, and $\{1\} = \pi_1 = \pi_2$, so $\{1\} = \pi$: X is simply connected.


Suppose that X is the bouquet of n circles, say joined at a point $x_0 \in X$. We define each X_1, \dots, X_n to be X with all but one of the circles punctured, and hence homotopy equivalent to a single circle. So $\pi_i = \pi_1(X_i, x_0) = \mathbb{Z}$. The various π_{ij} are all trivial: all circles are punctured. So $\pi = \mathbb{Z} * \dots * \mathbb{Z} = *^n \mathbb{Z}$.

If Y_a are path connected and locally simply connected spaces with marked points $x_a \in Y_a$, and let $Y = *Y_a$ be the join of them along those points, then we can take simply connected neighborhoods $U_a \subset Y_a$ of x_a , and let X_a be the join $Y_a *_{b \neq a} U_b$. Then $\pi_{ab} = \{1\}$ and so $\pi = *\pi_a$, i.e. the fundamental group of the join at points is the free product of the fundamental groups.

Puncture a torus, and gently widen the puncture, until you see a pair of “belts” attached; homotopy equivalent to a bouquet of two circles. A torus, say with fundamental group $\langle x, y | xy = yx \rangle$, becomes punctured, and becomes a bouquet of circles with fundamental group $\langle x, y \rangle$, included into the torus as $x, y \mapsto x, y$.

Suppose that X is a connected manifold of dimension at least 3; pick a point $p_0 \in X$. Take a chart around p_0 , with connected domain, and a point $x_0 \in X$ in the domain of the chart. Let X_2 be an open set identified by the chart with the interior of a closed ball. Cover X in the open set $X_1 = X - \{p_0\}$ and X_2 . They intersect in the set $X_0 = X_1 \cap X_2$, homeomorphic to a punctured ball according to the chart. Then $\pi_2 = \{1\}$ and $\pi_0 = \{1\}$, so $\pi = \pi_1$, i.e. $\pi_1(M - \{p_0\}, x_0) = \pi_1(M, x_0)$: deleting a point does not affect the fundamental group of a manifold of dimension 3 or more.

How does poking a hole affect the fundamental group of a surface? Suppose that X is a connected surface; pick a point $p_0 \in X$. Take a *chart* around p_0 : a homeomorphism from a connected neighborhood of p_0 to an open subset of the plane. Take a point $x_0 \in X$ in the domain of the chart. Let X_2 be an open set identified by the chart with the interior of a closed disk in the plane. Cover X in the open set $X_1 = X - \{p_0\}$ and X_2 . They intersect in the set $X_0 = X_1 \cap X_2$, homeomorphic to a punctured ball according to the chart. Then $\pi_2 = \{1\}$ and $\pi_0 = \mathbb{Z}$. so $\pi = \pi_1 *_{\mathbb{Z}} 1 = \pi_1/N$, where N is the normal subgroup generated by the puncture.



Take two tori, poke holes in them, and join together along a tube; let X be the resulting surface of genus 2. Each punctured torus is an open set $X_1, X_2 \subset X$, and the tube is their intersection $X_0 = X_1 \cap X_2$. The tube X_0 has fundamental group $\pi_0 = \mathbb{Z}$, and this includes into each fundamental group of each punctured torus as a small circle around the

puncture. On each torus, the circle around the puncture is the product $xyx^{-1}y^{-1}$, which of course vanishes once we close up the puncture. So if we glue two tori together, the resulting surface X of genus 2 has fundamental group

$$\langle x, y, X, Y | xyx^{-1}y^{-1} = XYX^{-1}Y^{-1} \rangle.$$

4.19 Prove that the fundamental group of any compact, path connected, and locally simply connected topological space is finitely presented.

Homotopy groups

For any topological space X with marked point x_0 , and any $n \geq 1$, let $\pi_n(X, x_0)$ be the set of all continuous maps $[0, 1]^n \rightarrow X$ taking every boundary point of $[0, 1]^n$ to x_0 , modulo homotopy through such maps. Each $\pi_n(X, x_0)$ has a distinguished point: the constant map.

There is a special case: $n = 0$. We define $\pi_0(X, x_0)$ to be the set of all components of X , with a special chosen component: that of x_0 . Make $\pi_0(X, x_0)$ into a topological space, by the quotient topology.

For $n \geq 1$, take $f, g: [0, 1]^n \rightarrow X$ each mapping the boundary of $[0, 1]^n$ to x_0 . Define $f * g$ to be the usual

$$(f * g)(t, x_2, \dots, x_n) = \begin{cases} f(2t, x_2, \dots, x_n), & \text{if } 0 \leq t \leq \frac{1}{2}, \\ g(2t - 1, x_2, \dots, x_n), & \text{if } \frac{1}{2} \leq t \leq 1, \end{cases}$$

and $\bar{f}(t, x_2, \dots, x_n) = f(1 - t, x_2, \dots, x_n)$. If $f: X \rightarrow Y$ is continuous, and $y_0 := f(x_0)$, define $f_*: \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$ by $f_*[g] = [f \circ g]$.

A *contraction* of a topological space X to a point $x_0 \in X$ is a continuous map $\varphi: X \times [0, 1] \rightarrow X$ so that $\varphi(x, 0) = x$ and $\varphi(x, 1) = x_0$ for any $x \in X$. A topological space X is *contractible* if it has a contraction.

4.20 Prove that every contractible space is connected and has trivial homotopy groups.

Given a continuous map $f: X \rightarrow Y$, a *lift* of a continuous map $g: Z \rightarrow Y$ is a continuous map $\hat{g}: Z \rightarrow X$ so that $f \circ \hat{g} = g$. A *Serre fibration* is a continuous map $f: X \rightarrow Y$ of topological spaces, so that for any box $Z = [0, 1]^n$ and continuous map $g: Z \times [0, 1] \rightarrow Y$, denoted $g_t(z) := g(z, t)$, every lift of g_0 extends to a lift of g .

Theorem 4.22. *If $f: X \rightarrow Y$ is a Serre fibration, then the obvious maps*

$$\pi_n(F, x_0) \rightarrow \pi_n(X, x_0) \rightarrow \pi_n(Y, y_0)$$

fit together into an exact sequence

$$\begin{array}{c}
\cdots \\
\curvearrowright \\
\pi_4(F, x_0) \longrightarrow \pi_4(X, x_0) \longrightarrow \pi_4(Y, y_0) \\
\curvearrowright \\
\pi_3(F, x_0) \longrightarrow \pi_3(X, x_0) \longrightarrow \pi_3(Y, y_0) \\
\curvearrowright \\
\pi_2(F, x_0) \longrightarrow \pi_2(X, x_0) \longrightarrow \pi_2(Y, y_0) \\
\curvearrowright \\
\pi_1(F, x_0) \longrightarrow \pi_1(X, x_0) \longrightarrow \pi_1(Y, y_0) \\
\curvearrowright \\
\pi_0(F, x_0) \longrightarrow \pi_0(X, x_0) \longrightarrow \pi_0(Y, y_0) \longrightarrow 0
\end{array}$$

Pick $x_0 \in X$, let $y_0 := f(x_0)$ and let $F := f^{-1}\{y_0\} \subseteq X$. The morphism $\pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ kills loops in X which lie entirely in a fiber $F = f^{-1}\{y_0\}$, i.e. those loops map to $1 \in \pi_1(Y, y_0)$. More generally, it kills the loops in X which admit a homotopy, fixing endpoints, to a loop in F . Any loop in Y starting and ending at y_0 lifts to a path in X , starting at x_0 , but perhaps not a loop. Since it is a loop in Y , its lift has ends in X lying in the fiber F . Under homotopy of the loop in Y , with fixed ends, the ends of the lift can only move inside F , so they have well defined components, a map

$$\pi_1(X, x_0) \rightarrow \pi_1(Y, y_0) \rightarrow \pi_0(F, x_0),$$

where $\pi_0(X_{y_0}, x_0)$ is the set of all path components of X_{y_0} , and contains a distinguished element, the path component that contains $x_0 \in X_{y_0}$. If the ends of that lift lie in the same path component, then we can draw those two ends together, obtaining the path in Y from a path in X , i.e. the image of the first map is the “kernel” of the second, i.e. the elements mapping to the distinguished component.

Take a continuous map $h: Z \times [0, 1] \rightarrow Y$, which we write as $h_t(z) = h(z, t)$. Suppose that $h_0(z) = y_0$ for all z , and $h_s(z) = y_0$ for all $z \in \partial Z$. Lift h_0 to the trivial map to x_0 . Lift h to a map $\hat{h}: Z \times [0, 1] \rightarrow X$ extending that lift. So $f \circ \hat{h}_t(z) = h_t(z)$, and $\hat{h}_0(z) = x_0$ for all $z \in Z$. Since $h_s(z) = y_0$ for $z \in \partial Z$, $\hat{h}_s(z) \in F$ for $z \in \partial Z$, for all s . In particular, $\hat{h}_s(z)$ stays in the component of x_0 in F , for $z \in \partial Z$, for all s . If we replace our choices at each step, still $\hat{h}_1: \partial Z \rightarrow F$ varies only by homotopy, giving an element of $\pi_{n-1}(F, x_0)$, a map

$$\pi_n(X, x_0) \rightarrow \pi_n(Y, y_0) \rightarrow \pi_{n-1}(F, x_0).$$

If this element vanishes, then \hat{h}_1 is nullhomotopic in F , and if we attach that homotopy to \hat{h} , and its composition with $X \rightarrow Y$ to h , then h continues to

map to $h_s(z) = y_0$. So h is an element of $\pi_n(Y, y_0)$ arising from an element of $\pi_n(X, x_0)$. In other words, in

$$\pi_n(X, x_0) \rightarrow \pi_n(Y, y_0) \rightarrow \pi_{n-1}(F, x_0).$$

the kernel of the second map is the image of the first.

Being a Serre fibration is a local property in Y :

Lemma 4.23. *A continuous map $f: X \rightarrow Y$ of topological spaces is a Serre fibration if and only if there is an open cover of Y by open sets $Y_a \subseteq Y$ so that, if $X_a := f^{-1}Y_a$ and $f_a := f|_{X_a}$, then each f_a is a Serre fibration.*

Proof. Take such an open cover, and some map $g: Z \times [0, 1] \rightarrow Y$ with a lift $\hat{g}_0: Z \rightarrow X$, where $Z = [0, 1]^n$. Cover the image of g in our open sets Y_a . Take a finite subcover. Divide up $Z \times [0, 1]$ into a grid of cubes small enough that each lies inside a single Y_a . Lift up one cube at a time. \square

A *projection* is a map $(x, z) \in X \times Z \mapsto z \in Z$; the *fiber* is Z . Two maps $X_0 \rightarrow Y_0$ and $X_1 \rightarrow Y_1$ are *isomorphic* if there are diffeomorphisms $X_0 \rightarrow X_1$

and $Y_0 \rightarrow Y_1$ making a commutative diagram

$$\begin{array}{ccc} X_0 & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ Y_0 & \longrightarrow & Y_1 \end{array}$$

A continuous

map is *trivial* if isomorphic to a projection. A continuous map $\pi: X \rightarrow Y$ is *locally trivial* if every point of Y lies in an open set $U \subseteq Y$ so that

$$\pi|_{\pi^{-1}U}: \pi^{-1}U \rightarrow U$$

is trivial. A *fiber bundle* is a continuous locally trivial map $\pi: X \rightarrow Y$ of topological spaces.

Lemma 4.24. *Every fiber bundle is a Serre fibration.*

Proof. Fiber bundles are locally trivial, so true locally; apply lemma 4.23. \square

Hints

1.3. On the empty set, the only topology is the one whose only open set is the empty set. On the set $X = \{0\}$, the only topology is the discrete topology, which is also the indiscrete topology. On the set $X = \{0, 1\}$, the possible topologies are

- a. discrete: open sets are: the empty set, $\{0\}$, $\{1\}$, $\{0, 1\}$;
- b. indiscrete: open sets are: the empty set, $\{0, 1\}$;
- c. the *Sierpinski topology*: open sets are: the empty set, $\{0\}$, $\{0, 1\}$;
- d. the *other Sierpinski topology*: open sets are: the empty set, $\{1\}$, $\{0, 1\}$.

1.4. Take some sets $\{C_a\}_{a \in \mathcal{A}}$, so that each $C_a \subseteq X$ is closed. The complement $U_a := X - C_a$ is open. So the union $U := \bigcup_a U_a$ is open. So its complement $C = X - U$ is closed. But C is the set of points of X not belonging to U , i.e. not belonging to any U_a , i.e. belonging to C_a instead of U_a for every a , i.e. $C = \bigcap_a C_a$.

1.40. The diagonal is closed just when its complement is open. Its complement is the set $U \subset X \times X$ of pairs of distinct points. This U is open just when every point of U lies inside a basis element which lies in U , for any basis, and in particular for the basis consisting of products $U_1 \times U_2$ of open sets $U_1, U_2 \subset X$. So the diagonal is closed just when, for any pair of distinct points $x_1 \neq x_2$ in X , the point (x_1, x_2) lies inside a subset $U_1 \times U_2 \subset X \times X$ which does not intersect the diagonal. In other words, just when $x_1 \in U_1$ and $x_2 \in U_2$ but $U_1 \cap U_2$ is empty. In other words, just when X is Hausdorff.

1.42. Take two points $(x_1, y_1), (x_2, y_2)$ in $X \times Y$. Take disjoint open sets $X_1, X_2 \subset X$ and $Y_1, Y_2 \subset Y$ so that x_1, x_2, y_1, y_2 belong to X_1, X_2, Y_1, Y_2 respectively. Then (x_1, y_1) is in $X_1 \times Y_1$ and so on.

1.46. Take X to be a set consisting of two points p, q with open sets: X , the empty set, and the set $\{p\}$. Then $\{q\}$ is the complement of $\{p\}$, so closed. But $\{p\}$ is *not* the complement of an open set, so is not closed. There are only 4 open sets, so every open cover is finite.

1.49. Take compact sets K_a and let K be their intersection. Take an open cover U_b of K . By lemma 1.5 on page 12, every K_a is closed. Therefore the intersection K is closed, so $X - K$ is open. The open sets $\{X - K\} \cup \{U_b\}$ cover X . So each set K_a is covered by these, and so covered by finitely many of these. But then those finitely many cover $K \subset K_a$. So K is compact. Suppose

that K is empty. The open sets $W_b := X - K_b$ cover X . Each set K_a is compact, so finitely many W_b cover K_a , say W_{b_1}, \dots, W_{b_n} . So $W_a, W_{b_1}, \dots, W_{b_n}$ covers X . So their complements $K_a, K_{b_1}, \dots, K_{b_n}$ intersect in an empty set.

1.50. Let $X = \{0, 1, 2, \dots\} \cup \{\infty, \infty'\}$, where a subset $U \subseteq X$ is open just when either $U \subseteq \{0, 1, 2, \dots\}$ or $X \setminus U$ is finite. Then $A = \{0, 1, 2, \dots\} \cup \{\infty\}$ and $B = \{0, 1, 2, \dots\} \cup \{\infty'\}$ are both compact. But $A \cap B = \{0, 1, 2, \dots\}$ has the discrete topology and is not compact.

1.67. Every point x of X lies in the interior U of a compact set $\bar{U} \subset X$. So K is covered by such interiors. Take a finite subcover U_1, U_2, \dots, U_k and let $V := U_1 \cup \dots \cup U_k$, so $\bar{V} \subset \bar{U}_1 \cup \dots \cup \bar{U}_k$ is compact.

1.68. If X has a basis of precompact open sets, then every point of X lies in one of them. If every point of X lies in a precompact open set, then those open sets cover X , so X is locally compact. Suppose that X is locally compact. Take a point $x_0 \in X$. So x_0 lies in one of our X_a open sets lying in a compact set, and hence (by Hausdorffness!) with compact closure \bar{X}_a . Every open set U around x_0 contains $U \cap X_a$, which has compact closure, hence a basis of precompact open sets.

1.69. We know the subsets of a Hausdorff space are Hausdorff. If $U \subseteq X$ is open, every point of U lies in an open set with compact closure inside U , so U is locally compact Hausdorff. If $A \subseteq X$ is closed, cover X in a basis of precompact open sets, and they intersect A in precompact open sets covering A .

2.20. Take an open cover $Y_a \subset Y$ of $f(K)$. Let $X_a = f^{-1}Y_a$. Then X_a form an open cover of $f^{-1}f(K)$, and so of K . Take a finite subcover of K , say $X_1, X_2, \dots, X_n \subset X$. Then the corresponding sets Y_1, Y_2, \dots, Y_n are an open cover of $f(K)$.

2.25. If X', X'' are two such spaces, define $f: X' \rightarrow X''$ to be the identity on X and $\infty \rightarrow \infty$. Take an open set $U \subseteq X'$. If ∞ is not in U , $f(U) = U$ is open since $X \subset X''$ is embedded. If $\infty \in U$, $C := X' - U = X - (U - \infty)$ is closed in X' so compact in X' , therefore in X since $X \subset X'$ is embedded, and so C is compact in X'' , and so closed since X'' is Hausdorff. Hence $U = X'' - C$ is open. Therefore f is an open map, and so is f^{-1} by the same argument, so both are homeomorphisms.

Following problem 2.24 on page 20, let X' be the one point compactification of X . We need to prove that X' is compact and Hausdorff and that $X \subset X'$ is embedded. Trivially any open set $U \subset X'$ intersects X in an open set, so the inclusion map $X \rightarrow X'$ is continuous. Conversely, any open set in X is open in X' , so the inclusion map is open, hence an embedding.

Take an open cover of X' by open sets X'_a . Each is either $X'_a = (X - C_a) \cup \{\infty\}$ for a compact set $C_a \subseteq X$ or $X'_a = X_a$ for open sets $X_a \subseteq X$. At least one of these X'_a contains ∞ , so is some $X - C_{a_0}$. The sets X_a or $X - C_a$ cover $X' - \{\infty\}$ so cover X . Because $C_{a_0} \subseteq X$ is compact, C_{a_0} is covered by finitely many of these. So then $X' = X \cup \{\infty\}$ is covered by those finitely many together with $(X - C_{a_0}) \cup \{\infty\}$.

To see that X' is Hausdorff, note that $X \subset X'$ is embedded and Hausdorff. Take two points $x, y \in X'$. If both are in X , then they are housed off. So we

can suppose that $y = \infty$ and x

in X . Because X is locally compact, we can choose a compact set $C \subseteq X$ containing an open set U around x , and then $U, (X - C) \cup \{\infty\}$ house x, ∞ .

If X' satisfying these three conditions exists, we need to prove that X is locally compact and Hausdorff. But $X \subset X'$ is embedded in a Hausdorff space, so Hausdorff. Pick a point $x \in X$ and houses in X' : open sets $U, V \subseteq X'$ so that $x \in U$ and $\infty \in V$ and $U \cap V$ is empty. Then $C = X' - V$ is closed in Y so compact, and so compact in X , and contains U . So X is locally compact.

2.26. If we write a point p of the sphere $S^n \subset \mathbb{R}^{n+1}$ as

$$p = \begin{pmatrix} a \\ b \end{pmatrix}$$

with $a \in \mathbb{R}^n$ and $b \in \mathbb{R}$, then map

$$y = \varpi(p) = \frac{a}{1 - b}.$$

The inverse map is

$$p = \varpi^{-1}(y) = \begin{pmatrix} 0 \\ 1 \end{pmatrix} + \frac{1}{1 + \|y\|^2} \begin{pmatrix} 2y \\ -2 \end{pmatrix}.$$

So

$$\varpi: S^n - \begin{pmatrix} 0 \\ -1 \end{pmatrix} \rightarrow \mathbb{R}^n$$

is a homeomorphism.

2.27. The image $f(A)$ of a closed set $A \subseteq X$ is closed in Y so closed $f(X)$. Hence we can replace Y by $f(X)$ and assume the map is a bijection. Closed sets have closed images. Open sets have open images by taking complements. But then for $f^{-1}: Y \rightarrow X$, the f^{-1} -preimages of open sets are just exactly the f -images, so open. So f^{-1} is continuous.

2.28. Suppose (a) and (b). Pick a compact set $K \subseteq Y$ and an open cover $U_\alpha \subset X$ of $f^{-1}K$. Pick a point $y_0 \in Y$. Choose finitely many open sets $U_{\alpha_i} \subseteq X$ covering $f^{-1}\{y_0\}$; let U be their union. Check that y_0 belongs to

$$W := Y - f(X - U)$$

which is open since f is closed. Note that $f^{-1}W \subset U$. For each $y_0 \in Y$, there is some such open set W . Cover K by finitely many such W , say W_j , each arising as

$$W_j = Y - f(X - U_j),$$

from some open set $U_j \subset X$ which itself is a finite union of open sets U_{α_i} on X . So $f^{-1}K$ lies inside the union of the various $f^{-1}W_j$, each of which lies in its U_j . So $f^{-1}K$ lies in a finite union of U_{α_i} sets.

2.29. Take a compact set $K \subseteq Y$. Since Y is Hausdorff, K is closed. Take an open cover of $f^{-1}K$ by open sets X_α ; add in one more open set $X_0 = X - f^{-1}K$

to give an open cover of X . Take a finite subcover of X ; throw away X_0 to give a finite subcover of $f^{-1}K$.

2.33. The first problem is to use linear algebra to prove that any matrix $A \in Y$ arises as a *Gram matrix*

$$A = \varphi(v_1, \dots, v_n),$$

where φ is the map giving $A_{ij} = \langle v_i, v_j \rangle$. For this, you can use induction on dimension. The second problem is to show that this Gram matrix uniquely determines the vectors v_1, \dots, v_n up to orthogonal linear transformation. Hence the map φ associating to vectors their Gram matrix is an injection modulo the action of the orthogonal group. The third problem is to prove that every point $A \in Y$ lies in the interior of the image of a compact set of choices of vectors v_1, \dots, v_n ; but this follows from taking all of the vectors to lie in some large enough ball in \mathbb{R}^n , so that every Gram matrix whose diagonals are bounded by the radius of that ball lies in the image of that set.

2.36. If we let $t = \pi|x|$ then $|f(x)|^2 = \cos^2 t + \sin^2 t = 1$, so f is valued in the unit sphere Y . Continuity is clear except when $|x| \rightarrow 0$, but then the fact that $\sin' t = \cos t$ tells us that

$$\frac{\sin t}{t}$$

is continuous with

$$1 = \lim_{t \rightarrow 0} \frac{\sin t}{t}$$

by L'Hôpital's rule. Write

$$f(x) = \begin{cases} \left(\cos(\pi|x|), \frac{\sin(\pi|x|)}{|x|}x \right), & \text{if } |x| \neq 0, \\ (1, 0), & \text{if } |x| = 0. \end{cases}$$

to see that $f(x) \rightarrow (1, 0)$ as $|x| \rightarrow 0$, continuity. Clearly $f(0) = (1, 0)$ and, if $|x| = 1$, then $f(x) = (-1, 0)$. Since $0 \leq \pi|x| \leq 1$, if $f(x) = (-1, 0)$ then $\cos(\pi|x|) = -1$ and $\sin(\pi|x|) = 0$ and so $\pi|x| = \pi$, i.e. $|x| = 1$. If $x, y \in X$ and $|x| < 1$ and $|y| < 1$ and if $f(x) = f(y)$ then

$$\cos(\pi|x|) = \cos(\pi|y|)$$

so $0 \leq |x| = |y| < 1$. But then also

$$\sin(\pi|x|)x = \sin(\pi|y|)y$$

and $\sin(\pi|x|) \neq 0$ so $x = y$. Hence f is continuous and 1-1. Since X is compact and Y is Hausdorff and f is onto, f identifies Y with the quotient of X by the map, i.e. with the closed ball with its boundary glued to a single point.

3.2. Let $F_s(t) = x((1-s)t + s\tau(t))$.

4.1. Suppose that $U_Y \subset Y$ is an evenly covered open set near $1 \in Y$. Write each point of Y as $w = u + iv \in Y$. Then U_Y contains a connected open set

near $1 \in Y$, say the “interval” $-\varepsilon < v < \varepsilon$. Since U_Y is evenly covered, so is this “interval”. The preimage of this “interval” is a union of 4 open intervals:

$$(0, \delta) \cup (1 - \delta, 1 + \delta) \cup (2 - \delta, 2 + \delta) \cup (3 - \delta, 3)$$

where

$$\delta = \frac{\sin^{-1} \varepsilon}{2\pi}.$$

Any sheet over our open set is homeomorphic so also path connected, so lies inside one of these 4 intervals. But no open subset of the first interval maps onto our “interval” as the map takes on values $u + iv$ with $0 < v < \varepsilon$ there.

4.3. Pick an evenly covered open set $U_{y_0} \subset Y$ containing y_0 . Every point $x_0 \in X_{y_0}$ lies in a sheet, say U_{x_0} , over U_{y_0} . Take two distinct points $x_0, x_1 \in X_{y_0}$. The sheets U_{x_0} and U_{x_1} are disjoint so x_1 is not in U_{x_0} and x_0 is not in U_{x_1} . Hence $X_{y_0} \cap U_{x_0} = \{x_0\}$. Take any subset $S \subset X_{y_0}$. Then

$$S = X_{y_0} \cap \bigcup_{x \in S} U_x$$

is the intersection of X_{y_0} with an open subset

$$\bigcup_{x \in S} U_x \subset X,$$

so is open inside X_{y_0} .

4.4. Suppose that Y is Hausdorff. Take distinct points x_1, x_2 of X . Let $y_1 = f(x_1)$ and $y_2 = f(x_2)$.

Suppose that $y_1 \neq y_2$. Take disjoint open sets $U_1, U_2 \subset Y$ so that $y_1 \in U_1$ and $y_2 \in U_2$. Then $f^{-1}U_1, f^{-1}U_2 \subset X$ are disjoint open sets so that $x_1 \in f^{-1}U_1$ and $x_2 \in f^{-1}U_2$.

Suppose that $y_1 = y_2$. Take an evenly covered open set $U \subset Y$ containing y_1 . Every point $x_1 \in X_{y_0}$ lies in a sheet, say U_{x_1} , over U . The sheets U_{x_1} and U_{x_2} are disjoint open sets.

Now instead suppose that X is Hausdorff. Take distinct points y_1, y_2 in Y . Take two points x_1, x_2 in X so that $y_1 = f(x_1)$ and $y_2 = f(x_2)$. Pick evenly covered open sets U_{y_1} and U_{y_2} around y_1 and y_2 . Pick disjoint open sets W_{x_1} and W_{x_2} around x_1 and x_2 . Let $U_{x_1} = W_{x_1} \cap f^{-1}U_{y_1}$. Then f is a homeomorphism of U_{x_1} to its image inside U_{y_1} ; call its image V_{y_1} . Similarly define V_{y_2} . Because f homeomorphically maps U_{x_1} to V_{y_1} , V_{y_1} is an open set in Y . The sets V_{y_1} and V_{y_2} are disjoint since they are images of disjoint open sets in X .

4.5. Take a point $q_0 \in Q$. Because f is proper, the fiber $P_{q_0} := f^{-1}\{q_0\}$ is compact. Because f is a local diffeomorphism, each point $p_0 \in P_{q_0}$ lies in an open set $U_{p_0} \subset P$ taken by f diffeomorphically to a neighborhood $U_{q_0} \subset Q$. The set P_{q_0} intersects U_{p_0} in a set taken by f to $\{q_0\}$, so just the single point p_0 . Therefore every point $p_0 \in P_{q_0}$ lies an open set U_{p_0} containing only p_0 , so

P_{q_0} is a discrete set of points. Being compact, P_{q_0} is therefore a finite set of points, say

$$P_{q_0} = \{ p_1, p_2, \dots, p_n \},$$

with each point p_j lying in an open set U_{p_j} taken diffeomorphically to some open set

$$f(U_{p_j}) \subset Q$$

around q_0 . Since there are finitely many such open sets, their intersection is open; call it U_{q_0} . Then replace U_{p_j} by

$$U_{p_j} \cap f^{-1}U_{q_0}$$

so we can arrange that f takes each of the open sets

$$U_{p_1}, U_{p_2}, \dots, U_{p_n}$$

diffeomorphically to U_{q_0} .

4.14. If X is a universal covering space, then the map $X \rightarrow Y$ lifts to a map $X \rightarrow Z$ by proposition 4.12 on page 47.

Take a covering map $(X, x_0) \rightarrow (Y, y_0)$ so that every other covering map $(Z, z_0) \rightarrow (Y, y_0)$ has a unique covering map $(X, x_0) \rightarrow (Z, z_0)$ making a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Z \\ & \searrow \quad \swarrow & \\ & Y & \end{array}$$

Take a universal covering map $(Z, z_0) \rightarrow (Y, y_0)$. Take the unique covering map described above. Any loop in X maps into Z , where it is homotopic to a constant map. Map that homotopy into Y , and then lift the curves of that homotopy up to curves in X , which are loops by taking the “endpoint of the lift” mapping. So then X is simply connected, i.e. a universal covering map.

4.15. By problem 4.14 on page 49, there are unique morphisms $(X_1, x_1) \rightarrow (X_2, x_2)$ and $(X_2, x_2) \rightarrow (X_1, x_1)$. The composition of these a morphism $X_1 \rightarrow X_1$. Again by problem 4.14 on page 49, there is a unique such morphism. But the identity map is a morphism. So $X_1 \rightarrow X_2$ composes with $X_2 \rightarrow X_1$ to give the identity map. Swapping the roles of X_1 and X_2 , the same is true for the composition in the other order. Therefore the two morphisms are inverses of each other.

4.16. By hypothesis, for any point $y_0 \in \mathbb{R}^n$, the fiber $f^{-1}\{y_0\}$ is bounded, but also closed, so compact. But around each point $x_0 \in f^{-1}\{y_0\}$, f is a local diffeomorphism, so x_0 is isolated in $f^{-1}\{y_0\}$. So $f^{-1}\{y_0\}$ is discrete and compact, so finite, say equal to $\{x_1, x_2, \dots, x_N\}$. Take a small enough open set around each x_i so that f is a diffeomorphism on that open set. Inside each, take a compact set containing a neighborhood of x_i , and take the images of those compact sets. Because f is a local diffeomorphism, each of these images contains some relatively compact open neighborhood of y_0 . So their intersection does as well. But then that open neighborhood is evenly covered.

4.17. Consider the fundamental group action, a covering group action on \mathbb{R} . Every orbit is discrete. Pick some $t_0 \in \mathbb{R}$. It has a discrete orbit: some $t_i \in \mathbb{R}$, which we can write in order $t_i < t_{i+1}$. Each element of the group is uniquely determined by how it maps any one point, since they act without fixed points. Define elements γ_i by $\gamma_i(t_0) = t_i$. Define a group operation on these t_i , isomorphic to the fundamental group, by $t_i + t_j = t_k$ if $\gamma_j(\gamma_i(t_0)) = \gamma_k(t_0)$.

Each element of the fundamental group is a homeomorphism of \mathbb{R} , so an increasing or decreasing proper map $\mathbb{R} \rightarrow \mathbb{R}$. If decreasing, it has a fixed point by the intermediate value theorem, so it is an increasing function, so preserves the ordering of the t_i . Hence t_j takes $t_0 \mapsto t_0 + t_j = t_j$, and so sends t_1 to t_{j+1} , and so on. Hence the fundamental group is isomorphic to \mathbb{Z} . Each interval $[t_i, t_{i+1}]$ is mapped to the quotient space injectively except at the ends, so quotients to S^1 , a bijective continuous map from a compact Hausdorff space, so a homeomorphism by theorem 2.5 on page 20.

Bibliography

- [1] Mark Anthony Armstrong, *Basic topology*, Undergraduate Texts in Mathematics, Springer-Verlag, New York-Berlin, 1983, Corrected reprint of the 1979 original. MR 705632 [iii](#)
- [2] Glen E. Bredon, *Topology and geometry*, Graduate Texts in Mathematics, vol. 139, Springer-Verlag, New York, 1997, Corrected third printing of the 1993 original. MR 1700700 [iii](#)
- [3] Allen Hatcher, *Algebraic topology*, Cambridge University Press, Cambridge, 2002. MR 1867354 [iii](#)
- [4] Morris W. Hirsch, *Differential topology*, Graduate Texts in Mathematics, vol. 33, Springer-Verlag, New York, 1994, Corrected reprint of the 1976 original. MR 1336822 [35](#)
- [5] Stephen Huggett and David Jordan, *A topological aperitif*, revised ed., Springer-Verlag London, Ltd., London, 2009, With a foreword by Roger Penrose. MR 2483686 [iii](#)
- [6] Klaus Jänich, *Topology*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 1984, With a chapter by Theodor Bröcker, Translated from the German by Silvio Levy. MR 734483 [iii](#)
- [7] John McCleary, *A first course in topology*, Student Mathematical Library, vol. 31, American Mathematical Society, Providence, RI, 2006, Continuity and dimension. MR 2218695 [iii](#)
- [8] John W. Milnor, *Topology from the differentiable viewpoint*, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1997, Based on notes by David W. Weaver, Revised reprint of the 1965 original. MR 1487640 (98h:57051) [iii](#)
- [9] James R. Munkres, *Topology*, Prentice Hall, Inc., Upper Saddle River, NJ, 2000, Second edition of [MR0464128]. MR 3728284 [iii](#)
- [10] V. V. Prasolov, *Elements of combinatorial and differential topology*, Graduate Studies in Mathematics, vol. 74, American Mathematical Society, Providence, RI, 2006, Translated from the 2004 Russian original by Olga Sipacheva. MR 2233951 [iii](#)

- [11] ———, *Elements of homology theory*, Graduate Studies in Mathematics, vol. 81, American Mathematical Society, Providence, RI, 2007, Translated from the 2005 Russian original by Olga Sipacheva. MR 2313004 [iii](#)
- [12] Volker Runde, *A taste of topology*, Universitext, Springer, New York, 2005. MR 2155623 [iii](#)
- [13] Hajime Sato, *Algebraic topology: an intuitive approach*, Translations of Mathematical Monographs, vol. 183, American Mathematical Society, Providence, RI, 1999, Translated from the 1996 Japanese original by Kiki Hudson, Iwanami Series in Modern Mathematics. MR 1679607 [iii](#)
- [14] John Stillwell, *Poincaré, geometry and topology*, Henri Poincaré: science et philosophie (Nancy, 1994), Publ. Henri-Poincaré-Arch., Akademie Verlag, Berlin, 1996, pp. 231–240, 580. MR 1384994 [iii](#)
- [15] Tammo tom Dieck, *Algebraic topology*, EMS Textbooks in Mathematics, European Mathematical Society (EMS), Zürich, 2008. MR 2456045 [iii](#)

Index

- action
 - covering, **40**
 - group, **40**
 - by isometries, **41**
 - free, **40**
 - proper, **40**
- Alexander's horned
 - ball, **35**
 - sphere, **35**
- algebra
 - fundamental theorem of, **34, 39**
- alphabet, **53**
- amalgamation, **56**
- arithmetic progression, **13**

- Baire category theorem, **15**
- Baire space, **15**
- basis
 - topology, **4**
- boundary, **5**
- bouquet of circles, **22**

- closed
 - locally, **9**
 - set, **3**
- closed map, **23**
- closure, **4**
- cofinite topology, **3**
- comeager, **14**
- commutative
 - diagram, **33**
- compact, **11**
 - locally, **15**
- complement, **3**
- complex
 - projective space, **25**
- component
 - connected, **13**
 - path, **13**
- concatenation, **53**
- connected, **13**
 - path, **30**
 - simply, **30**
- connected component, **13**
- continuous, **17**
- convergence, **10**
- cover
 - open, **11**
- covering
 - group action, **40**
- covering map, **37**
- covering space
 - isomorphism, **47**
 - morphism, **47**
 - regular, **51**
 - universal, **47**

- deck transformation, **50**
- dense, **6**
 - everywhere, **14**
 - nowhere, **14**
- diagonal, **10**
- diagram, **33**
 - commutative, **33**
- diffeomorphism, **32**
- discrete
 - topology, **2**
- disjoint union, **9**

- evenly covered, **37**
- everywhere dense, **14**
- exterior, **5**

- fiber, **39**
- fiber bundle, **61**
- finitely presented, **53**
- free
 - group, **53**
 - group action, **40**
 - product, **55**
- fundamental
 - group, **29**

- theorem of algebra, 34, 39
- generated, 5
- generic, 15
- Golomb topology, 13
- group
 - action
 - covering, 40
 - action, 40
 - by isometries, 41
 - free, 40
 - proper, 40
 - free, 53
- half-open topology, 7
- Hausdorff, 9
- homeomorphic, 19
- homeomorphism, 19
- homotopic, 27
- homotopy, 27
 - equivalence, 33
 - relative, 27
- housed off, 9
- indiscrete
 - topology, 2
- interior, 5
- isometry
 - action, 41
- isomorphism
 - of covering spaces, 47
- join, 22
- Klein bottle, 25, 43, 50
- lift, 44, 59
- locally
 - closed, 9
 - compact, 15
 - topological space, 20
- locally path connected, 14
- logarithm, 46
- loop, 29
- manifold, 28
- manifold with corners, 28
- map
 - closed, 23
 - open, 23
 - strict, 23
- maximal subtree, 31
- Möbius strip, 21, 25, 42
- monodromy, 46
- morphism
 - of covering spaces, 47
- neighborhood, 4
- nowhere dense, 14
- null homotopic, 27
- open
 - cover, 11
 - set, 1
 - Zariski, 1
- open map, 23
- orbit, 40
- path, 27
- path component, 13
- path connected, 13, 30
 - locally, 14
- point, 1
- pointed space, 22
- preimage, 17
- presentation, 53
- product
 - topology, 7
- projection, 18
- projection map, 61
- proper
 - group action, 40
 - map, 20
- quaternionic projective space, 25
- quotient, 23
 - map, 23
 - space, 40
 - topology, 21
- real
 - projective space, 25
- reduce, 53
- reduced word, 55
- regular
 - covering space, 51
- relative homotopy, 27
- reparameterisation, 27
- restriction, 18
- Serre fibration, 59

- set
 - closed, [3](#)
 - open, [1](#)
- sheet, [37](#)
- simply connected, [30](#)
- space
 - quotient, [40](#)
 - topological, [1](#)
 - universal covering, [47](#)
 - with marked point, [22](#)
- spanned, [5](#)
- star shaped, [30](#)
- strict map, [23](#)
- subspace
 - topology, [2](#), [8](#)
- surface, [28](#)
- theorem
 - Baire category, [15](#)
 - fundamental, of algebra, [34](#), [39](#)
 - van Kampen, [52](#), [56](#)
- topological space, [1](#)
 - locally compact, [20](#)
- topology, [1](#)
 - basis, [4](#)
 - cofinite, [3](#)
 - discrete, [2](#)
 - Golomb, [13](#)
 - half-open, [7](#)
 - indiscrete, [2](#)
 - product, [7](#)
 - quotient, [21](#)
 - subspace, [2](#), [8](#)
 - Zariski, [1](#), [17](#), [18](#)
- tree, [31](#)
- universal covering space, [47](#)
- van Kampen's theorem, [52](#), [56](#)
- word, [53](#), [55](#)
 - reduced, [55](#)
- Zariski
 - open set, [1](#)
 - topology, [1](#), [17](#), [18](#)