

Title	A simple evaluation of the benefit of combined kinetic analysis of multiple injection dynamic PET scans
Authors	Gu, Fengyun;O'Sullivan, Finbarr;Muzi, Mark;Mankoff, David A.
Publication date	2019-11
Original Citation	Gu, F., O'Sullivan, F., Muzi, M. and Mankoff, D. A. 'A Simple Evaluation of the Benefit of Combined Kinetic Analysis of Multiple Injection Dynamic PET Scans'. 2019 IEEE Nuclear Science Symposium and Medical Imaging Conference (NSS/MIC), Manchester, UK, 26 Oct.-2 Nov. 2019, 1-3. doi: 10.1109/NSS/MIC42101.2019.9060073
Type of publication	Conference item
Link to publisher's version	https://ieeexplore.ieee.org/document/9060073 - 10.1109/NSS/MIC42101.2019.9060073
Rights	© 2019 IEEE. Personal use of this material is permitted. Permission from IEEE must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collective works, for resale or redistribution to servers or lists, or reuse of any copyrighted component of this work in other works.
Download date	2026-08-17 18:32:21
Item downloaded from	https://hdl.handle.net/10468/10582



UCC

University College Cork, Ireland
 Coláiste na hOllscoile Corcaigh

A Simple Evaluation of the Benefit of Combined Kinetic Analysis of Multiple Injection Dynamic PET Scans

Fengyun Gu, Finbarr O'Sullivan, Mark Muzi and David A. Mankoff

Abstract—The multiple injection dynamic Positron Emission Tomography (PET) scanning is used in the clinical management of certain groups of cancer patients and in medical research. The analysis of such studies can be approached in one of two ways: analyze individual injections separately to recover tracer kinetic information, or concatenate data from separate injections and carry out a combined analysis. Separate analysis offers some simplicity but may not be as efficient statistically. The mixture technique is readily implemented in a separated or combined analysis mode. We evaluate these approaches in a 1-D simulation setting matched to the mathematical complexity of PET. These simulations are largely guided by experience with breast cancer flow-metabolism mismatch studies using ^{15}O -Water (H_2O) and ^{18}F -Fluorodeoxyglucose (FDG). An efficient implementation in the R (an open-source environment) is used to implement simulations. The simulations evaluate mean square error (MSE) characteristics, for separate and combined analysis, both as a function of dose. The relationship between MSE characteristics of the underlying source distribution is described and the combined analysis is found to reduce MSE by between 18.1% and 33.85%. The quantitative advantages of combined approach have been demonstrated.

Index Terms—Multiple Injections, Combined Kinetic Modelling, Simulation, Mixture analysis

I. INTRODUCTION

THE multiple injection Positron Emission Tomography (PET) scanning have the ability to image two or more tracers in a single scan. Usually one tracer can just provide one kind of information, like ^{15}O -H₂O for measuring blood flow and ^{18}F -FDG for glucose metabolism. Such multitracers PET imaging would provide a wealth of complementary information for tumor grading and prognosis[7]. This technique is used in the clinical management of certain groups of cancer patients and in medical research, for example, ^{18}F -FDG and ^{18}F -FLT for brain tumors, ^{15}O -H₂O and ^{18}F -FDG for breast cancer and six different tracers for risk characterization in Sarcoma[5]. Data in these studies can be approached in one of two ways: (i) data from individual injections can be separately analysed to recover kinetic information corresponding to individual tracers, or (ii), studies can be concatenated and a combined analysis of the resulting data carried out. Initial efforts at analysing

multiple-tracer PET image were performed in phantoms[6], dual-tracer brain imaging [10, 11] and there are a series reports by group in Uath [9, 16, 2, 7, 8, 17]. But most of work has focused on the first approach by applying the compartmental modelling, which is not as efficient (statistically) as a combined analysis.

Additive mixture models [13, 14, 15] can be implemented in the separate and combined approach and it has been applied to the flow-metabolism mismatch study [3, 4, 12] in a breast cancer patient using ^{15}O -H₂O and ^{18}F -FDG. Motivated by experience with mixture model and flow-metabolism mismatch study in breast cancer, this work conducts some numerical analysis matched with this study. Our objective is to measure the quantitative advantages of combined analysis for multiple injections evaluated by analyzing MSE of underlying source distribution as a function of dose.

The basic theory and methodology is developed in section II. Results are presented in Section III. The paper concludes with discussion.

II. METHODOLOGY

We examine the efficiency of the mixture model estimation process for combined analysis in 1D simulation. The focus is on evaluation of MSE calculated in combined and separated analysis. This process, matched to the flow-metabolism studies with ^{15}O -H₂O and ^{18}F -FDG[12].

A. Analysis Approaches

In the multiple-injection PET studies, two approaches can be applied and one dual-tracer study example is presented in (Fig. 1).




Fig. 1: A dual-tracer study example

Manuscript received December 11, 2019. This work was supported in part by Science Foundation Ireland Grant No. PI-11/1027, by the National Cancer Institute USA grant R33-CA225310 and NIH/NCI R50-CA211270.

Fengyun Gu and Finbarr O'Sullivan are with the Department of Statistics, University College Cork, Ireland; Mark Muzi is with the Department of Radiology, University of Washington, USA; David A. Mankoff is with the Department of Radiology, University of Pennsylvania, USA. (e-mail address for correspondence: 116106718@uemail.ucc.ie).

- 1) *Combined analysis*: concatenate all of the time courses from multiple injections and analyse them together.
- 2) *Separate analysis*: analyse each time course one by one.

B. Mixture Model

In a study involving two tracer injections, the full voxel-level time-course can be approximated by an additive mixture model [13, 14, 15]:

$$z(x, t) \approx z(x, t|\alpha) = \sum_{k=1}^K \alpha_k(x) \mu_k(t) \quad (1)$$

where z are the full voxel-level data, $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_K)$ are positive mixing coefficients and $\mu = (\mu_1, \mu_2, \dots, \mu_K)$ are underlying time courses (sub-TACs), $t = (t_1, t_2, \dots, t_T)$.

In the combined analysis, mixing coefficient estimator ($\hat{\alpha}_c$) is calculated by using the full time-course data for both tracers. In the separated analysis, mixing coefficient estimator for H₂O ($\hat{\alpha}_s^1$) and FDG ($\hat{\alpha}_s^2$) is from part time-course data separately.

C. 1-D Simulation

In our simulation study (Fig.2), 1-D numerical phantom was conducted to examine and compare the efficiency of combined and separated analysis in multiple injection dynamic PET scans.




Fig. 2: 1D Simulation Process

1-D mixture model with K known components is employed to recover the mixing coefficients(α) from simulated image data.

$$\lambda(x, t) = \sum_{k=1}^K \alpha_k(x) \mu_k(t) \quad (2)$$

where λ is the true tissue source distribution of PET acquisition. Mixing coefficient estimators from combined ($\hat{\alpha}_c$)

and separated analysis ($\hat{\alpha}_s^1; \hat{\alpha}_s^2$) are recovered as introduced in section II-B. Estimated distributions ($\hat{\lambda}$) in combined and separate analysis are $\hat{\lambda}_c = \hat{\alpha}_c \mu$ and $\hat{\lambda}_s = (\hat{\alpha}_s^1 \mu^1, \hat{\alpha}_s^2 \mu^2)$

MSE can be calculated as below:

$$\begin{cases} MSE1 = E[(\hat{\lambda}_s - \lambda)^2] \\ MSE2 = E[(\hat{\lambda}_c - \lambda)^2] \end{cases} \quad (3)$$

III. RESULTS

The MSE of the source distribution estimation as the function of the dose levels from combined and separated approach in 1-D simulation are presented in Fig.3. It illustrates errors from combined and separated analysis decrease with the dose. It is demonstrated that combined analysis is more efficient than the separate analysis and the magnitude of the improvements ranges from 18.10% to 33.85% for two injections with different dose levels. These results show combined analysis has quantitative advantages and it also validates the efficiency of mixture model approach for multiple injections.




Fig. 3: MSE Comparison in 1D Simulation Study

TABLE I: Improvements at different dose levels

dose levels	dose1	dose2	dose3	dose4	dose5	dose6	dose7
improvements(%)	18.10	21.78	24.60	26.56	30.16	32.67	33.85

IV. DISCUSSION

The quantitation of data from multiple-injection PET studies is enhanced by combining information from separate injections. Mixture modelling provides simple mechanism to realise the combined analysis with long time-frames. It has a potential to process large scale dynamic data with the announcement of the next generation of total-body PET scanners[1] for research and clinical practice. More detailed examination of the benefits of combined analysis in different settings would be helpful.

REFERENCES

- [1] R. D. Badawi, H. Shi, P. Hu, S. Chen, T. Xu, P. M. Price, Y. Ding, B. A. Spencer, L. Nardo, W. Liu *et al.*, "First human imaging studies with the explorer total-body pet scanner," *Journal of Nuclear Medicine*, vol. 60, no. 3, pp. 299–303, 2019.
- [2] N. F. Black, S. McJames, and D. J. Kadrmas, "Rapid multi-tracer pet tumor imaging with ^{18}F -fdg and secondary shorter-lived tracers," *IEEE Transactions on Nuclear Science*, vol. 56, no. 5, pp. 2750–2758, 2009.
- [3] L. K. Dunnwald, R. K. Doot, J. M. Specht, J. R. Gralow, G. K. Ellis, R. B. Livingston, H. M. Linden, V. K. Gadi, B. F. Kurland, E. K. Schubert *et al.*, "Pet tumor metabolism in locally advanced breast cancer patients undergoing neoadjuvant chemotherapy: value of static versus kinetic measures of fluorodeoxyglucose uptake," *Clinical Cancer Research*, vol. 17, no. 8, pp. 2400–2409, 2011.
- [4] L. K. Dunnwald, J. R. Gralow, G. K. Ellis, R. B. Livingston, H. M. Linden, J. M. Specht, R. K. Doot, T. J. Lawton, W. E. Barlow, B. F. Kurland, E. K. Schubert, and D. A. Mankoff, "Tumor metabolism and blood flow changes by positron emission tomography: relation to survival in patients treated with neoadjuvant chemotherapy for locally advanced breast cancer," *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, vol. 26, no. 27, pp. 4449–4457, 09 2008. [Online]. Available: <https://www.ncbi.nlm.nih.gov/pubmed/18626006>
- [5] J. F. Eary, J. M. Link, M. Muzi, E. U. Conrad, D. A. Mankoff, J. K. White, and K. A. Krohn, "Multiagent pet for risk characterization in sarcoma," *Journal of Nuclear Medicine*, vol. 52, no. 4, pp. 541–546, 2011.
- [6] S. C. Huang, R. E. Carson, E. J. Hoffman, D. E. Kuhl, and M. E. Phelps, "An investigation of a double-tracer technique for positron computerized tomography," *The Journal of Nuclear Medicine*, vol. 23, no. 9, pp. 816–822, 1982.
- [7] D. J. Kadrmas and J. M. Hoffman, "Methodology for quantitative rapid multi-tracer pet tumor characterizations," *Theranostics*, vol. 3, no. 10, p. 757, 2013.
- [8] —, "Methodology for quantitative rapid multi-tracer pet tumor characterizations," *Theranostics*, vol. 3, no. 10, p. 757, 2013.
- [9] D. J. Kadrmas and T. C. Rust, "Feasibility of rapid multitracers pet tumor imaging," *IEEE transactions on nuclear science*, vol. 52, no. 5, pp. 1341–1347, 2005.
- [10] R. Koeppe, E. Ficaro, D. Raffel, S. Minoshima, and M. Kilbourn, "Temporally overlapping dual-tracer pet studies," in *Quantitative functional brain imaging with positron emission tomography*. Elsevier, 1998, pp. 359–366.
- [11] R. A. Koeppe, D. M. Raffel, S. E. Snyder, E. P. Ficaro, M. R. Kilbourn, and D. E. Kuhl, "Dual-[^{11}C] tracer single-acquisition positron emission tomography studies," *Journal of Cerebral Blood Flow & Metabolism*, vol. 21, no. 12, pp. 1480–1492, 2001.
- [12] D. A. Mankoff, L. K. Dunnwald, J. R. Gralow, G. K. Ellis, E. K. Schubert, J. Tseng, T. J. Lawton, H. M. Linden, and R. B. Livingston, "Changes in blood flow and metabolism in locally advanced breast cancer treated with neoadjuvant chemotherapy," *Journal of Nuclear Medicine*, vol. 44, no. 11, pp. 1806–1814, 2003.
- [13] F. O'sullivan, "Locally constrained mixture representation of dynamic imaging data from pet and mr studies," *Biostatistics*, vol. 7, no. 2, pp. 318–338, 2005.
- [14] F. O'Sullivan, M. Muzi, D. A. Mankoff, J. F. Eary, A. M. Spence, and K. A. Krohn, "Voxel-level mapping of tracer kinetics in pet studies: a statistical approach emphasizing tissue life tables," *The annals of applied statistics*, vol. 8, no. 2, p. 1065, 2014.
- [15] F. O'sullivan, M. Muzi, A. M. Spence, D. M. Mankoff, J. N. O'sullivan, N. Fitzgerald, G. C. Newman, and K. A. Krohn, "Nonparametric residue analysis of dynamic pet data with application to cerebral fdg studies in normals," *Journal of the American Statistical Association*, vol. 104, no. 486, pp. 556–571, 2009.
- [16] T. Rust and D. Kadrmas, "Rapid dual-tracer psm+ atsm pet imaging of tumour blood flow and hypoxia: a simulation study," *Physics in Medicine & Biology*, vol. 51, no. 1, p. 61, 2005.
- [17] J. L. Zhang, A. M. Morey, and D. J. Kadrmas, "Application of separable parameter space techniques to multi-tracer pet compartment modeling," *Physics in Medicine & Biology*, vol. 61, no. 3, p. 1238, 2016.