

Title	Sugar-sweetened beverage consumption and association with weight status in Irish children: A cross-sectional study prior to the introduction of a government tax on sugar-sweetened beverages
Authors	Harrington, Janas M.;Perry, Catherine;Keane, Eimear;Perry, Ivan J.
Publication date	2020-05-28
Original Citation	Harrington, J. M., Perry, C., Keane, E. and Perry, I. J. (2020) 'Sugar-sweetened beverage consumption and association with weight status in Irish children: A cross-sectional study prior to the introduction of a government tax on sugar-sweetened beverages', Public Health Nutrition, 23(12), pp. 2234-2244. doi: 10.1017/ S1368980020000014
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1017/S1368980020000014
Rights	© 2020, the Authors. Published by Cambridge University Press on behalf of The Nutrition Society. This material is free to view and download for personal use only. Not for re-distribution, re-sale or use in derivative works.
Download date	2025-08-01 09:46:14
Item downloaded from	https://hdl.handle.net/10468/12273



University College Cork, Ireland Coláiste na hOllscoile Corcaigh NS Public Health Nutrition

Sugar-sweetened beverage consumption and association with weight status in Irish children: a cross-sectional study prior to the introduction of a government tax on sugar-sweetened beverages

Janas M Harrington^{1,*} , Catherine Perry², Eimear Keane³ and Ivan J Perry¹ ¹School of Public Health, University College Cork, Cork City T12 XF62, Ireland: ²Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, UK: ³Department of Public Health, HSE Mid-West, Ireland

Submitted 25 January 2019: Final revision received 9 December 2019: Accepted 2 January 2020: First published online 28 May 2020

Abstract

Objective: To provide baseline evidence of sugar-sweetened beverage (SSB) consumption in a sample of Irish children prior to the introduction of the SSB tax; to identify the energy contribution of SSB to daily energy intake; and to explore the association between SSB consumption and overweight/obesity.

Design: Cross-sectional study.

Setting: Primary schools in Cork, Ireland in 2012.

Participants: 1075 boys and girls aged 8–11 years. SSB consumption was assessed from 3-d food diaries. BMI was used to define obesity (International Obesity Taskforce definitions). Plausible energy reporters (n 724, 68% of total sample) were classified using Schofield equation.

Results: Eighty-two per cent of children with plausible energy intake consumed SSB. Mean energy intake from SSB was 485 kJ (6% of total kJ). Mean kilojoules from SSB increased with weight status from 443 kJ for normal-weight children to 648 kJ for children with overweight/obesity (5·8 and 7·6% of total kJ, respectively). Mean SSB intake was significantly higher in children with overweight/ obesity than normal-weight children (383 and 315 ml/d). In adjusted analyses, children consuming >200 ml/d had an 80% increased odds of overweight/obesity compared to those consuming <200 ml/d (OR 1·8, 95% CI 1·0, 3·5). Family socioeconomic status and lifestyle determinants, including frequency of takeaway consumption and TV viewing, were also significantly associated with SSB consumption.

Conclusions: SSB account for a substantial proportion of daily energy intake and are significantly associated with child overweight/obesity. This study provides baseline data from a sample of children from which the impact of the SSB tax can be benchmarked.

Keywords Child obesity Sugar-sweetened beverages Sugar tax Child weight

Ireland has an unacceptably high prevalence of childhood obesity, with an estimate that 25 % of 9-year-olds live with overweight/obesity⁽¹⁾. Though there are some indications that the prevalence may be stabilising⁽²⁾, it remains at a level that imparts significant health-related risk.

The current epidemic of overweight and obesity represents a public health crisis with the potential to reverse recent favourable trends in life expectancy^(2,3) and undermine the financial viability of health systems worldwide. While effective government policies and actions are essential to increase the healthiness of food environments, monitoring the degree of implementation of the policies and actions and impact of these policies is essential to ensuring progress towards better nutritional health. One policy measure introduced by the Irish government was a tax on sugar-sweetened beverages (SSB) from 1 May 2018. Though evidence from other sectors, including alcohol and tobacco, supports the use of fiscal measures as legitimate strategies in the public health toolkit, the support for and effectiveness of this tax is controversial and is a subject of debate both nationally and internationally^(4–6). Several countries have chosen to use taxes on foods and non-alcoholic beverages in an attempt to improve the quality of people's diets^(7,8) and reduce the prevalence of

obesity. Some recent evidence from a number of countries indicates positive population health and economic benefits of such a $tax^{(5,9-11)}$.

Essential to measuring the impact of this fiscal measure is a baseline from which to benchmark. The aim of the current article is to provide evidence of consumption levels of SSB in Ireland prior to the introduction of a tax; to identify the energy contribution of SSB to overall energy intake; and to explore the association between SSB consumption and weight in a cross-sectional sample of children living in Cork, Ireland. This will provide baseline data from which the tax impact on SSB consumption can be assessed in the short and long terms.

Methods

Study design and sample

Details of the Cork Children's Lifestyle Study (CCLaS) are described elsewhere⁽¹²⁾. In summary, the study recruited children in third and fourth classes (years 5 and 6 of enrolment to primary school) between April 2012 and June 2013. Schools from the urban area were recruited using probability proportionate-to-size and purposive sampling. All schools in the rural area were invited to participate. At the school level, twenty-seven out of forty-six schools participated (response rate 58.6%), and 1075 out of 1641 children participated (response rate 65.5%) in the study. The present study only includes children with completed food diaries and who had a plausible energy intake (n724).

Dietary assessment

Participating children completed a consecutive 3-d food diary developed for the purposes of this study⁽¹²⁾. Diaries were completed over weekdays (Monday–Friday) and weekend days (Saturday and Sunday). Overall, 45% (n 327) of children with plausible energy intake had at least one weekend data collection day. All data reported here refer to a daily average of 3-d consumption. Each food item the child reported in their food diary was entered into the nutritional analyses software, NetWISP version 4 (Tinuviel software). All diaries were completed and data entered according to a specifically developed standard operating procedure (SOP). Ten per cent of diaries were double-entered, and all data cleaning was conducted according to a study-specific SOP.

Assessment of sugar-sweetened beverage intake

SSB intake was assessed from food diaries. The type and quantity of SSB was reported by the child. The nutritional software's output provided information for quantity (ml), kilojoules and sugars per day from each food/drink type. SSB consumers were identified using NetWISP's food item categorisation. The McCance and Widdowson 7th edition⁽¹³⁾ and the Irish Food Composition nutrient databases⁽¹⁴⁾ were used for analysis in NetWISP. Diet carbonated soft drinks were disaggregated from the beverage category and excluded from this SSB category. The beverages that contributed to overall SSB are carbonated soft drinks, juices or cordials and energy drinks. Each beverage category was checked to verify that the drink items contained within it had added sugar. All beverages included contained 5 g sugar/100 ml. The volume (ml) of each beverage category was summed to generate an overall SSB amount, and this was then divided by the number of food diary days the child actually completed. SSB consumption was analysed as a continuous variable (ml/d), and for multivariate regression analyses, it was collapsed into a categorical variable: 'non consumers', those who had no reported SSB consumption over the food diary days; 'low consumers', less than one standard household glass (200 ml/d); and 'high consumers', those with reported consumption >200 ml/d. The Irish Healthy Eating Guidelines suggest a serving size of 200 ml cup to measure portion sizes. Furthermore, the Irish guidelines suggest that 200 ml is an identifiable portion size for the general population, which is useful for public health $messages^{(15)}$.

In this sample, the average amount of sugar (g) that a 200-ml cup of SSB provides is 10.3 g. For ease of interpretation, 200 ml was chosen rather than an uneven number. The WHO recommends an intake of free sugars of <10 % of total energy intake⁽¹⁶⁾. The Irish energy recommendation for 5–12-year-old children is 5858–9205 kJ/d⁽¹⁵⁾.

Energy misreporting

An energy intake to BMR ratio was calculated for each child to identify under- or plausible reporters. BMR was estimated using the method outlined by Schofield *et al.*⁽¹⁷⁾. Cut-off values for energy intake to BMR were defined by an equation developed by Goldberg *et al.*⁽¹⁸⁾, with updated cut-off points for children, defined by Black *et al.*⁽¹⁹⁾. Implausible reporters, based on Black's equation, were excluded from the analysis (*n* 348). Data presented are separated for the full study sample, plausible and under-reporters (Table 1), while subsequent tables are for plausible energy reporters only.

Anthropometric measurements and obesity definition

Children's height and weight were measured by trained researchers using standardised methods⁽¹²⁾. Age- and sex-specific International Obesity Taskforce (IOTF) definitions were used to categorise children as normal weight or overweight/obese^(20,21). Data for children's measured BMI were available for 99.3 % (*n* 1068) of the sample.

NS Public Health Nutrition

Table 1 Sociodemographic and lifestyle characteristic of the Cork Children's Lifestyle Study population by energy reporting

	Total po	pulation*	Plau energy	sible** reporters	Under-re		
Factor	n	%	<u> </u>	%	n	%	<i>P</i> -value
	1068		717	68.0	337	32.0	
SSB consumption**	1000		7.17	000	007	02.0	
Non-consumer	225	21.4	130	18.2	95	28.2	
Low consumer. <200 ml/d	339	32.2	208	29.1	131	38.9	
High consumer. >200 ml/d	489	46.4	378	52.8	111	32.9	0.000
Child characteristics							
Weight category							
Normal	797	74.6	586	81.7	201	59.6	
Overweight/obese	271	25.4	131	18.3	136	40.4	0.000
Sex							
Boy	620	58·1	427	59.6	184	54.6	
Girl	448	42.0	290	40.5	153	45.4	0.129
Age (years)							
8–9	560	52.4	379	52.9	175	51·9	
10–11	508	47.6	338	47.1	162	48·1	0.778
Parent-reported child TV viewing							
<1 h/d	224	22.1	158	23.1	63	19.3	
1–3 h/d	614	60.4	405	59.3	207	63·5	
>3 h/d	178	17.5	120	17.6	56	17.2	0.346
Child-reported TV viewing							
<1 h/d	516	48.9	341	48.0	168	50.0	
1–3 h/d	401	37.9	266	37.4	131	39.0	
>3 h/d	138	13.0	101	14.2	36	10.7	0.464
Meeting MVPA targets‡							
Yes	183	22.2	139	24.9	43	16.7	
No	643	77.9	421	75.2	215	83.3	0.009
Parent-reported family takeaways							
<once per="" td="" week<=""><td>847</td><td>84.8</td><td>570</td><td>84.8</td><td>270</td><td>84.4</td><td></td></once>	847	84.8	570	84.8	270	84.4	
≥Once per week	152	15·2	102	15·2	50	15.6	0.855
Parental characteristics							
Ethnicity							
Irish	828	87.2	575	89.0	247	82.9	
Other	122	12.8	71	11.0	51	17.1	0.009
Family type							
Single-parent	198	18·5	127	18.9	67	20.7	
Two-parent	803	75.2	544	81·1	257	79.3	0.513
Parental education							
Primary/lower secondary only	106	10.8	68	10.4	34	10.7	
Higher secondary	189	19.3	123	18.8	65	20.4	
Certificate/diploma	283	28.9	193	29.5	90	28.3	
Tertiary	401	41.0	271	41.4	129	40.6	0.928

SSB, sugar sweetened beverage.

*Excludes n 7; no BMI data.

**Excludes n 15; no food diary data.

 $\dagger \ensuremath{\textit{P}}\xspace$ value is the association between factors and energy reporting.

[‡]Total study population is 1075, however data presented in this table relate only for children with measured BMI data. Children missing data were excluded from this table (*n*7). Data presented for participants with valid objectively measured accelerometry data only.

Covariates

Additional socioeconomic and lifestyle behaviours were provided by participants in self-reported questionnaires and objective measures of physical activity. Self-reported child questionnaires were completed in school during class time, with researchers providing assistance where necessary. Parent-reported questionnaires were completed in the home and returned by the children.

Demographics

Sex (girl or boy) was recorded by trained researchers during physical measurements. Child's age was calculated using the date of physical measurement and date of birth (parent-reported). Parent-reported highest level of maternal education was used as a proxy measure of socioeconomic status. The variable was coded as primary/lower secondary education, higher secondary education, post-secondary education, tertiary education. Ethnic background was reported in the parental questionnaire. This was coded as a binary variable as Irish or non-Irish. Family type was recorded by the parental questionnaire and coded as single-parent or two-parent.

Parent-reported family takeaway consumption

The frequency of takeaway consumption was assessed from the parent-reported questionnaire. Parents were asked to report how often they ordered family takeaways.

This was coded as a categorical variable: ' \leq once a week' and '>once a week'.

Parent-reported child TV-viewing habits

Child TV-viewing habits were assessed from the parentreported and child-reported questionnaires.

Parents reported how much time the child spends watching TV on a 'normal weekday during term time'. This was coded as a categorical variable: '<1 h/d', '1–3 h/d' and ' \geq 3 h/d'. Children reported how much time they watch TV each day. This was coded as: '<1 h/d', '1–3 h/d' and ' \geq 3 h/d'.

Physical activity

Details of the objectively measured physical activity levels have been previously recorded^(22,23). In summary, freeliving physical activity was measured over 7 consecutive days using a wrist-worn validated tri-axial Geneactiv accelerometer^(24,25). The accelerometers were set to record at 100 Hz for 7 d using the 'on button press' setting of Geneactiv software, version 2.2. The accelerometers were placed on the non-dominant hand of each participating child. Children were asked to wear the accelerometers 24 h a day, each day.

Parent-reported questionnaire data were used to identify the most frequent waking time and bedtime on week days and weekend days. These data were then used to estimate the number of waking hours each day. The mean waking time was 14 h/d. To be included in the current analysis, children needed to have recorded ≥ 600 min of waking time data each day. For children included in the analysis who had missing wear time information, the data were scaled to full waking time. Non-wear time was determined using an algorithm by van Hees *et al.*⁽²⁶⁾. Children who engaged in ≥ 60 min of moderate to vigorous physical activity (MVPA) on each of the 7 d were categorised as meeting WHO MVPA recommendations⁽²⁷⁾.

Ethical considerations

All study procedures were approved by the local research ethics committee. Only children who provided assent and whose parents/guardians provided written informed consent participated in the study. Feedback on the physical measurements was provided to all parents of participating children in the form of a letter.

Data analysis

Statistical analyses were completed using STATA/IC 13.1 Descriptive results are presented as mean (SD) or median (IQR) for continuous data, and frequency (percentage) for categorical data. *t*-Tests and ANOVA were used to compare mean differences in continuous variables, while

 χ^2 tests were used to determine differences in categorical data.

Multivariate analyses

Unadjusted estimates of the association between SSB consumption and child weight status were performed using a binary logistic regression. Partially adjusted estimates were obtained by adjusting for sociodemographic variables and lifestyle behaviours separately in a binary logistic regression model. Covariates that were statistically significant in the partially adjusted logistic regression analyses were retained for the multivariate fully adjusted logistic analyses to assess the association between SSB consumption and child overweight/obesity for low consumers (reference group), non-consumers and high consumers.

Using BMI as a continuous variable, the distribution of BMI across SSB consumption quintiles was assessed using fully adjusted kernel density estimates at the upper cut-points of SSB consumption quintiles.

Results

Descriptive details of the CCLaS study population have been previously reported⁽¹²⁾. Table 1 shows the descriptive characteristics of all participants, participants with plausible energy intake and those classified as under-reporters with measured height, weight and BMI data (n 1068).

Overview of participants in this study

According to IOTF obesity classification, a quarter of children had overweight/obesity (25 %, n 271). The majority of participating parents were female (58%, n 620), Irish (87 %, *n* 828) and from two-parent families (75 %, *n* 803). Over 40% of the children's parents (n 401) had a tertiary qualification. Sixty per cent of parents (n 614) reported that their children watched between 1-3 h of TV per day, while 38% (*n* 401) of children self-reported watching 1–3 h TV per day. The majority of children (78%, n 643) did not meet the daily recommendation for MVPA, while 15% (*n* 152) of parents reported consuming a takeaway more than once a week (Table 1). Almost half of children (46%, n 489) were classified as high consumers of SSB (>200 ml/d). Significant differences were seen in energy reporting (plausible v. under-reporting) for SSB consumption status (P < 0.0001), weight category (P < 0.0001), meeting MVPA targets (P = 0.009) and ethnicity (P = 0.009)(Table 1).

Child weight status

Child's physical activity, parent-reported child TV viewing and level of SSB consumption were significantly associated with child weight status (Table 2). Significantly more children with overweight/obesity, compared to children with normal weight, watched TV >3 h daily (26 v. 17 % daily),

Public Health Nutrition

	Plausible reporters*‡		Normal weight*		Overweight/ obese*		<i>P</i> -value‡	Non- consumers§		Low consumers§		High consumers§		
	n	%	п	%	п	%		п	%	п	%	п	%	<i>P</i> -value
	586		471	80.4	115	19.6		130	18.2	208	29.1	378	52.8	
Gender					-				-		-			
Boy	352	60.1	285	60.5	67	58.3	0.659	75	57.7	122	58.7	230	60.9	0.772
Girl	234	39.9	186	39.4	48	41.7		55	42.3	86	41.4	148	39.2	
Weight category														
Normal	471	80.4	_		-		_	114	87.7	179	86.1	292	77.3	0.005
Overweight/obese	115	19.6	_		_			16	12.3	29	13.9	86	22.7	
Age														
8–9 years	307	52.4	247	52.4	60	52.2	0.959	72	55.4	108	51.9	199	52.7	0.841
10-11 years	279	47.6	224	47.6	55	47.8		58	44.6	100	48·1	179	47.4	
Parent-reported child TV viewing (h	/d)													
<1	[´] 118	21.2	101	22.5	17	15.5	0.042	39	31.5	54	27.7	64	17.6	0.005
1–3	335	60.0	271	60.5	64	58.2		70	56.5	109	55.9	226	62.3	
>3	105	18.8	76	17.0	29	26.4		15	12.1	32	16.4	73	20.1	
Child-reported TV viewing (h/d)														
<1	273	46.9	224	48.0	49	42.6	0.512	67	52.3	110	53.4	163	43.4	0.153
1–3	222	38.1	176	37.7	46	40.0		44	34.4	70	34.0	152	40.4	
>3	84	14.4	64	13.7	20	17.4		17	13.3	24	11.7	60	16.0	
Meeting MVPA targets	•••		0.					••						
Yes	110	24.1	98	26.4	12	14.0	0.015	29	28.4	49	30.3	61	20.7	0.050
No	347	75.9	273	73.6	74	86.1	0010	73	71.6	113	69.8	234	79.3	0 000
Parent-reported family takeaways	047	100	210	700	14	001		70	/10	110	000	204	100	
<pre>/Once per week</pre>	453	82.8	369	84.1	84	77.8	0.121	116	93.6	168	87.5	285	80.3	0.001
>Once per week	400 Q/	17.2	70	16.0	24	22.2	0.121	8	6.5	24	12.5	70	19.7	0.001
SSB consumption	54	17.2	70	10.0	24			0	0.0	24	12.0	70	10.7	
Non-consumer	_		_		_		0.010	_		_		_		_
< 200 m/d	200	25.5	174	26.0	29	24.4	0.010	_		_		_		_
\geq 200 ml/d	200	64.5	207	62.1	20	24.4		-		_		_		
>200 m/u	304	04.5	291	03.1	07	75.7		-		_		_		
Irich	175	00.9	200	00.7	05	01.4	0 0 0 0	00	01.0	160	00.4	215	01.6	0.007
Other	4/5	90.8	300	90.7	95	91.4	0.030	99	10.0	100	09·4	315	91.0	0.007
	40	9.2	39	9.3	9	0.1		23	18.9	19	10.0	29	ö ∙4	
	100	10.0	00	00 5	10	17.0	0.400	10	45.4	00	10.0	70	01 5	0 1 0 7
Single-parent	108	19.9	90	20.5	18	17.3	0.469	19	15.1	32	16.8	76	21.5	0.187
I wo-parent	436	80.2	350	79.6	86	12.1		107	84.9	159	83.3	211	78.5	
Parental education	00	4 4 -	50	11.0	40	10.1	0.004	0	4.0		44 -	10	44 7	0 0000
Primary/lower secondary only	62	11.7	50	11.6	12	12.1	0.801	6	4.8	22	11.7	40	11.7	0.0000
Higher secondary	105	19.8	82	19.0	23	23.2		18	14.5	32	17.0	73	21.4	
Certificate/diploma	165	31.1	136	31.6	29	29.3		28	22.6	51	27.1	114	33.3	
lertiary	198	37.4	163	37.8	35	35.4		72	58.1	83	44.2	115	33.6	
SSB ml*														
Mean		328.7		315.4		383.1	0.0054	-			113.8		447.0	0.0000
SD		234.4		222.0		273.9					49.9		210.7	,
Median		268.7		262.7		316.7					110.0		390.7	
IOR		312.7		293.0		326.7					92.7		290.7	

JM Harrington et al.

Public Health Nutrition

Table 2 Continued

	Plausible reporters*‡		Plausible reporters*‡ Normal we		Overweight/ ۱ weight* obese*		<i>P</i> -value‡	Non- consumers§		Low consumers§		High consumers§		19 DC
	n	%	п	%	n	%		n	%	n	%	n	%	P-value
SSB kJ*														
Mean		116.3		106.8		155.3	0.0005	-			43.2		156.6	0.0000
SD		134.1		98.4		224.6					44.6		149.1	
Median		84.3		78.0		99.7					34.2		123.3	
IQR		114.3		110.3		123.0					32.9		114.7	
SSB %kJ*														
Mean		6.1		5.8		7.6	0.0025	-			2.6		8.2	0.0000
SD		5.7		5.0		7.9					2.6		6.0	
Median		4.6		4.4		5.3					2.0		6.7	
IQR		6.3		6.2		7.5					2.2		6.2	
SSB sugars*														
Mean		27.1		24.8		36.6	0.0006	-			10.1		36.5	0.0000
SD		33.1		24.1		56.0					11.3		37.3	
Median		18.5		17.5		22.8					7.7		28.8	
IQR		28.3		28.4		32.0					10.1		28.1	
SSB %sugars*														
Mean		22.1		21.1		26.1	0.0049	-			11.0		28.2	0.0000
SD		17.0		16.3		19.3					10.5		16.8	
Median		18.8		18.0		21.4					8.6		24.7	
IQR		22.6		22.0		23.3					11.6		23.0	

MVPA, moderate to vigorous physical activity.

*Plausible energy reporters and SSB consumers only.

†SSB is defined as carbonated soft drinks (excluding diet drinks), cordials/squash/juices and energy/sports drinks. % denotes the percentage contribution of SSB to overall mean daily energy intake and mean daily sugar intake. ‡*P*-value is the association between SSB consumer category and the factors; χ^2 test for categorical data and *t* test for continuous data.

§Plausible energy reporters only.

||Low consumer <5 g sugars/100 ml SSB, high consumer \geq 5 g sugars/100 ml SSB; non-consumers *n* 130 (18·2 %).

nS

2240

did not meet MVPA targets (86 *v*. 74%) and consumed \geq 200 ml SSB (76 *v*. 63%). The volume of consumption differed statistically significantly in overweight/obese children compared to normal-weight children (383 *v*. 315 ml, P = 0.0054) (Table 2).

Child's sugar-sweetened beverage consumption

The majority of participants (82%) were consumers of SSB (Table 1). The average SSB consumption was 328 ml/d (a standard unit of SSB available for purchase is 330 ml). SSB contributed a mean of 6% of total energy intake and 22% of total sugar intake. No gender differences in SSB consumption were evident. However, consumption differed significantly by child's weight status. Mean energy intake from SSB increased from 448 to 649 kJ for children who were normal weight compared with those with overweight/obesity, equating to 5.8 and 7.6% of total energy intake, respectively. Mean intake volumes were significantly higher in children with overweight/obesity compared to normal-weight children. Average consumption was 315.5 and 383.1 ml/d for normal-weight children and children with overweight/obesity, respectively. Sugar intake from SSB and per cent contribution of sugars from SSB to total sugar intake was significantly higher in children with overweight/obesity compared to normal-weight children (Table 2).

Twenty-nine per cent of children (*n* 208) were 'low SSB consumers' (\leq 200 ml/d), 18% (*n* 130) 'non-consumers' and 53% (*n* 380) 'high SSB consumers' (>200 ml/d). Significantly more children with overweight/obesity compared to normal-weight children were high consumers of SSB (76 v. 63%) (Table 2). Child's SSB consumption status was significantly associated with parent-reported child TV viewing (P = 0.005), meeting physical activity recommendation (P = 0.005) and eating takeaway meals on a weekly basis (P = 0.001). Children whose parents reported they watched >3 h TV per day were more likely to be high SSB consumers (*n* 15, 12%). Of those reporting eating takeaway meals more than once per week, significantly more were high SSB consumers than low consumers

(20 v. 6 %). Ethnicity and parental education are associated with SSB consumption (Table 2). A higher proportion of 'Irish' participants were classified as high SSB consumers, while a higher proportion of 'other ethnic backgrounds' were classified as non- or low consumers (Table 2). The highest proportion of non-consumers of SSB were among children of parents with higher levels of education (58 v. 5% for tertiary education v. primary or secondary only) (Table 2).

Quantity of sugar-sweetened beverage consumption in plausible energy reporters

Compared to the total sample (detailed in Table 1), a higher proportion of plausible energy reporters were 'high SSB consumers' (Table 1) (53 v. 46%). Across weight categories, a higher percentage of plausible reporters were normal weight compared to the total sample. Plausible reporters were less likely to have overweight/obesity (18%) compared to under-reporters (40%). A subsequent analysis focused only on plausible energy reporters.

Table 3 displays the contribution of each SSB type to the overall SSB intake in millilitres, kilojoules and sugars. It also shows the pattern of consumption over weekdays and weekends. The mean contribution of fizzy drinks to overall SSB kilojoules differed between weekdays and weekends. Weekend contribution was 50.9 %, while the mean contribution during weekdays was 30.6 %.

Unadjusted and partially adjusted analysis

Crude unadjusted estimates of the association between SSB consumption and child's weight status are presented in Table 4. High SSB consumers have an increased odds of being overweight/obese (OR 1.8, 95% CI 1.1, 2.9) than low SSB consumers. This association remained stable in separate analyses after adjusting for individual characteristics (child's age and gender), family characteristics (parental ethnicity, maternal education, marital status and family type) and child's lifestyle behaviours (MVPA, takeaway consumption, TV viewing and average daily kJ intake) (Table 4).

Table 3 Patterns of SSB consumption by SSB components by weekday and weekends*

		Weekdays (<i>n</i> 586)						Weekends (n 274)						
	Carbonated fizzy drinks		Juices/ squash/ cordials		Energy drinks		Carbonated fizzy drinks		Juices/ squash/ cordials		Energy drinks			
	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD		
% contribution to overall SSB ml % contribution to overall SSB kJ % contribution to overall SSB sugars (g)	26·3 30·6 32·2	39·5 46·0 47·9	70·7 66·8 64·9	54∙4 57∙8 58∙8	5·3 5·0 5·6	21·3 20·0 24·0	48·0 50·9 53·1	69∙5 72∙8 74∙2	48 42·6 40·4	67∙1 66∙7 68∙5	3∙1 3∙0 3∙5	16·6 16·2 18·7		

*Weekdays, Monday–Friday; weekends, Saturday and Sunday. *n* is the number of children with food diary data from at least 1 weekday and the number of children with food diary data from at least 1 weekend day, and who were plausible energy reporters and SSB consumers.

Table 4 Unadjusted and partially adjusted logistic regression on the association between SSB consumption (ref: low consumers, <200 ml/d) and overweight/obesity

Variables in the modelOR95 % ClPUnadjusted-0.012Low consumers-0.012Non-consumers1.81.1, 2.9Partially adjusted model with different child characteristics-0.011Child's age, gender-0.011Low consumers-0.011Non-consumers0.90.4, 1.7High consumers-0.011Non-consumers0.90.4, 1.7High consumers-0.058Non-consumers0.60.3, 1.4Uw consumers0.60.3, 1.4Low consumers0.60.3, 1.4High consumers-0.022Non-consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilojule intake-0.022Non-consumers0.80.4, 1.7High consumers0.90.4, 1.8High consumers1.71.1, 2.8Parent characteristics-0.007Non-consumers0.90.4, 1.8Low consumers0.90.4, 1.8High consumers1.10.5, 2.2Non-consumers2.21.3, 3.7Parent characteristics-0.003Non-consumers0.90.4, 1.7High consumers2.21.3, 3.7Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers0.90.4, 1.7High consumers1.10.5, 2.2 <tr< th=""><th></th><th colspan="4">Overweight/obesity</th></tr<>		Overweight/obesity			
Unadjusted	Variables in the model	OR	95 % CI	Р	
Low consumers-0.012Non-consumers0.80.4, 1.6High consumers1.81.1, 2.9Partially adjusted model with different child characteristicsChild's age, genderChild's age, gender-0.011Non-consumers0.90.4, 1.7Non-consumers0.90.4, 1.7High consumers0.60.3, 1.4Low consumers0.60.3, 1.4Low consumers0.60.3, 1.4Low consumers0.80.4, 1.6High consumers0.80.4, 1.6High consumers0.80.4, 1.7Low consumers0.80.4, 1.6High consumers0.80.4, 1.7Low consumers0.80.4, 1.7Low consumers0.80.4, 1.7Low consumers0.90.4, 1.7Non-consumers0.90.4, 1.7Low consumers0.90.4, 1.7High consumers1.71.1, 2.8Parental ethnicity (linsh, other)-0.007Low consumers0.90.4, 1.8High consumers1.10.5, 2.2Non-consumers0.90.4, 1.7High consumers1.10.5, 2.2High consumers1.10.5, 2.2High consumers0.90.4, 1.7High consumers0.90.4, 1.7High consumers1.10.5, 2.2High consumers1.10.5, 2.2High consumers0.90.4, 1.7Low consumers0.90.	Unadjusted				
Non-consumers0.80.4, 1-6High consumers1.81.1, 2.9Parilally adjusted model with different child characteristics0.011Child's age, gender-0.011Low consumers0.90.4, 1.7High consumers1.81.1, 2.9MVPA targets achieved (yes, no)-0.058Low consumers0.60.3, 1.4High consumers0.60.3, 1.4High consumers0.60.3, 1.4High consumers1.71.0, 2.9Child-reported TV viewing (<1, 1–3, >3 h/d)-0.022Low consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilojoule intake-0.022Non-consumers0.80.4, 1.7High consumers0.80.4, 1.7Parent characteristics1.71.1, 2.8Parental ethnicity (frish, other)-0.007Low consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental ethnicity (frish, other)-0.007Low consumers-0.003Non-consumers1.10.5, 2.2High consumers1.3, 3.7Family type (single-parent, two-parent)-0.0011Low consumers-0.9Uow consumers-0.011Non-consumers1.81.2, 2.9Parental throity (high consumers)-0.011Non-consumers0.90.4, 1.7High consumers1.8	Low consumers	_		0.012	
High consumers 1-8 1-1, 2-9 Partially adjusted model with different child characteristics - 0-011 Child's age, gender - 0-011 Low consumers 0-9 0-4, 1.7 High consumers 0-9 0.4, 1.7 High consumers 0-9 0.4, 1.7 Low consumers 0-6 0.3, 1.4 Non-consumers 0.6 0.3, 1.4 High consumers 1.7 1.0, 2.9 Child-reported TV viewing (<1, 1-3, >3 h/d) - 0.022 Low consumers 0.8 0.4, 1.6 High consumers 1.7 1.1, 2.7 Daily kilojoule intake - 0.022 Low consumers 0.8 0.4, 1.7 Low consumers 0.8 0.4, 1.7 Low consumers 0.9 0.4, 1.8 Low consumers - 0.002 Non-consumers 0.9 0.4, 1.8 Low consumers - 0.007 Non-consumers 0.9 0.4, 1.8 Low consumers 2.2 1.3, 3.7 Parental ethnicity (firsh, other) </td <td>Non-consumers</td> <td>0.8</td> <td>0.4, 1.6</td> <td></td>	Non-consumers	0.8	0.4, 1.6		
Partially adjusted model with different child characteristics Child's age, gender Low consumers $-$ 0.911 Non-consumers 0.9 0.4, 1.7 High consumers $-$ 0.058 Non-consumers $-$ 0.66 0.3, 1.4 High consumers 0.66 0.3, 1.4 High consumers 0.66 0.3, 1.4 High consumers 0.66 0.3, 1.4 High consumers 0.66 0.4, 1.6 Child-reported TV viewing (<1, 1–3, >3 h/d) Low consumers 0.8 0.4, 1.6 High consumers 0.8 0.4, 1.7 Daily kilojoule intake $-$ 0.022 Non-consumers 0.8 0.4, 1.7 Daily kilojoule intake $-$ 0.022 Non-consumers 0.8 0.4, 1.7 High consumers 0.8 0.4, 1.7 High consumers 0.8 0.4, 1.7 High consumers 0.8 0.4, 1.7 High consumers 1.7 1.1, 2.8 Parent characteristics $-$ 0.007 Non-consumers 0.9 0.4, 1.8 High consumers 0.9 0.4, 1.8 High consumers 1.99 1.2, 3.3 Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) Low consumers 1.1 0.5, 2.2 High consumers $-$ 0.003 Non-consumers $-$ 0.9 Non-consumers 1.1 0.5, 2.2 High consumers $-$ 0.003 Non-consumers $-$ 0.9 Non-consumers $-$ 0.0011 Non-consumers $-$ 0.9 Non-consumers $-$ 0.9 Non-consumers $-$ 0.9 Non-consumers $-$ 0.011 Non-consumers $-$ 0.9 Non-consumers $-$ 0.011 Non-consumers $-$ 0.9 Non-consumers $-$ 0.9 Non-co	High consumers	1.8	1.1, 2.9		
Child's sige, gender Low consumers0-90.4, 1.7High consumers0.90.4, 1.7High consumers1.81.1, 2.9MVPA targets achieved (yes, no)0.0058Low consumers0.60.3, 1.4High consumers0.60.3, 1.4High consumers0.60.21, 1.4Low consumers0.680.4, 1.6Low consumers0.880.4, 1.6Low consumers0.880.4, 1.6High consumers0.880.4, 1.7Non-consumers0.880.4, 1.7Valuers1.71.1, 2.7Daily kilojoule intake0.022Non-consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers0.90.4, 1.8Parent al ethnicity (Irish, other)0.007Low consumers0.90.4, 1.8High consumers1.10.5, 2.2Parental ethnicity (Irish, other)0.007Low consumers1.10.5, 2.2Parent al ethnicity (Irish, other)0.003Low consumers1.10.5, 2.2Low consumers1.10.5, 2.2High consumers1.10.5, 2.2Low consumers0.90.4, 1.7High consumers1.10.5, 2.2Low consumers1.10.5, 2.2Low consumers1.10.5, 2.2Low consumers1.10.9Low consumers0.90.4, 1.7High consumers1.3, 3.7Family type (s	Partially adjusted model with different child characteristics				
Low consumers-0-011Non-consumers0.90.4, 1.7High consumers1.81.1, 2.9MVPA targets achieved (yes, no)-0.058Low consumers0.60.3, 1.4High consumers0.60.3, 1.4High consumers0.60.3, 1.4Child-reported TV viewing (<1, 1-3, >3 h/d)-0.022Low consumers0.80.4, 1.6High consumers0.80.4, 1.7Low consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers0.90.4, 1.8Parent characteristics-0.007Non-consumers0.990.4, 1.8High consumers1.991.2, 3.3Parental ethnicity (Irish, other)-0.003Low consumers1.991.2, 3.3Parental ethnicity (Irish, other)-0.003Low consumers1.10.5, 2.2High consumers1.10.5, 2.2High consumers1.10.5, 2.2High consumers-0.011Non-consumers1.81.2, 2.9Parent education (primary/lower secondary, complete secondary, tertiary level, postgraduate)-Low consumers-0.011Non-consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent exported takeaway food cons	Child's age, gender				
Non-consumers0.90.4, 1.7High consumers1.81.1, 2.9MVPA targets achieved (yes, no)0.058Low consumers-0.058Non-consumers0.60.3, 1.4High consumers0.60.3, 1.4High consumers0.60.3, 1.4Low consumers0.60.4, 1.6Low consumers0.80.4, 1.6Low consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilopule intake0.022Non-consumers0.80.4, 1.7Low consumers0.80.4, 1.7Non-consumers0.80.4, 1.7High consumers0.980.4, 1.7High consumers0.990.4, 1.7Non-consumers0.990.4, 1.8Parent characteristics-0.007Non-consumers0.990.4, 1.8High consumers-0.007Non-consumers-0.007Non-consumers0.990.4, 1.8High consumers1.10.5, 2.2Low consumers1.10.5, 2.2Low consumers0.990.4, 1.7High consumers1.81.2, 2.9Parent elevation (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.011Low consumers0.990.4, 1.7High consumers1.81.2, 2.9Parent reported takeaway food consumption (<once per="" td="" week)<="">0.011Low consumers0.990.4, 1.7High consum</once>	Low consumers	_		0.011	
High consumers1.81.1, 2.9MVPA targets achieved (yes, no)-0.058Non-consumers0.60.3, 1.4High consumers0.60.3, 1.4High consumers1.71.0, 2.9Child-reported TV viewing (<1, 1–3, >3 h/d)-0.022Low consumers0.80.4, 1.6High consumers0.80.4, 1.7Daily kilojoule intake-0.022Low consumers-0.022Non-consumers0.80.4, 1.7Daily kilojoule intake-0.022Low consumers-0.022Non-consumers0.80.4, 1.7High consumers1.71.1, 2.8Parental ethnicity (Irish, other)-0.007Low consumers0.90.4, 1.8High consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers-0.003Non-consumers-0.001Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers1.41.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once="">0.011Low consumers-0.011Non-consumers0.80.4, 1.6Non-consumers0.80.4, 1.6<!--</td--><td>Non-consumers</td><td>0.9</td><td>0.4, 1.7</td><td></td></once>	Non-consumers	0.9	0.4, 1.7		
$ \begin{array}{cccc} MVPA \mbox{targets achieved (yes, no)} & - & 0.058 \\ \mbox{Low consumers} & 0.6 & 0.3, 1.4 \\ \mbox{High consumers} & 1.7 & 1.0, 2.9 \\ \mbox{Child-reported TV viewing (<1, 1–3, >3 h/d)} & 0.022 \\ \mbox{Low consumers} & 0.8 & 0.4, 1.6 \\ \mbox{High consumers} & 0.8 & 0.4, 1.6 \\ \mbox{High consumers} & 0.8 & 0.4, 1.7 \\ \mbox{Daily kilojule intake} & 0.0022 \\ \mbox{Non-consumers} & 0.8 & 0.4, 1.7 \\ \mbox{High consumers} & 0.9 & 0.4, 1.8 \\ \mbox{Parental ethnicity (Irish, other)} & 0.9 & 0.4, 1.8 \\ \mbox{High consumers} & 1.99 & 1.2, 3.3 \\ \mbox{Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) \\ \mbox{Low consumers} & 0.9 & 0.4, 1.7 \\ \mbox{High consumers} & 1.1 & 0.5, 2.2 \\ \mbox{High consumers} & 2.2 & 1.3, 3.7 \\ \mbox{Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) \\ \mbox{Low consumers} & 0.9 & 0.4, 1.7 \\ \mbox{High consumers} & 1.1 & 0.5, 2.2 \\ \mbox{High consumers} & 0.9 & 0.4, 1.7 \\ \mbox{High consumers} & 1.1 & 0.5, 2.2 \\ \mbox{High consumers} & 0.9 & 0.4, 1.7 \\ High consu$	High consumers	1.8	1.1, 2.9		
Low consumers-0.058Non-consumers0.60.3, 1.4High consumers1.71.0, 2.9Child-reported TV viewing (<1, 1–3, >3 h/d)-0.022Low consumers-0.022Non-consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilojoule intake-0.022Non-consumers0.80.4, 1.7Low consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers0.90.4, 1.8Parent characteristics-0.007Non-consumers-0.007Non-consumers-0.007Non-consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers0.90.4, 1.7High consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">-Low consumers0.90.4, 1.6Non-consumers0.80.4, 1.6High consumers0.80.4, 1.6High consumers1.1Low consumers0.90.90.4, 1.6Non-consumers1.1Non-consumers0.90</once>	MVPA targets achieved (yes, no)				
Non-consumers0-60-3, 1.4High consumers1.71.0, 2.9Child-reported TV viewing (<1, 1–3, >3 h/d)0Low consumers-0.022Non-consumers0-80-4, 1-6High consumers1.71.1, 2.7Daily kilojoule intake-0.022Non-consumers0-80-4, 1.7High consumers0-80-4, 1.7Non-consumers0-80-4, 1.7High consumers0.71.1, 2.8Parent characteristics1.71.1, 2.8Parent al ethnicity (fish, other)-0.007Low consumers0-90.4, 1.8High consumers1.991.2, 3.3Parental ethnicity (fish, other)-0.003Low consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers0.90.4, 1.7Low consumers0.90.4, 1.7High consumers0.90.4, 1.7High consumers0.90.4, 1.7Non-consumers <td< td=""><td>Low consumers</td><td>_</td><td></td><td>0.058</td></td<>	Low consumers	_		0.058	
High consumers 1.7 1.0, 2.9 Child-reported TV viewing (<1, 1–3, >3 h/d) 0.022 Low consumers 0.8 0.4, 1.6 High consumers 0.8 0.4, 1.6 High consumers 0.8 0.4, 1.7 Daily kilojoule intake 0.022 0.022 Low consumers 0.8 0.4, 1.7 High consumers 0.8 0.4, 1.7 High consumers 0.8 0.4, 1.7 High consumers 0.9 0.4, 1.8 Parent characteristics 0.9 0.4, 1.8 Parental ethnicity (Irish, other) 0.9 0.4, 1.8 Low consumers 0.9 0.4, 1.8 High consumers 1.99 1.2, 3.3 Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) 0.003 Low consumers 1.1 0.5, 2.2 High consumers 1.1 0.5, 2.2 High consumers 0.9 0.4, 1.7 Non-consumers 0.9 0.4, 1.7 Low consumers 0.9 0.4, 1.7 High consumers 1.8 1.2, 2.9 <tr< td=""><td>Non-consumers</td><td>0.6</td><td>0.3, 1.4</td><td></td></tr<>	Non-consumers	0.6	0.3, 1.4		
Child-reported TV viewing (<1, 1–3, >3 h/d)Low consumers-0.022Non-consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilojoule intake-0.022Low consumers0.80.4, 1.7High consumers0.80.4, 1.7High consumers1.71.1, 2.8Parent characteristics-0.007Non-consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental ethnicity (Irish, other)-0.003Low consumers-0.003Non-consumers2.21.3, 3.7Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">-0.019Low consumers0.80.4, 1.6High consumers1.81.1, 2.9</once>	High consumers	1.7	1.0, 2.9		
Low consumers-0.022Non-consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilojoule intake-0.022Low consumers-0.022Non-consumers0.80.4, 1.7High consumers1.71.1, 2.8Parent characteristics-0.007Non-consumers0.90.4, 1.8High consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental etucation (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">0.019Low consumers0.80.4, 1.6High consumers1.81.1, 2.9</once>	Child-reported TV viewing $(<1, 1-3, >3 h/d)$				
Non-consumers0.80.4, 1.6High consumers1.71.1, 2.7Daily kilojoule intake	Low consumers	_		0.022	
High consumers1.7 $1.1, 2.7$ Daily kilojoule intake-0.022Low consumers0.80.4, 1.7High consumers1.71.1, 2.8Parent characteristics-0.007Parent characteristics-0.007Parental ethnicity (Irish, other)-0.007Low consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers-0.003Non-consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">0.019Low consumers0.90.4, 1.6High consumers1.80.4, 1.6High consumers0.980.4, 1.6High consumers1.81.2, 2.9</once>	Non-consumers	0.8	0.4, 1.6		
Daily kilojoule intake-0.022Non-consumers0.80.4, 1.7High consumers1.71.1, 2.8Parent characteristics-0.007Non-consumers-0.007Non-consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers-0.003Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)0.011Low consumers-0.011Non-consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once="">0.019Low consumers-0.019Non-consumers0.80.4, 1.6High consumers1.81.1, 2.9</once>	High consumers	1.7	1.1, 2.7		
Low consumers-0.022Non-consumers0.80.4, 1.7High consumers1.71.1, 2.8Parent characteristics-0.007Parental ethnicity (Irish, other)-0.09Low consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers-0.003Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">0.019Non-consumers0.80.4, 1.6High consumers-0.019</once>	Daily kilojoule intake				
Non-consumers0.80.4, 1.7High consumers1.71.1, 2.8Parent characteristicsParental ethnicity (Irish, other)0.007Low consumers-0.007Non-consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers-0.003Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)0.90.4, 1.7Low consumers-0.011Non-consumers0.90.4, 1.7High consumers0.90.4, 1.7Low consumers-0.011Non-consumers0.90.4, 1.7Low consumers-0.011Non-consumers0.90.4, 1.7Low consumers0.90.4, 1.7High consumers0.90.4, 1.7High consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">0.019Low consumers0.80.4, 1.6High consumers1.81.1, 2.9</once>	Low consumers	_		0.022	
High consumers 1.7 1.1 , 2.8 Parent characteristicsParental ethnicity (Irish, other) 0.9 0.4 , 1.8 Low consumers 0.9 0.4 , 1.8 1.99 1.2 , 3.3 Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) 0.003 0.003 Low consumers $ 0.003$ Non-consumers 1.1 0.5 , 2.2 High consumers 2.2 1.3 , 3.7 Family type (single-parent, two-parent) $ 0.011$ Low consumers 0.9 0.4 , 1.7 High consumers 1.8 1.2 , 2.9 Parent-reported takeaway food consumption (<once per="" td="" week)<="">0.019Non-consumers0.019Non-consumers$0.8$$0.4$, 1.6High consumers$1.8$$1.1$, 2.9</once>	Non-consumers	0.8	0.4, 1.7		
Parent characteristics Parental ethnicity (Irish, other) Low consumers – 0.007 Non-consumers 0.9 0.4, 1.8 High consumers 1.99 1.2, 3.3 Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) 0.003 Low consumers – 0.003 Non-consumers 1.1 0.5, 2.2 High consumers 2.2 1.3, 3.7 Family type (single-parent, two-parent)	High consumers	1.7	1.1, 2.8		
Parental ethnicity (Irish, other)0.007Low consumers-0.007Non-consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)0.003Low consumers-0.003Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once="">0.019Low consumers0.80.4, 1.6High consumers1.81.1, 2.9</once>	Parent characteristics				
Low consumers-0.007Non-consumers0.90.4, 1.8High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate)-0.003Low consumers-0.003Non-consumers1.10.5, 2.2High consumers2.21.3, 3.7Family type (single-parent, two-parent)-0.011Low consumers-0.011Non-consumers0.90.4, 1.7High consumers1.81.2, 2.9Parent-reported takeaway food consumption (<once per="" td="" week)<="">-0.019Low consumers0.80.4, 1.6High consumers1.81.1, 2.9</once>	Parental ethnicity (Irish, other)				
Non-consumers 0.9 $0.4, 1.8$ High consumers 1.99 $1.2, 3.3$ Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) $ 0.003$ Low consumers $ 0.003$ Non-consumers 1.1 $0.5, 2.2$ High consumers 2.2 $1.3, 3.7$ Family type (single-parent, two-parent) $ 0.011$ Low consumers $ 0.011$ Non-consumers 0.9 $0.4, 1.7$ High consumers 1.8 $1.2, 2.9$ Parent-reported takeaway food consumption (<once <math="" per="" week,="">\geqonce per week)0.019Low consumers0.019Non-consumers$0.8$$0.4, 1.6$High consumers$1.8$$1.1, 2.9$</once>	Low consumers	_		0.007	
High consumers1.991.2, 3.3Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) $ 0.003$ Low consumers $ 0.003$ Non-consumers 1.1 $0.5, 2.2$ High consumers 2.2 $1.3, 3.7$ Family type (single-parent, two-parent) $ 0.011$ Low consumers $ 0.011$ Non-consumers 0.9 $0.4, 1.7$ High consumers 1.8 $1.2, 2.9$ Parent-reported takeaway food consumption (<once per="" td="" week)<="">$-$Low consumers0.019Non-consumers$0.8$$0.4, 1.6$High consumers$1.8$$1.1, 2.9$</once>	Non-consumers	0.9	0.4. 1.8		
Parental education (primary/lower secondary, complete secondary, tertiary level, postgraduate) 0.003 Low consumers - 0.003 Non-consumers 1.1 0.5, 2.2 High consumers 2.2 1.3, 3.7 Family type (single-parent, two-parent) - 0.011 Low consumers - 0.011 Non-consumers 0.9 0.4, 1.7 High consumers 1.8 1.2, 2.9 Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once=""> 0.019 Low consumers - 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9</once>	High consumers	1.99	1.2.3.3		
Low consumers- 0.003 Non-consumers 1.1 $0.5, 2.2$ High consumers 2.2 $1.3, 3.7$ Family type (single-parent, two-parent)- 0.011 Low consumers- 0.9 Non-consumers 0.9 $0.4, 1.7$ High consumers 1.8 $1.2, 2.9$ Parent-reported takeaway food consumption (<once per="" td="" week)<="">-Low consumers-0.019Non-consumers$0.8$$0.4, 1.6$High consumers$1.8$$1.1, 2.9$</once>	Parental education (primary/lower secondary, complete seconda	rv. tertiarv level. postgra	duate)		
Non-consumers 1.1 $0.5, 2.2$ High consumers 2.2 $1.3, 3.7$ Family type (single-parent, two-parent) $ 0.011$ Low consumers $ 0.011$ Non-consumers 0.9 $0.4, 1.7$ High consumers 1.8 $1.2, 2.9$ Parent-reported takeaway food consumption (<once per="" td="" week)<="">$-$Low consumers0.019Non-consumers$0.8$$0.4, 1.6$High consumers$1.8$$1.1, 2.9$</once>	Low consumers		· · · · · ,	0.003	
High consumers 2.2 $1.3, 3.7$ Family type (single-parent, two-parent)– 0.011 Low consumers– 0.011 Non-consumers 0.9 $0.4, 1.7$ High consumers 1.8 $1.2, 2.9$ Parent-reported takeaway food consumption (<once <math="" per="" week,="">\geqonce per week)0.019Low consumers–0.019Non-consumers$0.8$$0.4, 1.6$High consumers$1.8$$1.1, 2.9$</once>	Non-consumers	1.1	0.5. 2.2		
Family type (single-parent, two-parent) - 0.011 Low consumers - 0.011 Non-consumers 0.9 0.4, 1.7 High consumers 1.8 1.2, 2.9 Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once=""> - 0.019 Low consumers - 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9</once>	High consumers	2.2	1.3. 3.7		
Low consumers - 0.011 Non-consumers 0.9 0.4, 1.7 High consumers 1.8 1.2, 2.9 Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once=""> - 0.019 Low consumers - 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9</once>	Family type (single-parent, two-parent)		-, -		
Non-consumers 0.9 0.4, 1.7 High consumers 1.8 1.2, 2.9 Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once=""> 0.019 Low consumers – 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9</once>	Low consumers	_		0.011	
High consumers 1.8 1.2, 2.9 Parent-reported takeaway food consumption (<once per="" td="" week)<="" week,="" ≥once=""> 0.019 Low consumers - 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9</once>	Non-consumers	0.9	0.4, 1.7		
Parent-reported takeaway food consumption (<once per="" week)<br="" week,="" ≥once="">Low consumers – 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9</once>	High consumers	1.8	1.2. 2.9		
Low consumers – 0.019 Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9	Parent-reported takeaway food consumption (<once per="" week.=""></once>	once per week)	,		
Non-consumers 0.8 0.4, 1.6 High consumers 1.8 1.1, 2.9	Low consumers			0.019	
High consumers 1.8 1.1, 2.9	Non-consumers	0.8	0.4. 1.6		
	High consumers	1.8	1.1, 2.9		

MVPA, moderate to vigorous physical activity.

Multivariate analysis

In multivariate logistic regression (Table 5), the associations observed in univariate analyses (Table 4) remained stable. In the fully adjusted model (model 5), high SSB consumers have 80 % increased odds of having overweight/ obesity compared to low SSB consumers (OR 1·8, 95 % CI $1\cdot0$, $3\cdot5$).

In separate fully adjusted regression analyses (Fig. 1), clear BMI distribution differences are evident across SSB consumption quintiles: a one unit difference in BMI score exists between the highest and the lowest quintiles of SSB consumption.

Discussion

This study provides evidence of SSB consumption levels in a sample of Irish children living in Cork city and county prior to the introduction of a sugar drinks tax. The Sugar Sweetened Beverage Tax came into effect on 1 May 2018 and applies to water- and juice-based drinks that have added sugar and a total sugar content of ≥ 5 g per 100 ml. Drink categories that are liable for the tax are flavoured waters, carbonated drinks, energy/sports drinks and juice-based drinks. The drink types are detailed using the Combined Nomenclature (CN) of the European Union (CN codes 2009 and 2202). There are four key findings. Firstly, the majority of participants consumed SSB (82%); secondly, participants consumed 328 ml SSB per day on average, which is equivalent to one standard commercial serving (330 ml); thirdly, SSB contributed to 6 and 8% of energy (kJ) intake and 22 and 26 % of sugar intake for normal-weight children and children with overweight/obesity, respectively; finally, SSB consumption >200 ml is associated with increased BMI. Clear positive distribution differences in BMI are evident across SSB consumption groups.

NS Public Health Nutrition

Table 5 Multivariate logistic regression of the association between SSB consumption (ref: <1 standard unit (200 ml/d)) and overweight/ obesity*

		Outcome: overweight/obesity											
	Low consumer				Non-consum	ner							
	OR	95 % CI	P-value	OR	95 % CI	P-value	OR	95 % CI	P-value	the model			
		1	_	0.9	0.5, 1.7	0.68	1.8	1.1, 2.9	0.012	716			
Model 1		1	-	1.1	0.5, 2.1	0.86	2.2	1.3, 3.7	0.003	626			
Model 2		1	-	0.8	0.3, 1.9	0.61	2.0	1.1, 3.7	0.022	499			
Model 3		1	_	0.7	0.3, 1.6	0.37	1.8	1.0, 3.3	0.05	488			
Model 4		1	-	0.6	0.2, 1.6	0.32	1.9	1.0, 3.5	<0.05	478			
Model 5		1	-	0.6	0.2, 1.6	0.34	1.8	1.0, 3.5	0.06	47			

*Model 1: adjusted for child's age and gender, parental education, family type and ethnicity. Model 2: model 1 + total daily MVPA. Model 3: model 2 + child-reported and parent-reported child TV viewing. Model 4: model 3 + parent-reported family takeaway consumption. Model 5: model 4 + average daily kJ intake.



Fig. 1. Kernel density distribution of BMI by quintiles of Sugar Sweetened Beverages consumption

Results in context

Evidence of an association between SSB consumption and increased weight is compelling^(28,29). Concurrent with increasing prevalence of global obesity, a surge in the availability of ultra-processed foods and beverages occurred⁽³⁰⁾. Included in this group of obesogenic foods and beverages are soft drinks (SSB). Our results are consistent with other published studies reporting that SSB consumption, which imparts no nutritional benefit on the individual, contributes significantly to the overall daily energy intake. US data indicate that SSB consumption contributes >2092 kJ/d in 5 % of 2–11-year-olds⁽³¹⁾. NHANES results show that between 2011 and 2014, six in ten youths (63 %) drank an SSB on a given day; this compares to 82 % in the current study. Further, evidence suggests that a high intake of these beverages is not accompanied by a reduction in food intake⁽¹⁰⁾,

and the lack of this compensation may contribute to the surplus daily energy consumed.

Strengths and limitations

This study is strengthened by the depth of data on lifestyle, diet and physical activity collected at an individual and family level, allowing for an in-depth exploration on the potential determinants of SSB consumption and its association with weight status. The study collected objectively measured height, weight and BMI data. Additionally, objectively measured physical activity data was collected under free-living conditions over a 7-d period. The thoroughly debriefed 3-d food diaries provided comprehensive data on dietary intake patterns and behaviours, including SSB consumption.

NS Public Health Nutrition

Limitations of the study include the cross-sectional design of the survey, the relatively modest response rate (67%) and the issue of measurement error in relation to the exposure (diet quality) and the self-reported nature of confounding variables. The study was conducted in 2012, and so it may not be possible to indicate how consumption levels changed between 2012 and the introduction of the tax in 2018. The study recruited children from only one region of Ireland, and the sample is predominantly urban; so caution needs to be exercised in making inferences for the rest of the Irish population. Further, active parental permission might depress response rates, and non-responding children (and parents) are likely to have less favourable BMI profiles. It was outside of the scope of this study to follow-up non-responders, and thus we cannot definitively state that the BMI profile of participants was similar to that of non-responders. By definition, we have to be cautious in making causal links in crosssectional analyses. However, the findings presented here are entirely consistent with available data from other studies. It should be noted that misclassification of exposures and outcomes due to random error would tend to underestimate the effect sizes, and it was highly likely that the magnitude of associations seen between SSB consumption and child's weight has been underestimated.

Conclusion

Public Health Nutrition

We provide a baseline from which the impact of the sugar drinks tax on SSB consumption can be benchmarked. We provide evidence of a very high level of SSB consumption among a sample of school-going children in Cork, Ireland, accounting for a significant proportion of daily energy and sugar intake. We further provide strong evidence of its association with overweight/obesity and suggest that differences in child's BMI are evident between high and low SSB consumers at a population level.

Population-based approaches to reduce SSB consumption, as part of a complex systems approach to tackling the problem of childhood obesity, are a public health necessity. While no single measure might reverse the current trends in obesity, a multi-component strategy, including a targeted approach towards improving the food environment, will be necessary.

Key points

- SSB impart no nutritional value to the overall diet, and high consumption levels are attributed to excess weight in children.
- To date, the magnitude of SSB consumption and its contribution to the overall daily energy and sugar intake of Irish children has not been documented.

- 2243
- The majority of children in our study consumed SSB, accounting for a significant proportion of daily energy and sugar intake.
- The study provides strong evidence of an association between SSB consumption and overweight and obesity.
- Our results provide a baseline from which the impact of the SSB tax, recently introduced in Ireland, on consumption levels of these drinks can be gauged.

Acknowledgements

Acknowledgements: The authors would like to acknowledge the contribution of all study participants and the participating schools for accommodating data collection. Financial support: The Cork Children's Lifestyle Study was funded by the National Children's Research Centre. Crumlin. The authors are affiliated with the HRB Centre for Health and Diet Research (HRC/2007/13). Conflict of interest: None. Authorship: J.M.H. is the PI and guarantor for this study; she designed the study, analysed the data, interpreted the results and drafted and revised the article. J.M.H. has access to the data and controls the decision to publish. C.P. conducted data collection, processed the food diary data, assisted with data analysis and reviewed and commented on manuscript drafts. E.K. conducted data collection, advised on results interpretation and commented on manuscript drafts. I.J.P. conceived and designed the study and helped draft the manuscript. Ethics of human subject participation: This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by the Clinical Research Ethics Committee of the Cork University Teaching Hospitals. Written informed consent was obtained from all subjects/patients.

References

- 1. Department of Health (2016) *A Healthy Weight for Ireland: Obesity Policy and Action Plan.* Dublin: Stationery Office.
- World Health Organisation (2015) Global Status Report on Noncommunicable Diseases 2014. http://apps.who.int/iris/ bitstream/10665/148114/1/9789241564854_eng.pdf (accessed June 2019).
- World Health Organization (2016) Report of the Commission on Ending Childbood Obesity. Geneva: World Health Organisation.
- Godlee F (2016) Sugar tax and screening: listen to the evidence. *BMJ* 352, i33.
- Sarlio-L\u00e4hteenkorva S & Winkler JT (2015) Could a sugar tax help combat obesity? *BMJ* 351, h4047.
- Sassi F (2016) An old idea with a new place in the fight against obesity? *BMJ* 352, h6904.
- 7. OECD (2012) OECD Obesity Update 2012. France: OECD Publishing.
- Redondo M, Hernández-Aguado I & Lumbreras B (2018) The impact of the tax on sweetened beverages: a systematic review. *Am J Clin Nutr* 108, 548–563.

- 2244
- Briggs ADM, Mytton OT, Kehlbacher A *et al.* (2013) Overall and income specific effect on prevalence of overweight and obesity of 20 % sugar sweetened drink tax in UK: econometric and comparative risk assessment modelling study. *BMJ* 347, f6189.
- 10. Colchero MA, Popkin BM, Rivera JA *et al.* (2016) Beverage purchases from stores in Mexico under the excise tax on sugar sweetened beverages: observational study. *BMJ* **352**, h6704.
- 11. Gulland A (2016) WHO calls for tax on sugary drinks to tackle child obesity. *BMJ* **352**, i475.
- 12. Keane E, Kearney PM, Perry IJ *et al.* (2014) Diet, physical activity, lifestyle behaviors, and prevalence of childhood obesity in Irish children: the Cork Children's Lifestyle Study Protocol. *JMIR Res Protoc* **3**, e44.
- 13. Food Standards Agency (2014) *McCance and Widdowson's the Composition of Foods*, 7th Summary ed. Cambridge: Royal Society of Chemistry.
- Black LJ, Ireland J, Møller A *et al.* (2011) Development of an on-line Irish food composition database for nutrients. *J Food Compos Anal* 24, 1017–1023.
- Food Safety Authority of Ireland (2019) Healthy Eating, Food Safety and Food Legislation. A Guide Supporting the Healthy Ireland Food Pyramid. Dublin: Food Safety Authority of Ireland.
- World Health Organisation (2015) Guideline: Sugars Intake for Adults and Children. Geneva: World Health Organisation.
- Schofield W, Schofield C & James W (1985) Predicting basal metabolic rate: new standards and review of previous work. *Hum Nutr Clini Nutr* **39**, Suppl. 1, 5–41.
- Goldberg G, Black A, Jegg S *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: derivation of cut-off limits to identify underrecording. *Eur J Clin Nutr* **45**, 569–581.
- Black AE, Prentice AM & Goldberg GR (1993) Measurements of total energy expenditure provide insights into the validity of dietary measurements of energy intake. *J Am Diet Assoc* 93, 572–579.

- 20. Cole TJ, Bellizzi MC, Flegal KM *et al.* (2000) Establishing a standard definition for child overweight and obesity world-wide: international survey. *BMJ* **320**, 1240.
- 21. Cole TJ & Lobstein T (2012) Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes* **7**, 284–294.
- 22. Keane E, Li X, Kearney P *et al.* (2017) Physical activity, sedentary behaviour and the risk of overweight and obesity in school aged children. *Paediatr Exerc Sci* **29**(3), 408–418.
- Li X, Kearney PM, Keane E *et al.* (2017) Levels and sociodemographic correlates of accelerometer-based physical activity in Irish children: a cross-sectional study. *J Epidemiol Community Health* **71**(6), 521–527.
- Esliger DW, Rowlands AV, Hurst TL *et al.* (2011) Validation of the GENEA accelerometer. *Med Sci Sports Exerc* 43, 1085–1093.
- 25. Phillips LR, Parfitt G & Rowlands AV (2013) Calibration of the GENEA accelerometer for assessment of physical activity intensity in children. *J Sci Med Sport* **16**, 124–128.
- van Hees VT, Renström F, Wright A *et al.* (2011) Estimation of daily energy expenditure in pregnant and non-pregnant women using a wrist-worn tri-axial accelerometer. *PLOS ONE* 6, e22922.
- 27. World Health Organisation (2010) *Global Recommendations* on *Physical Activity for Health*. Geneva: WHO.
- Malik VS, Schulze MB & Hu FB (2006) Intake of sugarsweetened beverages and weight gain: a systematic review. *Am J Clin Nutr* 84, 274–288.
- 29. Vartanian LR, Schwartz MB & Brownell KD (2007) Effects of soft drink consumption on nutrition and health: a systematic review and meta-analysis. *Am J Public Health* **97**, 667–675.
- Rodgers A, Woodward A, Swinburn B *et al.* (2018) Prevalence trends tell us what did not precipitate the US obesity epidemic. *Lancet Public Health* 3, e162–e163.
- Han E & Powell LM (2013) Consumption patterns of sugar sweetened beverages in the United States. J Acad Nutr Diet 113, 43–53.