

| | |
|-----------------------------|--|
| Title | Protein quality and the protein to carbohydrate ratio within a high fat diet influences energy balance and the gut microbiota in C57BL/6J mice |
| Authors | McAllan, Liam;Skuse, Peter;Cotter, Paul D.;O'Connor, Paula M.;Cryan, John F.;Ross, R. Paul;Fitzgerald, Gerald F.;Roche, Helen M.;Nilaweera, Kanishka N. |
| Publication date | 2014 |
| Original Citation | McAllan L, Skuse P, Cotter PD, Connor PO, Cryan JF, Ross RP, et al. (2014) Protein Quality and the Protein to Carbohydrate Ratio within a High Fat Diet Influences Energy Balance and the Gut Microbiota In C57BL/6J Mice. PLoS ONE 9(2): e88904. doi:10.1371/journal.pone.0088904 |
| Type of publication | Article (peer-reviewed) |
| Link to publisher's version | 10.1371/journal.pone.0088904 |
| Rights | © 2015 McAllan et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited - http://creativecommons.org/licenses/by/4.0/ |
| Download date | 2024-02-23 07:25:36 |
| Item downloaded from | https://hdl.handle.net/10468/2342 |



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Protein Quality and the Protein to Carbohydrate Ratio within a High Fat Diet Influences Energy Balance and the Gut Microbiota In C57BL/6J Mice

Liam McAllan^{1,2}, Peter Skuse^{1,3}, Paul D. Cotter^{1,4}, Paula O' Connor¹, John F. Cryan^{4,5}, R. Paul Ross¹, Gerald Fitzgerald³, Helen M. Roche⁶, Kanishka N. Nilaweera^{1*}

1 Food Biosciences Department, Teagasc, Fermoy, County Cork, Ireland, **2** Department of Pharmacology and Therapeutics, University College Cork, Cork, Ireland, **3** Department of Microbiology, University College Cork, Cork, Ireland, **4** Alimentary Pharmabiotic Centre, University College Cork, Cork, Ireland, **5** Department of Anatomy & Neuroscience, University College Cork, Cork, Ireland, **6** UCD Conway Institute of Biomolecular & Biomedical Research, University College Dublin, Dublin, Ireland

Abstract

Macronutrient quality and composition are important determinants of energy balance and the gut microbiota. Here, we investigated how changes to protein quality (casein versus whey protein isolate; WPI) and the protein to carbohydrate (P/C) ratio within a high fat diet (HFD) impacts on these parameters. Mice were fed a low fat diet (10% kJ) or a high fat diet (HFD; 45% kJ) for 21 weeks with either casein (20% kJ, HFD) or WPI at 20%, 30% or 40% kJ. In comparison to casein, WPI at a similar energy content normalised energy intake, increased lean mass and caused a trend towards a reduction in fat mass ($P=0.08$), but the protein challenge did not alter oxygen consumption or locomotor activity. WPI reduced HFD-induced plasma leptin and liver triacylglycerol, and partially attenuated the reduction in adipose FASN mRNA in HFD-fed mice. High throughput sequence-based analysis of faecal microbial populations revealed microbiota in the HFD-20% WPI group clustering closely with HFD controls, although WPI specifically increased *Lactobacillaceae/Lactobacillus* and decreased *Clostridiaceae/Clostridium* in HFD-fed mice. There was no effect of increasing the P/C ratio on energy intake, but the highest ratio reduced HFD-induced weight gain, fat mass and plasma triacylglycerol, non-esterified fatty acids, glucose and leptin levels, while it increased lean mass and oxygen consumption. Similar effects were observed on adipose mRNA expression, where the highest ratio reduced HFD-associated expression of UCP-2, TNF α and CD68 and increased the diet-associated expression of β 3-AR, LPL, IR, IRS-1 and GLUT4. The P/C ratio also impacted on gut microbiota, with populations in the 30/40% WPI groups clustering together and away from the 20% WPI group. Taken together, our data show that increasing the P/C ratio has a dramatic effect on energy balance and the composition of gut microbiota, which is distinct from that caused by changes to protein quality.

Citation: McAllan L, Skuse P, Cotter PD, Connor PO, Cryan JF, et al. (2014) Protein Quality and the Protein to Carbohydrate Ratio within a High Fat Diet Influences Energy Balance and the Gut Microbiota In C57BL/6J Mice. PLoS ONE 9(2): e88904. doi:10.1371/journal.pone.0088904

Editor: Darcy Johannsen, Pennington Biomed Research Center, United States of America

Received: July 23, 2013; **Accepted:** January 13, 2014; **Published:** February 10, 2014

Copyright: © 2014 McAllan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: KN is supported by the Teagasc Vision Programme on Obesity, which also funded the work detailed in this manuscript. LM is supported by a Teagasc PhD Walsh Fellowship. HMR is supported by SFI PI (11/PI/1119). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Paul D. Cotter is a PLOS ONE Editorial Board member. The authors also declare that this does not alter their adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: kanishka.nilaweera@teagasc.ie

Introduction

It is widely recognised that levels of obesity and related clinical conditions such as diabetes, stroke, hyperlipidemia and cardiovascular disease are increasing worldwide [1]. Importantly, the development of obesity increases the set point at which the body weight, more specifically body fat, is defended, thus making its reversal difficult to achieve [2,3]. As such, there is an increased research interest to develop effective treatments for this disease.

Dairy proteins belonging to the whey fraction (a by-product of cheese manufacture) have been increasingly tested for their potential anti-obesity effect, specifically for their ability to reduce high fat diet (HFD)-associated body weight and fat mass gain [4–6]. Shi *et al.*, [7] showed that replacing 5%, 50% or 100% of the dietary casein protein-derived energy content with a lactoperoxidase and lactoferrin-enriched whey protein isolate (WPI) caused a proportional suppression of body weight gain in HFD fed mice.

We have previously demonstrated that a WPI-related reduction in body weight and fat mass gain in HFD fed mice was accompanied by a normalisation of energy intake and complete or partial reversal of energy balance-related gene expression in the adipose tissue and the hypothalamus [8]. While these data suggest that whey proteins have specific-effects on energy balance, such effects appear to be modified by the macronutrient composition in the diet [9]. In the latter study, it was shown that increasing the lipid to carbohydrate ratio within a whey protein-rich diet significantly reduced energy intake and bodyweight gain in rats. Collectively, these data suggest that protein quality and macronutrient composition are important determinants of energy balance.

Interestingly, diet is also an important factor in determining the composition of the gut microbiota [10,11] and specific gut microbiota signatures are associated with obesity phenotypes in animals and humans [12–14]. Notably, studies have shown specific whey proteins to possess anti-microbial activity [15–17], and that

the digestive process itself facilitates the formation of potent antimicrobial whey-derived peptides, such as pepsin catalysed lactoferrin to lactoferricin [18]. A study by Sprong *et al.*, [19] demonstrated that in comparison to casein, whey protein intake increased levels of lactobacilli and bifidobacteria in a rat model of colitis. However, in a more recent study, whey protein intake was found to have no influence on gut microbiota composition in mice fed a HFD for 7 or 13 weeks [20]. Several key unanswered questions are; could whey proteins specifically influence the gut microbiota composition associated with prolonged high fat feeding, and would any changes relate to energy balance? Could changes to protein to carbohydrate ratio within a HFD vary the gut microbiota profile and energy balance in a different way to changes to protein quality?

To assess WPI specific effects on above parameters, we subjected male C57BL/6J mice to 21 weeks of either a low fat diet (LFD) with 20% kJ casein or a HFD with 20% kJ casein or WPI. In addition, using two additional HFD dietary groups on 30 or 40% kJ WPI, we evaluated the impact of increasing the protein to carbohydrate (P/C) ratio within the HFD on parameters of interest. Our data show that WPI has a specific effect on energy balance and gut microbiota, while increasing the P/C ratio within the HFD leads to dramatic alterations in energy balance, body composition, metabolic health and the composition of the gut microbiota.

Materials and Methods

Ethics Statement

All research involving mice was licensed under the Cruelty to Animal Act 1876 and received ethical approval from the University College Cork Animal Ethics Review Committee (#2011/005).

Animals

Male 3–4 week old C57BL/6J mice (Harlan, Oxon, UK) were group housed either 5 per cage (Study 1) or 4 per cage (Study 2) in individually ventilated cages and acclimatised for four weeks in a light (06:00–18:00), temperature ($21 \pm 1^\circ\text{C}$) and humidity (45–65%) controlled environment with free access to water and a low fat diet (LFD; 10% kJ fat and 20% kJ casein; #D12450, Research diets; New Brunswick, NJ, USA).

Experimental protocol

Two studies were performed to assess how the WPI-derived energy content within a HFD (study 1) or LFD (study 2) impacts on energy balance-related parameters in mice over a 21 week (study 1) or 7 week period (study 2).

Study 1: Following the acclimatisation period, weight-matched dietary groups were maintained on the LFD or switched to either a HFD (45% kJ fat and 20% kJ casein; #D12451) or a HFD with WPI (Alacentm 895 NZMP, New Zealand) at an energy content of 20% kJ (HFD-20% WPI), 30% kJ (HFD-30% WPI) or 40% kJ (HFD-40% WPI) (Table S1) ($n = 10$) for a total of 21 weeks. Body weights were measured weekly. Energy intake in group housed mice was measured by weighing the food hopper each week until week 16. During weeks 17–20, energy intake and metabolic activity in individual mice was measured using TSE Phenomaster cages (TSE systems, Bad Homburg, Germany). Following this analysis and prior to re-housing the mice in home cages, faecal pellets were collected from individual mice for examination of microbial composition via pyrosequencing and subsequent bioinformatic analysis. At the end of the experimental period, mice were fasted for 6 hours and the body composition was

measured using the Bruker minispec LF50H (Bruker optics, Ettlingen, Germany). Mice were then anaesthetised using ketamine (65 mg/kg bodyweight) and xylazine (13 mg/kg bodyweight). Blood was collected from anaesthetised mice into vacutainer EDTA tubes (BD, USA) and treated with Aprotinin (500,000 KIU/L final concentration; Sigma, Ireland) and Diprotin A (0.1 mM final concentration; Sigma, Ireland) to protect plasma peptides from proteolytic degradation. Plasma was isolated from blood by centrifugation at 2000 rpm at 4°C for 15 mins. Mice were sacrificed by cervical dislocation, and tissues of interest were dissected and snap frozen in liquid nitrogen (liver, adipose and stomach) or on dry ice (brain). Plasma and tissue samples were stored at -80°C until analysis.

Study 2: Weight matched mice were provided for 7 weeks with either the LFD or a LFD with WPI replacing the casein protein (LFD-WPI; 10% kJ fat and 20% kJ WPI) ($n = 8$). Body weights were measured weekly. Energy intake and metabolic activity in individual mice was measured during weeks 5 and 6 using the TSE Phenomaster system. After the analysis, mice were re-housed, as before, in the home cages and the experiment was terminated at the end of week 7.

Analysis of metabolic parameters

The TSE Phenomaster cages comprised an open-circuit indirect calorimetry system with gas sensing units to measure oxygen consumption (ml/h/kg) (VO_2) and CO_2 production (ml/h/kg) (VCO_2). The cages also contained high precision sensor associated-feeding baskets to accurately measure food intake (g), with a meal defined as intake over 0.01 g. A multi-dimensional infrared beam system allowed the measurement of locomotor activity, which was defined as the total number of infrared beam breaks in the X and Y axis. Mice were singly housed in TSE Phenomaster cages for a total of 3 days, with data collected during the final 24 hours, following a 2 day acclimatisation to the new cage environment. The acclimatisation period was established based on the data from our previous study [8]. Heat production (kcal/h/kg) in individual mice was calculated using the Weir equation ($3.941 \times \text{VO}_2 + 1.106 \times \text{VCO}_2$) [21]), and this was converted to kJ/h/kg using $1 \text{ kcal} = 4.184 \text{ kJ}$. The respiratory exchange ratio (RER) was calculated by VCO_2/VO_2 . Energy intake was calculated from food intake measurements using the energy content of the diets supplied by the manufacturer.

Microbial DNA extraction, amplification and high throughput DNA sequencing

Total metagenomic DNA was extracted from individual faecal samples using QJamp DNA Stool Mini Kit (Qjagen, Hilden, Germany), after an additional bead-beating step. Bacterial composition was determined by sequencing of 16S rRNA amplicons (V4-V5 region; 408nt long) generated by a separate PCR reaction for each sample (in triplicate) using universal 16S primers, where, the forward primer (5'-AYTGGGYD-TAAAGNG), with attached molecular identifier tags between 454 adapter sequence and target-specific primer sequence, and reverse primer V5 (5'-CCGTCAATTTTTRAGTTT) [22], were used along with Biomix Red (Bioline, London UK). The template DNA was amplified under the following PCR conditions for a total of 35 cycles: 94°C for 2 minutes and 1 minute respectively (initialization and denaturation), 56°C for 60 seconds (annealing) and 72°C for 60 seconds (elongation), proceeded by a final elongation stage of 2 minutes. Negative control reactions with PCR grade water in place of template DNA were used to confirm a lack of contamination. Amplicons were pooled and cleaned using the AMPure XP purification system (Beckman and Coulter,

Takeley, UK) and DNA concentration was determined using the NANODROP 3300 Fluorospectrometer (Thermo Scientific, USA) coupled with the Quant-it™ Picogreen® dsDNA Assay Kit (Invitrogen, Paisley, UK). Equal volumes of each sample were then pooled together and underwent a final cleaning and quantification stage. Amplicons were sequenced in-house on a Roche GS FLX Titanium platform.

Bioinformatics

Raw sequencing reads were ‘de-noised’ using traditional techniques implemented in the Ribosomal Database Project Pyrosequencing (RDP) Pipeline with ambiguous bases, non exact primer matches and reads shorter than 150 bp being excluded. Trimmed FASTA files were then BLASTed against a previously published 16S-specific database using default parameters. The resulting files were then parsed using the MEGAN software package, which assigns reads to the National Centre for Biotechnology Information (NCBI) taxonomies via the lowest common ancestor algorithm. Results were filtered prior to tree construction and summarization by the use of bit scores from within MEGAN where a cut-off bit score of 86 was employed [23,24]. The QIIME software suite was employed to achieve clustering of sequence reads into operational taxonomic units (OTUs) [25]. Chimeric OTUs were removed using the Chimera-Slayer program [26] and phylogenetic trees constructed using the FastTreeMP tool [27]. Beta diversity values were calculated based on Bray Curtis, weighted and unweighted UniFrac distances, and the KING viewer was used to visualise resulting PCoA plots [28,29]. Sequence reads were deposited in the European Nucleotide Archive (ENA) under the accession number PRJEB4636.

Plasma analysis

Colorimetric assays were used to measure plasma levels of glucose (Calbiochem, Darmstadt, Germany), triacylglycerol (TAG; Wako Chemicals, Richmond, VA, USA) and non-esterified fatty acids (NEFA; Abcam, Cambridge, UK). Commercially available ELISA kits were used to analyse plasma levels of insulin, leptin (Crystal Chem, Downers Grove, IL, USA), glucagon-like peptide 1 (GLP-1; Millipore, St. Charles, MO, USA) and corticosterone (Enzo Life sciences, Farmingdale, NY, USA). The homeostasis model assessment of insulin resistance (HOMA-IR) was determined using the formula: fasting plasma insulin ($\mu\text{U}/\text{ml}$) \times fasting plasma glucose (mmol/L)/22.5 [30]. To measure plasma amino acid levels, samples were first deproteinised by mixing with equal volumes of 24% (w/v) tri-chloroacetic acid. The samples were then allowed to stand for 10 minutes before being centrifuged at $14400 \times g$ (Microcentaur, MSE, UK) for 10 minutes. Supernatants were mixed with 0.2 M sodium citrate buffer, pH 2.2, and the plasma concentration of amino acids were quantified using a Jeol JLC-500/V amino acid analyser (Jeol Ltd., Garden city, UK) fitted with a Jeol Na^+ high performance cation exchange column.

Liver TAG analysis

Total lipids from liver samples (approx. 50 mg) were extracted as described previously [8] using the Folch extraction method [31]. Briefly, total lipids were extracted using 2:1 (v/v) chloroform:methanol solution, into which a 0.88% NaCl solution was added before centrifugation at 2000 rpm and 4°C for 30 mins. Aliquots of the organic phase were collected, dried and re-suspended in the LabAssay TAG reagent (Wako Chemicals, Richmond, VA, USA) to measure TAG levels using LabAssay TAG kit according to the manufacturer’s protocol.

Real-Time PCR analysis

Total RNA was isolated from tissues using RNeasy mini (liver and stomach) or RNeasy lipid mini (adipose and hypothalamus) kits (Qiagen, Hilden, Germany) according to manufacturers’ instructions. RNA was treated with DNase (Qiagen, Hilden, Germany) during RNA isolation to eliminate any possible genomic DNA contamination. Reverse transcription of 1 μg of RNA was performed using 2.5 ng/ μl random hexamer primers (Biolone, London, UK), 0.5 mM dNTP (Promega, Madison, WI, USA), 2 U/ μl RNase inhibitor (Promega, Madison, WI, USA), and the Superscript II first stand system (Invitrogen, Carlsbad, CA, USA) according to manufactures’ instructions. Gene expression was measured by the Roche Lightcycler 480 system (Rotkreuz, Switzerland) via amplification of 1 μl complementary DNA using the Lightcycler SYBR Green I Mastermix kit (Roche, Penzberg, Germany) and 2.5 μM gene specific primers (Eurofins MWG operon, Ebersberg, Germany) in a 10 μl total reaction volume. Primer sequences used are listed in Table S2. PCR conditions were; 10 mins at 95°C , followed by 50 cycles of 95°C for 10 s, $58\text{--}65^\circ\text{C}$ for 5 s and 72°C for 15 s. Authenticity of PCR products was determined by melting curve analysis and by automated sequencing. Crossing point (Cp) of fluorescence signals were used to calculate target gene expression by $2^{-\Delta\Delta\text{Cp}}$, following normalisation against housekeeping gene according to $\Delta\Delta\text{Cp} = \Delta\text{Cp target gene} - \Delta\text{Cp housekeeping gene}$. Housekeeping genes used were β -actin (liver, stomach and hypothalamus), YWHAZ (liver and hypothalamus), 18-S (adipose) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (adipose and stomach). Relative gene expression is shown compared to the LFD group.

Statistical analysis

Data are expressed as means \pm standard error of the mean (SEM). Differences between experimental dietary groups were analysed by one-way or two-way ANOVA followed by pairwise comparisons using *tukey* or *bonferroni* post hoc tests, respectively. Body weight trajectories were analysed by two-way repeated measures ANOVA with *bonferroni* post hoc tests. Non-parametric data was compared by Kruskal-Wallis ANOVA followed by *Dunn*’s pairwise comparisons. Significance levels were set at $P \leq 0.05$, and statistical analysis was performed using Graphpad prism (ver. 5.04; San Diego, CA, USA) and Minitab (ver.15; State College, PA, USA).

Results

WPI inclusion or increasing the P/C ratio within a HFD alters body composition and plasma amino acids

Fig. 1A–B demonstrates that body weight gain of HFD-20%WPI fed mice was similar to HFD controls. However, intake of WPI was seen to have a specific effect on body composition, with HFD-20% WPI fed mice having an increased lean mass (%) ($P < 0.05$), and a trend towards a reduction in fat mass (%) ($P = 0.08$) compared to the HFD control group. Increasing the WPI derived energy content in the HFD to 40% and proportionally reducing the carbohydrate energy content led to a significant reduction in body weight gain compared to all other HFD-WPI groups ($P < 0.001$), with observed values similar to that seen for the LFD group. This was accompanied by significantly reduced body fat mass and increased lean mass levels in the HFD-40% WPI fed mice compared to HFD control and other WPI diets groups ($P < 0.001$), while the body composition of HFD-30% WPI group did not differ from that of the HFD-20% WPI group (Fig. 1C).

Comparison of the plasma amino acid profiles including those that could influence lean and fat mass, revealed an impact of WPI




Figure 1. Impact of whey protein isolate and protein to carbohydrate ratio on body weight and composition. (A) shows the body weight trajectories of mice during 21 weeks of dietary treatment with a 10% kJ low fat diet (LFD), 45% kJ high fat diet (HFD) or a HFD with 20, 30 or 40% kJ whey protein isolate (WPI). Body weight gain (B) and body composition (C) of mice after 21 weeks on experimental diets are also shown. Data represent mean values \pm S.E.M. (n = 10 per group). Groups that do not share a common letter are significantly different at $P < 0.05$. doi:10.1371/journal.pone.0088904.g001

and the P/C ratio (Table 1). WPI specific effects were observed on glutamic acid, aspartic acid and glycine, which either decreased (glutamic acid and aspartic acid) or increased (glycine) compared

to HFD fed mice ($P < 0.05$) (Table 1). Changes in macronutrient ratio in HFD-40%WPI, decreased plasma histidine, phenylalanine, serine and threonine levels compared to the lowest P/C ratio (20% WPI) ($P < 0.01$) (Table 1).

WPI-enriched HFD normalised energy intake, while increasing the P/C ratio accentuated metabolism

The cumulative energy intake (MJ) for the dietary groups (2 cages/group, all with n = 5 mice), measured over the first 16 weeks did not differ between LFD, HFD and HFD-20% WPI groups (24.73 ± 2.70 vs. 27.69 ± 1.54 vs. 29.40 ± 0.31 , respectively). In contrast, data gathered by individually housing the mice in TSE Phenomaster cages in weeks 17–20 demonstrated that the energy intake of the HFD-WPI groups was greater than that of the HFD control group during both the light and dark phases ($P < 0.05$), while being similar to that of the LFD group (Fig. 2A). Increasing the P/C ratio had no significant effect on cumulative energy intake (MJ) in HFD fed mice up to week 16 (20%WPI, 29.40 ± 0.31 vs. 30%WPI, 30.01 ± 0.62 vs. 40%WPI, 26.71 ± 0.15). Energy intake measurements from TSE Phenomaster cages corroborated this data (Fig 2A). There was also no significant effect on meal number or meal size of altering the P/C ratio (i.e. between WPI groups; Fig 2B–C).

The HFD-20% WPI diet had no impact on VO_2 , heat production, locomotor activity or respiratory exchange ratio (RER) when compared to HFD fed mice (Fig. 3A–D). Increasing the P/C ratio was found to impact on energy expenditure with HFD-40% WPI fed mice having significantly increased levels of dark phase VO_2 compared to HFD-20 and 30% WPI fed mice ($P < 0.001$) (Fig 3A). A similar change in heat production was observed between the groups, albeit data was only significant between HFD-40% and HFD-30% WPI groups ($P < 0.05$) (Fig 3B). There was no effect of WPI or P/C ratio on locomotor activity (Fig 3C). RER values of all HFD groups were lower than the LFD group in both the light and dark phases, consistent with increased fat metabolism ($P < 0.001$) (Fig 3D).

Investigation of the above parameters in mice fed a LFD with WPI or casein for 7 weeks (study 2) revealed that WPI does not influence body weight, energy intake, VO_2 , locomotor activity or RER in a low fat background (Figure S1A–D).

Increasing the protein to carbohydrate ratio attenuated adverse metabolic impact of HFD

Specific effects of WPI and the P/C ratio were observed on lipid metabolism-related gene expression and on tissue lipid deposition. Firstly, the decrease in epididymal adipose tissue fatty acid synthase (FASN) mRNA expression with HFD feeding, was somewhat attenuated by WPI challenge ($P < 0.05$), with no added benefit of increasing the P/C ratio on expression of this gene (Fig 4A). Notably, the epididymal mRNA expression of a number of other genes were altered by the P/C ratio, specifically, fatty acid transporter 1 (FATP1), beta-3 adrenergic receptor (β_3 -AR), peroxisome proliferator-activated receptor gamma (PPAR γ), uncoupling protein 2 (UCP-2) and lipoprotein lipase (LPL) ($P \leq 0.05$) (Fig 4). In the liver, WPI specifically reduced TAG levels (Table 2) and the mRNA expression of fatty acid binding protein 1 (FABP1) compared to HFD fed mice (Table 3). The highest P/C ratio (40% WPI) significantly decreased mRNA levels of cluster of differentiation 36 (CD36) and PPAR γ ($P < 0.05$) (Table 3), dramatically reduced liver TAG levels compared to 20/30% WPI fed mice (Table 2), and normalised the elevated plasma levels of TAG and NEFA observed with HFD feeding ($P \leq 0.05$) (Table 2). Finally, the mRNA level of lipid metabolism-related

Table 1. Plasma amino acid levels ($\mu\text{mol/L}$) in mice fed a 45%kJ high fat diet (HFD) or HFD with 20%, 30% or 40% kJ whey protein isolate (WPI) for 21 weeks¹.

| | HFD | 20% WPI | 30% WPI | 40% WPI | P value |
|---------------|--------------------------------|----------------------------------|---------------------------------|--------------------------------|---------|
| Alanine | 170.91 \pm 8.82 ^a | 163.83 \pm 11.59 ^{ab} | 137.44 \pm 9.72 ^{ab} | 127.15 \pm 8.46 ^b | <.05 |
| Arginine | 41.74 \pm 4.82 | 38.41 \pm 7.16 | 56.39 \pm 7.60 | 39.08 \pm 4.34 | NS |
| Aspartic acid | 7.47 \pm 1.25 ^a | 3.63 \pm 0.51 ^b | 3.49 \pm 0.51 ^b | 4.85 \pm 1.02 ^{ab} | <.05 |
| Cysteine | 11.95 \pm 1.34 | 8.87 \pm 1.37 | 8.93 \pm 1.33 | 7.86 \pm 1.72 | NS |
| Glutamic acid | 90.76 \pm 3.42 ^a | 74.74 \pm 1.89 ^b | 75.25 \pm 1.86 ^b | 71.25 \pm 2.69 ^b | <.001 |
| Glycine | 106.22 \pm 4.28 ^a | 131.43 \pm 3.57 ^b | 116.45 \pm 4.52 ^{ab} | 131.92 \pm 2.69 ^b | <.001 |
| Histidine | 75.43 \pm 2.12 ^a | 74.93 \pm 2.98 ^a | 68.21 \pm 3.35 ^{ab} | 64.21 \pm 2.51 ^b | <.05 |
| Isoleucine | 213.81 \pm 8.87 | 207.87 \pm 9.09 | 195.27 \pm 5.74 | 186.78 \pm 10.17 | NS |
| Leucine | 91.86 \pm 9.66 | 81.36 \pm 6.65 | 66.93 \pm 6.69 | 69.71 \pm 10.21 | NS |
| Lysine | 113.81 \pm 5.88 | 108.18 \pm 8.30 | 102.57 \pm 8.39 | 88.29 \pm 4.07 | NS |
| Methionine | 23.94 \pm 1.77 | 23.65 \pm 0.75 | 21.05 \pm 0.97 | 20.40 \pm 1.08 | NS |
| Phenylalanine | 46.92 \pm 1.70 ^a | 41.13 \pm 1.13 ^{ab} | 38.23 \pm 1.52 ^{bc} | 33.30 \pm 2.35 ^c | <.001 |
| Proline | 118.26 \pm 10.14 | 127.63 \pm 9.63 | 106.04 \pm 11.02 | 104.39 \pm 11.93 | NS |
| Serine | 68.18 \pm 5.69 ^{ab} | 74.98 \pm 2.29 ^a | 62.42 \pm 2.57 ^{ab} | 55.73 \pm 2.12 ^b | <.01 |
| Threonine | 92.71 \pm 2.93 ^a | 87.63 \pm 2.49 ^a | 78.29 \pm 2.95 ^{ab} | 69.28 \pm 3.96 ^b | <.01 |
| Tyrosine | 40.29 \pm 2.47 ^a | 36.37 \pm 1.12 ^{ab} | 33.76 \pm 1.54 ^{ab} | 29.63 \pm 2.77 ^b | <.01 |
| Valine | 154.51 \pm 8.65 ^a | 130.59 \pm 5.99 ^{ab} | 118.40 \pm 6.69 ^b | 113.16 \pm 9.49 ^b | <.01 |

¹Data are means \pm SEM (n=7–10). In each row values without a common letter significantly differ, $P<0.05$; NS, non-significant.
doi:10.1371/journal.pone.0088904.t001

carnitine palmitoyltransferase 1a-c (CPT1a-c), fatty acid transport protein 5 (FATP5) and PPAR α , in tissues of interest, was not influenced by the dietary challenges (Tables 3–4).

Increasing the P/C ratio reduced plasma glucose levels, particularly in the 40% WPI group ($P<0.05$) (Table 2). In parallel HOMA-IR values were also reduced ($P<0.05$), but the change in plasma insulin concentration did not reach statistical significance (Table 2). At a cellular level, the highest P/C ratio normalised the HFD-induced reduction in adipose expression of insulin receptor (IR) and insulin receptor substrate 1 (IRS-1), and partially prevented the HFD-induced reduction in glucose transporter 4 (GLUT4) (Fig. 4B) ($P<0.001$). In the hypothalamus, WPI specifically increased IR mRNA expression ($P<0.05$), as did P/C ratio, with the highest P/C ratio having the greatest impact (Table 4). Epididymal adipose tissue mRNA expression of inflammatory markers, namely tumour necrosis factor (TNF)- α and cluster of differentiation (CD) 68 only responded to the highest P/C ratio, which significantly reduced the expression of both in a HF background (Fig. 4B) ($P<0.001$). In the hypothalamus, whilst TNF α mRNA was elevated by HFD feeding, neither WPI nor the P/C ratio influenced its levels, although there was a trend towards a decrease for the highest P/C ratio (40% WPI) (Table 4). None of the dietary challenges influenced hepatic glucose transporter 2 (GLUT2), IRS-1 and TNF- α mRNA expression (Table 3) or hypothalamic IRS-1 mRNA expression (Table 4).

The increased plasma leptin concentration in response to the HFD was significantly blunted by WPI intake with dramatic reductions seen at the highest P/C ratio ($P<0.001$) (Table 2). Yet, the hypothalamic expression of genes known to be responsive to plasma leptin levels were unaffected, specifically mRNA levels of leptin receptor (ObR), pro-opiomelanocortin (POMC), neuropeptide Y (NPY) and growth hormone secretagogue receptor (GHS-R) (Table 4). In addition, gastric mRNA expression for the orexigenic hormone ghrelin was not found to significantly differ between all dietary treatment groups (1.00 \pm 0.37, LFD vs. 1.11 \pm 0.40, HFD

vs. 0.73 \pm 0.35, 20%-WPI vs. 0.29 \pm 0.28, 30%-WPI vs. 1.07 \pm 0.45, 40%-WPI). Plasma corticosterone levels were also elevated with HF feeding, but were not influenced by protein source (WPI or casein) or P/C ratio (Table 2). Similarly, there was no effect of WPI on the HFD-associated suppression of adipose 11 β -hydroxysteroid dehydrogenase type 1 (11 β HSD1) (Fig 4B), or the HFD-induced increase in glucocorticoid receptor (GCCR) in the hypothalamus (Table 4).

WPI inclusion or increasing the P/C ratio within a HFD altered the composition of gut microbiota

A total of 251,395 V4–V5 16s rRNA sequence reads were generated which corresponded to an average of 50,279 reads per diet group or 5,130 reads per animal. α -diversity values were calculated for biodiversity (Shannon index), species richness (Chao1) and the number of species relative to the abundance in the sample (Simpson diversity index). When α -diversity values were compared by diet group, the only difference observed was a significantly higher microbial richness (Chao1) within the HFD microbiota compared to the HFD-30% WPI ($P=0.028$). Principal coordinate analysis (based on unweighted UniFrac distances) (Fig. 5) of the sequence data highlighted a clustering of the LFD, HFD and HFD-20% WPI group microbial populations, while HFD-30% and 40% WPI groups clustered in close proximity to each other and distinctly from the LFD, HFD and HFD-20% WPI group clusters. Indeed, the LFD diet with casein as a protein source clustered most closely with the HFD containing the casein protein.

Phylogenetic analysis revealed several significant microbial population shifts between the HFD control and WPI groups (Table 5). At the family level, all WPI diet groups had significantly increased proportions of *Lactobacillaceae* and significantly decreased proportions of *Clostridiaceae* compared to the HFD control group. *Bifidobacteriaceae* populations were increased in both the HFD-20%




Figure 2. Impact of whey protein isolate and protein to carbohydrate ratio on energy intake. Energy intake (A) and feeding behaviour (meal size & meal number) (B–C) was measured using TSE Phenomaster cages at 17–20 weeks for mice on either a 10% kJ low fat diet (LFD), 45% kJ high fat diet (HFD) or a HFD with 20, 30 or 40% kJ whey protein isolate (WPI). Experimental data collected from individual mice at 9 minute intervals over a 24 hour period are shown as mean values \pm SEM (n=8–10 per group) for light and dark phases. In light and dark phase, groups that do not share a common letter are significantly different at $P < 0.05$. doi:10.1371/journal.pone.0088904.g002

WPI and HFD-30% WPI diet groups compared to the HFD control, while in contrast they were significantly lower in the HFD-40% WPI group compared to the HFD control. The aforementioned patterns were also observed with respect to the corresponding genera (*Lactobacillus*, *Clostridium* and *Bifidobacterium* respectively) (Table 5). Also at genus level, proportions of *Rikenella*




Figure 3. Impact of whey protein isolate and protein to carbohydrate ratio on metabolic activity. Metabolic activity was

measured using TSE Phenomaster cages at 17–20 weeks for mice on either a 10% kJ low fat diet (LFD), 45% kJ high fat diet (HFD) or a HFD with 20, 30 or 40% kJ whey protein isolate (WPI). Experimental data for (A) oxygen consumption (VO_2), (B) heat production, (C) locomotor activity and (D) respiratory exchange ratio (RER), collected from individual mice at 9 minute intervals over a 24 hour period, are shown as mean values \pm SEM ($n=8-10$ per group) for light and dark phases. In light and dark phase, groups that do not share a common letter are significantly different at $P<0.05$.
doi:10.1371/journal.pone.0088904.g003

were significantly higher in the HFD-40% WPI group compared to HFD-20/30% WPI groups, while proportions of *Peptostreptococcus* were significantly higher in the HFD-40% WPI group than in any other diet group (Table 5). Specific comparison of the microbiota of the HFD-20% WPI and HFD control was deemed particularly important given that the changes occurring here reflected changes resulting specifically from the presence of whey protein, rather than casein, in the diet, and not simply a change in the P/C ratio in the diet. In addition to the changes in the *Lactobacillus*, *Clostridium* and *Bifidobacterium* populations (and associ-

ated families) referred to above, it was also noted that proportions of *Desulfovibrio* and *Mucispirillum* (genus) were increased in the HFD-20% WPI relative to HFD control animals.

Discussion

The key findings of this study are that WPI has a specific effect on HFD-induced energy intake, metabolic health and gut microbiota composition. Additionally, with the exception of energy intake, increasing the P/C ratio, by increasing WPI dietary content, was seen to dramatically alter the above parameters.

Energy intake

Cumulative energy intake measured up to week 16 did not significantly differ between LFD and HFD fed mice. While this is consistent with data reported elsewhere [20], HF feeding has also been shown to increase or decrease energy intake in rodents [32,33]. Differences between data reported may relate to variances in diet composition including fat source/composition, or it may be




Figure 4. Impact of whey protein isolate and protein to carbohydrate ratio on adipose cellular activity. Epididymal adipose tissue gene expression was investigated in mice after 21 weeks on a 10% kJ low fat diet (LFD), 45% kJ high fat diet (HFD), or HFD with 20, 30 or 40% kJ whey protein isolate (WPI). Relative mRNA expression is shown for (A) fatty acid synthase (FASN), fatty acid transporter 1 (FATP1), cluster of differentiation 36 (CD36), beta-3 adrenergic receptor (β 3-AR), uncoupling protein 2 (UCP-2), lipoprotein lipase (LPL) and carnitine palmitoyltransferase 1b (CPT1b), and (B) for peroxisome proliferator activated receptor gamma (PPAR γ), insulin receptor (IR), insulin receptor substrate 1 (IRS-1), glucose transporter 4 (GLUT4), 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1), tumour necrosis factor alpha (TNF- α) and cluster of differentiation 68 (CD68). Data represent mean values \pm SEM ($n=9-10$ per group). Gene expression is shown relative to the LFD control group set at 1.00. Groups that do not share a common letter are significantly different at $P<0.05$.
doi:10.1371/journal.pone.0088904.g004

Table 2. Tissue lipid parameters and plasma levels of hormones and metabolites in mice fed a 10%kJ low fat diet (LFD), 45%kJ high fat diet (HFD) or HFD with 20%, 30% or 40% kJ whey protein isolate (WPI) for 21 weeks¹.

| | LFD | HFD | 20% WPI | 30% WPI | 40%WPI | P value |
|-------------------------|-------------------------|--------------------------|--------------------------|--------------------------|--------------------------|---------|
| Liver TAG (mg/g tissue) | 71.47±9.17 ^a | 139.03±6.92 ^b | 104.94±5.88 ^c | 107.21±8.83 ^c | 70.82±7.64 ^a | <.001 |
| Leptin (ng/ml) | 4.41±0.75 ^a | 84.69±3.17 ^b | 53.78±5.67 ^c | 51.12±4.69 ^c | 15.01±4.07 ^a | <.001 |
| TAG (mg/dl) | 39.85±2.91 ^a | 56.90±6.31 ^b | 45.15±4.32 ^{ab} | 52.36±3.91 ^{ab} | 40.19±3.96 ^a | <.05 |
| NEFA (mmol/L) | 0.37±0.09 ^a | 0.67±0.07 ^b | 0.64±0.05 ^b | 0.63±0.06 ^b | 0.36±0.05 ^a | <.01 |
| Corticosterone (ng/ml) | 150.0±21.7 ^a | 319.0±48.6 ^b | 286.8±31.9 ^{ab} | 334.1±36.7 ^b | 277.1±35.5 ^{ab} | <.01 |
| GLP-1 (pM) | 25.03±2.47 | 33.15±2.83 | 28.07±2.3 | 27.88±1.98 | 24.03±2.25 | NS |
| Glucose (mmol/L) | 9.14±1.24 ^a | 15.68±0.83 ^b | 13.92±0.95 ^b | 14.08±1.05 ^b | 9.25±0.77 ^a | <.001 |
| Insulin (ng/ml) | 0.29±0.05 | 0.55±0.08 | 0.40±0.10 | 0.37±0.04 | 0.31±0.06 | NS |
| HOMA-IR | 2.48±0.67 ^a | 9.79±1.07 ^b | 6.81±2.58 ^{ab} | 6.20±0.95 ^{ab} | 3.04±0.64 ^a | <.01 |

¹Data are means ± SEM (n = 5–10). In each row results without a common letter significantly differ, $P \leq 0.05$ NS, non-significant. TAG, triacylglycerol. NEFA, non-esterified fatty acids. GLP-1, glucagon-like peptide 1. HOMA-IR, homeostasis model assessment of insulin resistance.
doi:10.1371/journal.pone.0088904.t002

due to differences in the palatability of the LFD used as the control [33,34].

In this study, there was a discrepancy in energy intake in HFD fed mice depending on the housing environment (single or group housed). In contrast to the group house environment up to week 16, HFD fed mice when individually housed in metabolic cages during weeks 17–20 showed a hypophagic response compared to LFD fed mice. It is possible that these differences may be related to the accuracy of the method used to measure food intake in group versus single housed mice, although if this was an error due to methodology, then it would likely to have influenced all dietary groups equally and not just the HFD group. Alternatively, the different behavioural responses could be result of social isolation, which has been shown to decrease energy intake and elevate plasma corticosterone levels [35,36]. However, socially isolated mice adapt to the new environment and consume similar amounts of food as pre-adapted singularly housed mice 6 hr post-novelty stress, but interestingly 24hr later, their food intake reduces significantly once again in the new environment, suggesting that stress of social isolation could extend up to 24 h [36]. We showed that group housed mice on a LFD when placed in isolation adapt to the new environment and continue to consume similar amounts of food by day 2 and 3 in the new location [8] and consequently used the day 3 time point to measure energy intake in this study. Rodents on a HFD have been shown to display increased anxiety [37], and have an over-active hypothalamic-pituitary-adrenal axis [38,39] resulting in elevated plasma corticosterone levels, as demonstrated here. This could explain why HFD fed mice are more susceptible to stress stimuli with more pronounced reductions in energy intake compared LFD fed counterparts subjected to the same stress levels [40–42]. Consequently, in a HFD background, it's possible that social isolation-induced stress responses could have had a greater impact on energy intake with effects lasting up to the 3 day housing period as used in this study. Given the finding that whey proteins such as lactoferrin and α -lactalbumin and native whey protein itself reduce stress [43–47], it is possible that replacing the casein protein with an equivalent WPI content caused normalisation of energy intake in HFD-fed mice by affecting a specific feeding behaviour related mechanism(s), with increasing the WPI protein-derived bioactives having no further effect. This WPI effect on energy intake appeared be specific to the HFD-induced neuroendocrine state because mice on the LFD with WPI showed similar energy intake to casein diet-

fed controls. Since neither WPI nor increasing the P/C ratio influenced plasma corticosterone levels, adipose expression of 11 β -HSD1 or hypothalamic expression of GCCR in HFD fed mice, it is possible that WPI may have affected other central mechanisms mediating stress responses not investigated in this study [48–50] either independently or in combination with key mechanisms regulating energy balance. Given that leptin decreases meal size and number [51–53], and WPI reduced the HFD-induced increase in plasma leptin levels, it's possible that WPI-derived bioactives could have specifically influenced circadian rhythm of leptin production and/or action within the neuroendocrine state of HFD-fed mice in a socially isolated environment. Additionally, the reduction in plasma amino acids associated with WPI intake (see below), could also have acted as a possible central trigger to increase energy intake in WPI groups compared to HFD control in the single house environment.

HFD feeding has been shown to cause a gain of weight in rats up to the duration of a test period lasting 76 weeks, with animal's body weight gain responding to changes to the dietary fat content introduced at various time points [54]. Lin *et al.*, [32] demonstrated that mice on a HFD for 19 weeks are responsive to intracerebroventricular administration of leptin. These data suggest that energy balance related mechanisms are able to respond to energetic challenges even after prolonged high fat intake. High protein intake within a HFD suppresses energy intake [55,56] albeit not consistently [4,57], and in our study, the P/C ratio did not alter energy intake in either housing environment. This could be a result of the quantity or composition of the macronutrient used in the test diets. Indeed, data from human trials showed that increasing protein dietary content (10/15% to 30%) only decreased energy intake when the carbohydrate content was kept constant [58,59]. This further highlights the importance of designing appropriate experimental diets with the correct macronutrient composition for uncovering the energy balance related impact of the dietary component under investigation.

Metabolic health

Replacement of the casein protein with an equivalent energy content of WPI (i.e. 20%) did not specifically alter metabolic activity, heat production or locomotor activity in HFD or LFD fed mice. In contrast, Acheson *et al.*, [60] showed that whey has a greater thermic effect than casein or soy in humans. These

Table 3. Relative hepatic gene expression in mice fed a 10%kJ low fat diet (LFD), 45%kJ high fat diet (HFD) or HFD with 20%, 30% or 40% kJ whey protein isolate (WPI) for 21 weeks¹.

| | LFD | HFD | 20% WPI | 30% WPI | 40% WPI | P value |
|---------------|-------------------------|------------------------|-------------------------|-------------------------|-------------------------|---------|
| CD36 | 1.00±0.20 ^a | 2.85±0.34 ^b | 2.98±0.45 ^b | 2.17±0.25 ^{ab} | 1.50±0.25 ^a | <.05 |
| PPAR γ | 1.00±0.12 ^a | 3.00±0.57 ^b | 2.30±0.49 ^b | 1.88±0.46 ^b | 0.85±0.13 ^a | <.001 |
| FABP1 | 1.00±0.09 ^{ab} | 1.27±0.10 ^b | 0.86±0.06 ^{ac} | 1.27±0.17 ^{bc} | 0.71±0.09 ^a | <.001 |
| IRS-1 | 1.00±0.15 ^a | 0.58±0.04 ^b | 0.69±0.05 ^{ab} | 0.68±0.06 ^{ab} | 0.79±0.06 ^{ab} | <.01 |
| CPT1a | 1.00±0.08 | 0.85±0.08 | 0.97±0.06 | 0.90±0.13 | 0.86±0.10 | NS |
| GLUT2 | 1.00±0.09 | 0.76±0.04 | 1.02±0.08 | 0.97±0.09 | 0.89±0.05 | NS |
| FATP5 | 1.00±0.21 | 0.89±0.07 | 0.88±0.07 | 1.16±0.07 | 1.24±0.12 | NS |
| FASN | 1.00±0.36 | 0.46±0.14 | 0.54±0.19 | 0.40±0.06 | 0.32±0.04 | NS |
| PPAR α | 1.00±0.04 | 1.13±0.05 | 1.21±0.05 | 1.17±0.10 | 1.28±0.11 | NS |
| TNF α | 1.00±0.26 | 0.84±0.15 | 1.12±0.18 | 0.80±0.09 | 0.71±0.10 | NS |

¹Data are means \pm SEM (n=7–10). In each row results without a common letter significantly differ, $P<0.05$; NS, non-significant. Gene expression shown relative to the LFD control group set at 1.00. CD36, cluster of differentiation 36; PPAR γ , peroxisome proliferator activated receptor gamma; FABP1, Fatty acid binding protein 1; IRS-1, Insulin receptor substrate 1; CPT1a, carnitine palmitoyltransferase 1a; GLUT2, Glucose transporter 2; FATP5, Fatty acid transporter 5; FASN, Fatty acid synthase; TNF- α , Tumour necrosis factor alpha.

doi:10.1371/journal.pone.0088904.t003

differences in data may be related to the fact that the latter study only investigated an acute post-prandial response to a defined test meal, or it may relate to how different species (humans versus mice) digest and metabolise dietary proteins. Shetzer *et al.* found that mice consuming a HFD and WPI-supplemented drinking water have enhanced oxygen consumption compared to mice drinking unsupplemented water [4]. In this instance, the increased metabolic activity may have arisen due to the increased protein intake (proteins from diet and from WPI supplemented water). In fact, this corroborates with the data presented here, which show that increasing the P/C ratio resulted in increased energy expenditure (VO₂ and heat production) and dark phase locomotor activity, resulting presumably from the increased catabolism of ingested dietary protein, coupled with thermic effects of WPI compared to casein [60] and/or due to increased deposition of lean mass with WPI content [61–63]. Interestingly, Zhang *et al.* [64] showed that HFD fed mice on leucine-supplemented drinking

water have reduced fasting plasma levels of aspartic acid, glutamic acid, and phenylalanine, as well as increased VO₂ and reduced adiposity compared to HFD controls. Given the influence of leucine on WPI-induced muscle hypertrophy [65,66] and its unique ability to regulate the translation of protein synthesis [67], it is possible that the elevated leucine content found normally in the WPI diets may have enhanced muscle protein synthesis by directing other amino acids towards protein synthesis and/or catabolism [68,69], with the required energy been derived possibly from fat catabolism [70]. Consistent with the latter suggestion, we found an increased lean mass and a trend towards a reduction in fat mass with decreased plasma levels of several amino acids, but not leucine, when the casein protein in a HFD was replaced with WPI or when the P/C ratio in the HFD was increased.

WPI intake appeared to cause a trend towards a reduction in fat mass, and in the liver this manifested as a WPI specific reduction

Table 4. Relative hypothalamic gene expression in mice fed a 10%kJ low fat diet (LFD), 45%kJ high fat diet (HFD) or HFD with 20%, 30% or 40% kJ whey protein isolate (WPI) for 21 weeks¹.

| | LFD | HFD | 20% WPI | 30% WPI | 40% WPI | P value |
|---------------|-------------------------|------------------------|-------------------------|-------------------------|-------------------------|---------|
| IR | 1.00±0.03 ^{ab} | 0.97±0.07 ^a | 1.22±0.04 ^{bc} | 1.37±0.04 ^{cd} | 1.55±0.08 ^d | <.05 |
| GCCR | 1.00±0.09 ^a | 1.40±0.10 ^b | 1.34±0.07 ^{ab} | 1.30±0.08 ^{ab} | 1.42±0.09 ^b | <.05 |
| TNF α | 1.00±0.05 ^a | 1.76±0.09 ^b | 1.57±0.13 ^b | 1.47±0.14 ^{ab} | 1.40±0.25 ^{ab} | <.01 |
| POMC | 1.00±0.08 | 0.99±0.06 | 1.01±0.06 | 0.98±0.06 | 1.00±0.08 | NS |
| NPY | 1.00±0.04 | 1.02±0.09 | 0.90±0.06 | 0.88±0.05 | 0.92±0.08 | NS |
| ObR | 1.00±0.11 | 0.92±0.05 | 0.99±0.05 | 1.07±0.07 | 0.95±0.05 | NS |
| GHS-R | 1.00±0.07 | 0.98±0.06 | 0.98±0.08 | 1.02±0.11 | 1.04±0.09 | NS |
| PPAR γ | 1.00±0.36 | 1.07±0.04 | 0.95±0.05 | 0.99±0.05 | 0.98±0.07 | NS |
| CPT1c | 1.00±0.09 | 0.92±0.04 | 0.91±0.01 | 0.97±0.06 | 0.86±0.04 | NS |
| IRS-1 | 1.00±0.04 | 0.99±0.04 | 0.97±0.02 | 1.06±0.08 | 1.00±0.06 | NS |

¹Data are means \pm SEM (n=5–10). In each row values without a common letter significantly differ, $P<0.05$; NS, non-significant. Gene expression shown relative to the LFD control group set at 1.00. IR, Insulin receptor; GCCR, Glucocorticoid receptor; TNF- α , Tumour necrosis factor alpha; POMC, Pro-opiomelanocortin; NPY, Neuropeptide Y; ObR, Leptin receptor; GHS-R, Growth hormone secretatgogue receptor; PPAR γ , peroxisome proliferator activated receptor gamma; CPT1c, carnitine palmitoyltransferase 1c; IRS-1, Insulin receptor substrate 1.

doi:10.1371/journal.pone.0088904.t004




Figure 5. Impact of whey protein isolate and protein to carbohydrate ratio on the gut microbiota composition. Principal Coordinate analysis (PCoA) of unweighted Unifrac distances of the 16srRNA sequences, demonstrating where sequences cluster according to diet group. Data were generated from analysis of faecal samples collected from mice on 10% kJ low fat diet (LFD, Δ) or 45% kJ high fat diet (HFD, \blacklozenge) or a HFD with 20% kJ whey protein isolate (HFD-20% WPI, \circ), 30% kJ WPI (HFD-30% WPI, \diamond) or 40% kJ WPI (HFD-40% WPI, \bullet) (n = 10). doi:10.1371/journal.pone.0088904.g005

in TAG levels, which was accompanied by the suppression of FABP1 mRNA expression, similar to previous findings [4,6,71,72]. In the epididymal adipose tissue, WPI prevented the HFD-induced FASN gene expression, albeit a recent study reported that WPI does not affect the weight of the epididymal tissue in HFD fed mice, but instead causes a reduction in subcutaneous fat pad weight [20]. These data suggest that WPI affects cellular activity in the liver and in specific adipose tissue depots. While it has been suggested that whey protein may facilitate enhanced postprandial chylomicron clearance via an alteration in LPL expression/activity [73,74], here we did not find a WPI specific effect on LPL expression or plasma TAG levels, but we did observe that intake of the highest P/C ratio (40%-WPI) led to an increase adipose tissue LPL mRNA expression which was accompanied by significant reduction in plasma TAG levels, and a complete reversal of genes involved in lipid accumulation (PPAR γ), fatty acid transport (FATP1), and lipolysis (β 3-AR). Given that HF feeding/obesity down-regulates β 3-AR mRNA expression [75], our data suggests an increased adipocyte lipolysis, and reduction in adipose TAG storage in HFD-40%WPI fed mice. Yet the endogenous CPT1b-associated β -oxidation pathway and the UCP-2-associated pathway in epididymal adipose tissue seem to be unaffected (CPT1b) or suppressed (UCP-2) by raising the P/C ratio. This data raises the possibility that the free fatty acids generated from the potentially increased availability of β 3-AR in the adipose may have been re-directed for utilisation by other physiological processes active in HFD-40% group, possibly leading to the increased metabolic activity (VO_2) observed in the animals. It is also noteworthy that WPI has been shown to increase faecal fat excretion compared to casein [5], which may have also contributed to the decreased plasma TAG and NEFA seen here with intake of the highest P/C ratio diet (HFD-40%WPI).

Given the link between HFD-induced obesity, low-grade inflammation and insulin resistance [76,77], one could argue that

the dramatic reduction in fat mass observed with the highest P/C ratio may underlie the effects on inflammatory markers in the adipose tissue (TNF α and CD68) and the hypothalamus (TNF α), along with simultaneous changes in expression of genes involved in insulin signalling (IR, IRS-1 and GLUT4 in the adipose, IRS-1 in liver and IR in the hypothalamus), and the reduction in plasma glucose in these mice. Improvements to insulin sensitivity with WPI have been reported previously [9,78], but our data suggested that only an increased P/C ratio in the HFD facilitated improvements to insulin signalling pathway associated gene expression, particularly in the adipose, in parallel with reduced HOMA-IR values.

Composition of gut microbiota

While many of the effects described above may be due to direct WPI or P/C ratio-host interactions, the effect of WPI and P/C ratio on the composition of the gut microbiota may also play an important role in adiposity and weight gain in these animals. Here, high throughput sequencing based analyses of faecal microbial populations revealed the clustering of the microbiota from animals in receipt of 30 and 40% WPI diets away from those in receipt of 20% kJ WPI or HFD-casein diets. Tranberg *et al* [20] recently suggested that the efficient absorption of dairy whey proteins in the small intestine may explain the absence of changes in the faecal microbiota. This may explain the clustering of the microbiota from animals fed 20% WPI or HFD-casein diets in our study. However, it is apparent that the high concentrations of WPI present in the 30 and 40% WPI diets employed in our study had a more profound effect, possibly due to additional whey proteins finding their way to the large intestine and/or the overall change in the P/C ratio in the diet. Consumption of the 30 and 40% WPI diets did not result in a shift in the microbiota toward that of the LFD animals and thus the effects on weight gain are not simply due to an overall

Table 5. Gut microbiota composition as % of reads in mice fed a 45%kJ high fat diet (HFD) or HFD with 20%, 30% or 40% kJ whey protein isolate (WPI) for 21 weeks¹.

| | HFD | 20% WPI | 30% WPI | 40% WPI |
|------------------------------|--------------------|--------------------|--------------------|--------------------|
| Phylum | | | | |
| <i>Proteobacteria</i> | 0.36 ^a | 0.63 ^b | 0.34 ^{ab} | 0.32 ^a |
| <i>Actinobacteria</i> | 0.63 ^a | 1.82 ^b | 3.79 ^b | 0.36 ^c |
| <i>Deferribacteres</i> | 0.57 ^a | 1.61 ^b | 1.56 ^{ab} | 2.03 ^b |
| Family | | | | |
| <i>Desulfovibrionaceae</i> | 0.12 ^a | 0.31 ^b | 0.21 ^{ab} | 0.23 ^{ab} |
| <i>Rikenellaceae</i> | 6.71 ^{ab} | 7.54 ^b | 3.9 ^a | 6.4 ^{ab} |
| <i>Bacteroidaceae</i> | 0.44 ^a | 0.42 ^a | 0.16 ^b | 0.21 ^b |
| <i>Lactobacillaceae</i> | 0.21 ^a | 3.03 ^b | 4.6 ^b | 2.14 ^b |
| <i>Bifidobacteriaceae</i> | 0.43 ^a | 1.71 ^b | 3.66 ^b | 0.22 ^c |
| <i>Deferribacteraceae</i> | 0.57 ^a | 1.59 ^b | 1.32 ^{ab} | 2.03 ^{ab} |
| <i>Peptostreptococcaceae</i> | 0.62 ^a | 1.79 ^a | 1.54 ^a | 8.01 ^b |
| <i>Succinivibrionaceae</i> | 0.13 ^a | 0.15 ^a | 0 ^b | 0 ^b |
| <i>Clostridiaceae</i> | 1.31 ^a | 0 ^b | 0 ^b | 0 ^b |
| <i>Veillonellaceae</i> | 0.02 ^a | 0.12 ^b | 0 ^a | 0 ^a |
| Genus | | | | |
| <i>Anaerobiospirillum</i> | 0.13 ^a | 0.15 ^a | 0 ^b | 0 ^b |
| <i>Desulfovibrio</i> | 0.07 ^a | 0.22 ^b | 0.17 ^{ab} | 0.15 ^{ab} |
| <i>Alistipes</i> | 4.33 ^{ab} | 4.41 ^a | 2.24 ^b | 3.76 ^{ab} |
| <i>Rikenella</i> | 1.04 ^{ab} | 0.49 ^b | 0.68 ^b | 1.08 ^b |
| <i>Bacteroides</i> | 0.44 ^a | 0.4 ^a | 0.16 ^b | 0.21 ^{ab} |
| <i>Oscillibacter</i> | 0.24 ^a | 0.67 ^{ab} | 0.42 ^{ab} | 0.52 ^b |
| <i>Lactobacillus</i> | 0.2 ^a | 3.03 ^b | 4.6 ^b | 2.39 ^b |
| <i>Bifidobacterium</i> | 0.43 ^a | 1.71 ^b | 3.66 ^b | 0.22 ^c |
| <i>Mucispirillum</i> | 0.57 ^a | 1.61 ^b | 1.56 ^{ab} | 1.92 ^{ab} |
| <i>Coprococcus</i> | 0.11 ^{ab} | 0.23 ^b | 0.06 ^a | 0.06 ^{ab} |
| <i>Turicibacter</i> | 0.56 ^a | 0.35 ^a | 0.15 ^{ab} | 0 ^b |
| <i>Clostridium</i> | 1.3 ^a | 0 ^b | 0 ^b | 0 ^b |
| <i>Peptostreptococcus</i> | 0.1 ^a | 0.14 ^a | 0.12 ^a | 0.78 ^b |

¹Data are means \pm SEM (n=10). Statistically significant differences generated using the Kruskal-Wallis algorithm. In each row values without a common letter significantly differ, $P \leq 0.05$.

doi:10.1371/journal.pone.0088904.t005

conversion to a LFD-like microbiota. Specific taxonomic changes were also noted in response to the different diets. In all cases dietary WPI resulted in significant increases in *Lactobacillaceae/Lactobacillus* and decreases in *Clostridiaceae/Clostridium*. Increased proportions of *Lactobacillus* have previously also been observed in a study of individuals following a regime of calorie restriction and exercise [79]. However, in contrast, increased proportions of *Lactobacillus* have also been noted in HFD fed rats [80] and diet-induced obese mice [81]. While specific species of *Lactobacillus* have been associated with both lean and obese gut microbiota profiles and also to play a role in obesity and immune response regulation [82–84], due to the length of the 16S sequences generated and the high degree of sequence homology, we cannot assess changes in proportions of *Lactobacillus* at the species level. An increase in the proportions of *Bifidobacteriaceae/Bifidobacterium* was also observed in both the HFD-20% WPI and HFD-30% WPI compared to the HFD group. This result, combined with the aforementioned increases in *Lactobacillaceae/Lactobacillus*, mirror those reported by

Sprong *et al* who suggest that whey proteins act as grow factors for certain species of bacteria by an amino acid composition mediated mechanism [19]. This pattern did not extend to the HFD-40% WPI group suggesting that, at these high protein levels, other factors are at play. Our observations are also consistent with previous findings that high proportions of the class *Clostridiales* are associated with the gut microbiota of animals fed a HFD [85], while fasting reduces the levels of *Clostridium* [86]. Notably, *Clostridiaceae* can produce short chain fatty acids as a product of their metabolism [87], which can play an important role in the regulation of immune cells and has been associated with inflammation and obesity [88]. These differences, as well as others in the *Proteobacteria Actinobacteria Deferribacteres* (phylum), *Desulfovibrionaceae Deferribacteraceae Veillonellaceae* (family), *Desulfovibrio* and *Mucispirillum* taxa in the HFD-20% whey protein relative to HFD controls (20% casein) are particularly notable as these reflect changes resulting from the specific presence of whey proteins in the diet, in place of casein, rather than changes in the P/C ratio. Changes in relative proportions may be attributed to (a) the ability of bacteria to utilise whey proteins as a growth medium, (b) the anti-microbial activity of whey protein/peptide components, (c) decreased competition as a result of the whey proteins/peptides antimicrobial activity or (d) whey protein mediated changes in the host. Ultimately, the question of cause versus effect remains unanswered, and so while the changes to the microbiota observed may contribute to the mechanisms involved in controlling weight gain, further studies with, for example germ free animals, will be required to determine this definitively.

WPI effects on energy balance from a whole animal context

Focusing on the experimental data gathered between weeks 17–21, during which we measured metabolic parameters, faecal microbial population, body composition and tissue and plasma level of energy balance related parameters, it is clear that WPI intake increased energy intake associated with the HFD, without altering energy expenditure, as measured by VO_2 and locomotor activity. However, the body composition and body weight in the HFD-WPI group does not appear to reflect a positive energy balance, as animals showed a trend towards a reduction in fat mass and increased lean mass. It is noteworthy in this regards that WPI has been shown to increase faecal fat excretion [5] and we also observed some subtle changes in the gut microbiota at a phylogenetic level that are associated with non-obese states, raising the possibility of a reduced intestinal TAG absorption in the HFD-WPI groups with increased energy intake, leading to similar body weight trajectories as HFD controls. Increasing the P/C ratio by changing WPI from 30 to 40% did not alter energy intake but significantly accentuated energy expenditure with a concurrent dramatic change in physiology.

In summary, our results show that WPI specifically normalises energy intake, increases lean mass and causes a trend towards a reduction in fat mass associated with prolonged high fat feeding. Raising the P/C ratio had no effect on energy intake but augmented metabolic activity and beneficially altered gene expression profiles for lipid metabolism, inflammation and insulin signalling, particularly in the adipose tissue. High throughput analysis of gut microbiota revealed distinct changes in microbial populations with increased P/C ratio causing clustering of 30/40% WPI groups together and distinct from those of HFD and 20% WPI groups, but with specific phylogenetic differences existing between the latter groups. These data indicate that changes to P/C ratio have a dramatic effect on energy balance and the composition of gut microbiota distinct from that seen with

changes to protein source. Future studies should focus on determining whether the effects demonstrated for highest P/C ratio are specific to the WPI content, a consequence of macronutrient change, or both.

Supporting Information

Figure S1 Effect of a 10%kJ low fat diet with 20%kJ casein (LFD) or 20%kJ whey protein isolate (LFD-WPI) on (A) body weight, (B) energy intake (C) oxygen consumption (VO₂) and (D) locomotor activity, which were measured in individual mice at 9 minute intervals over a 24 hour period using TSE Phenomaster cages.

(DOC)

References

- Caterson ID, Hubbard V, Bray GA, Grunstein R, Hansen BC, et al. (2004) Prevention Conference VII: Obesity, a worldwide epidemic related to heart disease and stroke: Group III: worldwide comorbidities of obesity. *Circulation* 110: e476–483.
- Guo J, Jou W, Gavrilova O, Hall KD (2009) Persistent diet-induced obesity in male C57BL/6 mice resulting from temporary obesigenic diets. *PLoS one* 4: e5370.
- Ryan KK, Woods SC, Seeley RJ (2012) Central nervous system mechanisms linking the consumption of palatable high-fat diets to the defense of greater adiposity. *Cell metabolism* 15: 137–149.
- Shertzer HG, Woods SE, Krishan M, Genter MB, Pearson KJ (2011) Dietary whey protein lowers the risk for metabolic disease in mice fed a high-fat diet. *The Journal of nutrition* 141: 582–587.
- Pilvi TK, Korpela R, Huttunen M, Vapaatalo H, Mervaala EM (2007) High-calcium diet with whey protein attenuates body-weight gain in high-fat-fed C57Bl/6J mice. *British Journal of Nutrition* 98: 900–907.
- Shi J, Tauriainen E, Martonen E, Finckenberg P, Ahlroos-Lehmus A, et al. (2011) Whey protein isolate protects against diet-induced obesity and fatty liver formation. *International Dairy Journal* 21: 513–522.
- Shi J, Ahlroos-Lehmus A, Pilvi TK, Korpela R, Tossavainen O, et al. (2012) Metabolic effects of a novel microfiltered native whey protein in diet-induced obese mice. *Journal of functional foods* 4: 440–449.
- McAllan L, Keane D, Schellekens H, Roche HM, Korpela R, et al. (2013) Whey protein isolate counteracts the effects of a high-fat diet on energy intake and hypothalamic and adipose tissue expression of energy balance-related genes. *The British journal of nutrition*: 1–13.
- Pichon L, Potier M, Tome D, Mikogami T, Laplaize B, et al. (2008) High-protein diets containing different milk protein fractions differently influence energy intake and adiposity in the rat. *Br J Nutr* 99: 739–748.
- Turnbaugh PJ, Ridaura VK, Faith JJ, Rey FE, Knight R, et al. (2009) The effect of diet on the human gut microbiome: a metagenomic analysis in humanized gnotobiotic mice. *Science translational medicine* 1: 6ra14.
- Wu GD, Chen J, Hoffmann G, Bittinger K, Chen YY, et al. (2011) Linking long-term dietary patterns with gut microbial enterotypes. *Science* 334: 105–108.
- Backhed F, Ding H, Wang T, Hooper LV, Koh GY, et al. (2004) The gut microbiota as an environmental factor that regulates fat storage. *Proceedings of the National Academy of Sciences of the United States of America* 101: 15718–15723.
- Murphy EF, Cotter PD, Healy S, Marques TM, O'Sullivan O, et al. (2010) Composition and energy harvesting capacity of the gut microbiota: relationship to diet, obesity and time in mouse models. *Gut* 59: 1635–1642.
- Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, et al. (2006) An obesity-associated gut microbiome with increased capacity for energy harvest. *Nature* 444: 1027–1031.
- Freedman DJ, Tackett CO, Delehanty A, Maneval DR, Nataro J, et al. (1998) Milk immunoglobulin with specific activity against purified colonization factor antigens can protect against oral challenge with enterotoxigenic *Escherichia coli*. *The Journal of infectious diseases* 177: 662–667.
- Shin K, Tomita M, Lonnerdal B (2000) Identification of lactoperoxidase in mature human milk. *The Journal of nutritional biochemistry* 11: 94–102.
- Yamauchi K, Tomita M, Giehl TJ, Ellison RT, 3rd (1993) Antibacterial activity of lactoferrin and a pepsin-derived lactoferrin peptide fragment. *Infection and immunity* 61: 719–728.
- Hoek KS, Milne JM, Grieve PA, Dionysius DA, Smith R (1997) Antibacterial activity in bovine lactoferrin-derived peptides. *Antimicrobial agents and chemotherapy* 41: 54–59.
- Sprong RC, Schonewille AJ, van der Meer R (2010) Dietary cheese whey protein protects rats against mild dextran sulfate sodium-induced colitis: role of mucin and microbiota. *Journal of dairy science* 93: 1364–1371.
- Tranberg B, Hellgren LI, Lykkesfeldt J, Sejrnsen K, Jeamet A, et al. (2013) Whey protein reduces early life weight gain in mice fed a high-fat diet. *PLoS one* 8: e71439.
- Weir JB (1949) New methods for calculating metabolic rate with special reference to protein metabolism. *The Journal of physiology* 109: 1–9.
- Claesson MJ, Wang Q, O'Sullivan O, Greene-Diniz R, Cole JR, et al. (2010) Comparison of two next-generation sequencing technologies for resolving highly complex microbiota composition using tandem variable 16S rRNA gene regions. *Nucleic acids research* 38: e200.
- Rea MC, Dobson A, O'Sullivan O, Crispie F, Foully F, et al. (2011) Effect of broad- and narrow-spectrum antimicrobials on *Clostridium difficile* and microbial diversity in a model of the distal colon. *Proceedings of the National Academy of Sciences of the United States of America* 108 Suppl 1: 4639–4644.
- Urich T, Lanzan A, Qi J, Huson DH, Schleper C, et al. (2008) Simultaneous assessment of soil microbial community structure and function through analysis of the meta-transcriptome. *PLoS one* 3: e2527.
- Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, et al. (2010) QIIME allows analysis of high-throughput community sequencing data. *Nature methods* 7: 335–336.
- Haas BJ, Gevers D, Earl AM, Feldgarden M, Ward DV, et al. (2011) Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons. *Genome research* 21: 494–504.
- Price MN, Dehal PS, Arkin AP (2010) FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS one* 5: e9490.
- Chen VB, Davis IW, Richardson DC (2009) KING (Kinemage, Next Generation): a versatile interactive molecular and scientific visualization program. *Protein science : a publication of the Protein Society* 18: 2403–2409.
- Huson DH, Richter DC, Rausch C, DeZulian T, Franz M, et al. (2007) Dendroscope: An interactive viewer for large phylogenetic trees. *BMC bioinformatics* 8: 460.
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, et al. (1985) Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 28: 412–419.
- Folch J, Lees M, Sloane Stanley GH (1957) A simple method for the isolation and purification of total lipides from animal tissues. *The Journal of biological chemistry* 226: 497–509.
- Lin S, Thomas TC, Storlien LH, Huang XF (2000) Development of high fat diet-induced obesity and leptin resistance in C57Bl/6J mice. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 24: 639–646.
- Sclafani A, Weiss K, Cardieri C, Ackroff K (1993) Feeding response of rats to no-fat and high-fat cakes. *Obesity research* 1: 173–178.
- West DB, York B (1998) Dietary fat, genetic predisposition, and obesity: lessons from animal models. *The American journal of clinical nutrition* 67: 505S–512S.
- Saegusa Y, Takeda H, Muto S, Nakagawa K, Ohnishi S, et al. (2011) Decreased plasma ghrelin contributes to anorexia following novelty stress. *American journal of physiology Endocrinology and metabolism* 301: E685–696.
- Yamada C, Saegusa Y, Nakagawa K, Ohnishi S, Muto S, et al. (2013) Rikkunshito, a Japanese Kampo Medicine, Ameliorates Decreased Feeding Behavior via Ghrelin and Serotonin 2B Receptor Signaling in a Novelty Stress Murine Model. *Biomed Res Int* 2013: 792940.
- Buchenaue T, Behrendt P, Bode FJ, Horn R, Brabant G, et al. (2009) Diet-induced obesity alters behavior as well as serum levels of corticosterone in F344 rats. *Physiol Behav* 98: 563–569.
- Dallman MF (2010) Stress-induced obesity and the emotional nervous system. *Trends in endocrinology and metabolism*: TEM 21: 159–165.
- Sharma S, Fulton S (2013) Diet-induced obesity promotes depressive-like behaviour that is associated with neural adaptations in brain reward circuitry. *Int J Obes (Lond)* 37: 382–389.

Table S1 The composition of the low fat diet (LFD), high fat diet (HFD) and HFD with 20% kJ, 30% kJ or 40% kJ whey protein isolate (WPI)¹.

(DOC)

Table S2 Sequences of mouse specific primers used in real-time PCR analysis¹.

(DOC)

Author Contributions

Conceived and designed the experiments: LM KNN JFC PDC PS RPR POC GF HMR. Performed the experiments: LM PS POC. Analyzed the data: LM KNN JFC PDC PS RPR POC GF HMR. Contributed reagents/materials/analysis tools: KNN PDC PS POC HMR. Wrote the paper: LM KNN JFC PDC PS RPR POC GF HMR. Ethical Approval: KNN JFC RPR GF HMR.

40. Finger BC, Dinan TG, Cryan JF (2011) High-fat diet selectively protects against the effects of chronic social stress in the mouse. *Neuroscience* 192: 351–360.
41. Harris RB, Zhou J, Youngblood BD, Rybkin II, Smagin GN, et al. (1998) Effect of repeated stress on body weight and body composition of rats fed low- and high-fat diets. *Am J Physiol* 275: R1928–R1938.
42. Zhou J, Yan X, Ryan DH, Harris RB (1999) Sustained effects of repeated restraint stress on muscle and adipocyte metabolism in high-fat-fed rats. *Am J Physiol* 277: R757–766.
43. Kamemori N, Takeuchi T, Hayashida K, Harada E (2004) Suppressive effects of milk-derived lactoferrin on psychological stress in adult rats. *Brain research* 1029: 34–40.
44. Markus CR, Olivier B, de Haan EH (2002) Whey protein rich in alpha-lactalbumin increases the ratio of plasma tryptophan to the sum of the other large neutral amino acids and improves cognitive performance in stress-vulnerable subjects. *The American journal of clinical nutrition* 75: 1051–1056.
45. Markus CR, Olivier B, Panhuysen GE, Van Der Gugten J, Alles MS, et al. (2000) The bovine protein alpha-lactalbumin increases the plasma ratio of tryptophan to the other large neutral amino acids, and in vulnerable subjects raises brain serotonin activity, reduces cortisol concentration, and improves mood under stress. *The American journal of clinical nutrition* 71: 1536–1544.
46. Takeuchi T, Hayashida K, Inagaki H, Kuwahara M, Tsubone H, et al. (2003) Opioid mediated suppressive effect of milk-derived lactoferrin on distress induced by maternal separation in rat pups. *Brain research* 979: 216–224.
47. Vekovisheva OY, Peuhkuri K, Backstrom P, Sihvola N, Pilvi T, et al. (2013) The effects of native whey and alpha-lactalbumin on the social and individual behaviour of C57BL/6J mice. *Br J Nutr* 110: 1336–1346.
48. Herman JP, Cullinan WE (1997) Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. *Trends Neurosci* 20: 78–84.
49. Ghosal S, Myers B, Herman JP (2013) Role of central glucagon-like peptide-1 in stress regulation. *Physiol Behav* 122: 201–207.
50. Viau V, Chu A, Soriano L, Dallman MF (1999) Independent and overlapping effects of corticosterone and testosterone on corticotropin-releasing hormone and arginine vasopressin mRNA expression in the paraventricular nucleus of the hypothalamus and stress-induced adrenocorticotrophic hormone release. *J Neurosci* 19: 6684–6693.
51. Bady I, Marty N, Dallaporta M, Emery M, Gyger J, et al. (2006) Evidence from Glut2-null mice that glucose is a critical physiological regulator of feeding. *Diabetes* 55: 988–995.
52. Hulsey MG, Lu H, Wang T, Martin RJ, Baile CA (1998) Intracerebroventricular (i.c.v.) administration of mouse leptin in rats: behavioral specificity and effects on meal patterns. *Physiology & behavior* 65: 445–455.
53. Brown LM, Clegg DJ, Benoit SC, Woods SC (2006) Intraventricular insulin and leptin reduce food intake and body weight in C57BL/6J mice. *Physiology & behavior* 89: 687–691.
54. Peckham SC, Entenman C, Carroll HW (1962) The influence of a hypercaloric diet on gross body and adipose tissue composition in the rat. *The Journal of nutrition* 77: 187–197.
55. Freudenberg A, Petzke KJ, Klaus S (2012) Comparison of high-protein diets and leucine supplementation in the prevention of metabolic syndrome and related disorders in mice. *The Journal of nutritional biochemistry* 23: 1524–1530.
56. Pichon L, Huneau JF, Fromentin G, Tome D (2006) A high-protein, high-fat, carbohydrate-free diet reduces energy intake, hepatic lipogenesis, and adiposity in rats. *The Journal of nutrition* 136: 1256–1260.
57. Schwarz J, Tome D, Baars A, Hooiveld GJ, Muller M (2012) Dietary protein affects gene expression and prevents lipid accumulation in the liver in mice. *PLoS One* 7: e47303.
58. Weigle DS, Breen PA, Matthys CC, Callahan HS, Meeuws KE, et al. (2005) A high-protein diet induces sustained reductions in appetite, ad libitum caloric intake, and body weight despite compensatory changes in diurnal plasma leptin and ghrelin concentrations. *The American journal of clinical nutrition* 82: 41–48.
59. Blatt AD, Roe LS, Rolls BJ (2011) Increasing the protein content of meals and its effect on daily energy intake. *Journal of the American Dietetic Association* 111: 290–294.
60. Acheson KJ, Blondel-Lubrano A, Oguey-Araymon S, Beaumont M, Emady-Azar S, et al. (2011) Protein choices targeting thermogenesis and metabolism. *Am J Clin Nutr*.
61. Kanda A, Nakayama K, Fukasawa T, Koga J, Kanegae M, et al. (2013) Post-exercise whey protein hydrolysate supplementation induces a greater increase in muscle protein synthesis than its constituent amino acid content. *The British journal of nutrition*: 1–7.
62. Pennings B, Boirie Y, Senden JM, Gijzen AP, Kuipers H, et al. (2011) Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men. *The American journal of clinical nutrition* 93: 997–1005.
63. Tipton KD, Elliott TA, Cree MG, Aarsland AA, Sanford AP, et al. (2007) Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *American Journal of Physiology-Endocrinology and Metabolism* 292: E71–E76.
64. Zhang Y, Guo K, LeBlanc RE, Loh D, Schwartz GJ, et al. (2007) Increasing dietary leucine intake reduces diet-induced obesity and improves glucose and cholesterol metabolism in mice via multimechanisms. *Diabetes* 56: 1647–1654.
65. Norton LE, Wilson GJ, Layman DK, Moulton CJ, Garlick PJ (2012) Leucine content of dietary proteins is a determinant of postprandial skeletal muscle protein synthesis in adult rats. *Nutrition & Metabolism* 9.
66. Rieu I, Balage M, Sornet C, Debras E, Ripes S, et al. (2007) Increased availability of leucine with leucine-rich whey proteins improves postprandial muscle protein synthesis in aging rats. *Nutrition* 23: 323–331.
67. Norton LE, Layman DK (2006) Leucine regulates translation initiation of protein synthesis in skeletal muscle after exercise. *The Journal of nutrition* 136: 533S–537S.
68. Bender DA (2012) The metabolism of “surplus” amino acids. *The British journal of nutrition* 108 Suppl 2: S113–121.
69. Garlick PJ, McNurlan MA, Patlak CS (1999) Adaptation of protein metabolism in relation to limits to high dietary protein intake. *European journal of clinical nutrition* 53 Suppl 1: S34–43.
70. McAllan L, Cotter PD, Roche HM, Korpela R, Nilaweera KN (2013) Impact of leucine on energy balance. *Journal of physiology and biochemistry* 69: 155–163.
71. Hamad EM, Taha SH, Abou Dawood AG, Sitohy MZ, Abdel-Hamid M (2011) Protective effect of whey proteins against nonalcoholic fatty liver in rats. *Lipids in health and disease* 10: 57.
72. Pilvi TK, Seppanen-Laakso T, Simolin H, Finckenberg P, Huotari A, et al. (2008) Metabolomic changes in fatty liver can be modified by dietary protein and calcium during energy restriction. *World Journal of Gastroenterology* 14: 4462–4472.
73. Mortensen LS, Hartvigsen ML, Brader LJ, Astrup A, Schrezenmeir J, et al. (2009) Differential effects of protein quality on postprandial lipemia in response to a fat-rich meal in type 2 diabetes: comparison of whey, casein, gluten, and cod protein. *The American journal of clinical nutrition* 90: 41–48.
74. Pal S, Ellis V, Ho S (2010) Acute effects of whey protein isolate on cardiovascular risk factors in overweight, post-menopausal women. *Atherosclerosis* 212: 339–344.
75. Collins S, Daniel KW, Rohlf EM (1999) Depressed expression of adipocyte beta-adrenergic receptors is a common feature of congenital and diet-induced obesity in rodents. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* 23: 669–677.
76. Gustafson B, Gogg S, Hedjazifaz S, Jenndahl L, Hammarstedt A, et al. (2009) Inflammation and impaired adipogenesis in hypertrophic obesity in man. *American journal of physiology Endocrinology and metabolism* 297: E999–E1003.
77. McGillicuddy FC, Harford KA, Reynolds CM, Oliver E, Claessens M, et al. (2011) Lack of interleukin-1 receptor I (IL-1RI) protects mice from high-fat diet-induced adipose tissue inflammation coincident with improved glucose homeostasis. *Diabetes* 60: 1688–1698.
78. Belobrajdic DP, McIntosh GH, Owens JA (2004) A high-whey-protein diet reduces body weight gain and alters insulin sensitivity relative to red meat in wistar rats. *J Nutr* 134: 1454–1458.
79. Santacruz A, Marcos A, Warnberg J, Marti A, Martin-Matillas M, et al. (2009) Interplay between weight loss and gut microbiota composition in overweight adolescents. *Obesity* 17: 1906–1915.
80. Parnell JA, Reimer RA (2012) Prebiotic fibres dose-dependently increase satiety hormones and alter Bacteroidetes and Firmicutes in lean and obese JCR:LA-cp rats. *The British journal of nutrition* 107: 601–613.
81. Clarke SF, Murphy EF, O’Sullivan O, Ross RP, O’Toole PW, et al. (2013) Targeting the microbiota to address diet-induced obesity: a time dependent challenge. *PLoS one* 8: e65790.
82. Armougom F, Henry M, Vialettes B, Raccach D, Raoult D (2009) Monitoring bacterial community of human gut microbiota reveals an increase in Lactobacillus in obese patients and Methanogens in anorexic patients. *PLoS one* 4: e7125.
83. Erickson KL, Hubbard NE (2000) Probiotic immunomodulation in health and disease. *The Journal of nutrition* 130: 403S–409S.
84. Ouwehand AC, Kirjavainen PV, Shortt C, Salminen S (1999) Probiotics: mechanisms and established effects. *International Dairy Journal* 9: 43–52.
85. Hildebrandt MA, Hoffmann C, Sherrill-Mix SA, Keilbaugh SA, Hamady M, et al. (2009) High-fat diet determines the composition of the murine gut microbiome independently of obesity. *Gastroenterology* 137: 1716–1724 e1711–1712.
86. Sonoyama K, Fujiwara R, Takemura N, Ogasawara T, Watanabe J, et al. (2009) Response of gut microbiota to fasting and hibernation in Syrian hamsters. *Applied and environmental microbiology* 75: 6451–6456.
87. Sousa DZ, Pereira MA, Smidt H, Stams AJ, Alves MM (2007) Molecular assessment of complex microbial communities degrading long chain fatty acids in methanogenic bioreactors. *FEMS microbiology ecology* 60: 252–265.
88. Schwiertz A, Taras D, Schafer K, Beijer S, Bos NA, et al. (2010) Microbiota and SCFA in lean and overweight healthy subjects. *Obesity* 18: 190–195.