

Title	Compactification for asymptotically autonomous dynamical systems: Theory, applications and invariant manifolds
Authors	Wieczorek, Sebastian;Xie, Chun;Jones, Chris K.R.T.
Publication date	2021-05-13
Original Citation	Wieczorek, S., Xie, C. and Jones, C. K.R.T. (2021) 'Compactification for asymptotically autonomous dynamical systems: Theory, applications and invariant manifolds', <i>Nonlinearity</i> , 34(5), pp. 2970-3000. doi: 10.1088/1361-6544/abe456
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1088/1361-6544/abe456
Rights	© 2021, IOP Publishing Ltd & London Mathematical Society. This is an author-created, un-copyedited version of an article accepted for publication in <i>Nonlinearity</i> . The publisher is not responsible for any errors or omissions in this version of the manuscript or any version derived from it. The Version of Record is available online at https://doi.org/10.1088/1361-6544/abe456
Download date	2024-04-16 04:35:09
Item downloaded from	https://hdl.handle.net/10468/11487



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Compactification for Asymptotically Autonomous Dynamical Systems: Theory, Applications and Invariant Manifolds

Sebastian Wieczorek*, Chun Xie,* and Chris K.R.T. Jones†

June 24, 2021

Abstract

We develop a general compactification framework to facilitate analysis of nonautonomous ODEs where nonautonomous terms decay asymptotically. The strategy is to compactify the problem: the phase space is augmented with a bounded but open dimension and then extended at one or both ends by gluing in flow-invariant subspaces that carry autonomous dynamics of the limit systems from infinity. We derive the weakest decay conditions possible for the compactified system to be continuously differentiable on the extended phase space. This enables us to use equilibria and other compact invariant sets of the limit systems from infinity to analyse the original nonautonomous problem in the spirit of dynamical systems theory. Specifically, we prove that solutions of interest are contained in unique invariant manifolds of saddles for the limit systems when embedded in the extended phase space. The uniqueness holds in the general case, that is even if the compactification gives rise to a centre direction and the manifolds become centre or centre-stable manifolds. A wide range of problems including pullback attractors, rate-induced critical transitions (R-tipping) and nonlinear wave solutions fit naturally into our framework, and their analysis can be greatly simplified by the compactification.

Keywords: asymptotically autonomous differential equations, compactification, nonautonomous dynamical systems, rate-induced tipping, nonlinear waves, centre manifold, centre-stable manifold.

Contents

1	Introduction	3
1.1	The Basic Setting	3
1.2	Motivating Examples	5
1.2.1	Pullback Attractors	5
1.2.2	Rate-Induced Tipping	5
1.2.3	Radial Steady States	5
1.2.4	Stability of Nonlinear Waves	6
2	Compactification	7
2.1	Two-Sided Compactification for Bi-Asymptotically Constant $\Gamma(t)$	8
2.1.1	Compactified System: Transformation Conditions	8
2.1.2	Compactified System: Optimal Existence Criteria	11
2.2	One-Sided Compactification for Asymptotically Constant $\Gamma(t)$	13
2.2.1	Right-Sided Compactification	14
2.2.2	Left-Sided Compactification	14

*School of Mathematical Sciences, University College Cork, Western Road, Cork, T12 XF62, Ireland

†Mathematics Department, University of North Carolina, Chapel Hill, NC 27599, USA

3	Compactified System Dynamics	15
3.1	Attractors and Repellers	16
3.2	Saddles and Invariant Manifolds	16
3.2.1	Dynamical Structure from the Future Limit System	17
3.2.2	Dynamical Structure from the Past Limit System	20
3.3	Invariant Manifolds and Applications	21
3.3.1	Pullback Attractors	22
3.3.2	Rate-Induced Tipping	22
3.3.3	Radial Steady States	22
3.3.4	Stability of Nonlinear Waves	22
3.4	Generalisations to More Complicated Compact Invariant Sets	23
4	Examples of Compactification	23
4.1	Γ -Compactification	23
4.2	Exponential Compactification	25
4.3	Algebraic Compactification	26
4.4	Other Compactification Types and Decay Limitation	28
5	Conclusion	28

1 Introduction

Let U and V be open subsets $U \subseteq \mathbb{R}^n$ and $V \subseteq \mathbb{R}^d$. Consider a *nonlinear nonautonomous* differential equation

$$\frac{dx}{dt} := \dot{x} = f(x, \Gamma(t)), \quad (1)$$

with the dependent state variable $x \in U$, independent variable $t \in \mathbb{R}$, C^1 -smooth nonautonomous term $\Gamma : \mathbb{R} \rightarrow V$, and C^1 -smooth vector field $f : U \times V \rightarrow \mathbb{R}^n$.

For an arbitrary $\Gamma(t)$, the theory of nonautonomous dynamical systems is presented in [19] which summarises work on the problem, discusses useful concepts such as pullback attractors, and gives some very general results on attraction and stability. The main obstacle to the analysis of nonautonomous system (1) is the absence of compact invariant sets such as equilibria, limit cycles or invariant tori. This obstacle becomes evident when the system is augmented with $\nu = t$ as an additional dependent variable to obtain the usual *autonomous extended system*¹

$$\begin{aligned} \dot{x} &= f(x, \Gamma(\nu)), \\ \dot{\nu} &= 1, \end{aligned} \quad (2)$$

that is defined on the phase space $U \times \mathbb{R}$, which is an open subset of \mathbb{R}^{n+1} .

Our strategy is to consider asymptotically decaying $\Gamma(t)$ and compactify the system (2). Compactification is not a new strategy and has been exploited in many applications cited in Sec. 1.2 below. However, in the context of decaying $\Gamma(t)$, it has been implemented only on a case-by-case basis for specific applications and specific types of decay [17, 1, 7, 36, 35, 2]. Our viewpoint here is to develop a general framework and derive in Theorem 2.2 the weakest decay conditions possible for which the compactified system is sufficiently smooth to enable the construction of invariant manifolds (stable, unstable, centre, etc.) in the compactified phase space.

The invariant manifolds are key in the applications we envision in Sec. 1.2. Addressing the issue of finding optimal conditions for compactification raises the problem that the manifolds of interest may involve a centre direction and need not be pure stable or unstable manifolds. This, in turn, raises the possibility that the manifolds of interest may not be unique. We show in Theorems 3.2 and 3.4 that uniqueness holds in spite of these centre directions.

1.1 The Basic Setting

The main point of this work is that one can make further progress on problem (1) if $\Gamma(t)$ limits to a (vector) constant as t tends to positive or negative infinity. Specifically,

Definition 1.1. *We say $\Gamma(t)$ is bi-asymptotically constant with future limit Γ^+ and past limit Γ^- if*

$$\lim_{t \rightarrow \pm\infty} \Gamma(t) = \Gamma^\pm \in \mathbb{R}^d.$$

We say $\Gamma(t)$ is asymptotically constant if it has a future limit but not necessarily a past limit, or if it has a past limit but not necessarily a future limit.

The main simplification is that nonautonomous system (1) becomes *asymptotically autonomous* in the terminology of [28]:

$$f(x, \Gamma(t)) \rightarrow f(x, \Gamma^+) \text{ as } t \rightarrow +\infty, \text{ or } f(x, \Gamma(t)) \rightarrow f(x, \Gamma^-) \text{ as } t \rightarrow -\infty,$$

¹We note that in the special case $\Gamma(t)$ is generated by a known set of autonomous ODEs, say $\dot{\Gamma} = w(\Gamma)$, the original system can be augmented with Γ to obtain the autonomous system $\dot{x} = f(x, \Gamma)$, $\dot{\Gamma} = w(\Gamma)$, defined on $U \times V \subseteq \mathbb{R}^{n+d}$. However, $w(\Gamma)$ is not always known and this approach may be impractical for analysis if d is large.

and we can define the autonomous *future limit system*

$$\dot{x} = f(x, \Gamma^+), \quad (3)$$

or the autonomous *past limit system*

$$\dot{x} = f(x, \Gamma^-). \quad (4)$$

The autonomous dynamics of the future (3) or past (4) limit systems will typically include equilibria and other compact invariant sets. Note, however, that the flow of (1) does not contain the autonomous dynamics of (3) or (4) because they only appear as t tends to positive and negative infinity.

The key idea is to overcome the main obstacle by using the autonomous dynamics and compact invariant sets of the limit systems to analyse the nonautonomous system (1). Specifically, we augment the nonautonomous system (1) with an additional dimension that is bounded but open, and then glue to its open ends the future limit system (3) from $+\infty$, the past limit system (4) from $-\infty$, or both. In this way, we combine systems (1), (3) and (4) into one *compactified system* that is autonomous, but not necessarily C^1 -smooth. The compactification approach presented in this paper is akin to the Poincaré-type phase-space compactification (of the x -dimensions) that enables analysis of dynamical behaviour at infinity [10, 21, 13, 31, 29, 16], collisions in many-body problems [30], stability of nonlinear waves [17, 1], as well as higher-codimension bifurcations [12] and canard solutions in slow-fast systems [14, 24, 43, 20, 25, 23] via blow up of singularities of vector fields. The difference between our work and these studies is twofold. Firstly, we compactify the augmented ν -dimension, but not necessarily the x -dimensions. Secondly, we consider a general case with arbitrary decay of $\Gamma(t)$ and

- Derive in Theorem 2.2 the optimal conditions for the compactification: the *weakest decay of $\dot{\Gamma}(t)$ possible* that allows us to construct a C^1 -smooth compactified system.
- Prove in Theorems 3.2 and 3.4 that the geometric object of interest, namely the set of orbits tending to a saddle at infinity, is a *unique invariant manifold* in the compactified phase space, even if a centre direction arises from non-exponential decay of $\Gamma(t)$.

Our framework enables analysis of asymptotically autonomous dynamical systems [28] in terms of unique invariant manifolds of saddles from infinity in a general setting, where the decay of $\Gamma(t)$ ranges from super-exponential to sub-logarithmic. Thus, it provides an alternative tool and complements the existing approaches based on pullback attractors [26, 37] and asymptotic equivalence of the nonautonomous system and the autonomous limit systems [45, 40].

The paper is organised as follows. In Sec. 1.2 we describe different problems from applications that provide direct motivation for this study and fit naturally into the framework of (1) with asymptotically constant $\Gamma(t)$. In Sec. 2 we develop the general compactification technique and introduce the underpinning coordinate transformation. To be more specific, in Sec. 2.1.1 we construct an autonomous compactified system on a suitably augmented and extended phase space. We then give two simple transformation conditions for the compactified system to be continuously differentiable on the extended phase space. In Sec. 2.1.2 we introduce the concept of “reference envelopes” to derive the weakest decay conditions possible for the existence of the desired compactification transformation. In Sec. 2.2 we distinguish between two-sided compactification for bi-asymptotically constant $\Gamma(t)$ and one-sided compactification for asymptotically constant $\Gamma(t)$. In Sec. 3 we relate the nonautonomous dynamics of (1) to the autonomous dynamics of the compactified system. Our focus is on the extrapolation of the dynamical structure from the limit systems. This leads us to the construction of stable, unstable and centre invariant manifolds of compact invariant sets for the limit systems when embedded in the extended

phase space of the compactified system. The geometric shape of these manifolds ‘encodes’ the nonautonomous profile of $\Gamma(t)$, which can greatly simplify analysis of (1). In Sec. 4 we construct examples of exponential and algebraic compactifications that are useful in practice.

1.2 Motivating Examples

Our work is directly motivated by a range of problems from applications that can be fitted into the framework of Eq. (1) with asymptotically constant $\Gamma(t)$, and become simpler to analyse after compactification.

1.2.1 Pullback Attractors

Consider a nonautonomous dynamical system (1) with time t and asymptotically constant $\Gamma(t)$ with a past limit

$$\Gamma(t) \rightarrow \Gamma^- \text{ as } t \rightarrow -\infty,$$

and refer to [19, Ch.3] for the notion of a local pullback attractor. Each asymptotically stable compact invariant set A^- for the past limit system (4) can be associated with a local pullback attractor of the nonautonomous system (1) as shown in [37, Lem.6.2], [6, Th.2.2] and [2, Th.II.2]. When embedded in the extended phase space of a compactified system, A^- gains one repelling direction and becomes a saddle. Thus, the process of compactification transforms the local pullback attractor associated with A^- into the unstable or centre invariant manifold of A^- , which can greatly facilitate analysis of the nonautonomous problem. Relating pullback attractors to unstable invariant manifolds gives an alternative approach to the existing nonautonomous stability theory [19].

1.2.2 Rate-Induced Tipping

A nonlinear nonautonomous dynamical system (1) with time t and bi-asymptotically constant

$$\Gamma(t) = \Lambda(rt) \rightarrow \lambda^\pm \in \mathbb{R}^d \text{ as } t \rightarrow \pm\infty,$$

is often used to describe nonlinear dynamics of open systems subject to finite-time external disturbances, growing or decaying external trends, or simply time-varying external inputs $\Lambda(rt)$. A rate-induced critical transition, known as R-tipping [7], occurs when the ‘rate’ $r > 0$ of the external input exceeds some critical value and the system transitions to a different state [46, 7, 32, 36, 38, 18, 3, 33, 47, 48]. This genuine nonautonomous bifurcation is of great interest to natural scientists but cannot, in general, be explained using classical bifurcation analysis of the autonomous frozen system

$$\dot{x} = f(x, \lambda),$$

with a fixed-in-time input parameter λ . The process of compactification transforms the nonautonomous R-tipping problem into a connecting heteroclinic orbit problem, which facilitates both numerical analysis [48] and the derivation of rigorous criteria for R-tipping [47]. Here, the closure of an image of $\Gamma(t)$ has an important meaning: it is a parameter path in the multi-dimensional parameter space $V \subseteq \mathbb{R}^d$ of the frozen system that is traced out by the time-varying external input.

1.2.3 Radial Steady States

The steady states of nonlinear wave, diffusion or Schrödinger equations with a potential $\mathcal{V}(x)$ on \mathbb{R}^n satisfy a semilinear elliptic equation:

$$\Delta_x u + f(u) + \mathcal{V}(x)u = 0.$$

Under the condition that $\mathcal{V}(x) = \mathcal{V}(\|x\|)$, spherically symmetric solutions satisfy the semilinear elliptic boundary value problem (BVP), where $r > 0$ is the radial direction [17, 41]:

$$u_{rr} + \frac{n-1}{r} u_r + f(u) + \mathcal{V}(r)u = 0, \quad \begin{aligned} \lim_{r \rightarrow 0} u_r(r) &= 0, \\ \lim_{r \rightarrow +\infty} u(r) &= 0, \end{aligned} \quad (5)$$

where $u_r = du/dr$. If the potential $\mathcal{V}(r)$ is asymptotically constant with a future limit², the BVP (5) fits naturally into the framework of (1). More precisely, by using r as the independent variable and introducing $v(r) = u_r(r)$ as an additional dependent variable so that

$$x(r) = \begin{pmatrix} u(r) \\ v(r) \end{pmatrix} \quad \text{and} \quad \Gamma(r) = \begin{pmatrix} \Gamma_1(r) \\ \Gamma_2(r) \end{pmatrix} = \begin{pmatrix} (-n+1)/r \\ -\mathcal{V}(r) \end{pmatrix} \rightarrow \begin{pmatrix} 0 \\ -\mathcal{V}^+ \end{pmatrix} \quad \text{as } r \rightarrow +\infty,$$

we can rewrite (5) as

$$x' = f(x, \Gamma(r)) = \begin{pmatrix} v \\ \Gamma_1(r)v + \Gamma_2(r)u - f(u) \end{pmatrix}, \quad \begin{aligned} \lim_{r \rightarrow 0} v(r) &= 0, \\ \lim_{r \rightarrow +\infty} u(r) &= 0, \end{aligned}$$

where $' = d/dr$, and $f(0) = 0$.

The future limit system corresponds to the problem with constant potential and is easily analysed as a Hamiltonian system in the plane [17]. Under common assumptions, the analysis reveals a saddle equilibrium η^+ at the origin. The full BVP is then solved by finding appropriate trajectories on the stable or centre-stable invariant manifold of η^+ in the compactified system.

There are other examples of compactification used for the study of particular solutions of nonlinear equations, such as the lens transform [44, Ch.2], which is a special case of pseudoconformal compactification [8] for the nonlinear Schrödinger equation.

1.2.4 Stability of Nonlinear Waves

Traveling wave solutions of a reaction-diffusion equation in one space dimension

$$U_t = U_{xx} + F(U), \quad (6)$$

where $U \in \mathbb{R}^m$, satisfy a second-order system of ordinary differential equations [1]:

$$U'' + cU' + F(U) = 0, \quad (7)$$

where $' = d/dz$, $z = x - ct$ and c is the speed of the wave. Of interest is commonly a wave $U(z)$ that decays to end-states, $U(z) \rightarrow U_{\pm}$ as $z \rightarrow \pm\infty$. The key in an analysis of the stability of such a wave is the eigenvalue problem for the linearisation of (6) at the wave [1]:

$$L(U)P = P'' + cP' + dF(U(z))P = \lambda P, \quad (8)$$

where $dF(U)$ denotes the derivative of $F(U)$ with respect to U . We can rewrite the second equality in (8) as a nonautonomous system on \mathbb{R}^{2m} :

$$P' = Q \quad (9)$$

$$Q' = -cQ + \lambda P - dF(U(z))P, \quad (10)$$

with bi-asymptotically constant $dF(U(z))$. Setting

$$x(z) = \begin{pmatrix} P(z) \\ Q(z) \end{pmatrix} \quad \text{and} \quad \Gamma(z) = dF(U(z)) \rightarrow dF(U_{\pm}) \quad \text{as } z \rightarrow \pm\infty,$$

²This includes the special case $\mathcal{V}(r) = 0$.




Figure 1: Examples of the coordinate transformation $s = g(t)$ from (a) Assumption 2.1, (b) Assumption 2.2 and (c) Assumption 2.3.

we can fit (9)–(10) into our general framework

$$x' = f(x, \Gamma(z)) = \begin{pmatrix} Q \\ -cQ + \lambda P - \Gamma(z)P \end{pmatrix}.$$

Note that the limit systems here are exactly the eigenvalue problems for the linearisation of (6) at the constant (time and space independent) solutions U_{\pm} .

The Evans Function is the primary tool used to capture these eigenvalues. The construction of it in [1], where the first general formulation was given, involves a two-sided compactification of system (9)–(10).

2 Compactification

The aim of this section is to reformulate the nonautonomous system (1) into an autonomous system so that:

- The new system contains the autonomous flow and compact invariant sets of the future limit system (3) and/or the past limit system (4).
- The dimension of the vector field increases just by one, independently of the dimension and monotonicity of the nonautonomous term $\Gamma(t)$.

We show that this can be achieved under quite general assumptions on $\Gamma(t)$ by augmenting system (1) with $s = g(t)$ depicted in Fig. 1 as an additional dependent variable. In other words, instead of having a problem with the additional dimension being unbounded as in the usual extended system (2), we augment system (1) so that the additional dimension becomes a compact interval. Specifically, we:

- Restrict to asymptotically and bi-asymptotically constant $\Gamma(t)$.
- Compactify the real ν -line into the compact s -interval $[-1, 1]$ if $\Gamma(t)$ is bi-asymptotically constant.
- Compactify the half ν -line $[\nu_-, +\infty)$ into the compact s -interval $[s_-, 1]$ if $\Gamma(t)$ is asymptotically constant with a future limit, or compactify the half ν -line $(-\infty, \nu_+]$ into the compact s -interval $[-1, s_+]$ if $\Gamma(t)$ is asymptotically constant with a past limit.

2.1 Two-Sided Compactification for Bi-Asymptotically Constant $\Gamma(t)$

We reformulate the nonautonomous system (1) with a bi-asymptotically constant $\Gamma(t)$ into a compactified system that is autonomous and contains the flow and compact invariant sets of the future (3) and past (4) limit systems. This is achieved via two-sided compactification that uses the coordinate transformation $s = g(t)$ depicted in Fig. 1(a). More precisely, we assume

Assumption 2.1. *A coordinate transformation $s = g(t)$ maps the real t -line onto the finite s -interval $(-1, 1)$, is at least C^2 -smooth, bi-asymptotically constant with future limit 1 and past limit -1 , and strictly increasing with vanishing first derivative as t tends to $\pm\infty$:*

$$g: \mathbb{R} \rightarrow (-1, 1), \quad g \in C^{k \geq 2}, \quad \lim_{t \rightarrow \pm\infty} g(t) = \pm 1, \quad \dot{g}(t) > 0 \text{ for } t \in \mathbb{R} \text{ and } \lim_{t \rightarrow \pm\infty} \dot{g}(t) = 0. \quad (11)$$

Remark 2.1. *In practice, we introduce compactification parameter(s) to control the rate/order of asymptotic decay of $g(t)$, and work with parametrised compactifications; see Sec. 4 ahead. The dependence on the compactification parameter(s) does not affect, and is thus left out of, general statements of this section.*

2.1.1 Compactified System: Transformation Conditions

Compactification is a three-step process. The first step is to make the additional dimension bounded by augmenting the asymptotically autonomous system (1) with $s = g(t)$ as an additional dependent variable

$$\dot{x} = f(x, \Gamma(g^{-1}(s))), \quad (12)$$

$$\dot{s} = \dot{g}(g^{-1}(s)), \quad s(t_0) = g(t_0). \quad (13)$$

Since the practical implementation of the compactification requires the inverse coordinate transformation $t = h(s) := g^{-1}(s)$, we reformulate (12)–(13) in terms of $h(s)$ alone:

$$\dot{x} = f(x, \Gamma(h(s))), \quad (14)$$

$$\dot{s} = \gamma(s), \quad (15)$$

where

$$\gamma(s) = \dot{g}(h(s)) = \frac{1}{h'(s)},$$

is the augmented component of the vector field, $\dot{\cdot} = d/dt$, and $' = d/ds$. System (14)–(15) is defined on $U \times (-1, 1)$, which is an open subset of \mathbb{R}^{n+1} . The second step is to make the s -interval closed by including $s = \pm 1$ ($t = \pm\infty$) and extending the augmented by (15) vector field to subspaces $\{s = \pm 1\}$:

$$f(x, \Gamma(h(s))) = \begin{cases} f(x, \Gamma(h(s))) & \text{for } s \in (-1, 1), \\ f(x, \Gamma^-) & \text{for } s = -1, \\ f(x, \Gamma^+) & \text{for } s = 1, \end{cases} \quad (16)$$

$$\gamma(s) = \begin{cases} 1/h'(s) & \text{for } s \in (-1, 1), \\ 0 & \text{for } s = \pm 1. \end{cases} \quad (17)$$

This gives an autonomous *compactified system* (14)–(17) that is defined on the extended phase space $U \times [-1, 1]$. Most importantly, subspaces $\{s = 1\}$ and $\{s = -1\}$ are flow-invariant and carry the autonomous dynamics and compact invariant sets of the future (3) and past (4) limit systems, respectively. However, it is not generally guaranteed that the extended vector field is differentiable at the added invariant subspaces $\{s = \pm 1\}$. The third step is to give testable criteria for the extended vector field to be continuously differentiable. Examining first-order partial derivatives of the extended vector field (16)–(17) leads to the following result:




Figure 2: (a)–(b) Example of transformation-induced loss of differentiability of $\Gamma(h(s))$ at $s = \pm 1$ when condition (18) is violated. (c)–(d) Example of $\gamma(s)$ that is non-differentiable at $s = \pm 1$ when condition (19) is violated.

Proposition 2.1. (Transformation Conditions.) *Consider a nonautonomous system (1) on U with C^1 -smooth f and Γ , and bi-asymptotically constant $\Gamma(t)$. For a chosen coordinate transformation $g(t)$ from Assumption 2.1, with the inverse $h(s) = g^{-1}(s)$, the ensuing compactified system (14)–(17) is C^1 -smooth on $U \times [-1, 1]$ if and only if*

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{g}(t)} = \lim_{s \rightarrow \pm 1^\mp} \frac{d}{ds} \Gamma(h(s)) \text{ exist,} \quad (18)$$

$$\lim_{t \rightarrow \pm\infty} \frac{\ddot{g}(t)}{\dot{g}(t)} = - \lim_{s \rightarrow \pm 1^\mp} \frac{h''(s)}{(h'(s))^2} \text{ exist.} \quad (19)$$

The one-dimensional examples from Fig. 2 give insight into transformation conditions (18) and (19), which ensure differentiability of $\Gamma(h(s))$ and $\gamma(s)$, respectively, at $s = \pm 1$. Recall that $\dot{g}(t)$ limits to zero as $t \rightarrow \pm\infty$ and note the following. The slope of $\Gamma(h(s))$ in Fig. 2(b) is given by $d\Gamma/ds = d\Gamma/dg = \dot{\Gamma}/\dot{g}$. Thus, the derivative of $\Gamma(h(s))$ becomes undefined at $s = \pm 1$ if $\dot{\Gamma}(t)$ does not limit to zero or if $\dot{\Gamma}(t)$ approaches zero slower than $\dot{g}(t)$ as $t \rightarrow \pm\infty$; see Sec. 2.1.2 for optimal conditions on the asymptotic decay of $\dot{\Gamma}(t)$. In other words, coordinate transformations that decay faster than the nonautonomous term violate condition (18) and cause transformation-induced loss of differentiability of $\Gamma(h(s))$ at $s = \pm 1$. Similarly, the slope of $\gamma(s)$ in Fig. 2(d) is given by $d\gamma/ds = d\dot{g}/ds = \ddot{g}/\dot{g}$. Thus, the derivative of $\gamma(s)$ becomes undefined at $s = \pm 1$ if $\ddot{g}(t)$ does not limit to zero or if $\ddot{g}(t)$ approaches zero slower than $\dot{g}(t)$ as $t \rightarrow \pm\infty$. In other words, coordinate transformations whose speed tends to zero faster than acceleration violate condition (19) and make $\gamma(s)$ non-differentiable at $s = \pm 1$. The following remark gives additional insight into condition (19).

Remark 2.2. *Transformation condition (19) excludes coordinate transformations $g(t)$ that decay faster than exponentially. To see that, consider*

$$g(t) \sim \begin{cases} 1 - \exp(-t^k) & \text{as } t \rightarrow +\infty, \\ -1 + \exp(-(-t)^k) & \text{as } t \rightarrow -\infty, \end{cases}$$

for $k > 0$, and note that

$$\lim_{t \rightarrow \pm\infty} \frac{\ddot{g}(t)}{\dot{g}(t)} = \mp k \lim_{t \rightarrow \pm\infty} (\pm t)^{k-1} = \begin{cases} \mp\infty & \text{if } k > 1, \\ \mp 1 & \text{if } k = 1, \\ 0 & \text{if } 0 < k < 1. \end{cases}$$

This can be understood intuitively via behaviour near invariant subspaces: solutions of ODEs cannot approach invariant subspaces faster than exponentially unless they blow up and cease to exist.

Proof of Proposition 2.1

Note that $h'(s) = 1/\dot{g}(h(s)) > 0$ and $h(s)$ is C^2 -smooth for $s \in (-1, 1)$ by the (general) inverse function theorem [42], and choose the following form of the augmented component of the vector field $\gamma(s) = \dot{g}(h(s))$ for $s \in (-1, 1)$. Then, C^1 -smoothness of the extended vector field (16)–(17) on $U \times (-1, 1)$ follows from C^1 -smoothness of f on U , C^2 -smoothness of Γ on \mathbb{R} , C^2 -smoothness of g on \mathbb{R} , and C^1 -smoothness of h on $(-1, 1)$. What needs to be examined is C^1 -smoothness of the extended vector field at the added right and left invariant subspaces $\{s = \pm 1\}$.

It follows from Definition 1.1 and Assumption 2.1 that $\Gamma(h(s))$ and $\gamma(s)$ are continuous at $s = \pm 1$:

$$\begin{aligned} \lim_{s \rightarrow \pm 1^\mp} h(s) &= \lim_{t \rightarrow \pm\infty} t, \\ \lim_{s \rightarrow \pm 1^\mp} \Gamma(h(s)) &= \lim_{t \rightarrow \pm\infty} \Gamma(t) = \Gamma^\pm = \Gamma(h(\pm 1)), \\ \lim_{s \rightarrow \pm 1^\mp} \gamma(s) &= \lim_{s \rightarrow \pm 1^\mp} \dot{g}(h(s)) = \lim_{t \rightarrow \pm\infty} \dot{g}(t) = 0 = \gamma(\pm 1). \end{aligned} \tag{20}$$

Thus, the extended vector field is continuous on $U \times [-1, 1]$. The first derivative of the extended vector field is continuous at $\{s = \pm 1\}$ if the left- and right-sided limits $s \rightarrow \pm 1^\mp$ exist for all first-order partial derivatives. To check this, consider the Jacobian of the extended vector field

$$J(x, s) = \begin{pmatrix} \left(\frac{\partial f}{\partial x}\right)_{n \times n} & \left(\frac{\partial f}{\partial s}\right)_{n \times 1} \\ (0)_{1 \times n} & \left(\frac{d\gamma}{ds}\right)_{1 \times 1} \end{pmatrix}, \tag{21}$$

where the subscripts indicate the size of the matrix components of $J(x, s)$, and

$$\left(\frac{\partial f}{\partial s}\right)_{n \times 1} = \left(\frac{\partial f}{\partial \Gamma}\right)_{n \times d} \left(\frac{d}{ds}\Gamma(h(s))\right)_{d \times 1}. \tag{22}$$

Using the chain rule and the inverse function theorem gives

$$\frac{d}{ds}\Gamma(h(s)) = \dot{\Gamma}(h(s)) h'(s) = \frac{\dot{\Gamma}(h(s))}{\dot{g}(h(s))}. \tag{23}$$

Differentiate $s = g(h(s))$ twice with respect to s to obtain

$$\ddot{g}(h(s)) (h'(s))^2 = -\dot{g}(h(s)) h''(s), \tag{24}$$

which together with the chain rule and the inverse function theorem gives

$$\frac{d}{ds}\gamma(s) = \frac{d}{ds}\dot{g}(h(s)) = \ddot{g}(h(s)) h'(s) = \frac{\ddot{g}(h(s))}{\dot{g}(h(s))} = -\frac{h''(s)}{(h'(s))^2}. \tag{25}$$




Figure 3: Consider C^1 -smooth bumps of equal amplitudes whose (shaded) areas form a convergent series as $t \rightarrow +\infty$. Then, $\|\dot{\Gamma}(t)\|$ is set to consist of such bumps with bump amplitudes remaining constant or decaying arbitrarily slowly to zero as $t \rightarrow +\infty$. Since the total area below $\|\dot{\Gamma}(t)\|$ is finite by construction, we have $\Gamma(t) \rightarrow \Gamma^+ \in \mathbb{R}$ as $t \rightarrow +\infty$ in spite of the envelope $\dot{A}(t)$ remaining constant or decaying arbitrarily slowly to zero.

It follows from the continuity of $\Gamma(h(s))$ at $s = \pm 1$ that $(\partial f / \partial x)_{n \times n}$ and $(\partial f / \partial \Gamma)_{n \times d}$ are continuous at $s = \pm 1$. It then follows from Eqs. (20), (22) and (23) that $(\partial f / \partial s)_{n \times 1}$ is continuous at $s = \pm 1$ if and only if the first transformation condition (18) is satisfied. It follows from Eqs. (20) and (25) that the $J_{n+1, n+1}$ component of the Jacobian is continuous at $s = \pm 1$ if and only if the second transformation condition (19) is satisfied. Thus, the first derivative of the extended vector field is continuous on $U \times [-1, 1]$ if and only if transformation conditions (18) and (19) are satisfied. \square

2.1.2 Compactified System: Optimal Existence Criteria

Here we derive optimal criteria on $\dot{\Gamma}(t)$ that guarantee a suitable coordinate transformation $g(t)$ that satisfies conditions (18) and (19) from Proposition 2.1 exists. The derivation of the existence criteria is guided by the observation that ‘normal’ examples of bi-asymptotically constant $\Gamma(t)$ have two additional and desirable properties. Firstly, $\dot{\Gamma}(t)$ limits to zero as t tends to positive and negative infinity. Secondly, the asymptotic approach of $\dot{\Gamma}(t)$ towards zero is not slower than the asymptotic approach of $\Gamma(t)$ towards Γ^\pm .

However, $\Gamma(t) \rightarrow \Gamma^\pm \in \mathbb{R}^d$ does not imply $\dot{\Gamma}(t) \rightarrow 0$ in general, meaning that there exist ‘pathological’ examples of $\Gamma(t) \rightarrow \Gamma^\pm$ whose derivatives do not have a future or past limit, or approach zero arbitrarily slowly as t tends to positive or negative infinity. One example are damped oscillations with increasing frequency, where the frequency increase ‘beats’ the amplitude decay. For example, $\Gamma(t) \sim \sin(t^2)/t$ as $t \rightarrow +\infty$ has a future limit but its first derivative $\dot{\Gamma}(t) \sim 2 \cos(t^2) - \sin(t^2)/t^2$ does not. Another example is depicted in Fig. 3 where $\Gamma(t) \rightarrow \Gamma^+$ but $\dot{\Gamma}(t)$ may not have a future limit or may approach zero arbitrarily slowly. Conversely, $\dot{\Gamma}(t) \rightarrow 0$ as $t \rightarrow \pm\infty$ does not imply $\Gamma(t) \rightarrow \Gamma^\pm \in \mathbb{R}^d$ either. For example, $\Gamma(t) \sim \ln(t)$ as $t \rightarrow +\infty$ does not have a future limit even though its first derivative $\dot{\Gamma}(t) \sim 1/t$ limits to zero.

To exclude ‘pathological’ examples of bi-asymptotically constant $\Gamma(t)$ it is convenient to work with a monotone envelope of $\|\dot{\Gamma}(t)\|$, denoted by $\dot{A}(t)$ as depicted in Fig. 3, and ask about the slowest-decaying $\dot{A}(t) \rightarrow 0$ so that $A(t)$ has a future and past limits. Formulating this question

in terms of the integral

$$\lim_{t \rightarrow +\infty} A(t) = \int_{t_0}^{\infty} \dot{A}(\mu) d\mu = A^+ \in \mathbb{R}, \quad (26)$$

shows that such a slowest-decaying envelope does not exist: given any $\dot{A}(t) \rightarrow 0$ that satisfies (26) for some $t_0 \in \mathbb{R}$, one can construct a slower-decaying one that also satisfies (26). Nonetheless, it is possible to work with a parametrised family of envelopes that satisfy (26) and can be chosen to decay sufficiently slowly for the problem at hand. For brevity, we introduce \exp^m and \ln^m to denote a composition of m exponential and logarithmic functions, respectively,

$$\exp^m(t) = \underbrace{\exp(\exp(\dots \exp(t)\dots))}_{m \text{ times}} \quad \text{and} \quad \ln^m(t) = \underbrace{\ln(\ln(\dots \ln(t)\dots))}_{m \text{ times}}.$$

Then, we consider

$$A(t, m) = -\frac{1}{\ln^m(|t|)},$$

together with its derivative

$$\dot{A}(t, m) = \left(t \ln^m(|t|) \prod_{k=1}^m \ln^k(|t|) \right)^{-1}, \quad (27)$$

parametrised by a non-negative integer $m \in \mathbb{N}_0$, where $|t| > \exp^{m-2}(e)$.

Definition 2.1. We call $\dot{A}(t, m)$ from Eq. (27) the parametrised reference envelope.

The reference envelope formula (27) looks rather technical but one can gain further insight by writing out examples of A and \dot{A} with $m = 2$ that are defined for $|t| > e$:

$$A(t, 2) = -\frac{1}{\ln(\ln(|t|))} \quad \text{and} \quad \dot{A}(t, 2) = \frac{1}{t \ln(|t|) (\ln(\ln(|t|)))^2}.$$

More generally, one can verify the following properties

$$\lim_{t \rightarrow \pm\infty} A(t, m) = 0, \quad \dot{A}(t, m) \begin{cases} > 0 & \text{for } t > \exp^{m-2}(e), \\ < 0 & \text{for } t < -\exp^{m-2}(e), \end{cases} \quad \text{and} \quad \lim_{t \rightarrow \pm\infty} \dot{A}(t, m) = 0. \quad (28)$$

We now use the reference envelope concept to restrict to ‘normal’ examples of bi-asymptotically constant $\Gamma(t)$ and ensure the existence of a continuously differentiable compactified system.

Definition 2.2. We call a bi-asymptotically constant $\Gamma(t)$ normal if there is an m such that

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{A}(t, m)} \text{ exist.} \quad (29)$$

Theorem 2.2. (Existence of a C^1 -smooth Compactified System.) *Consider a nonautonomous system (1) with C^1 -smooth f and Γ . If $\Gamma(t)$ is normal bi-asymptotically constant then there exists a coordinate transformation $s = g(t)$ from Assumption 2.1 such that the extended vector field (16)–(17) of the compactified system is C^1 -smooth on $U \times [-1, 1]$.*

In other words, the normal bi-asymptotic constant condition (29) gives sufficient existence criteria that are both *optimal* and *easily testable*. To be more precise, the criteria are optimal in the sense that they identify the weakest decay of $\dot{\Gamma}(t)$ possible to eliminate ‘pathological’ bi-asymptotically constant $\Gamma(t)$ and enable construction of a continuously differentiable compactified system, while still allowing super-exponential, exponential, algebraic, logarithmic or even sub-logarithmic decay of $\Gamma(t)$.

Proof of Theorem 2.2

Use the parametrised reference envelope from Def. 2.1 to construct a parametrised by m coordinate transformation $g_{(m)} : \mathbb{R} \rightarrow (-1, 1)$ that is at least C^2 -smooth, strictly increasing and

$$g_{(m)}(t) \sim \begin{cases} 1 + A(t, m) & \text{as } t \rightarrow +\infty, \\ -1 - A(t, m) & \text{as } t \rightarrow -\infty. \end{cases}$$

It follows from the asymptotic properties of $A(t, m)$ that

$$\lim_{t \rightarrow \pm\infty} g_{(m)}(t) = \pm 1 \quad \text{and} \quad \lim_{t \rightarrow \pm\infty} \dot{g}_{(m)}(t) = 0.$$

Thus, $g_{(m)}(t)$ satisfies Assumption 2.1 by construction. By Definition 2.2, for any normal bi-asymptotically constant $\Gamma(t)$ there is an m such that

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{g}_{(m)}(t)} = \pm \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{A}(t, m)},$$

exist, meaning that $g_{(m)}(t)$ satisfies the first transformation condition (18) by construction. Finally, one can verify by induction that

$$\lim_{t \rightarrow \pm\infty} \frac{\ddot{g}_{(m)}(t)}{\dot{g}_{(m)}(t)} = \lim_{t \rightarrow \pm\infty} \frac{\ddot{A}(t, m)}{\dot{A}(t, m)} = 0,$$

for any non-negative $m \in \mathbb{N}_0$, meaning that $g_{(m)}(t)$ satisfies the second transformation condition (19) by construction. It then follows from Proposition 2.1 that the extended vector field (16)–(17) is C^1 -smooth on $U \times [-1, 1]$. \square

2.2 One-Sided Compactification for Asymptotically Constant $\Gamma(t)$

Here we discuss briefly two one-sided subcases of the two-sided compactification from Sec 2.1. The discussion does not require new analysis, but may be helpful to readers interested in problems with asymptotically constant $\Gamma(t)$. For asymptotically constant $\Gamma(t)$ with a future limit, we reformulate the nonautonomous system (1) into a compactified system that is autonomous and contains the flow and compact invariant sets of the future limit system (3). This is achieved via one-sided compactification that uses coordinate transformation $s = g(t)$ depicted in Fig. 1(b). For asymptotically constant $\Gamma(t)$ with a past limit, we reformulate the nonautonomous system (1) into a compactified system that is autonomous and contains the flow and compact invariant sets of the past limit system (4). This is achieved via one-sided compactification that uses coordinate transformation $s = g(t)$ depicted in Fig. 1(c).

Similar to Def. 2.2, we restrict to ‘normal’ examples of asymptotically constant $\Gamma(t)$:

Definition 2.3. *We call an asymptotically constant $\Gamma(t)$ with a future limit normal if there is an m such that the $t \rightarrow +\infty$ limit in (29) exists. We call an asymptotically constant $\Gamma(t)$ with a past limit normal if there is an m such that the $t \rightarrow -\infty$ limit in (29) exists.*

2.2.1 Right-Sided Compactification

Consider nonautonomous system (1) with asymptotically constant $\Gamma(t)$ with future limit Γ^+ , and assume that

Assumption 2.2. *A coordinate transformation $s = g(t)$ maps the half t -line $[t_-, +\infty)$ onto the finite s -interval $[s_-, 1)$, is at least C^2 -smooth, asymptotically constant with future limit 1, and strictly increasing with vanishing first derivative as $t \rightarrow +\infty$:*

$$g : [t_-, +\infty) \rightarrow [s_-, 1), \quad g \in C^{k \geq 2}, \quad \lim_{t \rightarrow +\infty} g(t) = 1, \quad \dot{g}(t) > 0 \text{ for } t \geq t_- \text{ and } \lim_{t \rightarrow +\infty} \dot{g}(t) = 0. \quad (30)$$

Then consider the following autonomous *compactified system*

$$\dot{x} = f(x, \Gamma(h(s))), \quad (31)$$

$$\dot{s} = \gamma(s), \quad (32)$$

$$f(x, \Gamma(h(s))) = \begin{cases} f(x, \Gamma(h(s))) & \text{for } s \in [s_-, 1), \\ f(x, \Gamma^+) & \text{for } s = 1, \end{cases} \quad (33)$$

$$\gamma(s) = \begin{cases} 1/h'(s) & \text{for } s \in [s_-, 1), \\ 0 & \text{for } s = 1, \end{cases} \quad (34)$$

that is defined on the extended phase space $U \times [s_-, 1]$ with flow-invariant subspace $\{s = 1\}$ that carries the autonomous dynamics and compact invariant sets of the future limit system (3).

Proposition 2.1 and Theorem 2.2 apply to compactified system (31)–(34) after we leave out the limits $t \rightarrow -\infty$ ($s \rightarrow -1^+$) and replace: “bi-asymptotically constant” with “asymptotically constant with a future limit”, “compactified system (14)–(17)” with “compactified system (31)–(34)”, “phase space $U \times [-1, 1]$ ” with “phase space $U \times [s_-, 1]$ ” and “Assumption 2.1” with “Assumption 2.2”.

2.2.2 Left-Sided Compactification

Similarly, consider nonautonomous system (1) with asymptotically constant $\Gamma(t)$ with past limit Γ^- , and assume that

Assumption 2.3. *A coordinate transformation $s = g(t)$ maps the half t -line $(-\infty, t_+]$ onto the finite s -interval $(-1, s_+]$, is at least C^2 -smooth, asymptotically constant with past limit -1 , and strictly increasing with vanishing first derivative as $t \rightarrow -\infty$:*

$$g : (-\infty, t_+] \rightarrow (-1, s_+], \quad g \in C^{k \geq 2}, \quad \lim_{t \rightarrow -\infty} g(t) = -1, \quad \dot{g}(t) > 0 \text{ for } t \leq t_+ \text{ and } \lim_{t \rightarrow -\infty} \dot{g}(t) = 0. \quad (35)$$

Then consider the following autonomous *compactified system*

$$\dot{x} = f(x, \Gamma(h(s))), \quad (36)$$

$$\dot{s} = \gamma(s), \quad (37)$$

$$f(x, \Gamma(h(s))) = \begin{cases} f(x, \Gamma(h(s))) & \text{for } s \in (-1, s_+], \\ f(x, \Gamma^-) & \text{for } s = -1, \end{cases} \quad (38)$$

$$\gamma(s) = \begin{cases} 1/h'(s) & \text{for } s \in (-1, s_+], \\ 0 & \text{for } s = -1, \end{cases} \quad (39)$$

that is defined on the extended phase space $U \times [-1, s_+]$ with flow-invariant subspace $\{s = -1\}$ that carries the autonomous dynamics and compact invariant sets of the past limit system (4).

Proposition 2.1 and Theorem 2.2 apply to compactified system (36)–(39) after we leave out the limits $t \rightarrow +\infty$ ($s \rightarrow 1^-$) and replace: “bi-asymptotically constant” with “asymptotically constant with a past limit”, “compactified system (14)–(17)” with “compactified system (36)–(39)”, “phase space $U \times [-1, 1]$ ” with “phase space $U \times [-1, s_+]$ ” and “Assumption 2.1” with “Assumption 2.3”.

3 Compactified System Dynamics

We now relate we relate the nonautonomous dynamics of system (1) with bi-asymptotically constant $\Gamma(t)$ and the autonomous dynamics of the compactified system (14)–(17). The main focus is on the extrapolation of dynamical structure from (one of) the limit systems (3) or (4). We start with a general remark on modified stability of compact invariant sets for the limit system when embedded in the extended phase space of the compactified system, and follow with rigorous statements that can greatly simplify analysis of (1). In the discussion, we distinguish between equilibria and more general compact invariant sets:

- A denotes a general compact invariant set.
- η denotes a compact invariant set that is an equilibrium point.

Remark 3.1. *In the compactified system (14)–(17), the time evolution of $s(t)$ is not influenced by the time evolution of $x(t)$. Owing to this special skew-product structure:*

- (i) *Any compact invariant set A^+ for the future limit system (3) gains one attracting direction, and any compact invariant set A^- for the past limit system (4) gains one repelling direction when embedded in the extended phase space of (14)–(17), in the sense of monotone increasing $s(t)$:*

$$\dot{s}(t) = \dot{g}(t) > 0.$$

- (ii) *A regular³ compact invariant set A^\pm of one of the limit systems can be viewed as gaining an additional Lyapunov exponent⁴, denoted l_s^\pm , when embedded in the extended phase space. The additional Lyapunov exponent quantifies linear stability in the s -direction, is independent of x , and is given by the $(n + 1, n + 1)$ -th element of the Jacobian (21) at $\{s = \pm 1\}$, or by the limit in the second transformation condition (19):*

$$l_s^\pm = \gamma'(s) \Big|_{s=\pm 1} = - \frac{h''(s)}{(h'(s))^2} \Big|_{s=\pm 1} = \lim_{t \rightarrow \pm\infty} \frac{\ddot{g}(t)}{\dot{g}(t)}, \quad (40)$$

where $' = d/ds$ and the second equality follows from Proposition 2.1. We note that l_s^\pm is zero when $g(t)$ decays slower than exponentially; see Sec. 4.3.

- (iii) *The Lyapunov vector corresponding to the additional Lyapunov exponent is independent of x . The vector is normal to $\{s = \pm 1\}$ if the top n elements in the last column of the Jacobian (21) at $\{s = \pm 1\}$ are zero, or if the first transformation condition (18) is zero*

$$\frac{\partial f}{\partial s} \Big|_{s=\pm 1} = \frac{\partial f}{\partial \Gamma} \Big|_{s=\pm 1} \frac{d}{ds} \Gamma(h(s)) \Big|_{s=\pm 1} = \frac{\partial f}{\partial \Gamma} \Big|_{s=\pm 1} \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{g}(t)} = 0,$$

where the second equality follows from Proposition 2.1.

³A compact invariant set is *regular* if the full Lyapunov spectrum exists.

⁴Referred to as the *normal Lyapunov exponent* in the terminology of [5].

3.1 Attractors and Repellers

In the particular case of an attractor in the future limit system (and a repeller in the past limit system), this can be stated more topologically. We say that a compact invariant set A is an attractor if it is the ω -limit set of a neighborhood of itself, i.e., there is an open set D with $A \subset D$, so that $\omega(D) = A$. With the notation that $\psi(t, y_0)$ is the flow evolution for time t of the initial condition y_0 , which also applies to a set of initial conditions $\psi(t, D) = \{\psi(t, y) : y \in D\}$, the ω -limit set is given by:

$$\omega(D) = \bigcap_{T>0} \overline{\{\psi(t, D) : t > T\}},$$

where \bar{S} denotes the closure of S . Similarly, the α -limit set is given by

$$\alpha(D) = \bigcap_{T<0} \overline{\{\psi(t, D) : t < T\}},$$

and a set A is a repeller if it is the α -limit set of a neighborhood of itself. The following proposition is then readily proved for the future and past limits.

Proposition 3.1. *Suppose that (1) and $\Gamma(t)$ are chosen such that Proposition 2.1 applies. If A^+ is an attractor for the future limit system (3), then $\tilde{A}^+ = \{(x, 1) : x \in A^+\} \subset \{s = 1\}$ is also an attractor for the compactified system (14)–(17) when considered in the extended phase space. If A^- is a repeller for the past limit system (4), then $\tilde{A}^- = \{(x, -1) : x \in A^-\} \subset \{s = -1\}$ is also a repeller for (14)–(17) when considered in the extended phase space.*

Proof of Proposition 3.1

It suffices to show the statement for an attractor as that for a repeller follows by reversing time.

From the theory of isolating blocks, see [9], we can find a neighborhood of A^+ inside $\{s = 1\}$, which we denote B , so that the flow takes points on the boundary ∂B to the interior of B for $t > 0$. Moreover, since this is stable under the continuous perturbation of vector fields and $\dot{s} > 0$ for $s \in (-1, 1)$, we can construct a neighborhood of \tilde{A}^+ , namely $\tilde{B} = B \times (1 - \delta, 1]$ for some $\delta > 0$ that is forward invariant.

Since A is the maximal invariant set in B and $\omega(\tilde{B}) \subset B$ we can conclude that $\omega(\tilde{B}) = \tilde{A}^+$ and thus \tilde{A}^+ is an attractor in the full compactified system. \square

3.2 Saddles and Invariant Manifolds

It follows from Rmk. 3.1 that repellers for the future limit system (3) and attractors for the past limit system (4) become saddles for the compactified system (14)–(17) when considered in the extended phase space. Furthermore, saddles for the future limit system (3) remain saddles but gain one stable direction. Before we discuss the role of saddles and their invariant manifolds in the compactified system, we recall basic facts about invariant manifolds in general.

In the following, it will be important to distinguish between the stable set and stable manifold of a compact invariant set (similarly the unstable set and unstable manifold.) For any compact invariant set A , we define its stable set $w^s(A)$ and unstable set $w^u(A)$ as

$$w^s(A) = \{y : \omega(y) \subseteq A\}, \quad w^u(A) = \{y : \alpha(y) \subseteq A\}.$$

The extra condition for the stable manifold is that the decay is exponential, i.e., the stable manifold $W^s(A)$ and unstable manifold $W^u(A)$ are given by

$$\begin{aligned} W^s(A) &= \{y \in w^s(A) : d(\psi(y, t), A) \leq Ke^{\beta t}, \text{ for some } K > 0, \beta < 0 \text{ and all } t > 0\}, \\ W^u(A) &= \{y \in w^u(A) : d(\psi(y, t), A) \leq Ke^{\beta t}, \text{ for some } K > 0, \beta > 0 \text{ and all } t < 0\}. \end{aligned}$$

The distance function d is between a point and a set, and is defined as the greatest lower bound in the Euclidean distance between the point on the trajectory and any point in the set A :

$$d(\psi(y, t), A) = \inf_{x \in A} \|\psi(y, t) - x\|.$$

We will also need local versions of the above sets/manifolds. Based on an open superset N of a compact invariant set A , we define the local stable set of A :

$$w_{loc}^s(A) = \{y : y \in w^s(A) \text{ and } \psi(y, t) \in N \text{ for all } t > 0\},$$

and a local stable manifold of A :

$$W_{loc}^s(A) = \{y : y \in W^s(A) \text{ and } \psi(y, t) \in N \text{ for all } t > 0\}.$$

The local unstable set of A , denoted $w_{loc}^u(A)$, and the local unstable manifold of A , denoted $W_{loc}^u(A)$, are defined similarly for $t < 0$.

Due to the possibility that $l_s^\pm = 0$, we need to consider compact invariant sets with centre directions. For now, we focus on equilibria and later on, in Sec. 3.4, generalise to more complicated compact invariant sets.

The corresponding invariant manifolds cannot be described in terms of decay rates so straightforwardly. They are characterised as the graphs of functions over the relevant subspaces from the linearised system. Specifically, let E^c , E^s and E^u be the subspaces based at η and spanned by the sets of eigenvectors corresponding to centre (zero real-part) eigenvalues, stable (negative real-part) eigenvalues and unstable (positive real-part) eigenvalues, respectively. These subspaces are referred to as the centre (E^c), stable (E^s) and unstable (E^u) eigenspaces, and are invariant under the linearised system at η . A (local) centre manifold is given as the graph of a Lipschitz function h^c that is flow-invariant relative to some chosen neighborhood of the equilibrium η . More precisely,

$$W_{loc}^c(\eta) = \{\text{gr}(h^c) \text{ where } h^c : E^c \cap N \rightarrow E^s \oplus E^u\},$$

where $W_{loc}^c(\eta)$ is flow-invariant relative to N and tangent to E^c at η . The (local) centre-stable manifold is defined similarly, as the graph of a Lipschitz function h^{cs} that is flow-invariant relative to some chosen neighborhood of the equilibrium η , but with a different domain space. More precisely,

$$W_{loc}^{cs}(\eta) = \{\text{gr}(h^{cs}) \text{ where } h^{cs} : E^c \oplus E^s \cap N \rightarrow E^u\},$$

where $W_{loc}^{cs}(\eta)$ is flow-invariant relative to N and tangent to $E^c \oplus E^s$ at η .

In general, the centre and centre-stable manifold may not be unique in that there may well be other functions whose graphs satisfy the tangency conditions and are invariant relative to N with the corresponding domains and ranges. It turns out that such manifolds can not only appear as a result of compactification, but they can also play a crucial role in simplifying analysis of the original nonautonomous system (1). Therefore, we now address the question of occurrence and uniqueness of centre and centre-stable manifolds in the compactified system (14)–(17).

3.2.1 Dynamical Structure from the Future Limit System

Of particular interest in both the R-tipping and radial steady state problems mentioned in Sec.1.2 is the situation where a hyperbolic saddle equilibrium, denoted η^+ , is present for the future limit system. In the extended phase space of the compactified system, this saddle becomes

$$\tilde{\eta}^+ = (\eta^+, 1) \in \{s = 1\}.$$

If the decay of $\dot{\Gamma}(t)$ to zero as $t \rightarrow +\infty$ is exponential or faster, then it is possible to construct a transformation $s = g(t)$ [e.g. transformation (48)] so the saddle gains an exponentially stable direction and remains hyperbolic with a higher-dimensional stable manifold when embedded in the extended phase space of (14)–(17); see Corollary 4.1. In this case, it follows from the stable manifold theorem that $W^s(\tilde{\eta}^+)$ is unique and

$$W^s(\tilde{\eta}^+) = w^s(\tilde{\eta}^+).$$

However, this may not be true in general.

An interesting situation occurs when the decay of $\dot{\Gamma}(t)$ as $t \rightarrow +\infty$ is slower than exponential. In this case, $l_s^+ = 0$ and the saddle gains a neutrally stable centre direction when embedded in the extended phase space of (14)–(17), meaning that $W^s(\tilde{\eta}^+) \neq w^s(\tilde{\eta}^+)$ because $W^s(\tilde{\eta}^+) \subset \{s = 1\}$; see Corollary 4.2 and Fig. 4(a)–(b). Nonetheless, we can work with a local centre-stable manifold $W_{loc}^{cs}(\tilde{\eta}^+)$ instead. This, however, raises the question of uniqueness of the manifold of interest.

Because of the special structure inherent in the compactified system, $s(t)$ tends monotonically to 1 as $t \rightarrow +\infty$, $W_{loc}^{cs}(\tilde{\eta}^+)$ is forward invariant by construction, and we can ensure that $W_{loc}^{cs}(\tilde{\eta}^+) \subset w^s(\tilde{\eta}^+)$. What is more, it turns out that the centre-stable manifold of $\tilde{\eta}^+$ actually comprises its entire local stable set, and is thus unique, which is very useful in applications where the decay is slower than exponential. We note that the hyperbolicity of the equilibrium within $\{s = 1\}$ is crucial here as any centre directions in the limit system itself would compromise the uniqueness of the centre-stable manifold. This follows from the following theorem.

Theorem 3.2. *Consider a hyperbolic saddle equilibrium η^+ for the future limit system (3) that becomes a non-hyperbolic saddle $\tilde{\eta}^+ = (\eta^+, 1)$ in the compactified system (14)–(17). Then, $W_{loc}^{cs}(\tilde{\eta}^+)$ is unique relative to some chosen neighborhood N of $\tilde{\eta}^+$, and*

$$W_{loc}^{cs}(\tilde{\eta}^+) = w_{loc}^s(\tilde{\eta}^+),$$

in the extended phase space of the compactified system (14)–(17).

Proof of Theorem 3.2

Without loss of generality, we set $\eta^+ = 0$ and introduce $\sigma = s - 1$. We can choose coordinates $x = (y, z)$ so that, in a neighborhood of $(x, \sigma) = (0, 0)$, (31)–(34) becomes

$$\dot{y} = M^+ y + f_1(y, z, \sigma), \quad (41)$$

$$\dot{z} = M^- z + f_2(y, z, \sigma), \quad (42)$$

$$\dot{\sigma} = \beta(\sigma), \quad (43)$$

where $\beta(\sigma) = \gamma(\sigma + 1)$, and $\beta(0) = \beta'(0) = 0$. The spectra of M^+ and M^- are in the right and left half planes respectively. Set

$$N = \{(y, z, \sigma) : \|y\| \leq \epsilon, \|z\| < \epsilon, -\delta < \sigma \leq 0\}.$$

Since the terms f_i are higher order and $\dot{\sigma} > 0$, we can choose ϵ and δ so that the vector field of (41)–(43) points into N on $\|z\| = \epsilon$ and $\sigma = -\delta$ while it points out of N on $\|y\| = \epsilon$, and the last face $\sigma = 0$ ($s = 1$) is invariant.

The variables z and σ represent the stable and centre directions respectively and so we put them together and introduce $\zeta = (z, \sigma)$. We also denote $\xi = (y, \zeta)$ and $\psi(\xi, t) = (y(t), \zeta(t))$. Note that if $\xi \in w_{loc}^s(\tilde{\eta}^+)$ then $\psi(\xi, t) \in N$ for all $t > 0$ by construction of $w_{loc}^s(\tilde{\eta}^+)$. If $\xi \in W_{loc}^{cs}(\tilde{\eta}^+)$ then $\psi(\xi, t) \in N$ for all $t > 0$ by construction of $W_{loc}^{cs}(\tilde{\eta}^+)$ and the direction of the vector field on the faces of N . To see this, note that $W_{loc}^{cs}(\tilde{\eta}^+)$ cannot intersect $\|y\| = \epsilon$ because it is invariant, and the graph of a Lipschitz function $y = h^{cs}(\zeta)$ on which $\dot{\sigma} > 0$.

Suppose that $\xi_1 = (y_1, \zeta_1)$ satisfies $\psi(\xi_1, t) \in N$ for all $t \geq 0$. Let $\xi_2 = (y_2, \zeta_1)$ be the point on W_{loc}^{cs} with the same ζ coordinate as ξ_1 , which will also stay in N for $t > 0$. Under these conditions, the following lemma describes the fate of the hypothesized trajectory from ξ_1 .




Figure 4: (a) A hyperbolic saddle η^+ for the future limit system (3) gains a neutrally stable centre direction and becomes (b) a non-hyperbolic saddle $\tilde{\eta}^+ = (\eta^+, 1)$ in the extended phase space of the compactified system (14)–(17). (c) A hyperbolic stable focus η^- for the past limit system (4) gains a neutrally unstable centre direction and becomes (d) a non-hyperbolic saddle-focus $\tilde{\eta}^- = (\eta^-, -1)$ with one-dimensional centre manifold in the extended phase space of (14)–(17).

Lemma 3.3. *There are coordinates for y and ζ so that the following holds for all $t \geq 0$:*

$$\|y_1(t) - y_2(t)\| \geq \|\zeta_1(t) - \zeta_2(t)\|.$$

Furthermore, there is an $\eta > 0$ so that

$$\|y_1(t) - y_2(t)\| \geq \|y_1(0) - y_2(0)\| \exp(\eta t)$$

for all $t \geq 0$.

If $y_1(0) \neq y_2(0)$ then Lemma 3.3 implies that $\xi_1(t)$ must leave N for some $t > 0$, which is a contradiction. In other words, if we suppose $\xi_1 \in W_{loc}^{cs}(\tilde{\eta})$ then the contradiction gives a unique function $y = h^{cs}(\zeta)$ whose graph satisfies the tangency at $\tilde{\eta}$ and flow-invariance relative to N conditions. If we suppose $\xi_1 \in w_{loc}^s(\tilde{\eta})$ then the contradiction gives $\xi_1 \in W_{loc}^{cs}$ and $W_{loc}^{cs}(\tilde{\eta}) = w_{loc}^s(\tilde{\eta})$. Once we prove Lemma 3.3 the proof of Theorem 3.2 is complete.

Proof of Lemma 3.3

It will be convenient to combine (42) with (43), and write

$$\dot{\zeta} = M^0 \zeta + \tilde{f}_2(y, \zeta). \quad (44)$$

The spectrum of M^0 , denoted $\sigma(M^0)$, contains one neutral and otherwise stable eigenvalues, while that of M^+ contains unstable eigenvalues. Therefore we can find α and β so that, with suitable coordinates, $\langle M^+y, y \rangle \geq \beta \langle y, y \rangle$ and $\langle M^0\zeta, \zeta \rangle \leq \alpha \langle \zeta, \zeta \rangle$, where $0 < \alpha < \beta$, and $\langle \cdot, \cdot \rangle$ denotes the inner product in those coordinates.

Now set $\Delta y = y_1 - y_2$ and $\Delta \zeta = \zeta_1 - \zeta_2$. The functions f_1 and \tilde{f}_2 being higher order means that $Df_1(0, 0, 0) = Df_2(0, 0, 0) = 0$. From the continuity of the first derivatives of these nonlinear terms, given $\rho > 0$, we can find $\epsilon > 0$ and $\delta > 0$ so that

$$\|f_1(y_1, \zeta_1) - f_1(y_2, \zeta_2)\| \leq \rho(\|\Delta y\| + \|\Delta \zeta\|).$$

A similar estimate holds for \tilde{f}_2 . The following inequality can then be shown to hold:

$$\frac{d}{dt}\|\Delta y\|^2 \geq \beta\|\Delta y\|^2 - \rho(\|\Delta y\| + \|\Delta \zeta\|)\|\Delta y\|, \quad (45)$$

$$\frac{d}{dt}\|\Delta \zeta\|^2 \leq \alpha\|\Delta \zeta\|^2 + \rho(\|\Delta y\| + \|\Delta \zeta\|)\|\Delta \zeta\|. \quad (46)$$

Subtracting (46) from (45), we obtain:

$$\frac{d}{dt}(\|\Delta y\|^2 - \|\Delta \zeta\|^2) \geq \beta\|\Delta y\|^2 - \alpha\|\Delta \zeta\|^2 - \rho(\|\Delta y\| + \|\Delta \zeta\|)^2.$$

Initially, $\|\Delta y\| > \|\Delta \zeta\|$ and if they ever became equal, we can conclude from this inequality that

$$\frac{d}{dt}(\|\Delta y\|^2 - \|\Delta \zeta\|^2) \geq (\beta - \alpha - 4\rho)\|\Delta y\|^2.$$

and so, if ρ is sufficiently small, $\frac{d}{dt}(\|\Delta y\|^2 - \|\Delta \zeta\|^2) \geq 0$, and hence $\|\Delta y(t)\| > \|\Delta \zeta(t)\|$ for all $t > 0$ so long as both trajectories stay in N , which they do by hypothesis.

It is now straightforward to obtain the second inequality of the lemma from (45) using $\|\Delta y(t)\| > \|\Delta \zeta(t)\|$ and setting $\eta = \beta - 2\rho$. \square

3.2.2 Dynamical Structure from the Past Limit System

A result completely analogous to Theorem 3.2 can be formulated for the past limit system where s (stable) is replaced by u (unstable) in the manifolds. In this section, however, we will elaborate on an important special case.

Of particular interest in the pullback attractor and R-tipping problems mentioned in Sec.1.2 is the situation where a hyperbolic sink, denoted η^- , is present for the past limit system. In the extended phase space of the compactified system, this equilibrium becomes

$$\tilde{\eta}^- = (\eta^-, -1) \in \{s = -1\}.$$

If the decay of $\dot{\Gamma}(t)$ to zero as $t \rightarrow -\infty$ is exponential or faster, then it is possible to construct a transformation $s = g(t)$ [e.g. transformation (48)] so the sink gains an exponentially unstable direction and becomes a hyperbolic saddle with a one-dimensional unstable manifold when embedded in the extended phase space of (14)–(17); see Corollary 4.1. In this case, it follows from the stable manifold theorem that $W^u(\tilde{\eta}^-)$ is unique and

$$W^u(\tilde{\eta}^-) = w^u(\tilde{\eta}^-).$$

If the decay of $\dot{\Gamma}(t)$ as $t \rightarrow -\infty$ is slower than exponential then the sink gains a neutrally unstable centre direction and becomes a non-hyperbolic saddle when embedded in the extended

phase space of (14)–(17), meaning that $W^u(\tilde{\eta}^-) \neq w^u(\tilde{\eta}^-)$ because $W^u(\tilde{\eta}^-) = \emptyset$; see Corollary 4.2 and Fig. 4(c)–(d). Nonetheless, we can work with a local centre manifold $W_{loc}^c(\tilde{\eta}^-)$. This again raises the question of uniqueness of the manifold of interest.

Because of the special structure inherent in the compactified system, $s(t)$ tends monotonically to -1 as $t \rightarrow -\infty$, $W_{loc}^c(\tilde{\eta}^-)$ is backward invariant by construction, and we can ensure that $W_{loc}^c(\tilde{\eta}^-) \subset w^u(\tilde{\eta}^-)$. As in the considerations underlying Theorem 3.2, the hyperbolicity of the sink within $\{s = -1\}$ and the centre direction being an unstable one entails that the centre manifold of $\tilde{\eta}^-$ comprises its entire local unstable set. This is encapsulated in the following theorem.

Theorem 3.4. *Consider a hyperbolic sink η^- for the past limit system (4) that becomes a non-hyperbolic saddle $\tilde{\eta}^- = (\eta^-, -1)$ in the compactified system (14)–(17). Then, $W_{loc}^c(\tilde{\eta}^-)$ is unique relative to some chosen neighborhood N of $\tilde{\eta}^-$, and*

$$W_{loc}^c(\tilde{\eta}^-) = w_{loc}^u(\tilde{\eta}^-),$$

in the extended phase space of the compactified system (14)–(17).

Proof of Theorem 3.4

This follows immediately from Theorem 3.2 by reversing time and seeing that $W_{loc}^{cu}(\tilde{\eta}^-) = w_{loc}^u(\tilde{\eta}^-)$. But, in this case, there is no unstable direction and so $W_{loc}^{cu} = W_{loc}^c$. \square

3.3 Invariant Manifolds and Applications

The key point of Sec. 3 is that the unique invariant manifolds extrapolated from the future limit system, from the past limit system, or from both, can be used to analyse the nonautonomous system (1) with exponential or non-exponential decaying $\Gamma(t)$. Specifically, the benefits of compactification when studying nonlinear dynamics are the following:

- (i) In contrast to the original nonautonomous system (1), the compactified system (14)–(17) is autonomous and may contain equilibria and more complicated compact invariant sets, e.g. limit cycles, invariant tori or strange attractors.
- (ii) In the extended phase space of (14)–(17), the temporal or spatial variation of the nonautonomous term $\Gamma(t)$ becomes in a certain sense ‘encoded’ in the geometric shape of:
 - The stable or centre-stable invariant manifold of a saddle $\tilde{\eta}^+$.
 - The unstable or centre invariant manifold of a saddle $\tilde{\eta}^-$.
 - The basin of attraction of an attractor \tilde{A}^+ .
- (iii) In the extended phase space of (14)–(17):
 - Orbits contained either in the basin of attraction of an attractor \tilde{A}^+ , or in the stable or centre-stable invariant manifold of a saddle $\tilde{\eta}^+$, correspond to solutions of the nonautonomous system (1) that remain bounded as $t \rightarrow +\infty$.
 - Orbits contained in the unstable or centre invariant manifold of a saddle $\tilde{\eta}^-$ correspond to solutions of the nonautonomous system (1) that remain bounded as $t \rightarrow -\infty$.
 - Connecting heteroclinic orbits from a saddle $\tilde{\eta}^-$ to either a saddle $\tilde{\eta}^+$ or an attractor \tilde{A}^+ correspond to solutions of the nonautonomous system (1) that remain bounded as $t \rightarrow \pm\infty$.

- (iv) The compactification enables numerical detection and parameter continuation of solutions to the nonautonomous system (1) that remain bounded as $t \rightarrow \pm\infty$. Since such solutions become connecting heteroclinic orbits in the compactified system (14)–(17), they can be detected and continued numerically, for example using the implementation of the Lin’s method [27] in the numerical continuation software AUTO [11, 34, 22, 48].

These benefits are best illustrated using the three motivating examples from Sec. 1.2.

3.3.1 Pullback Attractors

A local pullback attractor for a nonautonomous system that is asymptotically constant as $t \rightarrow -\infty$ will emanate from a normally hyperbolic attractor A^- for the past limit system. For example, consider a sink η^- for the past limit system, which becomes a saddle $\tilde{\eta}^- = (\eta^-, -1)$ when embedded in the extended phase space of the compactified system (14)–(17). Then, the local pullback point attractor of (1) associated with η^- is transformed into the unstable or centre invariant manifold of the saddle $\tilde{\eta}^-$ in the extended phase space of (14)–(17).

3.3.2 Rate-Induced Tipping

The compactification lays foundations for genuine advances in R-tipping, which is of great interest to the nonlinear science community [7]. In the R-tipping problems, the objective is to find thresholds for the tipping. We claim these are naturally anchored in the future limit system by a saddle or repeller that is normally hyperbolic. We call such a state for the future limit system an *R-tipping edge state*; see [47] where this terminology is fully explained. The challenge is to pin down, in the extended phase space of (14)–(17), what we call *R-tipping thresholds* [47] separating solutions of (1) that R-tip from those that do not. The key point is that these can be naturally characterised as the n -dimensional stable or centre-stable manifolds of an appropriate saddle edge state. What is more, the critical rates of change of $\Gamma(t)$ at which the system transitions to a different state correspond to heteroclinic connections from a saddle $\tilde{\eta}^-$ to the saddle edge state. Thus, the compactification: (i) simplifies rigorous analysis such as derivation of testable criteria for the occurrence of R-tipping, as demonstrated in [47], and (ii) enables numerical detection and parameter continuation of genuine nonautonomous R-tipping bifurcations, as demonstrated in [48, Ch.4] for a number of canonical examples.

3.3.3 Radial Steady States

For the construction of radial steady states as in (5), the invariant manifolds constructed above will provide the sets of solutions satisfying the boundary conditions. We would be interested in saddles for the future limit system. This is because, in the extended phase space of (14)–(17), the (family of) solutions to the radial steady state problem (1) must decay as $r \rightarrow +\infty$. This would then use a right-sided compactification. A separate construction is needed for the boundary condition at $r = 0$. When this has been carried out, the problem becomes one of finding heteroclinic connections given by transverse intersections of the unstable or centre manifold of a saddle $\tilde{\eta}^-$ and the stable or centre-stable manifold of a saddle $\tilde{\eta}^+$, see [17].

3.3.4 Stability of Nonlinear Waves

The Evans Function is defined using sets of solutions of (9)–(10) which decay at $\pm\infty$, denoted $Y_{\pm}(z)$. Using exterior algebra, the Evans Function is defined as a generalised angle between these subspaces when compared at a fixed $z \in (-\infty, +\infty)$. These subspaces are realised in the compactified system as (at $-\infty$) a centre-unstable manifold of the zero solution for the left-sided

compactification and, analogously, (at $+\infty$) a centre-stable manifold of the zero solution for the right-sided compactification; see [1].

In most circumstances, the decay of the underlying wave to its end-states U_{\pm} is exponential. Nevertheless, the theory presented here would afford a construction of the subspaces $Y_{\pm}(z)$, and hence the Evans Function, when the decay were much weaker than exponential.

3.4 Generalisations to More Complicated Compact Invariant Sets

We anticipate that it will be possible to extend the results for equilibria from Secs. 3.2.1 and 3.2.2 to cover more complicated compact invariant sets in the limit systems and their invariant manifolds in the extended phase space of the compactified system. It is worth distinguishing two separate cases here.

If the compact invariant set A is a submanifold of $\{s = 1\}$, then the relevant invariant manifold theorems can be inferred from the results in Fenichel [15]. Theorems 3.2 and 3.4 will follow in a straightforward manner with the use of sub-bundles of the relevant tangent spaces as opposed to subspaces.

Arnol'd et al. [4, Sec.8.2] define a strange attractor as an attractor that is not the union of smooth submanifolds. Similarly, we might define a strange compact invariant set as one that cannot be cast as the union of submanifolds. The idea from the previous paragraph should extend to the case of a union of submanifolds, and so all non-strange compact invariant sets will be covered. This will obviously include important invariant sets such as periodic orbits and tori.

Invariant manifold theorems do exist for general compact invariant sets, including strange ones, see the book by Robinson [39] for instance. Nevertheless, it is not apparent how to generalise the arguments in the proofs of Theorems 3.2 and 3.4 as these proofs require a nicer local structure than might hold near a more complicated compact invariant set or strange attractor. We formulate the general result as a conjecture.

Conjecture 3.5. *The statements for equilibria in Theorems 3.2 and 3.4 also hold for general normally hyperbolic compact invariant sets A^{\pm} of the limit systems.*

4 Examples of Compactification

In this section we construct actual examples of coordinate transformations $g(t)$, given by (11), (30) and (35) and depicted in Fig. 1(a)–(c), that are useful in practice. This raises the question of the ‘rate’ of compactification relative to the rate of asymptotic decay of the nonautonomous term. To address this question, we introduce in Secs. 4.2 and 4.3 compactification parameters and work with *parametrised compactifications*. The additional freedom acquired in a parametrisation of $g(t)$ allows us to:

- (i) Fulfill the compactification conditions (18)–(19) by a suitable choice of the compactification parameter(s).
- (ii) Control the direction of the additional Lyapunov vector transverse to invariant subspaces $\{s = \pm 1\}$ to facilitate analysis.

Moreover, we consider different types of asymptotic decay of $g(t)$ giving rise to different behaviour of $s(t)$ near $\{s = \pm 1\}$.

4.1 Γ -Compactification

In some cases, a compactification transformation can be constructed in terms of (one component of) the nonautonomous term $\Gamma(t) \in \mathbb{R}^d$ itself; see [7, 36, 35, 38, 2] for examples. Let $\tilde{\Gamma}(t) =$

($\dot{\Gamma}_1(t), \dots, \dot{\Gamma}_d(t)$) be ordered such that $\dot{\Gamma}_1(t)$ decays not faster than the other components

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}_i(t)}{\dot{\Gamma}_1(t)} = L^\pm \in \mathbb{R} \text{ for } i = 2, \dots, d.$$

If $\Gamma_1(t)$ satisfies Assumption 2.1 with future limit Γ_1^+ and past limit Γ_1^- , we can construct a coordinate transformation (11) in terms of $\Gamma_1(t)$ as follows

$$g(t) = \frac{2\Gamma_1(t) - \Gamma_1^+ - \Gamma_1^-}{\Gamma_1^+ - \Gamma_1^-}. \quad (47)$$

One can verify that such $g(t)$ satisfies the first transformation condition (18):

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}_i(t)}{\dot{g}(t)} = \begin{cases} (\Gamma_1^+ - \Gamma_1^-)/2 \neq 0 & \text{for } i = 1, \\ (\Gamma_1^+ - \Gamma_1^-)/2 \neq 0 \text{ or } 0 & \text{for } i = 2, \dots, d, \end{cases}$$

and satisfies the second transformation condition (19):

$$\lim_{t \rightarrow \pm\infty} \frac{\ddot{g}(t)}{\dot{g}(t)} = \lim_{t \rightarrow \pm\infty} \frac{\ddot{\Gamma}_1(t)}{\dot{\Gamma}_1(t)} \text{ exists,}$$

if $\Gamma_1(t)$ satisfies (19). Similarly, one can construct a coordinate transformation (30):

$$g(t) = \frac{\Gamma_1(t) - \Gamma_1(0)}{\Gamma_1^+ - \Gamma_1(0)},$$

if $\Gamma_1(t)$ satisfies Assumption 2.2 with future limit Γ_1^+ , or a coordinate transformation (35):

$$g(t) = -\frac{\Gamma_1(t) - \Gamma_1(0)}{\Gamma_1^- - \Gamma_1(0)},$$

if $\Gamma_1(t)$ satisfies Assumption 2.3 with past limit Γ_1^- .

For example, one can use [7, 36]:

$$g(t) = \tanh(t) \quad \text{if } \Gamma_1(t) = \tanh(t).$$

This approach can be extended to certain non-monotone inputs that can be expressed in terms of a monotone growth that satisfies Assumption 2.1, such as

$$\Gamma_1(t) = \operatorname{sech}(t) = \sqrt{1 - \tanh(t)^2}.$$

Constructing $g(t)$ in terms of $\Gamma_1(t)$ is not always possible, for example it usually does not work for non-monotone $\Gamma_1(t)$. When possible, it has certain limitations. For example, there may be no algebraic formula for the inverse $h(s)$ required for the augmented component (15) of the vector field. Moreover, to facilitate analysis, one may wish the additional Lyapunov vectors to be normal to invariant subspaces $\{s = \pm 1\}$, which does not normally occur for the Γ -compactification because the resulting Jacobian (21) is not block-diagonal. Therefore, more universal compactification transformations will be required in general.

4.2 Exponential Compactification

Guided by Assumption 2.1 and Proposition 2.1, we construct an example of a *parametrised coordinate transformation* (11) with exponential asymptotic decay as proposed in [1, Eq.(3.2)]:

$$s = g_{(\alpha)}(t) = \tanh\left(\frac{\alpha t}{2}\right), \quad t = g_{(\alpha)}^{-1}(s) := h_{(\alpha)}(s) = \frac{1}{\alpha} \ln \frac{1+s}{1-s}, \quad (48)$$

where the *compactification parameter* $\alpha > 0$ parametrises the *rate* of exponential decay of $\dot{g}_{(\alpha)}(t)$. One can verify that this transformation satisfies Assumption 2.1 and the second transformation condition (19):

$$\lim_{t \rightarrow \pm\infty} \frac{\ddot{g}_{(\alpha)}(t)}{\dot{g}_{(\alpha)}(t)} = -\alpha \lim_{t \rightarrow \pm\infty} \tanh\left(\frac{\alpha t}{2}\right) = \mp\alpha. \quad (49)$$

The augmented component $\gamma_{(\alpha)}(s)$ of the vector field is obtained by computing and then inverting $h'_{(\alpha)}(s)$, so that the compactified system (14)–(15) becomes

$$\left. \begin{aligned} \dot{x} &= f(x, \Gamma(h_{(\alpha)}(s))), \\ \dot{s} &= \frac{\alpha}{2} (1 - s^2). \end{aligned} \right\} \quad (50)$$

We now examine properties of (50) with dependence on α and the asymptotic decay of $\dot{\Gamma}(t)$.

Corollary 4.1. (Compactification for $\dot{\Gamma}(t)$ with exponential or faster decay.) *Consider a nonautonomous system (1) with C^1 -smooth f and Γ , and bi-asymptotically constant $\Gamma(t)$. Moreover, suppose there is a $\rho > 0$ such that*

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{e^{\mp\rho t}} \text{ exist.} \quad (51)$$

Then, given the coordinate transformation (48) with any $\alpha \in (0, \rho]$, the ensuing autonomous compactified system (50) is C^1 -smooth on the extended phase space $U \times [-1, 1]$.

The distance between any trajectory of (50) and $\{s = \pm 1\}$ decays exponentially at a rate α as $t \rightarrow \pm\infty$. A regular compact invariant set of the future limit system gains one negative Lyapunov exponent $l_{s,(\alpha)}^+ = -\alpha < 0$, and a regular compact invariant set of the past limit system gains one positive Lyapunov exponent $l_{s,(\alpha)}^- = \alpha > 0$, when embedded in the extended phase space of (50).

In other words, exponentially or faster decaying $\dot{\Gamma}(t)$ is sufficient for (50) to be C^1 -smooth and for regular compact invariant sets of the limit systems to gain a non-zero Lyapunov exponent when embedded in the extended phase space of (50). This gives rise to pure stable and unstable manifolds in the extended phase space.

Proof of Corollary 4.1

Given (51), it follows from the asymptotic properties of $\dot{g}_{(\alpha)}(t)$:

$$\dot{g}_{(\alpha)}(t) \sim 2\alpha e^{\mp\alpha t} \text{ as } t \rightarrow \pm\infty,$$

and from the algebraic limit theorem

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{g}_{(\alpha)}(t)} = \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{e^{\mp\rho t}} \frac{e^{\mp\rho t}}{\dot{g}_{(\alpha)}(t)} = \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{e^{\mp\rho t}} \lim_{t \rightarrow \pm\infty} \frac{e^{\mp\rho t}}{\dot{g}_{(\alpha)}(t)} = \frac{1}{2\alpha} \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{e^{\mp\rho t}} \lim_{t \rightarrow \pm\infty} e^{\mp(\rho-\alpha)t},$$

that the first transformation condition (18) is satisfied if $0 < \alpha \leq \rho$. It follows from (49) that the second transformation condition (19) is satisfied for $\alpha > 0$. It then follows from Proposition 2.1

that the compactified system (50) is C^1 -smooth on the extended phase space $U \times [-1, 1]$ for any $0 < \alpha \leq \rho$.

The Hausdorff semi-distance $d(p(t), S^\pm)$ between a point $p(t) = (x(t), s(t))$ on a trajectory and invariant subspace $S^\pm = \{(x, s) : s = \pm 1\}$ in the extended phase space is the distance between $s(t)$ and ± 1 . Thus, exponential decay of $d(p(t), S^\pm)$ follows from exponential approach of $s(t)$ towards ± 1 as $t \rightarrow \pm\infty$:

$$d(p(t), S^\pm) = \inf_{y \in S^\pm} \|p(t) - y\| = 1 \mp s(t) = \frac{2}{e^{\pm\alpha t} + 1}.$$

It follows from Eq. (40) that the additional Lyapunov exponent $l_{s,(\alpha)}^\pm$ is given by Eq. (49). \square

Similarly, one can discuss one-sided subcases of the two-sided compactification above, using an example of parametrised coordinate transformation (30) with exponential decay in the form

$$s = g_{(\alpha)}(t) = 1 - e^{-\alpha t}, \quad t = h_{(\alpha)}(s) = -\frac{1}{\alpha} \ln(1 - s), \quad (52)$$

or example of parametrised coordinate transformation (35) with exponential decay in the form

$$s = g_{(\alpha)}(t) = -1 + e^{\alpha t}, \quad t = h_{(\alpha)}(s) = \frac{1}{\alpha} \ln(1 + s). \quad (53)$$

4.3 Algebraic Compactification

Guided by Assumption 2.1 and Proposition 2.1, we construct an inverse of the *parametrised compactification transformation* (11) with algebraic asymptotic decay in the form akin to stereographic projection

$$t = g_{(\alpha)}^{-1}(s) := h_{(\alpha)}(s) = \frac{s}{(1 - s^2)^{\frac{1}{\alpha}}}, \quad (54)$$

where the *compactification parameter* $\alpha > 0$ parametrises the *order* of algebraic decay of $\dot{g}_{(\alpha)}(t)$. There is no formula for the corresponding compactification transformation $s = g_{(\alpha)}(t)$ in the general case $\alpha > 0$. However, there is one in the special case $\alpha = 1$:

$$s = g(t) = \frac{-1 + \sqrt{1 + 4t^2}}{2t}, \quad t = h(s) = \frac{s}{1 - s^2}. \quad (55)$$

One can verify that transformation (54) satisfies Assumption 2.1 and the second transformation condition (19):

$$\lim_{s \rightarrow \pm 1} \frac{h_{(\alpha)}''(s)}{(h_{(\alpha)}'(s))^2} = \lim_{s \rightarrow \pm 1} \frac{2(2 - \alpha)s^3 + 6\alpha s}{[\alpha(1 - s^2) + 2s^2]^2} (1 - s^2)^{\frac{1}{\alpha}} = 0. \quad (56)$$

The augmented component $\gamma_{(\alpha)}(s)$ of the vector field is obtained by computing and then inverting $h_{(\alpha)}'(s)$, so that the compactified system (14)–(15) becomes

$$\left. \begin{aligned} \dot{x} &= f(x, \Gamma(h_{(\alpha)}(s))), \\ \dot{s} &= \frac{\alpha(1 - s^2)^{1 + \frac{1}{\alpha}}}{\alpha(1 - s^2) + 2s^2}. \end{aligned} \right\} \quad (57)$$

We now examine properties of (57) with dependence on α and the asymptotic decay of $\dot{\Gamma}(t)$.

Corollary 4.2. (Compactification for $\dot{\Gamma}(t)$ with algebraic or faster decay.) *Consider a nonautonomous system (1) with C^1 -smooth f and Γ , and bi-asymptotically constant $\Gamma(t)$. Moreover, suppose there is an $m > 1$ such that*

$$\lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{t^{-m}} \text{ exist.} \quad (58)$$

Then, given the coordinate transformation (48) with any $\alpha \in (0, m-1]$, the ensuing autonomous compactified system (57) is C^1 -smooth on the extended phase space $U \times [-1, 1]$.

The distance between any trajectory of (57) and $\{s = \pm 1\}$ decays algebraically as $t \rightarrow \pm\infty$. A regular compact invariant set of a limit system gains one zero Lyapunov exponent $l_{s,(\alpha)}^\pm = 0$ when embedded in the extended phase space of (57).

In other words, algebraically or faster decaying $\dot{\Gamma}(t)$ is sufficient for (57) to be C^1 -smooth, but regular compact invariant sets for the limit systems gain one zero Lyapunov exponent when embedded in the extended phase space of (57). This gives rise to centre-stable and centre manifolds in the extended phase space.

Proof of Corollary 4.2

Given (58), it follows from the asymptotic properties of $h_{(\alpha)}(t)$ and from the algebraic limit theorem

$$\begin{aligned} \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{\dot{g}_{(\alpha)}(t)} &= \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{t^{-m}} \frac{t^{-m}}{\dot{g}_{(\alpha)}(t)} = \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{t^{-m}} \lim_{t \rightarrow \pm\infty} \frac{t^{-m}}{\dot{g}_{(\alpha)}(t)} = \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{t^{-m}} \lim_{s \rightarrow \pm 1} \frac{(h_{(\alpha)}(s))^{-m}}{\gamma_{(\alpha)}(s)} \\ &= \lim_{t \rightarrow \pm\infty} \frac{\dot{\Gamma}(t)}{t^{-m}} \lim_{s \rightarrow \pm 1} \frac{1}{\alpha s^m} \left(\alpha(1-s^2)^{\frac{m-1}{\alpha}} + 2s^2(1-s^2)^{\frac{m-1-\alpha}{\alpha}} \right), \end{aligned}$$

that the first transformation condition (18) is satisfied if both $(1-s^2)$ terms above have non-negative powers, or if $0 < \alpha \leq m-1$. It follows from (56) that the second transformation condition (19) is satisfied for $\alpha > 0$. It then follows from Proposition 2.1 that the compactified system (57) is C^1 -smooth on the extended phase space $U \times [-1, 1]$ for any $0 < \alpha \leq m-1$.

The Hausdorff semi-distance $d(p(t), S^\pm)$ between a point $p(t) = (x(t), s(t))$ on a trajectory and invariant subspace $S^\pm = \{(x, s) : s = \pm 1\}$ in the extended phase space is the distance between $s(t)$ and ± 1 . Thus, algebraic decay of $d(p(t), S^\pm)$ follows from the algebraic approach of $s(t)$ towards ± 1 as $t \rightarrow \pm\infty$:

$$d(p(t), S^\pm) = \inf_{y \in S^\pm} \|p(t) - y\| = 1 \mp s(t).$$

It follows from Eq. (40) that the additional Lyapunov exponent $l_{s,(\alpha)}^\pm$ is given by Eq. (56). \square

Similarly, one can discuss one-sided subcases of the two-sided compactification above, using an example of a parametrised inverse coordinate transformation (30) with algebraic decay in the form

$$t = h_{(\alpha)}(s) = \frac{s}{(1-s)^{\frac{1}{\alpha}}}, \quad (59)$$

with the special case $\alpha = 1$:

$$s = g(t) = \frac{t}{1+t}, \quad t = h(s) = \frac{s}{1-s}, \quad (60)$$

used in the radial steady state problem [17], or example of parametrised inverse coordinate transformation (35) with algebraic decay in the form

$$t = h_{(\alpha)}(s) = \frac{s}{(1+s)^{\frac{1}{\alpha}}}, \quad (61)$$

with the special case $\alpha = 1$:

$$s = g(t) = \frac{t}{1-t}, \quad t = h(s) = \frac{s}{1+s}. \quad (62)$$

4.4 Other Compactification Types and Decay Limitation

In Secs. 4.2 and 4.3 we constructed examples of exponential (48) and algebraic (54) transformations, respectively, with the same type and rate of asymptotic decay as $t \rightarrow \pm\infty$. Transformations with different rates or different types of asymptotic decay as $t \rightarrow +\infty$ and $t \rightarrow -\infty$ are also possible. However, there is one important limitation, namely that the second compactification condition (19) excludes coordinate transformations with faster than exponential decay; see Remark 2.2.

In case of bi-asymptotically constant $\Gamma(t)$ whose speed $\|\dot{\Gamma}(t)\|$ has the same type and rate of asymptotic decay as $t \rightarrow \pm\infty$, the natural choice is transformation (11) whose derivative $\dot{g}_{(\alpha)}(t)$ has one type and rate of asymptotic decay. Such a transformation works also for $\dot{\Gamma}(t)$ with different rates or types of asymptotic decay.

Nonetheless, in the case of bi-asymptotically constant $\Gamma(t)$ whose speed $\|\dot{\Gamma}(t)\|$ has the same type but different rates of asymptotic decay as $t \rightarrow +\infty$ and $t \rightarrow -\infty$, one may wish to construct a transformation (11) that, unlike examples (48) and (54), has the same type but different rates of asymptotic decay as $t \rightarrow +\infty$ and $t \rightarrow -\infty$. For example, the exponential transformation (48) can be generalised to

$$t = h_{(\alpha_-, \alpha_+)}(s) = \frac{1}{\alpha_-} \ln(1+s) - \frac{1}{\alpha_+} \ln(1-s), \quad (63)$$

where the *compactification parameters* α_- and $\alpha_+ > 0$ parametrise the *rates* of exponential decay of $\dot{g}_{(\alpha_-, \alpha_+)}(t)$ as $t \rightarrow -\infty$ and $t \rightarrow +\infty$, respectively. Similarly, algebraic transformation (54) can be generalised to

$$t = h_{(\alpha_-, \alpha_+)}(s) = \frac{s}{(1+s)^{\frac{1}{\alpha_-}} (1-s)^{\frac{1}{\alpha_+}}}, \quad (64)$$

where the *compactification parameters* α_- and $\alpha_+ > 0$ parametrise the *orders* of algebraic decay of $\dot{g}_{(\alpha_-, \alpha_+)}(t)$ as $t \rightarrow -\infty$ and $t \rightarrow +\infty$, respectively. In the case of bi-asymptotically constant $\Gamma(t)$ whose speed $\|\dot{\Gamma}(t)\|$ has different types of asymptotic decay as $t \rightarrow +\infty$ and $t \rightarrow -\infty$, one may wish to construct a ‘hybrid’ transformation (11) that, unlike examples (63) and (64), has different types of asymptotic decay as $t \rightarrow +\infty$ and $t \rightarrow -\infty$. For each option discussed above, the first transformation condition (18) is satisfied by choosing the compactification parameter(s) so that $\dot{g}_{(\alpha_-, \alpha_+)}(t)$ does not decay faster than $\dot{\Gamma}(t)$.

5 Conclusion

We have developed a general framework to facilitate stability and nonlinear dynamics analysis in nonautonomous dynamical systems where the nonautonomous terms decay asymptotically. We refer to such systems as asymptotically autonomous dynamical systems [28]. The theoretical work is directly motivated by a wide range of problems from applications including pullback attractors, rate-induced tipping and nonlinear wave solutions, all of which fit naturally into the asymptotically autonomous setting. The analysis is based on a suitable compactification technique developed in Section 2, in conjunction with a dynamical system approach to study invariant manifolds of saddles in the ensuing autonomous compactified system.

The main obstacle to the analysis of the original nonautonomous system is the absence of compact invariant sets, such as equilibria, limit cycles or invariant tori, in its phase space. The key idea to overcome this obstacle is twofold:

- Introduce compact invariant sets to the problem: The original nonautonomous system is reformulated into an *autonomous compactified system* by augmenting the phase space with an additional bounded but open dimension, which is then extended at one or both ends by bringing in flow-invariant subspaces that carry autonomous dynamics and compact invariant sets of the limit systems from infinity.
- Use autonomous dynamics and compact invariant sets of the limit systems from infinity to analyse the original nonautonomous system: it turns out that solutions of interest form unique invariant manifolds of saddles for the limit systems when embedded in the extended phase space of the compactified system.

We describe a general case with an arbitrary decay of nonautonomous terms and construct the compactified system for this general case. The main results can be summarised as follows:

- In Theorem 2.2 we derive the weakest decay conditions possible for the existence of a C^1 -smooth compactified system to enable the construction of invariant manifolds (stable, unstable, centre, etc.) in the compactified phase space.
- We describe the compactified system dynamics topologically in terms of invariant manifolds of saddles in the extended phase space. We show in Theorems 3.2 and 3.4 that the manifolds of interest are unique and comprise the entire local stable or unstable set of the saddle, even if the saddle gains a centre direction due to non-exponential decay of the nonautonomous term(s) and the manifolds are not pure stable or unstable manifolds.
- We construct examples of parametrised (exponential and algebraic) compactifications that can be implemented in actual practice.

Our general compactification framework provides an alternative tool for the analysis of asymptotically autonomous dynamical systems that complements the existing approach based on asymptotic equivalence of the nonautonomous system and the autonomous limit systems. Extending our framework to other problems such as exponential dichotomies or infinite dimensions for PDEs remains an interesting research question for future study.

Acknowledgments

The authors would like to thank P. Ashwin, T. Carroll, F. Holland, M. Krupa and M. Rasmussen for interesting discussions. C.X. and S.W. received funding from the the CRITICS Innovative Training Network via the European Union’s Horizon 2020 research and innovation programme under Grant Agreement No. 643073. CJ acknowledges support from the US Office of Naval Research under grant number N00014-18-1-2204 and the US National Science Foundation under grant number DMS-1722578.

References

- [1] J. Alexander, R. Gardner, and C. K. R. T. Jones. A topological invariant arising in the stability analysis of travelling waves. *J. Reine Angew. Math*, **410**:167–212, 1990.
- [2] H. Alkhayuon and P. Ashwin. Rate-induced tipping from periodic attractors: Partial tipping and connecting orbits. *Chaos*, **28**, 033608, 2018.
- [3] H. Alkhayuon, P. Ashwin, L.C. Jackson, C. Quinn, and R.A. Wood. Basin bifurcations, oscillatory instability and rate-induced thresholds for atlantic meridional overturning circulation in a global oceanic box model. *Proceedings of the Royal Society A*, **475**(2225):20190051, 2019.

- [4] V.I. Arnold, V.S. Afrajmovich, Yu.S. Il'yashenko, and L.P. Shil'nikov. *Dynamical systems V: bifurcation theory and catastrophe theory*, volume 5. Springer Science & Business Media, 2013.
- [5] P. Ashwin, J. Buescu, and I. Stewart. From attractor to chaotic saddle: a tale of transverse instability. *Nonlinearity*, **9**(3):703–737, 1996.
- [6] P. Ashwin, C. Perryman, and S. Wieczorek. Parameter shifts for nonautonomous systems in low dimension: bifurcation- and rate-induced tipping. *Nonlinearity*, **30**(6):2185–2210, 2017.
- [7] P. Ashwin, S. Wieczorek, R. Vitolo, and P. Cox. Tipping points in open systems: bifurcation, noise-induced, and rate-dependent examples in the climate system. *Phil. Trans. Roy. Soc. A*, **370**(1962):1166–1184, 2012.
- [8] R. Carles. Critical nonlinear Schrödinger equations with and without harmonic potential. *Mathematical Models and Methods in Applied Sciences*, **12**(10):1513–1523, 2002.
- [9] C. Conley and R. Easton. Isolated invariant sets and isolating blocks. *Transactions of the American Mathematical Society*, **158**(1):35–61, 1971.
- [10] J. Delgado, E.A. Lacombe, J. Llibre, and E. Pérez. *Poincaré Compactification of Hamiltonian Polynomial Vector Fields*. In: *Dumas H.S., Meyer K.S., Schmidt D.S. (eds) Hamiltonian Dynamical Systems*. The IMA Volumes in Mathematics and its Applications, vol 63. Springer, New York, 1995.
- [11] Eusebius J Doedel, Thomas F Fairgrieve, Björn Sandstede, Alan R Champneys, Yuri A Kuznetsov, and Xianjun Wang. Auto-07p: Continuation and bifurcation software for ordinary differential equations. 2007.
- [12] F. Dumortier. *Techniques in the theory of local bifurcations: Blow-up, normal forms, nilpotent bifurcations, singular perturbations*. Springer, Dordrecht, 19–73, 1993.
- [13] F. Dumortier. Compactification and desingularization of spaces of polynomial Liénard equations. *J. Differential Equations*, **224**:296–313, 2006.
- [14] F. Dumortier and R. Roussarie. Canard cycles and center manifolds. *Memoirs of the American Mathematical Society*, 121(577), 1996.
- [15] N. Fenichel and J.K. Moser. Persistence and smoothness of invariant manifolds for flows. *Indiana University Mathematics Journal*, **21**(3):193–226, 1971.
- [16] A. Giraldo, B. Krauskopf, and H.M. Osinga. Cascades of Global Bifurcations and Chaos near a Homoclinic Flip Bifurcation: A Case Study. *SIAM Journal on Applied Dynamical Systems*, **17**(4):2784–2829, 2018.
- [17] C. Jones and A. Küpper. On the infinitely many solutions of a semilinear elliptic equation. *SIAM Journal on Mathematical Analysis*, **17**(4):803–835, 1986.
- [18] C. Kiers and C.K.R.T. Jones. On conditions for rate-induced tipping in multi-dimensional dynamical systems. *arXiv:1810.02808*, 2018.
- [19] P. Kloeden and M. Rasmussen. Nonautonomous dynamical systems. *AMS Mathematical Surveys and Monographs*, **176**, 2011.

- [20] I. Kosiuk and P. Szmolyan. Scaling in singular perturbation problems: blowing up a relaxation oscillator. *SIAM Journal on Applied Dynamical Systems*, **10**(4):1307–1343, 2011.
- [21] B. Krauskopf. Bifurcations at ∞ in a model for 1:4 resonance. *Ergod. Th. & Dynam. Sys.*, **17**:899–931, 1997.
- [22] Bernd Krauskopf and Thorsten Rieß. A lin’s method approach to finding and continuing heteroclinic connections involving periodic orbits. *Nonlinearity*, **21**(8):1655, 2008.
- [23] K.U. Kristiansen. Blowup for flat slow manifolds. *Nonlinearity*, **30**:2138–2184, 2017.
- [24] M. Krupa and P. Szmolyan. Extending geometric singular perturbation theory to nonhyperbolic points—fold and canard points in two dimensions. *SIAM Journal on Mathematical Analysis*, **33**(2):286–314, 2001.
- [25] C. Kuehn. Normal hyperbolicity and unbounded critical manifolds. *Nonlinearity*, **27**:1351–1366, 2014.
- [26] J.A. Langa, J.C. Robinson, and A. Suárez. Forwards and pullback behaviour of a non-autonomous lotka–volterra system. *Nonlinearity*, **16**(4):1277–1293, 2003.
- [27] Xiao-Biao Lin. Using melnikov’s method to solve silnikov’s problems. *Proc. Roy. Soc. Edinburgh*, **116**(3-4):295–325, 1990.
- [28] L. Markus. Asymptotically autonomous differential systems. *Contributions to the Theory of Nonlinear Oscillations (Annals of Mathematics Studies, no. 36)*, vol. **3**, Princeton, NJ:Princeton University Press 17–29, 1956.
- [29] K. Matsue. On Blow-Up Solutions of Differential Equations with Poincaré-Type Compactifications. *SIAM Journal on Applied Dynamical Systems*, **17**(3):2249–2288, 2018.
- [30] R. McGehee. Triple collision in the collinear three-body problem. *Inventiones mathematicae*, **27**(3):191–227, 1974.
- [31] M. Messaias. Dynamics at infinity and the existence of singularly degenerate heteroclinic cycles in the Lorenz system. *J. Phys. A: Math. Theor.*, **42**:115101, 2009.
- [32] J. Mitry, M. McCarthy, N. Kopell, and M. Wechselberger. Excitable neurons, firing threshold manifolds and canards. *Journal of Mathematical Neuroscience*, **3**(12), 2013.
- [33] P.E. O’Keeffe and S. Wicczorek. Tipping phenomena and points of no return in ecosystems: Beyond classical bifurcations. *SIAM Journal on Applied Dynamical Systems*, **19**(4):2371–2402, 2020.
- [34] Bart E Oldeman, Alan R Champneys, and Bernd Krauskopf. Homoclinic branch switching: A numerical implementation of lin’s method. *International Journal of Bifurcation and Chaos*, **13**(10):2977–2999, 2003.
- [35] C. G. Perryman. *How fast is too fast? Rate-induced bifurcations in multiple time-scale systems*. Ph.D. thesis, University of Exeter, 2015.
- [36] C. G. Perryman and S. Wicczorek. Adapting to a changing environment: Non-obvious thresholds in multi-scale systems. *Proc. Roy. Soc. A*, **470**:20140226, 2014.
- [37] M. Rasmussen. Bifurcations of asymptotically autonomous differential equations. *Set-Valued Analysis*, **16**:821–849, 2008.

- [38] P. Ritchie and J. Sieber. Early-warning indicators for rate-induced tipping. *Chaos*, **26**(9):093116, 2016.
- [39] C. Robinson. *Dynamical systems: stability, symbolic dynamics, and chaos*. CRC press, 1998.
- [40] J.C. Robinson. The asymptotic completeness of inertial manifolds. *Nonlinearity*, **9**:1325–1340, 1996.
- [41] A. Scheel. Radially symmetric patterns of reaction-diffusion systems. *MEM. AMER. MATH. SOC*, **165**(786), 2003.
- [42] M. Spivak. *Calculus on Manifolds: A Modern Approach to Classical Theorems of Advanced Calculus*. Advanced book program. Avalon Publishing, 1965.
- [43] P. Szmolyan and M. Wechselberger. Canards in \mathbb{R}^3 . *Journal of Differential Equations*, **177**(2):419–453, 2001.
- [44] T. Tao. A pseudoconformal compactification of the nonlinear schrödinger equation and applications. *New York J. Math.*, **15**:265–282, 2009.
- [45] H. R. Thieme. Asymptotically autonomous differential equations in the plane. *Rocky Mountain J. Math.*, **24**(1):351–380, 1993.
- [46] S. Wieczorek, P. Ashwin, C.M. Luke, and P.M. Cox. Excitability in ramped systems: the compost-bomb instability. *Proc. Roy. Soc. A*, **467**:1215–1242, 2011.
- [47] S. Wieczorek, P. Ashwin, and Ch. Xie. Rate-Induced Tipping: Regular Thresholds, Edge States and Testable Criteria. *in preparation*, 2021.
- [48] Chun Xie. *Rate-induced critical transitions*. PhD thesis, University College Cork, 2020.