

Title	Microbial succession and flavor production in the fermented dairy beverage kefir
Authors	Walsh, Aaron M.;Crispie, Fiona;Kilcawley, Kieran N.;O'Sullivan, Orla;O'Sullivan, Maurice G.;Claesson, Marcus J.;Cotter, Paul D.
Publication date	2016-10-04
Original Citation	Walsh, A. M., Crispie, F., Kilcawley, K., O'Sullivan, O., O'Sullivan, M. G., Claesson, M. J. and Cotter, P. D. (2016) 'Microbial succession and flavor production in the fermented dairy beverage kefir', mSystems 1(5), e00052-16 (16pp). doi:10.1128/mSystems.00052-16
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://msystems.asm.org/content/2/1/e00003-17 - 10.1128/mSystems.00052-16
Rights	© 2016, the Authors. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International license. - https://creativecommons.org/licenses/by/4.0/
Download date	2023-10-03 01:33:05
Item downloaded from	https://hdl.handle.net/10468/3850



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Table S3. Correlations between the relative abundances of microbial genera and the levels of volatile compounds

Genus	Compound(s)	R-value	Uncorrected p-value
Acetobacter	Acetic acid	0.76	<0.01
	2-methyl-1-butanol	0.65	0.01
	2,3-butanedione	0.67	<0.01
Kazachstania	Acetic acid	0.52	0.05
	2-methyl-1-butanol	0.53	0.04
	2,3-butanedione	0.85	<0.01
	2,3-pentanedione	0.68	<0.01
	2,3-hexanedione	0.72	<0.01
Lactobacillus	Carboxylic acids	0.6	0.02
	Esters	0.59	0.02
	3-methyl-1-butanol	0.58	0.02
Leuconostoc	2,3-butanedione	0.68	0.005
Saccharomyces	Carboxylic acids	0.71	<0.01
	Esters	0.78	<0.01