

Title	Detection and prevention of adverse drug reactions in multi-morbid older patients
Authors	Jennings, Emma L. M.;Gallagher, Paul F.;O'Mahony, Denis
Publication date	2018-09-12
Original Citation	Jennings, E., Gallagher, P. and O'Mahony, D. (2018) 'Detection and prevention of adverse drug reactions in multi-morbid older patients', Age and Ageing, 48(1), pp. 10-13. doi: 10.1093/ageing/afy157
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1093/ageing/afy157
Rights	© 2018, the Authors. Published by Oxford University Press on behalf of the British Geriatrics Society. All rights reserved. This is a pre-copyedited, author-produced PDF of an article accepted for publication in Age and Ageing following peer review. The version of record is available online at: https://doi.org/10.1093/ageing/afy157
Download date	2023-10-03 18:08:52
Item downloaded from	https://hdl.handle.net/10468/7480



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Table 1. Comparison of ADR Risk Prediction Models by Included Population Studied, Risk Predictive Variables and Area Under the Receiver Operator Curve

	<i>Studies as reported by Stevenson et al. [6] Systematic Review</i>				
	O'Mahony et al.[4] 2018 ADRROP	Tangiisuran et al. [7] 2014 BARDI Model	McElnay et al. [3] 1997	Onder et. al [13] 2010 GerontoNet	Trivalle et al. [8] 2011
Study Demographics	n = 2217, 467 ADRs (21%) Validation n = 530 Acute hospital (medical and surgical)	n = 690, 86 ADRs (12.5%) Validation n = 483 Acute hospital (elderly care and stroke wards)	n = 929; 149 ADEs (16%) Validation n = 204 Acute hospital (medical and surgical)	n = 5936, 383 ADRs (6.5%) Validation n = 483 Acute and community hospitals	n = 576, 152 ADRs (39%) Validation n = NR 16 geriatric rehabilitation centres
<i>Age (in years)</i>	>80% >70 years	100% ≥ 85 (81-89)*	100% ≥ 65; range 65-98	>90% ≥ 80; 78.0 ± 7.2**	100% ≥ 65, 83.6 ± 7.9**
<i>Polypharmacy</i>	>40% 6-10 medications	5 (3-7)*	4 (1-15) medications	>65% >5 medications	9 (6-12) medications*
<i>Multi-morbidity</i>	>75% 4+ co-morbidities	8 (6-10)*	-	>50% 4+ co-morbidities	-
Risk Predictive Variables	Female versus male Age > 70 eGFR < 30 ml/min BSA Needs assistance ≥ 1 ADL 4+ co-morbidities Liver Failure STOPP Category: 1 versus 0 STOPP PIMs ≥ 2 versus 0 STOPP PIMs ≥ 1 Fall in last year	≥ 8 drugs Hyperlipidaemia Raised WCC Use of anti-diabetic agents LOS ≥ 12 days	Prescribed antidepressants Prescribed digoxin Gastrointestinal problems Abnormal potassium level Patient belief drugs were responsible Experiences angina Experiences COAD	≥ 4 Comorbid conditions Heart Failure Liver Disease No. of Drugs – ≤ 5 – 5-7 – ≥ 8 Previous ADR Renal Failure	Number of medications – 0–6 – 7–9 – 10–12 – ≥ 13 Antipsychotic treatment Recent anticoagulant
Area Under Receiver Operator Curve (AUROC) Statistic	Development 0.63 [95% CI 0.60-0.67] Validation 0.59 [95% CI 0.53-0.65]	Internal Validation 0.74 [95% CI 0.68-0.79] External Validation 0.73 [95% CI 0.66-0.80]	No AUROC reported – Accuracy 63% – Sensitivity 40.5% – Specificity 69%	Development 0.71 [95% CI 0.68-0.73] Validation 0.70 [95% CI 0.63-0.78] External Validation O'Connor et. al [5] 0.62 [95% CI 0.57-0.68]	Overall 0.70 [95% CI 0.65-0.74]
*median (IQR); **mean ± standard deviation ADR – Adverse Drug Reaction (reported as number of patients); ADE – Adverse Drug Event; NR – Not recorded; WCC – White blood cell count; LOS – Length of stay; COAD Chronic Obstructive Airways Disease					