

Title	Nuclear membrane-localised NOX4D generates pro-survival ROS in FLT3-ITD-expressing AML
Authors	Moloney, Jennifer N.;Jayavelu, Ashok Kumar;Stanicka, Joanna;Roche, Sarah L.;O'Brien, Rebecca L.;Scholl, Sebastian;Böhmer, Frank D.;Cotter, Thomas G.
Publication date	2017
Original Citation	Moloney, J. N., Jayavelu, A. K., Stanicka, J., Roche, S. L., O'Brien, R. L., Scholl, S., Böhmer, F.-D. and Cotter, T. G. (2017) 'Nuclear membrane-localised NOX4D generates pro-survival ROS in FLT3-ITD-expressing AML', <i>Oncotarget</i> , 8(62), pp. 105440-105457. doi: 10.18632/oncotarget.22241
Type of publication	Article (peer-reviewed)
Link to publisher's version	<a href="http://www.oncotarget.com/index.php?journal=oncotarget&amp;page=article&amp;op=view&amp;path[]=22241&amp;path[]=70409-10.18632/oncotarget.22241">http://www.oncotarget.com/index.php?journal=oncotarget&amp;page=article&amp;op=view&amp;path[]=22241&amp;path[]=70409-10.18632/oncotarget.22241</a>
Rights	© 2017, Moloney et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License 3.0 (CC BY 3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. - <a href="https://creativecommons.org/licenses/by/3.0/">https://creativecommons.org/licenses/by/3.0/</a>
Download date	2024-12-04 04:00:32
Item downloaded from	<a href="https://hdl.handle.net/10468/6339">https://hdl.handle.net/10468/6339</a>




# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

# Nuclear membrane-localised NOX4D generates pro-survival ROS in FLT3-ITD-expressing AML

## SUPPLEMENTARY MATERIALS



**Supplementary Figure 1: NOX4 antibody specificity.** Abcam NOX4 antibody (Ab109225) specificity. Western blot analysis of NOX4 67 kDa protein levels in 32D/FLT3-ITD whole cell lysates at 48 h following NOX4 siRNA and shRNA transfection (**A**).  $\beta$ -actin was used as a loading control. Novus Biologicals NOX4 antibody (NB110-58849) specificity. HEK293-T cells were transfected with EV-HA or pCMV3-C-HA encoding NOX4 using Calcium Phosphate. Western blot analysis of NOX4 67 kDa and NOX4D 28 kDa protein expression in HEK 293-T whole cell lysates following transfection with EV-HA or NOX4-HA for 48 h (**B**). Total protein was used as a loading control. Blots are representative of three independent experiments.