

Title	Dietary energy density: estimates, trends and dietary determinants for a nationally representative sample of the Irish population (aged 5-90 years)
Authors	O'Connor, Laura;Walton, Janette;Flynn, Albert
Publication date	2014-11-17
Original Citation	O'Connor, L., Walton, J. and Flynn, A. (2015) 'Dietary energy density: estimates, trends and dietary determinants for a nationally representative sample of the Irish population (aged 5-90 years)', <i>British Journal of Nutrition</i> , 113(1), pp. 172-180. doi:10.1017/S0007114514003420
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1017/S0007114514003420
Rights	© The Authors 2014. Published by Cambridge University Press (CUP) on behalf of The Nutrition Society.
Download date	2025-03-28 08:24:27
Item downloaded from	https://hdl.handle.net/10468/3719



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh



Dietary energy density: estimates, trends and dietary determinants for a nationally representative sample of the Irish population (aged 5–90 years)

Laura O'Connor^{1,2*}, Janette Walton¹ and Albert Flynn¹

¹School of Food and Nutritional Science, University College Cork, Cork, Republic of Ireland

²MRC Epidemiology Unit, School of Clinical Medicine, Institute of Metabolic Sciences, University of Cambridge, Box 285, Cambridge Biomedical Campus, Cambridge CB2 0QQ, UK

(Submitted 10 June 2014 – Final revision received 3 September 2014 – Accepted 1 October 2014 – First published online 17 November 2014)

Abstract

Higher dietary energy density (DED) has been reported to be associated with weight gain, obesity and poorer dietary quality, yet nationally representative estimates that would allow tracking of secular trends and inter-country comparisons are limited. The aims of the present study were to calculate DED estimates for the Irish population and to identify dietary determinants of DED. Weighed/semi-weighed food records from three cross-sectional surveys (the National Children's Food Survey, the National Teens' Food Survey and the National Adult Nutrition Survey) were collated to estimate habitual dietary intakes for a nationally representative sample of the Irish population, aged 5–90 years (n 2535). DED estimates, calculated using the total diet method, the food only method and a novel method, including foods and solids in beverages, were 3·70 (SD 1·09), 7·58 (SD 1·72) and 8·40 (SD 1·88) kJ/g, respectively. Determinants of DED did not vary by the calculation method used. Variation in the intakes of fruit, vegetables and sugar-sweetened beverages (SSB) across consumer groups contributed to the largest variance in DED estimates, followed by variation in the intakes of potatoes, fresh meat, bread, chips, ready-to-eat breakfast cereals, and confectionery. DED estimates were inversely associated with age group and consistently lower for females than for males. The inverse association of DED with age group was explained by higher intakes of vegetables, fruit, fish, potatoes, fresh meat and brown bread and lower intakes of SSB, chocolate confectionery, ready-to-eat breakfast cereals and savoury snacks in older age groups. Females consumed, on average, 1·5 times more fruit and vegetables combined when compared with males, largely explaining the sex differences in DED estimates. Current DED estimates for adults were similar to those calculated in a previous survey, carried out 10 years earlier. These estimates and determinants serve as a baseline for comparison for other works and public health campaigns.

Key words: Energy density: Dietary determinants: Nationally representative estimates: Trends: Children: Adolescents: Adults

A solid body of evidence exists from experimental investigations and intervention studies for the association of dietary energy density (DED) with weight gain, increased energy intake (EI) and obesity^(1–3). Body weight status⁽⁴⁾, BMI⁽⁵⁾, waist circumference⁽⁶⁾, obesity and the metabolic syndrome⁽⁷⁾ have all been found to be positively associated with DED in cross-sectional studies, providing evidence that the observations extend to free-living populations. A relationship between DED and the nutritional quality of the diet has also been highlighted; lower DED has been reported to be associated with higher dietary quality in Irish children and teenagers, American adults, Spanish adults and elderly Spanish adults^(8–11) and with intakes of foods and nutrients related to better dietary quality in Mediterranean, Australian and American adults^(12–15). DED has also been proposed

as a potential proxy for dietary quality⁽¹⁶⁾, and low DED has been reported to be associated with a trend towards a healthier lifestyle, including lower alcohol consumption, smoking and physical inactivity, in a representative sample of Spanish adults⁽¹⁰⁾.

To date, there have been no estimations of DED in an Irish population. As DED is associated with dietary quality and with obesity, it is worth estimating DED values for the population. This will allow tracking of secular trends and enable comparison with those of other countries. It is also worth distinguishing the dietary components that influence DED in free-living populations to explain differences in DED estimates and to ascertain potential targets for public health messages to lower DED estimates. The National Children's Food Survey (NCFS), the National Teens' Food Survey (NTFS) and the

Abbreviations: %TE, percentage of total energy; DED, dietary energy density; EI, energy intake; NANS, National Adult Nutrition Survey; NCFS, National Children's Food Survey; NSIFCS, North/South Ireland Food Consumption Survey; NTFS, National Teens' Food Survey; RTEBC, ready-to-eat breakfast cereals; SSB, sugar-sweetened beverages; UR, under-reporters for energy.

* **Corresponding author:** L. O'Connor, email laura.oconnor@mrc-epid.cam.ac.uk



National Adult Nutrition Survey (NANS) together form a nationally representative sample of the Irish population aged 5–90 years that provides fully disaggregated habitual food consumption data at the level of the individual from which calculation of such estimates is possible.

The aims of the present study were to calculate and present DED estimates for a nationally representative sample of the Irish population and to identify dietary determinants of DED in a free-living population.

Methods

Study sample and dietary intake assessment

The NCFS (2003–4), the NTFS (2005–6) and the NANS (2008–10) were carried out to establish databases of habitual food and drink consumption in representative samples of Irish children, adolescents and adults aged 5–12, 13–17 and 18–90 years, respectively. A 7 d weighed food record was used to collect dietary intake data from 594 children (293 boys and 301 girls), a 7 d semi-weighed food record was used to collect dietary intake data from 441 adolescents (224 boys and 217 girls), and a 4 d semi-weighed food record was used to collect dietary intake data from 1500 adults (760 females and 740 males). Analyses of dietary intake data were carried out using WISP[®] (Tinuviel Software), which contains McCance and Widdowson's *The Composition of Foods*, 6th edition⁽¹⁷⁾, 5th edition⁽¹⁸⁾ plus supplemental volumes^(19–28) and the Irish Food Composition Database⁽²⁹⁾. Ethical approval was obtained from St James' Hospital and Federated Dublin Voluntary Hospitals Joint Research Ethics Committee for the NCFS and the University College Cork Clinical Research Ethics Committee of the Cork Teaching Hospitals for the NTFS and the NANS. Written consent was obtained from the participants or from the parents/guardians for participants aged <18 years. A full description of methodologies is given elsewhere (<http://www.iuna.net>).

Estimation of dietary energy density

DED was estimated as kJ/g and was calculated using three different methods by taking the following into account: (1) energy and weight of the total diet; (2) energy and weight including only foods and excluding all beverages (i.e. all solid foods and liquid-like foods such as yogurts and soups included and all beverages excluded); (3) energy and weight of all foods and solids in beverages (values for solids in beverages were calculated by removing the water portion of each food code).

Estimation of dietary energy density accounting for under-reporters for energy

DED was estimated both including and excluding under-reporters for energy (UR). For each child and teenager, BMR was predicted from standard equations using body weight and height⁽³⁰⁾. Minimum EI cut-off points, calculated as multiples of BMR⁽³¹⁾, were used to identify UR. In the study

sample, 32% of children and 64% of teenagers were classified as UR.

In adults, body composition was measured using the Tanita body composition analyser BC-420MA, which calculated BMR through multiple regression analysis using fat-free mass values (Tanita Limited). EI is expressed as a ratio of BMR (EI:BMR). The cut-off limits proposed by Goldberg *et al.*⁽³²⁾ to classify individuals as UR were used. The cut-off point relevant to the NANS was 1.10; 30% of the participants had an EI:BMR_{est} below this value.

Statistical analyses

Statistical analyses were conducted using SPSS[®] version 15.0 for Windows[™] (SPSS, Inc.). DED was estimated and expressed as mean, median and standard deviation by sex and age group. An independent-samples *t* test and a two-way between-groups ANOVA were conducted to evaluate differences in DED estimates by sex and age group.

The dietary determinants of DED were identified by categorising the participants by tertile of energy, macronutrient and food group intakes into low-, medium- and high- or non-, low- and high-consumer groups of each. The participants were categorised into consumers and non-consumers for some less consumed food groups. A one-way ANOVA or an independent-samples *t* test was used to test for significant differences ($P < 0.05$) in means across consumer groups. Significant differences between the groups were determined using ANOVA followed by Tukey's honestly significant difference or Hochberg's GT2 *post hoc* test where appropriate.

Post hoc tests were chosen depending on similarity between sample sizes and homogeneity of variance between the groups (as determined by Levene's test). The effect size of the significant differences between consumer groups was calculated as follows: $\eta^2 = \text{sum of squares between groups} / \text{total sum of squares}$ (values ≥ 0.01 indicate a small effect, ≥ 0.06 a medium effect and ≥ 0.14 a large effect, as classified by Cohen⁽³³⁾).

Food intake patterns that may explain the difference in DED estimates between sexes and with age were estimated by examining food group intakes across age groups in males and females separately. For this purpose, the participants were categorised into three age groups: 5–18; 19–50; 51–90 years. ANOVA was used to test for significant differences ($P < 0.05$) in mean intakes across age groups. Significant differences between the groups were determined using ANOVA followed by either Tukey's honestly significant difference or Hochberg's GT2 *post hoc* test. A one-way between-groups ANCOVA was used to test for significant differences between sexes adjusted for age.

Secular trends and generational effects

The DED estimates of a subsample from the NANS (those aged 18–64 years, $n = 1274$) were compared with estimates calculated for the participants of the North/South Ireland Food Consumption Survey (NSIFCS), 2001 ($n = 1379$). The NSIFCS was carried out 10 years before the NANS and included a

representative sample of adults aged 18–64 years living in Ireland. Dietary intake data were collected using a 7 d food record. A full description of the survey design and methodology of the NSIFCS has been published previously⁽³⁴⁾. Differences in DED estimates calculated for the NANS and NSIFCS were examined using a one-way between-groups ANCOVA. This comparison also allowed to assess the possibility of a generational effect on trends observed between DED and age groups in the cross-sectional analysis.

Results

For each of the three methods, DED estimates were inversely associated with age group (P linear trend <0.001) and females had lower mean DED estimates than males ($P < 0.001$) (Table 1). DED estimates were lower for females in all age groups ($P < 0.001$).

Mean DED estimates calculated excluding UR (see online supplementary Table S1) were similar to those calculated for the total population and trended similarly by sex and across

age groups. Exclusion of UR resulted in estimates similar to those calculated for the total population regardless of the calculation method used.

Determinants of DED did not vary by the calculation method. Determinants are thus only presented for DED calculated by including energy and weight of all foods and solids in beverages.

Higher mean DED estimates were found to be associated with higher intake of energy (kJ) and percentage of total energy (%TE) from fat, carbohydrates and total sugar ($P < 0.001$) and lower intakes of protein and dietary fibre ($P < 0.001$) (Table 2). Higher consumption or being a consumer of certain foods was found to be both positively and inversely associated with DED. DED estimates were significantly lower in those with higher intakes (g/10 MJ) of vegetables, fruit, potatoes (boiled, baked or mashed), fresh meat, brown bread, fish, vegetable and pulse dishes, eggs, pulses, soup, cooked breakfast cereals and nuts ($P < 0.001$). Mean DED estimates were significantly higher in those with higher intakes (g/10 MJ) of white bread, chips, ready-to-eat

Table 1. Dietary energy density (DED; kJ/g) estimates for the Irish population, aged 5–90 years, by age and sex (n 2535) (Number of participants and medians; mean values and standard deviations)

Sex	Age group	n	DED (kJ/g) total diet			DED (kJ/g) food only			DED (kJ/g) including foods and solids in beverages		
			Mean	Median	SD	Mean	Median	SD	Mean	Median	SD
Males	5–90	1257	3.85	3.84	1.05	7.86	7.84	1.62	8.75	8.76	1.77
	5–10	218	4.59	4.59	0.71	8.31	8.17	1.60	9.33	9.18	1.63
	11–15	212	4.53	4.52	0.84	8.44	8.33	1.51	9.39	9.37	1.59
	16–20	139	4.05	3.94	0.98	8.20	8.27	1.44	9.41	9.46	1.48
	21–25	72	3.39	3.22	0.92	8.33	8.47	1.61	9.52	9.79	1.91
	26–30	86	3.19	3.01	0.98	7.89	7.94	1.41	8.83	9.03	1.58
	31–35	66	3.39	3.24	1.06	7.71	7.55	1.55	8.50	8.27	1.69
	36–40	63	3.37	3.19	0.97	7.84	7.78	1.68	8.65	8.56	1.81
	41–45	66	3.53	3.48	1.04	7.76	7.65	1.33	8.48	8.39	1.47
	46–50	76	3.49	3.42	0.83	7.40	7.33	1.53	8.05	7.96	1.65
	51–55	64	3.26	3.22	0.83	7.29	7.04	1.30	8.05	7.87	1.31
	56–60	49	3.42	3.35	0.84	6.97	6.96	1.28	7.66	7.76	1.26
	61–65	46	3.13	2.96	0.90	6.71	6.83	1.55	7.28	7.40	1.54
66–70	37	3.66	3.65	0.99	6.93	6.91	1.72	7.44	7.06	1.77	
71–75	31	3.34	3.33	0.87	6.55	6.10	1.82	7.13	6.96	1.85	
> 75	32	3.37	3.40	0.88	6.52	6.48	1.43	7.01	6.88	1.45	
$P_{\text{trend}}\dagger$			<0.001			<0.001			<0.001		
Females	5–90	1278	3.55*	3.49	1.11	7.31*	7.33	1.77	8.07*	8.17	1.93
	5–10	228	4.56	4.54	0.69	8.11	8.08	1.29	9.07	9.04	1.37
	11–15	211	4.31	4.29	0.76	8.30	8.20	1.38	9.20	9.08	1.30
	16–20	115	3.73	3.84	0.96	7.88	7.79	1.64	8.81	8.83	1.73
	21–25	76	3.33	3.21	1.22	7.74	7.68	1.93	8.60	8.57	2.12
	26–30	76	3.13	3.04	1.10	7.36	7.36	1.59	8.09	8.25	1.80
	31–35	67	3.13	2.87	1.04	7.44	7.40	1.76	8.12	8.16	1.83
	36–40	59	2.95	3.02	0.75	7.16	7.12	1.32	7.86	7.73	1.54
	41–45	87	2.87	2.80	0.83	7.04	6.99	1.73	7.67	7.74	1.81
	46–50	86	2.95	2.86	0.89	6.55	6.56	1.65	7.18	7.21	1.75
	51–55	63	2.90	2.76	0.97	6.22	5.96	1.58	6.72	6.59	1.65
	56–60	48	2.93	2.82	0.82	5.80	5.84	1.47	6.24	6.19	1.58
	61–65	52	2.65	2.63	0.73	5.50	5.24	1.42	5.93	5.65	1.47
66–70	43	2.79	2.81	0.73	5.67	5.67	1.36	6.23	6.25	1.53	
71–75	32	2.96	2.73	1.08	6.07	6.03	1.58	6.50	6.43	1.66	
> 75	35	2.99	2.85	0.70	5.82	5.77	1.35	6.42	6.43	1.56	
P_{trend}			<0.001			<0.001			<0.001		

* Mean value was significantly different from that of males as calculated using an independent-samples t test ($P < 0.05$).

† P linear trend across age groups as calculated using two-way ANOVA.

Table 2. Dietary energy density (DED) estimates by tertile of mean daily energy intake (kJ), macronutrient intakes (percentage of total energy, %TE), and dietary fibre and food group intakes (g/10 MJ/d) in the Irish population (*n* 2535)

(Number of participants, mean values and standard deviations)

	DED (kJ/g) including foods and solids in beverages									<i>P</i> *	η^2		Mean intake
	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD				
	Low consumer			Medium consumer			High consumer						
Energy	845	7.96 ^a	1.97	845	8.47 ^b	1.89	845	8.87 ^c	1.69	0.000	0.04	↑	8094 kJ
Protein	845	9.47 ^a	1.71	845	8.53 ^b	1.60	845	7.30 ^c	1.70	0.000	0.22	↓	15.85 %TE
Fat	845	7.94 ^a	1.96	845	8.52 ^b	1.77	845	8.83 ^c	1.83	0.000	0.04	↑	34.67 %TE
Carbohydrates	845	8.30 ^a	1.89	845	8.34 ^a	1.87	845	8.65 ^b	1.89	0.000	0.01	↑	46.06 %TE
Total sugar	845	8.32 ^a	1.72	845	8.23 ^a	1.89	845	8.74 ^b	2.01	0.000	0.01	↑	19.12 %TE
Dietary fibre	845	9.76 ^a	1.49	845	8.61 ^b	1.41	845	6.93 ^c	1.57	0.000	0.38	↓	21.22 g/10 MJ
	Low consumer			Medium consumer			High consumer						
Vegetables (excluding pulses and composite meals)	845	9.67 ^a	1.63	845	8.57 ^b	1.48	845	7.05 ^c	1.56	0.000	0.32	↓	79.3 g/10 MJ
Fruit	845	9.37 ^a	1.79	845	8.64 ^b	1.62	845	7.29 ^c	1.64	0.000	0.21	↓	109.8 g/10 MJ
Potatoes (boiled, baked or mashed)	845	9.08 ^a	1.98	845	8.56 ^b	1.72	845	7.66 ^c	1.68	0.000	0.10	↓	88.4 g/10 MJ
White bread and rolls	845	7.64 ^a	1.93	845	8.74 ^b	1.85	845	8.92 ^b	1.62	0.000	0.09	↑	70.0 g/10 MJ
Chipped, fried and roasted potatoes	845	7.72 ^a	1.96	845	8.56 ^b	1.77	845	9.02 ^c	1.70	0.000	0.08	↑	54.2 g/10 MJ
Fresh meat and meat dishes	845	9.04 ^a	2.10	845	8.48 ^b	1.67	845	7.77 ^c	1.66	0.000	0.07	↓	133.1 g/10 MJ
Ready-to-eat breakfast cereals (including milk)	845	7.89 ^a	1.99	845	8.33 ^b	1.79	845	9.08 ^c	1.68	0.000	0.07	↑	224.4 g/10 MJ
Processed meat	845	7.96 ^a	1.99	845	8.56 ^b	1.75	845	8.77 ^b	1.84	0.000	0.03	↑	64.7 g/10 MJ
Biscuits (including crackers)	845	8.30 ^a	1.93	845	8.53 ^b	1.85	845	8.46 ^{a,b}	1.89	0.040	0.00	↑	16.1 g/10 MJ
Butters, spreading fats and oils	845	8.43 ^a	1.91	845	8.54 ^{a,b}	1.88	845	8.31 ^{a,c}	1.88	0.045	0.00	↔	15.0 g/10 MJ
Sugars, syrups, preserves and sweeteners	845	8.41	1.91	845	8.54	1.82	845	8.34	1.93	0.097	0.00	–	11.2 g/10 MJ
	Non-consumer			Low consumer			High consumer						
Sugar-sweetened beverages	979	7.40 ^a	1.77	777	8.81 ^b	1.60	779	9.34 ^c	1.68	0.000	0.20	↑	319.4 g/10 MJ
Savoury snacks (e.g. crisps and popcorn)	1162	7.71 ^a	1.90	686	8.84 ^b	1.66	687	9.24 ^c	1.62	0.000	0.13	↑	11.0 g/10 MJ
Brown bread	995	9.07 ^a	1.76	770	8.51 ^b	1.81	770	7.53 ^c	1.78	0.000	0.11	↓	46.6 g/10 MJ
Fish and fish dishes	1287	8.84 ^a	1.86	624	8.66 ^a	1.63	624	7.37 ^b	1.80	0.000	0.11	↓	41.8 g/10 MJ
Chocolate confectionery	953	7.77 ^a	1.90	791	8.50 ^b	1.73	791	9.16 ^c	1.76	0.000	0.09	↑	16.2 g/10 MJ
Non-chocolate confectionery	1448	8.05 ^a	1.92	543	8.87 ^b	1.69	544	9.02 ^b	1.74	0.000	0.06	↑	9.5 g/10 MJ
Vegetable and pulse dishes	1644	8.66 ^a	1.92	445	8.49 ^a	1.63	446	7.54 ^b	1.78	0.000	0.05	↓	16.7 g/10 MJ
Savouries (e.g. pasta dishes and pizzas)	1116	7.98 ^a	2.01	709	8.76 ^b	1.79	710	8.81 ^b	1.63	0.000	0.04	↑	39.6 g/10 MJ
Eggs and egg dishes	1340	8.56 ^a	1.90	597	8.77 ^a	1.79	598	7.80 ^b	1.82	0.000	0.04	↓	26.4 g/10 MJ
Pulses (excluding composite dishes)	1134	8.61 ^a	2.03	700	8.68 ^a	1.68	701	7.90 ^b	1.76	0.000	0.03	↓	19.3 g/10 MJ
Rice, pasta and noodles	1193	8.52 ^a	2.05	671	8.75 ^b	1.63	671	7.95 ^c	1.75	0.000	0.03	↔	39.1 g/10 MJ
Alcoholic beverages	1657	8.57 ^a	1.88	438	7.78 ^b	1.90	440	8.54 ^a	1.79	0.000	0.02	↔	542.6 g/10 MJ
Creams, ice creams and chilled desserts	1110	8.36 ^a	2.00	712	8.73 ^b	1.75	713	8.24 ^a	1.82	0.000	0.01	↔	28.2 g/10 MJ
Yogurts	1292	8.45 ^a	2.00	621	8.67 ^b	1.81	622	8.14 ^c	1.70	0.000	0.01	↔	42.4 g/10 MJ
Cakes, pastries and buns	1302	8.39 ^a	1.96	616	8.70 ^b	1.75	617	8.26 ^a	1.85	0.000	0.01	↔	17.4 g/10 MJ
Cheeses	948	8.41	2.04	793	8.52	1.76	794	8.36	1.84	0.244	0.00	–	14.2 g/10 MJ
Juices and smoothies	1247	8.34	2.02	644	8.49	1.83	644	8.54	1.68	0.060	0.00	–	197.0 g/10 MJ
	Non-consumer			Consumer									
Soups	1760	8.66	1.89	775	7.91	1.80				0.000	0.03	↓	32.3 g/10 MJ
Cooked breakfast cereals (e.g. porridge)	1953	8.54	1.90	582	8.07	1.82				0.000	0.01	↓	37.0 g/10 MJ
Potato products	2015	8.34	1.94	520	8.80	1.64				0.000	0.01	↑	7.0 g/10 MJ
Nuts	2258	8.48	1.89	277	8.02	1.84				0.000	0.01	↓	1.5 g/10 MJ

↑ and ↓, the direction of change in DED as the dietary intakes increase from low to high; ↔, a non-directional association of DED with dietary intakes; –, no significant difference in DED with change in dietary intake.

^{a,b,c} Mean values with unlike superscript letters were significantly different between the consumer groups (*P* < 0.05).

* As calculated using a one-way ANOVA with Tukey's honestly significant difference or Hochberg's GT2 *post hoc*, or an independent-samples *t* test.

Table 3. Dietary energy density (DED) estimates and mean daily energy intake (kJ), macronutrient intakes (percentage of total energy, %TE), and dietary fibre and food group intakes (g/10 MJ) in the Irish population by age group split by sex (*n* 2535)

(Mean values and standard deviations)

	Males							Females								
	5–18 years (<i>n</i> 534)		19–50 years (<i>n</i> 464)		51–90 years (<i>n</i> 259)		<i>P</i> *	5–18 years (<i>n</i> 527)		19–50 years (<i>n</i> 478)		51–90 years (<i>n</i> 273)		<i>P</i> *		
	Mean	SD	Mean	SD	Mean	SD		Mean	SD	Mean	SD	Mean	SD			
DED (kJ/g) including foods and solids in beverages	9.38 ^a	1.60	8.71 ^b	1.71	7.51 ^c	1.53	0.000	↓	9.07 ^a	1.40	7.95 ^b	1.89	6.34 ^c	1.58	0.000	↓
Energy (kJ)	8386 ^a	2303	10266 ^b	2777	8911 ^c	2563	0.000	↔	6864 ^a	1586	7302 ^b	2094	6819 ^a	1701	0.000	↔
Protein (%TE)	14.3 ^a	2.5	17.1 ^b	4.0	17.4 ^b	3.6	0.000	↑	13.9 ^a	2.3	16.4 ^b	3.7	17.9 ^c	3.4	0.000	↑
Fat (%TE)	34.3	4.9	34.2	6.1	34.7	7.1	0.618	–	35.0	4.4	35.0	6.0	34.8	6.1	0.858	–
Carbohydrates (%TE)	50.6 ^a	5.7	41.6 ^b	7.3	42.2 ^b	7.2	0.000	↓	50.5 ^a	4.8	43.6 ^b	6.4	44.4 ^b	6.3	0.000	↓
Total sugar (%TE)	22.3 ^a	5.7	15.6 ^b	5.6	16.2 ^b	6.3	0.000	↓	22.3 ^a	5.3	17.3 ^b	5.5	18.7 ^c	5.6	0.000	↑
Dietary fibre (g/10 MJ)	18.2 ^a	5.0	20.9 ^b	7.4	23.2 ^c	7.4	0.000	↑	18.2 ^a	4.6	23.3 ^b	7.5	28.1 ^c	8.6	0.000	↑
Vegetables (excluding pulses and composite meals) (g/10 MJ)	41.1 ^a	42.7	69.5 ^b	64.8	104.7 ^c	86.8	0.000	↑	46.8 ^a	49.8	113.5 ^b	101.1	149.9 ^c	109.8	0.000	↑
Fruit (g/10 MJ)	89.9 ^a	136.7	83.4 ^a	95.8	114.7 ^b	128.9	0.000	↑	103.0 ^a	103.5	117.2 ^a	135.3	189.3 ^b	171.3	0.000	↑
Fish and fish dishes (g/10 MJ)	12.9 ^a	22.2	41.2 ^b	69.4	88.6 ^c	109.7	0.000	↑	13.5 ^a	24.9	58.6 ^b	89.4	80.2 ^c	98.6	0.000	↑
Potatoes (boiled, baked or mashed) (g/10 MJ)	80.8 ^a	72.0	80.5 ^a	83.0	134.4 ^b	106.9	0.000	↑	78.1 ^a	70.8	73.7 ^a	79.9	118.4 ^b	90.6	0.000	↑
Meat and meat dishes (g/10 MJ)	112.0 ^a	71.7	157.3 ^b	110.3	144.0 ^b	95.1	0.000	↑	110.8 ^a	71.5	136.8 ^b	101.1	159.5 ^c	107.9	0.000	↑
Brown bread (g/10 MJ)	19.5 ^a	38.6	55.3 ^b	58.8	79.5 ^c	80.0	0.000	↑	19.9 ^a	32.9	60.3 ^b	61.5	81.0 ^c	66.3	0.000	↑
Sugar-sweetened beverages (g/10 MJ)	280.5 ^a	226.7	167.8 ^b	284.2	58.2 ^c	174.0	0.000	↓	271.2 ^a	253.0	188.6 ^b	364.2	77.7 ^c	287.6	0.000	↓
Chocolate confectionery (g/10 MJ)	22.8 ^a	22.5	11.1 ^b	16.5	5.1 ^c	10.5	0.000	↓	25.4 ^a	23.1	14.6 ^b	19.9	7.8 ^c	15.9	0.000	↓
Ready-to-eat breakfast cereals (including milk) (g/10 MJ)	415.3 ^a	252.3	127.2 ^b	130.1	84.0 ^c	100.2	0.000	↓	352.4 ^a	250.5	108.2 ^b	127.7	105.3 ^b	124.9	0.000	↓
Savoury snacks (e.g. crisps and popcorn) (g/10 MJ)	14.2 ^a	15.1	8.1 ^b	13.6	1.8 ^c	8.2	0.000	↓	20.1 ^a	18.1	10.4 ^b	15.3	2.1 ^c	6.1	0.000	↓
Chipped, fried and roasted potatoes (g/10 MJ)	65.2 ^a	53.6	52.2 ^b	54.0	37.5 ^c	47.9	0.000	↓	67.7 ^a	56.5	48.9 ^b	56.2	35.4 ^c	46.9	0.000	↓
White bread (g/10 MJ)	87.6 ^a	50.6	62.6 ^b	56.9	69.0 ^b	72.4	0.000	↓	80.8 ^a	47.6	57.9 ^b	56.9	49.1 ^b	62.6	0.000	↓

↑ and ↓, the direction of change in DED as the nutrient or food group intakes increase from low to high; ↔, a non-directional association of DED with nutrient or food group intakes; –, no significant difference in DED with change in dietary intake.

^{a,b,c} Mean values with unlike superscript letters were significantly between the age groups within a sex (*P* < 0.05).

* As calculated using a one-way ANOVA.

breakfast cereals (RTEBC), processed meat, sugar-sweetened beverages (SSB), savoury snacks, chocolate and non-chocolate confectionery, savouries, alcohol and potato products ($P < 0.001$). Variation in the intakes of vegetables, fruit and SSB (g/10MJ) was found to be associated with the largest differences in mean DED estimates between consumer groups (η^2 : vegetables 0.32, fruit 0.21 and SSB 0.20). Differences in the intakes (g/10MJ) of potatoes, fresh meat, brown bread, bananas, fish, white bread, chips, RTEBC, and chocolate and non-chocolate confectionery also contributed to the differences in DED estimates, but to a lesser extent (η^2 : 0.06–0.13).

To explain the lower DED estimates obtained for females and the inverse association of DED with age, energy-adjusted intake patterns of macronutrients and dietary fibre and of the food groups that contributed to the largest variation in mean DED estimates were examined across age groups, split by sex (Table 3), and between sexes, adjusted for age (Table 4). Higher intakes of protein, total sugar (%TE) and dietary fibre (g/10MJ) and lower intakes of carbohydrates (%TE) were found to be associated with older age groups ($P < 0.001$). The intakes (g/10MJ) of vegetables, fruit, fish, potatoes, fresh meat and brown bread were significantly higher ($P < 0.001$) in older age groups in males and females, while the intakes (g/10MJ) of SSB, chocolate and non-chocolate confectionery, RTEBC, savoury snacks, chips and white bread were significantly lower ($P < 0.001$). The intakes of fat ($P = 0.007$), carbohydrates, total sugar (%TE) and dietary fibre (g/10MJ) ($P < 0.001$) were higher in females than in males, while those of energy (kJ) ($P < 0.001$) and protein (%TE) ($P = 0.003$) were lower. The intakes of vegetables, fruit, chocolate confectionery and savoury snacks were significantly higher in females than in males ($P < 0.001$) and those of

potatoes ($P = 0.020$), RTEBC ($P = 0.001$) and white bread ($P < 0.001$) were significantly higher in males than in females.

The DED (calculated including energy and weight of all foods and solids in beverages) estimates of the NSIFCS participants were also inversely associated with age group ($P < 0.001$) and females had lower DED estimates than males ($P < 0.001$). Mean DED was 8.50 (SD 1.55) kJ/g for males and 7.10 (SD 1.59) kJ/g for females. Comparison of DED estimates of the sample from the NSIFCS and the comparable subsample from the NANS, split by age and sex, is shown in Fig. 1. The mean DED estimate obtained for women, adjusted for age, in the NSIFCS did not differ significantly from that obtained for the subsample in the NANS ($P = 0.427$). However, DED estimates obtained for men in the NSIFCS were significantly lower than those obtained for men in the NANS ($P < 0.001$); over the 10 years, differences were apparent in only two age groups: 41–45 ($P = 0.015$) and 56–60 ($P = 0.019$) years.

Discussion

In the nationally representative free-living Irish population aged 5–90 years included in the present study, DED estimates were higher for males than for females and were inversely associated with age group. Little difference was observed in DED estimates calculated for adults between the two surveys carried out 10 years apart. The intakes of fruit, vegetables and SSB were the major dietary determinants of DED in this population.

Many methods have been proposed for the calculation of DED and there is little consensus as to which is the most appropriate method for free-living populations^(35–37). Part of the difficulty lies in the high dependency of DED on the amount of water in the diet. The treatment of beverages is

Table 4. Dietary energy density (DED) estimates and mean daily energy intake (kJ), macronutrient intake (percentage of total energy, %TE), and dietary fibre and food group intakes (g/10MJ) in the Irish population by sex adjusted for age (n 2535)

(Mean values with their standard errors)

	Males		Females		P*
	Mean	SE	Mean	SE	
DED (kJ/g) including foods and solids in beverages	8.73	0.05	8.09	0.05	0.000
Energy (kJ)	9190	64	7016	64	0.000
Protein (%TE)	16.0	0.1	15.7	0.1	0.003
Fat (%TE)	34.4	0.2	35.0	0.2	0.007
Carbohydrates (%TE)	45.5	0.2	46.6	0.2	0.000
Total sugar (%TE)	18.5	0.2	19.7	0.2	0.000
Dietary fibre (g/10 MJ)	20.3	0.2	22.1	0.2	0.000
Vegetables (excluding pulses and composite meals) (g/10 MJ)	65.4	2.1	93.0	2.1	0.000
Fruit (g/10 MJ)	93.0	3.6	126.3	3.6	0.000
Fish and fish dishes (g/10 MJ)	39.6	2.0	44.0	2.0	0.106
Potatoes (boiled, baked or mashed) (g/10 MJ)	92.2	2.3	84.7	2.3	0.020
Meat and meat dishes (g/10 MJ)	135.7	2.6	130.5	2.6	0.159
Brown bread (g/10 MJ)	45.6	1.5	47.5	1.5	0.376
Sugar-sweetened beverages (g/10 MJ)	191.1	7.7	200.9	7.6	0.363
Chocolate confectionery (g/10 MJ)	14.7	0.6	17.7	0.5	0.000
Ready-to-eat breakfast cereals (including milk) (g/10 MJ)	238.1	5.7	210.8	5.6	0.001
Savoury snacks (e.g. crisps and popcorn) (g/10 MJ)	9.3	0.4	12.8	0.4	0.000
Chipped, fried and roasted potatoes (g/10 MJ)	54.4	1.5	54.0	1.5	0.841
White bread (g/10 MJ)	74.3	1.6	65.7	1.6	0.000

* As calculated using a one-way ANCOVA.




Fig. 1. Comparison of dietary energy density (DED) estimates (kJ/g) calculated including solids in beverages for the Irish adult population aged 18–64 years, by age and sex ((a) females and (b) males) in the North/South Ireland Food Consumption Survey (—) and the National Adult Nutrition Survey (---). Error bars: ± 2 sd.

therefore important, as the inclusion of beverages has a lowering effect on the resulting DED estimates⁽³⁷⁾. Variation in the associations of dietary and anthropometric factors with DED, depending on the method of computation, has been acknowledged^(9,14,35). Johnson *et al.*⁽³⁸⁾ proposed calculating DED by excluding drinks and including energy contributed by drinks as a covariate in analyses as the most reliable method for testing the relationship between DED and weight gain in free-living humans. The energy derived from beverages may be an important factor in the causation of obesity, in particular, from SSB, which have been reported to be independently associated with obesity⁽³⁹⁾. In the present study, nationally representative estimates of DED calculated using three different calculation methods were evaluated. Although the DED estimates differed, the determinants and age–sex trends did not. As such, the primary analysis was focused on DED estimates calculated using a novel method: including energy and weight of all foods and solids in beverages. This approach aims to remove the bias that including beverages can introduce in DED estimates while taking into account the contribution of beverages to EI and as such has potential as a standardised method for further publications.

There was little difference in mean DED estimates on including and excluding UR. Trends in DED estimates across sexes and age groups were constant irrespective of accuracy in reporting energy. Other studies have also reported energy under-reporting as having no substantial effects on DED estimates^(10,16). Thus, the trends in DED estimates with population demographics and associations of dietary intake with DED were investigated by including all participants.

It is difficult to compare these estimates with published estimates from other populations as the methods used for

calculating DED vary considerably, as discussed above. This further highlights the need for a standardised method. The DED estimates calculated for the Irish population using the food only method were found to be remarkably similar to those calculated for a sample from Northern Ireland⁽⁴⁰⁾, higher than those for a sample from Spain⁽¹⁰⁾ and lower than those for a sample from the USA⁽³⁷⁾. Some consistent trends were observed across the data; females have been reported to have lower DED estimates in child and adult groups in a number of studies^(7,10,14,16,37); similarly, Irish DED estimates were lower for females than for males. In keeping with the Irish data, a decrease in DED with age has been observed; Mendoza *et al.*⁽⁷⁾ reported a peak in DED at the age of 7–8 years and a decline thereafter in US children, while Kant & Graubard⁽¹⁴⁾ reported a lower mean DED in US adults aged >50 years than in those aged ≤ 50 years, and Ledikwe *et al.*⁽³⁷⁾ found that those aged ≥ 70 years had significantly lower DED estimates than other age groups.

DED is primarily determined by the water and fat contents of the diet⁽⁴¹⁾. Studies have shown a positive association of EI with DED⁽²⁾ and associations of lower DED with higher fat, added sugar and dietary fibre intakes and lower protein and carbohydrate intakes^(14,42). Fruit and vegetables are the food groups most commonly associated with lower DED in the literature^(8,14,16). In the present study, higher intakes of energy (kJ), fat (%TE), carbohydrates (%TE) and total sugar (%TE) were found to be associated with higher DED estimates, while higher intakes of protein (%TE) and dietary fibre (g/10MJ) were found to be associated with lower DED estimates. Variation in the intakes of fruit, vegetables and SSB had the greatest effect on mean DED estimates, followed

by that in the intakes of potatoes, fresh meat, brown bread, white bread, chips, RTEBC, and chocolate and non-chocolate confectionery. Higher intakes of fruit, vegetables, potatoes, fresh meat and brown bread were associated with lower DED estimates, while higher intakes of SSB, white bread, chips, RTEBC, and chocolate and non-chocolate confectionery were associated with higher DED estimates.

In an attempt to explain the lower DED estimates obtained for females and the inverse association of DED estimates with age group, energy-adjusted intake patterns of nutrients and the food groups that explained large amounts of variance in DED estimates in the population as a whole were examined across sexes and age groups. As DED decreased across age groups, intakes of all dietary components examined that were associated with higher DED estimates in the total population decreased and those associated with lower DED estimates increased. In particular, the intakes of fruit and vegetables increased and those of RTEBC decreased considerably across age groups, with the intakes of vegetables in 5- to 18-year-olds more than doubling when compared with those in the 51–90 years age group and those of RTEBC decreasing 3-fold. The disparity in DED estimates between sexes is less clearly explained by differences in consumption patterns. Females consumed significantly more fat, carbohydrates, total sugar, dietary fibre, fruit, vegetables, chocolate confectionery and savoury snacks and less energy, protein, potatoes and RTEBC when compared with males. Of particular interest is that females consumed 1.5 times more fruit and vegetables combined when compared with males.

The downward trend in DED estimates with age group should not be interpreted as longitudinal variations in DED, as the present analysis is based on cross-sectional data and generational effects are possible. However, to examine the potential generational effect, estimates calculated for adults from the NSIFCS were compared with those calculated for a subsample, comparable for age and sex, of adults from the NANS. Little difference was observed in Irish DED estimates over time, adding credibility to the presence of an inverse association between DED and age group.

In conclusion, differences in food choices, in particular, fruit, vegetables and RTEBC, explain the variation in DED estimates by age group and sex. The intakes of fruit, vegetables and SSB appear to have the greatest influence on DED in the Irish population and are potential targets for public health interventions. DED estimates calculated for Irish adults have not changed appreciably in the last 10 years. The DED estimates calculated in the present study are the first nationally representative estimates for an Irish population and are reported by sex and age group to serve as a baseline for future studies and public health interventions.

Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0007114514003420>

Acknowledgements

The authors thank the participants and their families for taking part in the surveys.

The present study was supported by the Irish Government under the National Development Plan (2000–6) and the Department of Agriculture, Fisheries & Food under the Food for Health Research Initiative (2007–12).

The authors' contributions are as follows: L. O'C carried out the analysis, drafted the manuscript and is responsible for the integrity of the data; J. W. is the co-ordinator of the national nutrition surveys and contributed to the manuscript content; A. F. is a principal investigator of the national nutrition surveys and provided critical input for the analysis. All authors critically reviewed the manuscript and approved the final version submitted for publication.

None of the authors has any conflicts of interest to declare.

References

1. Yao M & Roberts SB (2001) Dietary energy density and weight regulation. *Nutr Rev* **59**, 247–258.
2. Rolls BJ (2009) The relationship between dietary energy density and energy intake. *Physiol Behav* **97**, 609–615.
3. Ledikwe JH, Rolls BJ, Smiciklas-Wright H, *et al.* (2007) Reductions in dietary energy density are associated with weight loss in overweight and obese participants in the PREMIER trial. *Am J Clin Nutr* **85**, 1212–1221.
4. Vernarelli JA, Mitchell DC, Hartman TJ, *et al.* (2011) Dietary energy density is associated with body weight status and vegetable intake in U.S. children. *J Nutr* **141**, 2204–2210.
5. Howarth NC, Murphy SP, Wilkens LR, *et al.* (2006) Dietary energy density is associated with overweight status among 5 ethnic groups in the multiethnic cohort study. *J Nutr* **136**, 2243–2248.
6. Murakami K, Sasaki S, Takahashi Y, *et al.* (2007) Dietary energy density is associated with body mass index and waist circumference, but not with other metabolic risk factors, in free-living young Japanese women. *Nutrition* **23**, 798–806.
7. Mendoza JA, Drewnowski A & Christakis DA (2007) Dietary energy density is associated with obesity and the metabolic syndrome in US adults. *Diabetes Care* **30**, 974–979.
8. Ledikwe JH, Blanck HM, Khan LK, *et al.* (2006) Low-energy-density diets are associated with high diet quality in adults in the United States. *J Am Diet Assoc* **106**, 1172–1180.
9. O'Connor L, Walton J & Flynn A (2013) Dietary energy density and its association with the nutritional quality of the diet of children and teenagers. *J Nutr Sci* **2**, e10.
10. Schroder H, Covas M, Elosua R, *et al.* (2008) Diet quality and lifestyle associated with free selected low-energy density diets in a representative Spanish population. *Eur J Clin Nutr* **62**, 1194–1200.
11. Schroder H, Vila J, Marrugat J, *et al.* (2008) Low energy density diets are associated with favorable nutrient intake profile and adequacy in free-living elderly men and women. *J Nutr* **138**, 1476–1481.
12. Cuco G, Arijia V, Marti-Henneberg C, *et al.* (2001) Food and nutritional profile of high energy density consumers in an adult Mediterranean population. *Eur J Clin Nutr* **55**, 192–199.
13. Crowe TC, Fontaine HL, Gibbons CJ, *et al.* (2004) Energy density of foods and beverages in the Australian food

- supply: influence of macronutrients and comparison to dietary intake. *Eur J Clin Nutr* **58**, 1485–1491.
14. Kant AK & Graubard BI (2005) Energy density of diets reported by American adults: association with food group intake, nutrient intake, and body weight. *Int J Obes (Lond)* **29**, 950–956.
 15. Rolls BJ & Bell EA (1999) Intake of fat and carbohydrate: role of energy density. *Eur J Clin Nutr* **53**, Suppl. 1, S166–S173.
 16. Patterson E, Warnberg J, Poortvliet E, *et al.* (2010) Dietary energy density as a marker of dietary quality in Swedish children and adolescents: the European Youth Heart Study. *Eur J Clin Nutr* **64**, 356–363.
 17. Food Standards Agency (2002) *McCance and Widdowson's The Composition of Foods*, 6th ed. Cambridge, UK: Royal Society of Chemistry.
 18. Holland B, Welch AA, Unwin ID, *et al.* (1995) *McCance & Widdowson's The Composition of Foods*, 5th ed. London: HMSO. (Royal Society of Chemistry and Ministry of Agriculture, Fisheries and Food).
 19. Holland B, Brown J & Buss DH (1993) *Fish and Fish Products. Third Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 20. Holland B, Unwin ID & Buss DH (1992) *Fruit and Nuts. First Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 21. Holland B, Unwin ID & Buss DH (1991) *Vegetables, Herbs and Spices. Fifth Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 22. Holland B, Unwin ID & Buss DH (1989) *Milk Products and Eggs. Fourth Supplement to McCance and Widdowson's The Composition of Foods*, 4th ed. London: HMSO.
 23. Holland B, Unwin ID & Buss DH (1988) *Cereal and Cereal Products. Third Supplement to McCance and Widdowson's The Composition of Foods*, 4th ed. London: HMSO.
 24. Holland B, Welch AA & Buss DH (1992) *Vegetable Dishes. Second Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 25. Chan DS, Lau R, Aune D, *et al.* (2011) Red and processed meat and colorectal cancer incidence: meta-analysis of prospective studies. *PLoS One* **6**, e20456.
 26. Chan W, Brown J & Buss DH (1994) *Miscellaneous Foods. Fourth Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 27. Chan W, Brown J, Church SM, *et al.* (1996) *Meat Products and Dishes. Sixth Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 28. Chan W, Brown J, Lee S, *et al.* (1996) *Meat, Poultry and Game. Fifth Supplement to McCance and Widdowson's The Composition of Foods*, 5th ed. London: HMSO.
 29. Black LJ, Ireland J, Moller A, *et al.* (2011) Development of an on-line Irish food composition database for nutrients. *J Food Comp Anal* **24**, 1017–1023.
 30. Schofield WN (1985) Predicting basal metabolic rate, new standards and review of previous work. *Hum Nutr Clin Nutr* **39**, Suppl. 1, 5–41.
 31. Torun B, Davies PS, Livingstone MB, *et al.* (1996) Energy requirements and dietary energy recommendations for children and adolescents 1 to 18 years old. *Eur J Clin Nutr* **50**, Suppl. 1, S37–S80 (discussion S80–S81).
 32. Goldberg GR, Black AE, Jebb SA, *et al.* (1991) Critical-evaluation of energy-intake data using fundamental principles of energy physiology. 1. Derivation of cutoff limits to identify under-recording. *Eur J Clin Nutr* **45**, 569–581.
 33. Cohen JW (1998) The analysis of variance. In *Statistical Power Analysis for the Behavioural Sciences*, chapter 8, 2nd ed. Hillsdale, NJ: Lawrence Erlbaum Associates.
 34. Harrington KE, Robson PJ, Kiely M, *et al.* (2001) The North/South Ireland Food Consumption Survey: survey design and methodology. *Public Health Nutr* **4**, 1037–1042.
 35. Cox DN & Mela DJ (2000) Determination of energy density of freely selected diets: methodological issues and implications. *Int J Obes Relat Metab Disord* **24**, 49–54.
 36. Vernarelli JA, Mitchell DC, Rolls BJ, *et al.* (2013) Methods for calculating dietary energy density in a nationally representative sample. *Procedia Food Sci* **2**, 68–74.
 37. Ledikwe JH, Blanck HM, Khan LK, *et al.* (2005) Dietary energy density determined by eight calculation methods in a nationally representative United States population. *J Nutr* **135**, 273–278.
 38. Johnson L, Wilks DC, Lindroos AK, *et al.* (2009) Reflections from a systematic review of dietary energy density and weight gain: is the inclusion of drinks valid? *Obes Rev* **10**, 681–692.
 39. Malik VS, Pan A, Willett WC, *et al.* (2013) Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *Am J Clin Nutr* **98**, 1084–1102.
 40. McCaffrey TA, Rennie KL, Kerr MA, *et al.* (2008) Energy density of the diet and change in body fatness from childhood to adolescence; is there a relation? *Am J Clin Nutr* **87**, 1230–1237.
 41. Stubbs J, Ferres S & Horgan G (2000) Energy density of foods: effects on energy intake. *Crit Rev Food Sci Nutr* **40**, 481–515.
 42. Ledikwe JH, Blanck HM, Kettel Khan L, *et al.* (2006) Dietary energy density is associated with energy intake and weight status in US adults. *Am J Clin Nutr* **83**, 1362–1368.