

Title	Pilot scale production of a phospholipid-enriched dairy ingredient by means of an optimised integrated process employing enzymatic hydrolysis, ultrafiltration and super-critical fluid extraction
Authors	Barry, Kate M.;Dinan, Timothy G.;Kelly, Philip M.
Publication date	2017-04-08
Original Citation	Barry, K. M., Dinan, T. G. and Kelly, P. M. (2017) 'Pilot scale production of a phospholipid-enriched dairy ingredient by means of an optimised integrated process employing enzymatic hydrolysis, ultrafiltration and super-critical fluid extraction', Innovative Food Science & Emerging Technologies, 41, pp. 301-306. doi: 10.1016/j.ifset.2017.04.004
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.ifset.2017.04.004
Rights	© 2017 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license. - <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
Download date	2024-04-15 23:14:45
Item downloaded from	<a href="https://hdl.handle.net/10468/3958">https://hdl.handle.net/10468/3958</a>




# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

## Accepted Manuscript

Pilot scale production of a phospholipid-enriched dairy ingredient by means of an optimised integrated process employing enzymatic hydrolysis, ultrafiltration and super-critical fluid extraction

Kate M. Barry, Timothy G. Dinan, Philip M. Kelly



PII: S1466-8564(17)30167-4  
DOI: doi: [10.1016/j.ifset.2017.04.004](https://doi.org/10.1016/j.ifset.2017.04.004)  
Reference: INNFOO 1742

To appear in: *Innovative Food Science and Emerging Technologies*

Received date: 9 February 2017  
Revised date: 6 April 2017  
Accepted date: 7 April 2017

Please cite this article as: Kate M. Barry, Timothy G. Dinan, Philip M. Kelly , Pilot scale production of a phospholipid-enriched dairy ingredient by means of an optimised integrated process employing enzymatic hydrolysis, ultrafiltration and super-critical fluid extraction. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Innfoo(2017), doi: [10.1016/j.ifset.2017.04.004](https://doi.org/10.1016/j.ifset.2017.04.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Pilot scale production of a phospholipid-enriched dairy ingredient by means of an optimised integrated process employing enzymatic hydrolysis, ultrafiltration and super-critical fluid extraction.**

**Kate M. Barry<sup>ab</sup>, Timothy G. Dinan<sup>b</sup>, & Philip M. Kelly<sup>a</sup>**

*<sup>a</sup> Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland*

*<sup>b</sup> Department of Psychiatry, University College Cork, Cork, Ireland*

**Corresponding Author**

Dr. Philip M. Kelly,

Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork

Tel: +353 (0)87 7735352

Email: [phil.kelly@teagasc.ie](mailto:phil.kelly@teagasc.ie)

**Abstract**

Pilot scale production of a dairy ingredient enriched in phospholipids (PLs) was generated from a buttermilk powder (BMP) substrate utilising a combined process of targeted enzymatic hydrolysis of the innate milk proteins followed by ultrafiltration with a 50 kDa membrane. An 8.5 fold increase in PL was achieved in the 50 kDa retentate (50 R) compared to the BMP,  $11.05 \pm 0.02$  % and  $1.30 \pm 0.00$  % total PL, respectively. Simultaneously, total lipid content in the retentate increased 8.7 fold with reference to the BMP,  $60.07 \pm 0.54$  % and  $6.84 \pm 0.17$  % total lipid respectively. Protein reduced to  $10.58 \pm 0.09$  % (50 R) from  $31.40 \pm 0.57$  % in BMP. Supercritical CO<sub>2</sub> fluid extraction (SFE) was employed to generate a purified lipid fraction. SFE with ethanol as a co-solvent yielded a purified lipid extract with enriched PLs level of  $56.24 \pm 0.07$  % on a dry matter basis.

**Industrial relevance**

PLs have many associated health and nutritional benefits including those related to cognitive development and repair. Generation of an ingredient enriched in dairy PLs would be advantageous from an industrial view to allow fortification of nutritionals, both infant and geriatric, in promoting brain health. The present work demonstrates a novel pilot scale process for the generation of a PL enriched ingredient from a BMP substrate. Utilising a combined process of targeted protein hydrolysis followed by selective removal by ultrafiltration of the smaller molecular weight peptide material, an ingredient with 8.5 fold increase in PL material was achieved. SFE technology was utilised to generate a purified lipid extract with greater PL levels for future applications in biological assays to determine these pathways. The need to investigation and further develop the knowledge relating to the modes of action of these bioactive compounds would be beneficial from a nutritional perspective.

Key words;

Pilot-scale, buttermilk powder (BMP), phospholipids (PLs), hydrolysis, ultrafiltration, supercritical CO<sub>2</sub> fluid extraction (SFE).

## 1. Introduction

The milk fat globule membrane (MFGM) is a unique biophysical system (Lopez, 2011; Singh, 2006) surrounding milk fat globules that facilitates their dispersal in milk. This surface active multi-layered membrane is derived from the apical membrane of the secretory cell in the mammary gland (Heid & Keenan, 2005) and is composed of numerous bioactive molecules of which 90 % of MFGM dry weight is represented by proteins and polar lipids (Lopez, 2011). Phospholipids (PLs), a complex class of polar lipids, whose structure comprises of a hydrophilic head group and a hydrophobic fatty acid tail (Contarini & Povolo, 2013), are amphiphilic in nature. This distinctive polar property is responsible for the structural architecture of MFGM which, in turn, aids in the emulsification of fat in milk (Barry, Dinan, Murray, & Kelly, 2016; Contarini & Povolo, 2013; Sánchez-Juanes, Alonso, Zancada, & Hueso, 2009).

Buttermilk, an aqueous serum phase released during cream de-emulsification/churning, possesses PL concentrations approx. 15x that of whole milk (Barry et al., 2016; Rombaut, van Camp, & Dewettinck, 2005) and, hence, is a preferred substrate for the PL enrichment. Dairy PLs are of considerable interest due to their particular prominence of sphingomyelin (SM) and phosphatidylserine (PS) (Burling & Graverholt, 2008). The technological and nutritional functionalities of PLs have been extensively studied (Contarini & Povolo, 2013; Dewettinck, Rombaut, Thienpont, Le, Messens, & van Camp, 2008). PLs, due to their emulsification and stabilisation properties afforded by their inherent amphiphilic nature are widely utilised by the food and dairy industry for their associated

physiochemical roles in dairy products that include emulsification, heat stability, wettability and viscosity (Contarini & Povo 2013; Corredig & Dalgleish 1997; Corredig & Dalgleish 1998; Dewettinck et al., 2008).

Likewise, there have been numerous health and nutritional benefits associated with PLs e.g. reduction in the incidence of cardiovascular disease, tolerance of depression and stress and myelination of the central nervous system (Contarini & Povo 2013; Dewettinck et al., 2008; Oshida, Shimizu, Takase, Tamura, Shimizu & Yamashiro, 2003). Sphingomyelin (SM), mainly through its sphingosine and ceramide metabolites, has been shown to affect anticancer activity notably in the suppression of colon tumours (Berra, Colombo, Scottocotnola, & Giacose, 2002; Hertevig, Nilsson, Cheng, & Duan, 2003; Kuchta-Noctor, Murray, Stanton, Devery, & Kelly, 2016; Spitsberg, 2005). The biological activity of PLs has also been associated with cognitive development and repair. Phosphatidylserine, pre-eminently, has been implicated in the treatment of Alzheimer's disease (Burling & Graverholt, 2008; Contarini & Povo, 2013; Dewettinck et al., 2008). In more recent years, studies on infant milk formula fortified with PLs have proved positive in terms of immunological defences (Timby, Hernell, Vaarala, Melin, Lönnerdal, & Domellöf, 2015), and also infant neurodevelopment (Timby, Domellöf, Hernell, Lönnerdal, & Domellöf, 2014). Thus, there is considerable interest and demand for the generation of PL concentrates from dairy sources.

Technological approaches utilised to concentrate PLs in buttermilk (BM) have been primarily membrane filtration based, particularly microfiltration (MF) (Astaire, Ward, German, & Jiménez- Flores, 2003; Corredig, Roesch, & Dalgleish, 2003; Holzmüller & Kulozik, 2016; Morin, Pouliot, & Jiménez-Flores, 2006; Morin, Britten, Jiménez-Flores, & Pouliot, 2007) due to its greater porosity. However, similarity in the size of the MFGM components and casein (Morin et al., 2007) limited the effectiveness of MF as a selective

separation process. Complementary measures adopted to enhance MF include: cream washing prior to ultrafiltration (UF) (Morin et al., 2007), rennet induced coagulation of the caseins prior to MF (Sachdeva & Buchheim, 1997), and disruption of the casein micelle with citrate prior to MF (Corredig et al., 2003; Roesch, Rincon, & Corredig, 2004; Rombaut, Dejonckheere, & Dewettinck, 2006). A more recent avenue of investigation employed the use of enzymes to pre-digest the milk proteins prior to UF (Barry, Dinan, & Kelly, 2017). Hydrolysis of milk proteins prior to UF enables selective transmission of smaller peptides through the membrane, thus, selectively concentrating PLs in the retentate. Barry et al., (2017) reported that PLs losses through the UF membrane using this novel process were non-detectable.

A more advanced separation technology in the form of supercritical fluid extraction (SFE) has been periodically investigated to selectively extract unique higher value lipids, in particular from primary food substrates including BM sources (Astaire et al., 2003; Costa, Elias-Argote, & Jiménez-Flores, 2010; Spence, Jiménez-Flores, Qian, & Goddik, 2009). Astaire et al., (2003) were able to selectively remove non-polar lipids from the buttermilk by coupling MF with SFC, thus, allowing concentration of the polar lipids in the buttermilk. Spence et al., (2009) concentrated dairy PLs 5-fold when applying SFC to MF pre-processed samples.

The objective of the present study was to apply SFE while upscaling a laboratory-based enzymatic hydrolysis/ultrafiltration process previously outlined by the authors (Barry et al., 2017), in order to produce a purified milk lipid extract with elevated levels of PLs relative to non-polar milk lipids.

## 2. Materials and methods

### 2.1. Materials

Spray dried buttermilk powder (BMP) was sourced from Tipperary Co-operative (Tipperary, Ireland) with compositional analysis by standard IDF protocols determining 6.84 % fat, 31.40 % protein, 48.00 % lactose, and 7.40 % minerals. The digestive enzyme Alcalase (E.E. 3.4.21.62) was procured from Novozymes (Bagsværd, Denmark). Alcalase is a serine type endoprotease with esterase activity enabling it to catalyse the stereoselective hydrolysis of amino esters and selective esters. Alcalase has a minimum activity of 2.4 Anson units per gram of composition (2.4 AU-A g<sup>-1</sup>) at pH 7.50, with optimal conditions for alcalase activity within 35- 60°C and pH 7-9. Synder PES spiral wound membranes with a nominal molecular weight cut off (MWCO) of 50 kDa were purchased from David Kellett & Partners Ltd. (Hereford, UK). Acetonitrile was procured from ThermoFisher Scientific (Waltham, MA, USA) and all other chemicals were HPLC grade, > 99.9 %, and purchased from Sigma Aldrich (Arklow, Ireland).

### 2.2. Enzymatic hydrolysis of BMP substrate protein fraction

The degree of hydrolysis (% DH) is interpreted as the number of peptide bonds (*h*) that are cleaved as a percentage of the total peptide bonds (*htot*) all of which is related to the consumption of base as a result of proton release during the hydrolysis reaction according Adler-Nissen's (1986) equation (Eq. 1).

$$\% DH = 100 \times B \times N_B \times \frac{1}{\alpha} \times \frac{1}{MP} \times \frac{1}{h_{tot}} \quad (1)$$

Where, B is the volume of base in mL, N<sub>B</sub> is the normality of the base, 1/α is the average degree of dissociation of the α-NH groups (=1.13 at pH 8.00, 50°C), MP is the molecular



weight of the protein and  $htot$  is the number of peptide bonds in the protein substance (=8.2 for casein/whey mix).

For all hydrolysis reactions, BMP was reconstituted to 10 % total solids in reverse osmosis (RO) water, 360 Kg total weight ( 37.29 Kg BMP and 322.7005 Kg RO water), and allowed to hydrate over night at 4 °C with gentle agitation. The BMP dispersions were heated gently to 50 °C with a water-jacket (to avoid burn on of the BMP to the walls of the tank) and subsequently pH corrected with 4 M NaOH to pH 8.00 (original pH 6.45). The hydrolysis reaction was initiated by addition of alcalase dissolved in RO water at an enzyme to substrate ratio (E:S) of 1: 100, w w<sup>-1</sup>, (1 %) on a protein equivalent basis with the temperature, 50 °C, and pH, pH 8.00, maintained throughout the reaction. The pH was controlled by addition of 4M NaOH and the reaction was agitated throughout utilizing the inherent tank impeller. The hydrolysis reaction was terminated by means of pH inactivation with 2 M citric acid until a pH of 6.5 was achieved. All hydrolysis reactions were performed in duplicate.

Determination of the end-point DH was measured by modification of the trinitrobenzenesulfonic acid (TNBS) reaction outlined by Adler-Nissen (1979) which involves measuring the reaction of liberated  $\alpha$ -amino groups ( $\alpha$ -NH<sub>2</sub>) with 2, 4, 6-trinitrobenzene 1- sulfonic acid. Briefly, samples with a protein concentration of 3 mg mL<sup>-1</sup> were prepared in 1 % SDS, 1 % SDS alone serving as a blank, and 0.2125 M phosphate buffer was added. 2 mL of TNBS was added to both the samples and blank with light omitted, and the reaction was incubated for 1 hr at 50 °C. 4 mL of 0.1 M HCl was added to terminate the reaction and the samples were cooled for 30 min prior to absorbance at 314 nm being read using a Varian Cary 1 dual beam UV- visible spectrometer (Varian Ltd., Walton-on-Thames, U.K). UV readings were determined against a standard curve generated using L-

leucine at concentrations ranging from 20-250 mg L<sup>-1</sup>. DH values were calculated using the following equation (Spadaro, Draghetta, del Lama, Camargo, & Greene, 1979) (Eqn. 2)

$$\% DH = \frac{AN_2 - AN_1}{N_{pb}} \times 100 \quad (2)$$

Where, AN<sub>1</sub> represents the amino nitrogen content of the protein substrate (unhydrolysed) (mg g<sup>-1</sup> protein), AN<sub>2</sub> represents the amino nitrogen content of the hydrolysed protein substrate and N<sub>pb</sub> represents the nitrogen content of the peptide bonds on the protein substrate (mg g<sup>-1</sup> protein).

### 2.3. Ultrafiltration of the BMP hydrolysate

The BMP hydrolysate produced was subjected to membrane ultrafiltration utilising a GEA Model F unit (GEA Process Engineering A/S, Skanderborg, Denmark). UF was accomplished with two Synder PES spiral wound membranes (David Kellet & Partners, Ltd.) having a nominal molecular weight cut off (MWCO) of 50 kDa and a total membrane area of 6.69 m<sup>2</sup>. UF was carried out at 50 °C to a volume concentration factor of 11 by addition of 4 x diafiltration steps (3x 200Kg RO water and 1 x 100Kg RO water) during the membrane process. A feed recirculation rate of 100 L h<sup>-1</sup> at 1 bar and a membrane inlet pressure of 4.2 bar were maintained throughout processing. UF was carried out in duplicate. All 50 kDa retentates (50 R) and 50 kDa permeates (50 P) were evaporated to 40 % total solids on a Tetra Scheffers™ falling film single-stage evaporator (Tetra Pak, Gorredijk, The Netherlands) and subsequently spray-dried on a pilot scale Anhydro Lab 3 spray drier (SPX Flow Technology A/S, Soeborg, Denmark) with an inlet temperature ranging from 185 to 190 °C and an outlet temperature ranging from 85 to 90 °C during spray drying. UF was carried out in duplicate.

#### **2.4. Chromatographic characterisation of the BMP hydrolysate and UF fractions protein.**

Observation of the loss of native protein and molecular weight characterisation of the BMP hydrolysate and UF fractions (50 R and 50 P), size exclusion chromatography (SEC), according to the method of O'Loughlin et al., (2013), was carried out using a Waters 2695 separation module coupled with a Waters 2487 dual wavelength absorbance detector with Waters Empower software (Milford, MA, USA). All column eluates were monitored at 214 nm unless otherwise stated. Briefly, samples with a  $2.5 \text{ mg mL}^{-1}$  protein concentration were injected,  $20 \text{ }\mu\text{L}$ , onto a TSK Gel G2000SW,  $7.8 \text{ mm} \times 600 \text{ mm}$  column (Tosoh Bioscience GmbH, Stuttgart, Germany). An isocratic gradient of 30 % acetonitrile with 0.1 % TFA was maintained for a run time of 60 min. Sample chromatograms were compared with a molecular weight curve generated from standards,  $2.5 \text{ mg mL}^{-1}$  protein, of known molecular weights;  $\alpha$ -lactalbumin,  $\beta$ -lactoglobulin A and B, Bovine serum albumin (BSA), lactoferrin and CMP procured from Sigma Aldrich and Ribonuclease A, Cytochrome C, Aprotinin, Bacitracin, His- Pro- Arg- Tyr, Leu- Tyr,- Met- Arg, Bradykinin, Leu- Phe and Tyr- Glu procured from Bachem AG (Bubendorf, Switzerland).

#### **2.5. Chromatographic quantification of BMP phospholipids.**

Extraction of the crude lipid from the BMP hydrolysate and UF fractions (50 R and 50 P) was carried out according to the method of Folch (Folch, Lees, & Stanley, 1957). Briefly, 4 g of powder was dissolved in 75 mL chloroform: methanol (2:1, v v<sup>-1</sup>), and refrigerated at 4 °C with agitation for 1 hr. The sample was then subjected to centrifugation at  $4000 \times g$  for 5 min following which the bottom  $\text{CHCl}_3$  layer was collected and the top MeOH layer was washed with 50 mL  $\text{CHCl}_3$  and refrigerated with agitation for another hour. After centrifugation of

the sample both  $\text{CHCl}_3$  layers were pooled together and evaporated using a Buechi rotary evaporator (BUCHI, Flawil, Switzerland), and dried to dryness under  $\text{N}_2$ .

Separation and quantification of the BMP hydrolysate and UF fraction PLs was carried out according to the method of Barry et al., (2016). Briefly, the crude lipid extract was re-solved in  $\text{CHCl}_3$ : MeOH (88:12, v v<sup>-1</sup>). 25  $\mu\text{L}$  of the sample was applied to 3  $\mu\text{m}$  particle diameter Prevail Silica column (Labquib Dublin, Ireland) with separation carried using a Waters 2695 separation module (Waters) coupled to a Corona Charged Aerosol Detector (CAD) (Thermo Scientific, Hemel Hempstead, U.K) under the following linear gradient elution conditions with a flow rate of 0.50  $\text{mL min}^{-1}$  and ratio in volume of solvent A to B; 96:4 at T0, 25:75 at T20, 6:94 at T21 and 96:4 at T22 and maintained until T27. All lipid extractions were performed in triplicate and were pre- filtered through 0.20  $\mu\text{m}$  Hydrophobic Fluoropore™ (PTFE) membrane filters (Merek Millipore Ltd., Cork, Ireland) prior to HPLC. PL recovery was calculated by quantification with calibration curves that had been generated for each individual PL species.

## **2.5. Supercritical carbon dioxide fluid extraction (SFE) of 50 kDa retentate.**

SFE was carried out on the 50 R that was generated from the combined hydrolysis and 50 kDa membrane ultrafiltration process. SFE was carried out at The Biocomposites Centre in Bangor University (Wales, UK) on a Thar laboratory rig, Thar SFC-1000 (Thar Energy LLC, Pittsburg, PA, USA), with a 1000 mL extractor vessel, with the following extraction parameters; 40 °C extraction temperature and automated back pressure at 300 bar. The system was controlled manually with a computer-based Windows software. The initial defatting stage involved subjecting the 50 R sample to supercritical  $\text{CO}_2$  only, to remove non-polar lipids. The subsequent secondary PL purification stage utilised  $\text{CO}_2$  with ethanol as a co-solvent at different percentages, 10 % and 20 %, to increase the polarity of the extraction

solvent to enable maximum separation of the PLs from the 50 R. The entire extraction took approximately 13 h and the resulting PL extract was concentrated using a Buechi rotary evaporator and dried to dryness under N<sub>2</sub> prior to analysis by HPLC as mentioned above (Section 2.4). SFE was carried out in duplicate.

## **2.6. Compositional analysis**

Compositional analysis was performed on all samples using the standard IDF/ISO or AOAC methods for determination of % protein (ISO 8968-1, 2014), % total solids/moisture and ash (ISO 6731, 2010) and lactose (AOAC 896.01, 1896). % fat was determined using the Folch method described by Folch et al., (1957).

## **2.7. Statistical analysis.**

Statistical analysis was performed with Minitab, version 15 (Minitab Ltd, Coventry, UK), with one- way ANOVAs and Tukey Tests for both intra- and inter-sample variance.

## **3. Results and discussion**

### **3.1. Enzymatic hydrolysis of BMP substrate protein fraction**

A previous study carried out by the authors determined that a 19 % degree of hydrolysis (%DH) sufficiently digested milk proteins in order to enable permeation through a 50 kDa ultrafiltration (UF) membrane. In this study, the authors additionally highlight the importance of mode of inactivation utilised to terminate the hydrolytic reaction at pilot scale level i.e., thermal versus pH inactivation. Furthermore, it was also necessary to consider the implications that the selected inactivation step may have on the final hydrolysate e.g. risk of aggregate formation upon thermal treatment (80°C for 20 min) of the hydrolysate. Hence, pH-based inactivation was adopted as the method for termination of the enzymatic reaction.

Hydrolysis to 19 % DH of the BMP starting material resulted in a shift in the molecular mass distribution of the starting material (Table 1). Hydrolysis resulted in a higher percentage of smaller molecular weight material, > 50 kDa in size, compared with that of the BMP,  $0.05 \pm 0.03\%$  and  $99.95\%$  (summation of M.W > 50 kDa) respectively (Table 1). Size exclusion chromatography (SEC) chromatograms (Fig. 1) illustrate the larger molecular weight material present in the starting BMP material (Fig. 1a) and the presence of a large body of smaller molecular weight peptides following extensive digestion (Fig. 1b). These results are in agreement with those obtained previously at the laboratory scale (Barry et al., 2017).

### 3.2. Ultrafiltration of the BMP hydrolysate

Previous work carried out by the authors (Barry et al., 2017) determined that laboratory-based ultrafiltration (UF) using a cartridge-type membrane with a nominal molecular weight cut off of 50 kDa permitted maximum permeation of the digested proteinaceous material. In this pilot scale study, UF was carried out on the BMP hydrolysate utilising a GEA Model F unit fitted with two Synder PES 50 kDa spiral wound membranes in series. To ensure optimum enrichment of PLs in the retentate, 4 diafiltration steps were carried out during the 3h membrane process to allow maximum washing of peptide material in the retentate through the membrane, thus enabling a volume concentration factor of 11 to be achieved. A feed recirculation rate of  $100 \text{ L h}^{-1}$  at 1 bar was maintained throughout the membrane process with steady stage permeate flux, this indicating that there was no evidence of membrane fouling during 3h filtration (supplementary fig. 1).

An 8.7-fold increase in lipid content of the 50 kDa retentate (50 R),  $60.07 \pm 0.54\%$  total lipid on a dry matter basis (DM), with reference to the starting BMP,  $6.84 \pm 0.17\%$  total lipid (DM) (Table 2.) was achieved. Concurrent with the increase in lipid concentration, a 3-fold protein decrease in the starting material relative to 50 R was achieved,  $31.40 \pm 0.57\%$  and

10.58 ± 0.09 % respectively (Table 2.) due to permeation of peptide material < 50 kDa (Table 1.) through the UF membrane. The deproteinising effect of the combined enzymatic hydrolysis and UF process was also confirmed by SEC molecular size distribution profiling of the 50 R and the 50 kDa permeate (50 P) (Fig. 2) e.g. a maximum absorbance of 0.15 mV for 50 R (Fig. 2a) was recorded compared to an absorbance of 2.81 mV for 50 P (Fig. 2b). UF was carried out in duplicate with no significant difference between each run ( $p < 0.01$ , one-way ANOVA).

### **3.3 Phospholipid (PL) enrichment of the 50 kDa retentate via enzymatic hydrolysis and UF.**

Determination of the degree of PL enrichment achieved utilising the combined enzymatic hydrolysis and UF process was confirmed through quantification (Barry et al., 2016) of the PL chromatographic profiles generated for the BMP starting material, the 50 R and the 50 P. The crude lipid fractions of each substrate were extracted via Folch (Folch et al., 1957) with, as mentioned in section 3.2., an 8.7-fold increase in the total lipid content on a dry matter basis between the 50 R and the original BMP, 60.07 ± 0.54 % total lipid (DM) and 6.84 ± 0.17 % total lipid (DM) respectively, was achieved (Table 2.). Subsequent to the crude lipid extraction, the extracts were analysed to determine their relative PL contents by HPLC with a charged aerosol detection (CAD) system. Due to permeation of the smaller molecular weight peptides (<50 kDa) and the concurrent increase in lipid content, a 8.5 fold increase in PLs was achieved in the 50 R, 11.05 ± 0.02 % total PL (summation of PI, PE, GluCer PS, PC and SM), compared to the starting BMP, 1.30 ± 0.00 % total PL (Table 2.). Thus, the 8.7-fold increase achieved at pilot scale level exceeds the 7.8-fold increase that the authors previously achieved during the development phase on a laboratory scale where a 50 kDa UF membrane cassette was employed to process the enzymatically hydrolysed buttermilk (Barry et al.,

2017). This enrichment is also greater than that reported in other work utilising similar hydrolysis and/or filtration technologies. Morin et al., (2006) achieved a 3-fold enrichment of PL during MF. However, lower filtration temperatures appear to have negatively impacted separation performance (Astaire et al., 2003; Morin et al., 2006). A 2.8-fold increase in PL was obtained by Morin et al., (2007) during combined MF and DF of washed cream buttermilk. A 2.5 % increase in PL was achieved by Konrad et al., (2013) following peptic hydrolysis and UF.

The higher sensitivity and precision of the CAD detector system (Barry et al., 2016) is largely responsible for the higher levels of the more acidic PLs, PI and PS, than those previously reported,  $10.84 \pm 0.04$  % PI and  $9.54 \pm 0.01$  % PS in the BMP and  $10.42 \pm 0.04$  % and  $9.88 \pm 0.07$  % in the 50 R. The previously reported improvements to the Folch crude lipid extraction protocol (Barry et al., 2016) is also likely to be a contributory factor.

Well defined peaks representing the five major PLs, namely phosphatidylinositol (PI), phosphatidylethanolamine (PE), phosphatidylserine (PS), phosphatidylcholine (PC) and sphingomyelin (SM) along with glucosylceramide (GluCer) were demonstrated in the HPLC-CAD chromatograms generated for the BMP and 50 R fractions (Fig. 3). Also present are the sub-peaks of both PC (PC 1, PC 2) and SM (SM 1, SM 2), which represent different molecular subspecies of the PLs as previously described (Barry et al., 2016, Fagan & Wijesundera, 2004).

The PL profile representing the 50 R (Fig. 3b) demonstrates the higher concentration of PLs when compared to the PL profile of the BMP (Fig. 3a),  $11.05 \pm 0.02$  % total PL and  $1.30 \pm 0.00$  % total PL, respectively (Table 2.). While there was some permeation of lipid material during the membrane filtration,  $0.29 \pm 0.07$  % total lipid in the 50 P, losses of PL during the combined hydrolysis and UF process were non-detectable,  $0.00 \pm 0.09$  % total PL



in the 50 P (Table 2.). This is in general agreement with that previously reported by the authors (Barry et al., 2017).

### 3.4 Supercritical CO<sub>2</sub> Extraction (SFE)

After UF of the BMP hydrolysate, the 50 R was evaporated to 40 % solids and dried on a pilot scale Anhydro Lab 3 spray drier. In order to maximise PL enrichment further than that previously achieved at the laboratory scale by the authors (Barry et al., 2017), the 50 R powder obtained was subjected to further 2-stage separation utilising SFE- An initial defatting stage that removed non-polar lipid material utilising SFE alone, followed by a secondary purified PL extraction stage using CO<sub>2</sub> with ethanol (EtOH) as a co-solvent to create a more polar environment for extraction of PLs. Duplicate SFE runs involving 20% EtOH as co-solvent increased PL content to  $56.24 \pm 0.07$  % (Table 2.) compared to  $11.05 \pm 0.02$  % for the original 50 R powder. From the chromatogram obtained for the SFE sample (Fig. 4), it is noted that only the peaks representing PE, PC and SM are present. This may be due to the more polar nature of the PI and PS and, hence, it may be necessary to increase the polarity of the extraction solvent further with supercritical H<sub>2</sub>O. A 10 % ratio of EtOH was also investigated; however there were non-detectable levels of PLs in this fraction (data not shown). While the authors were unsuccessful in extracting all of the PL species, they were successful in generating a fraction where more than half the weight,  $56.24 \pm 0.07$  %, on a dry matter basis is represented by PLs. To our knowledge, the use of SFE technology in this manner to generate a purified PL fraction free of protein or lactose has not been previously accomplished. This purified fraction is a useful substrate with which to undertake various bioassay studies in order to properly investigate the numerous effects of PLs and the pathways by which their effects are believed to be implicated. Spence et al., (2009) achieved a PL concentration factor of between 3.49 and 4.24 when performing SFE on sweet and whey cream buttermilks following MF (Spence et al., 2009). A 4-fold increase in PLs was achieved

by Costa et al., (2010) in a whey buttermilk source treated with SFE after UF (Costa et al., 2010). However, in both cases the enrichment was carried out by reducing the non-polar lipid content, thereby, increasing both PL and residual protein content, simultaneously.

#### 4. Conclusion

This study demonstrated a number of key observations with the aim of the work to generate a novel ingredient enriched in dairy PLs. Firstly, successful scalability of a laboratory based process was achieved. With some adaptation of the original combined enzymatic hydrolysis and UF process, the authors succeeded in producing a PL enriched 50 kDa retentate (50 R) at pilot scale that surpassed their previously reported laboratory-based results (Barry et al., 2017). An 8.7-fold increase in lipid content achieved in the 50 R following the combined pilot scale process compared to the starting BMP ( $60.07 \pm 0.54$  % v.  $6.84 \pm 0.17$  %) using spiral-wound UF membranes is very favourable for industrial scale exploitation at this of enrichment. Parallel to this increase in lipid content, an 8.5-fold increase in PL material resulted comparing the BMP starting substrate and the 50 R,  $1.30 \pm 0.00$  % and  $11.05 \pm 0.02$  %, respectively. Concurrent with the increases in both total lipid content and PL content, a 3-fold decrease in proteinaceous material was observed,  $31.40 \pm 0.57$  % in the BMP and  $10.58 \pm 0.09$  % in the 50 R. The increase in PLs was greater than that previously reported using similar hydrolytic and separation technologies. The association of PLs in cognitive development and repair in the case of Alzheimer's disease could open up avenues for utilisation of this enriched PL fraction to fortify infant or geriatric nutritional with PLs.

Secondly, to complement the PL enrichment mentioned above, generation of a purified PL fraction was achieved through the utilisation of SFE. Following SFE, a fraction with a  $56.24 \pm 0.07$  % total PL content was achieved, with PLs representing more than half the dry matter. Generation of a purified PL fraction, depleted of proteinaceous and lactose

material, has to the best of our knowledge not been accomplished to this extent in previous published work and, hence, presents an excellent opportunity with which to pursue further *invitro* and *in vivo* studies on the putative health benefits associated with PLs.

### **Acknowledgements**

The authors would like to acknowledge that the work herein was funded by the Food Institutional Research Measure (FIRM), project number: 10RD/ TMFRC/709 administered by the Irish Department of Agriculture, Food & Marine. K. M. Barry was a Teagasc PhD Walsh Fellow supported by this FIRM project.

The authors would also like to thank and express their appreciation for the help and expertise provided by Professor Ray Marriott and Dr Luis Martin of The Biocomposites Centre in Bangor University when performing the supercritical carbon dioxide extraction process.

## References

- Adler-Nissen, J. (1979). Determination of the degree of hydrolysis of food protein hydrolysates by trinitrobenzenesulfonic acid. *Journal of Agriculture and Food Chemistry*, 27, 1256- 1262.
- Adler-Nissen, J. (1986). *Enzymatic Hydrolysis of Food Proteins*. Elsevier Applied Science Publishers Ltd, New York.
- AOAC (1896). Lactose in milk. Polarimetric method, method no. 896.01. In Latimer (Ed) Official methods of analysis of AOAC International (19th Ed). Gaithersburg, MD, USA: AOAC International, 2012.
- Astaire, J. C., Ward, R., German, J. B., & Jiménez- Flores, R. (2003). Concentration of polar MFGM lipids from buttermilk by microfiltration and supercritical fluid extraction. *Journal of Dairy Science*, 86, 2297-2307.
- Barry, K. M., Dinan, T. G., Murray, B. A., & Kelly, P. M. (2016). Comparison of dairy phospholipid preparative extraction protocols in combination with analysis by high performance liquid chromatography coupled to a charged aerosol detector. *International Dairy Journal*, 56, 179-185.
- Barry, K. M., Dinan T. G., & Kelly, P. M. (2017). Selective enrichment of dairy phospholipids in a buttermilk substrate through investigation of enzymatic hydrolysis of milk proteins in conjunction with ultrafiltration. *International Dairy Journal*, 68, 80-87.
- Berra, B., Colombo, I., Scottocornola, E., & Giacosa, A. (2002). Dietary sphingolipids in colorectal cancer prevention. *European Journal of Cancer Prevention*, 1, 198-197.

- Burling, H., & Graverholt, G. (2008). Milk- a new source for bioactive phospholipids for use in food formulations. *Lipid Technology*, 20, 229-231.
- Contarini, G., & Povolo, M. (2013). Phospholipids in milk fat: composition, biological and technological significance, and analytical strategies. *International Journal of Molecular Science*, 14, 2808-2831.
- Corredig, M., & Dalgleish, D. G. (1997). Isolates from industrial buttermilk: emulsifying properties of materials derived from the milk fat globule membrane. *Journal of Agricultural and Food Chemistry*, 45, 4595-4600.
- Corredig, M., & Dalgleish, D. G. (1998). Buttermilk properties in emulsions with soybean oil as affected by fat globule membrane derived proteins. *Journal of Food Science*, 63, 466- 480.
- Corredig, M., Roesch, R. R., & Dalgleish, D. G. (2003). Production of a novel ingredient from buttermilk. *Journal of Dairy Science*, 86, 2744-2750.
- Costa, M. R., Elias- Argote, X. E., & Jiménez- Flores, R. (2010). Use of ultrafiltration and supercritical fluid extraction to obtain a whey buttermilk powder enriched in milk fat globule membrane phospholipids. *International Dairy Journal*, 20, 598- 602.
- Dewettinck, K., Rombaut, R., Thienpont, N., Le, T. T., Messens, K., & van Camp, J. (2008). Nutritional and technological aspects of milk fat globule membrane material. *International Dairy Journal*, 18, 436-457.
- Fagan, P., & Wijesundera, C. (2004). Liquid chromatographic analysis of milk phospholipids with on- line pre- concentration. *Journal of Chromatography A*, 1054, 241-509.

- Folch, J., Lees, M., & Stanley, G. H. S. (1957). A simple method for the isolation and purification of total lipides from animal tissue. *Journal of Biological Chemistry*, 226, 497- 509.
- Heid, H. W., & Keenan, T. W. (2005). Intracellular origin and secretion of milk fat globules. *European Journal of Cell Biology*, 84, 245- 258.
- Hertervig, E., Nilsson, A., Cheng, Y., & Duan, R. D. (2003). Purified intestinal alkaline sphingomyelinase inhibits proliferation without inducing apoptosis in HT-29 colon carcinoma cells. *Journal of Cancer Research and Clinical Oncology*, 129, 577-582.
- Holzmüller, W., & Kulozik, U. (2016). Technical difficulties and future challenges in isolating membrane material from milk fat globules in industrial settings- a critical review. *International Dairy Journal*, 61, 51-66.
- ISO (2010). *Milk, cream and evaporated milk- determination of total solids content. ISO 6731*. Geneva, Switzerland: International Organisation for Standardization.
- ISO (2014). *Milk and milk products- determination of nitrogen content. Part 1: Kjeldahl principle & crude protein. ISO 8968-1*. Geneva, Switzerland: International Organisation for Standardization.
- Konrad, G., Kleinschmidt, T., & Lorenz, C. (2013). Ultrafiltration of whey buttermilk to obtain a phospholipid concentrate. *International Dairy Journal*, 30, 39-40.
- Kuchta- Noctor, A. M., Murray, B. M., Stanton, C., Devery, R., & Kelly, P. M. (2016). Anticancer Activity of Buttermilk Against SW480 Colon Cancer Cells is Associated with Caspase- Independent Cell Death and Attenuation of Wnt, Akt, and ERK Signaling. *Nutrition and Cancer*, 68, 1234-1246.

- Lopez, C. (2011). Milk fat globules enveloped by their biological membrane: unique colloidal assemblies with a specific composition and structure. *Current Opinions in Colloid and Interface Science*, 16, 391- 404.
- Morin, P., Pouliot, Y., & Jiménez- Flores, R. (2006). A comparative study of the fractionation of regular buttermilk and whey buttermilk by microfiltration. *Journal of Food Engineering*, 77, 521-528.
- Morin, P., Britten, M., Jiménez- Flores, R., & Pouliot, Y. (2007). Microfiltration of buttermilk and washed cream buttermilk for concentration of milk fat globule membrane components. *Journal of Dairy Science*, 90, 2132-2140.
- O'Loughlin, I. B., Murray, B. A., Brodkorb, A., FitzGerald, R. J., Robinson, A. A., Holton, T. A., & Kelly, P. M. (2013). Whey protein isolate polydispersity affects enzymatic hydrolysis outcomes. *Food Chemistry*, 141, 2334-2342.
- Oshida, K., Shimizu, T., Takase, M., Tamura, Y., Shimizu, T., & Yamashiro, Y. (2003). Effects of dietary sphingomyelin on central nervous system myelination in developing rats. *Paediatric Research*, 53, 589-593.
- Roesch, R. R., Rincon, A., & Corredig, M. (2004). Emulsifying properties of fractions prepared from commercial buttermilk by microfiltration. *Journal of Dairy Science*, 87, 4080-4087.
- Rombaut, R., Dejonckheere, V., & Dewettinck, K. (2006). Microfiltration of butter serum upon casein micelle destabilisation. *Journal of Dairy Science*, 89, 1915-1925.
- Rombaut, R., van Camp, J., & Dewettinck, K. (2005). Analysis of phospho- and sphingolipids in dairy products by a new HPLC method. *Journal of Dairy Science*, 88, 482-488.

- Sachdeva, S., & Buchheim, W. (1997). Recovery of phospholipids from buttermilk using membrane processing. *Kieler Milchwirtschaftliche Forschungsberichte*, *49*, 47-68.
- Sánchez- Juanes, F., Alonso, J. M., Zancada, L., & Hueso, P. (2009). Distribution and fatty acid content of phospholipids from bovine milk and bovine milk fat globule membranes. *International Dairy Journal*, *19*, 273- 278.
- Spence, A. J., Jiménez- Flores, R., Qian, M., & Goddik, L. (2009). Phospholipid enrichment in sweet and whey cream buttermilk powders using supercritical fluid extraction. *Journal of Dairy Science*, *92*, 2373-2381.
- Singh, H. (2006). The milk fat globule membrane-a biophysical system for food systems. *Current Opinions in Colloid and Interface Science*, *11*, 154- 163.
- Spadaro, A. C. C., Draghetta, W., del Lama, S. N., Camargo, A. C. M., & Greene, L. J. (1979). A convenient manual trinitrobenzenesulfonic acid method for monitoring amino acids and peptides in chromatographic column effluents. *Analytical Biochemistry*, *96*, 371-321.
- Spitsberg, V. L. (2005). Invited review: bovine milk fat globule membrane as a potential nutraceutical. *Journal of Dairy Science*, *88*, 2289-2294.
- Timby, N., Domellöf, E., Hernell, O., Lönnerdal, B., & Domellöf, M. (2014). Neurodevelopment, nutrition, and growth until 12 mo of age in infants fed a low-energy, low-protein formula supplemented with bovine milk fat globule membranes: a randomized controlled trial. *American Journal of Clinical Nutrition*, *99*, 860-868.
- Timby, N., Hernell, O., Vaarala, O., Melin, M., Lönnerdal, B., & Domellöf, M. (2015). Infections in infants fed formula supplemented with bovine milk globule membranes. *Journal of Pediatric Gastroenterology and Nutrition*, *60*, 348-389.



**Figure Legends**

**Figure 1.** SEC overlay chromatogram representing the size distribution profile of the BMP powders where A; BMP and B; 19 % DH BMP hydrolysate.

**Figure 2.** SEC overlay chromatogram representing molecular weight distribution profiles of the ultrafiltration fractions where A; 50 kDa retentate and B; 50 kDa permeate.

**Figure 3.** HPLC- CAD chromatogram representing phospholipid composition of the BMPs where A; BMP and B; 50 kDa retentate.

**Figure 4.** HPLC- CAD chromatogram representing phospholipid profiles of the supercritical fluid treated samples where A; CO<sub>2</sub> extraction only and B; CO<sub>2</sub>: 20% Ethanol.

**Tables and Figures****Table 1.** Molecular weight (M.W.) distributions of BMP and BMP hydrolysate. M. W. distribution quantified with respect to calibration with known M.W. standards.

Enzyme	M.W. Distributions					
	>100 kDa (%)	100-75 kDa (%)	75-50 kDa (%)	50-30 kDa (%)	30-10 kDa (%)	< 10 kDa (%)
<b>BMP</b>	92.10 ± 0.03	7.30 ± 0.11	0.55 ± 0.11	0.05 ± 0.03	-	-
<b>19 % DH</b>	-	-	-	20.66 ± 0.07	42.51 ± 0.12	36.83 ± 0.07

Data are expressed as the average (+ % RSD) of two replicate hydrolysis reactions of the % of total area under the curve.

**Table 2.** Gross compositional analysis of Buttermilk powder (BMP), 50 kDa retentate (50 kDa R) and 50 kDa permeate (50 kDa P) with respect to % Fat (Folch, Barry et al., 2016), % protein (Kjeldahl ISO 8968-1, 2014), and % total phospholipid (PL) determined by summation of each individual PL with respect to that extracted from the sample..

<b>Compositional analysis</b>			
	<b>Fat (%)<sup>a</sup></b>	<b>Protein (%)<sup>b</sup></b>	<b>Total Phospholipid (PL) (% of sample<sup>a</sup>)</b>
<b>BMP</b>	6.84 ± 0.17	31.40 ± 0.57	1.30 ± 0.00
<b>BMP hydrolysate 50 kDa retentate (50 kDa R)</b>	60.07 ± 0.54	10.58 ± 0.09	11.05 ± 0.02
<b>BMP hydrolysate 50 kDa permeate (50 kDa P)</b>	0.29 ± 0.07	52.06 ± 0.16	0.00 ± 0.09
<b>CO2 sample</b>	72.66 ± 0.51	-	0.00 ± 0.03
<b>CO2: 20 % EtOH extract</b>	99.3 ± 0.01	-	56.24 ± 0.07

<b>Individual PL (% of total PL)</b>			
<b>PL Species</b>	<b>BMP</b>	<b>50 kDa R</b>	<b>CO2: EtOH treated sample</b>
<b>Phosphatidylinositol (PI)</b>	10.84 ± 0.04	10.42 ± 0.04	
<b>Phosphatidylethanolamine (PE)</b>	21.66 ± 0.07	20.48 ± 0.01	11.26 ± 0.07
<b>Glucosylceramide (GluCer)</b>	0.07 ± 0.32	0.08 ± 0.11	
<b>Phosphatidylserine (PS)</b>	9.54 ± 0.01	9.88 ± 0.07	
<b>Phosphatidylcholine (PC)</b>	31.31 ± 0.01	31.75 ± 0.12	82.22 ± 0.16
<b>Sphingomyelin (SM)</b>	26.57 ± 0.04	27.39 ± 0.17	6.52 ± 0.02

<sup>a</sup>Data are expressed as the average ( $\pm$  % RSD) of three repeats.

<sup>b</sup>Data are expressed as the average ( $\pm$  % RSD) of two repeats.

<sup>c</sup>Data are expressed as % of total PL, average ( $\pm$  % RSD) of duplicate injections of each sample in triplicate.

Compositional data for BMP previously recorded in Barry et al., 2017.

ACCEPTED MANUSCRIPT

Figure 1.




Figure 2.




Figure 3.





Figure 4.



**Highlights**

- A novel pilot scale process generates high levels of buttermilk phospholipids (PLs)
- Targeted protein hydrolysis reduces co-separation with lipids at ultrafiltration
- Features a non-thermal method of enzyme inactivation to counter membrane fouling
- An 8.7-fold PL enrichment is highest known to date for membrane separation
- Supercritical CO<sub>2</sub> extraction increases PL concentration further to 56.24 %

ACCEPTED MANUSCRIPT